



## Topobathymetric LiDAR point cloud processing and landform classification in a tidal environment

Andersen, Mikkel Skovgaard; Al-Hamdani, Zyad K.; Steinbacher, Frank; Larsen, Laurids Rolighed; Ernstsen, Verner Brandbyge

*Published in:*  
Geophysical Research Abstracts

*Publication date:*  
2017

*Document version*  
Publisher's PDF, also known as Version of record

*Citation for published version (APA):*  
Andersen, M. S., Al-Hamdani, Z. K., Steinbacher, F., Larsen, L. R., & Ernstsen, V. B. (2017). Topobathymetric LiDAR point cloud processing and landform classification in a tidal environment. *Geophysical Research Abstracts*, 19, [EGU2017-5571].



## **Topobathymetric LiDAR point cloud processing and landform classification in a tidal environment**

Mikkel Skovgaard Andersen (1), Ziyad Al-Hamdani (2), Frank Steinbacher (3), Laurids Rolighed Larsen (4), and Verner Brandbyge Ernstsens (1)

(1) Department of Geosciences and Natural Resource Management, University of Copenhagen, Copenhagen, Denmark, (2) Geological Survey of Denmark and Greenland (GEUS), Copenhagen, Denmark, (3) Airborne Hydro Mapping GmbH, Innsbruck, Austria, (4) NIRAS, Allerød, Denmark

Historically it has been difficult to create high resolution Digital Elevation Models (DEMs) in land-water transition zones due to shallow water depth and often challenging environmental conditions. This gap of information has been reflected as a “white ribbon” with no data in the land-water transition zone. In recent years, the technology of airborne topobathymetric Light Detection and Ranging (LiDAR) has proven capable of filling out the gap by simultaneously capturing topographic and bathymetric elevation information, using only a single green laser. We collected green LiDAR point cloud data in the Knudedyb tidal inlet system in the Danish Wadden Sea in spring 2014. Creating a DEM from a point cloud requires the general processing steps of data filtering, water surface detection and refraction correction. However, there is no transparent and reproducible method for processing green LiDAR data into a DEM, specifically regarding the procedure of water surface detection and modelling. We developed a step-by-step procedure for creating a DEM from raw green LiDAR point cloud data, including a procedure for making a Digital Water Surface Model (DWSM) (see Andersen et al., 2017). Two different classification analyses were applied to the high resolution DEM: A geomorphometric and a morphological classification, respectively. The classification methods were originally developed for a small test area; but in this work, we have used the classification methods to classify the complete Knudedyb tidal inlet system.

### References

Andersen MS, Gergely Á, Al-Hamdani Z, Steinbacher F, Larsen LR, Ernstsens VB (2017). Processing and performance of topobathymetric lidar data for geomorphometric and morphological classification in a high-energy tidal environment. *Hydrol. Earth Syst. Sci.*, 21: 43-63, doi:10.5194/hess-21-43-2017.

### Acknowledgements

This work was funded by the Danish Council for Independent Research | Natural Sciences through the project “Process-based understanding and prediction of morphodynamics in a natural coastal system in response to climate change” (Steno Grant no. 10-081102) and by the Geocenter Denmark through the project “Closing the gap! – Coherent land-water environmental mapping (LAWA)” (Grant no. 4-2015).