



Electrophysiologic effects of the IK1 inhibitor PA-6 are modulated by extracellular potassium in isolated guinea pig hearts

Hoeker, Gregory S; Skarsfeldt, Mark A; Jespersen, Thomas; Poelzing, Steven

Published in:
Physiological Reports

DOI:
[10.14814/phy2.13120](https://doi.org/10.14814/phy2.13120)

Publication date:
2017

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](#)

Citation for published version (APA):
Hoeker, G. S., Skarsfeldt, M. A., Jespersen, T., & Poelzing, S. (2017). Electrophysiologic effects of the IK1 inhibitor PA-6 are modulated by extracellular potassium in isolated guinea pig hearts. *Physiological Reports*, 5(1), [e13120]. <https://doi.org/10.14814/phy2.13120>

ORIGINAL RESEARCH

Electrophysiologic effects of the I_{K1} inhibitor PA-6 are modulated by extracellular potassium in isolated guinea pig hearts

Gregory S. Hoeker¹, Mark A. Skarsfeldt², Thomas Jespersen² & Steven Poelzing¹

¹ Biomedical Engineering and Mechanics, Center for Heart and Regenerative Medicine, Virginia Tech, Virginia Tech Carilion Research Institute, Roanoke, Virginia

² Department of Biomedical Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

Keywords

Action potential, conduction velocity, inward rectifier current, pentamidine, potassium, repolarization.

Correspondence

Steven Poelzing, Virginia Tech Carilion Research Institute, 2 Riverside Circle, Roanoke, VA 24016.
Tel: 540-526-2108
Fax: 540-989-3373
E-mail: poelzing@vtc.vt.edu

Funding Information

This work was supported in part by the Novo Nordisk Foundation (TJ) and the National Institutes of Health R01 (SP, R01-HL102298).

Received: 7 December 2016; Accepted: 14 December 2016

doi: 10.14814/phy2.13120

Physiol. Rep., 5 (1), 2017, e13120,
doi: 10.14814/phy2.13120

Abstract

The pentamidine analog PA-6 was developed as a specific inward rectifier potassium current (I_{K1}) antagonist, because established inhibitors either lack specificity or have side effects that prohibit their use in vivo. We previously demonstrated that BaCl₂, an established I_{K1} inhibitor, could prolong action potential duration (APD) and increase cardiac conduction velocity (CV). However, few studies have addressed whether targeted I_{K1} inhibition similarly affects ventricular electrophysiology. The aim of this study was to determine the effects of PA-6 on cardiac repolarization and conduction in Langendorff-perfused guinea pig hearts. PA-6 (200 nm) or vehicle was perfused into ex-vivo guinea pig hearts for 60 min. Hearts were optically mapped with di-4-ANEPPS to quantify CV and APD at 90% repolarization (APD₉₀). Ventricular APD₉₀ was significantly prolonged in hearts treated with PA-6 ($115 \pm 2\%$ of baseline; $P < 0.05$), but not vehicle ($105 \pm 2\%$ of baseline). PA-6 slightly, but significantly, increased transverse CV by 7%. PA-6 significantly prolonged APD₉₀ during hypokalemia (2 mmol/L [K⁺]_o), although to a lesser degree than observed at 4.56 mmol/L [K⁺]_o. In contrast, the effect of PA-6 on CV was more pronounced during hypokalemia, where transverse CV with PA-6 (24 ± 2 cm/sec) was significantly faster than with vehicle (13 ± 3 cm/sec, $P < 0.05$). These results show that under normokalemic conditions, PA-6 significantly prolonged APD₉₀, whereas its effect on CV was modest. During hypokalemia, PA-6 prolonged APD₉₀ to a lesser degree, but profoundly increased CV. Thus, in intact guinea pig hearts, the electrophysiologic effects of the I_{K1} inhibitor, PA-6, are [K⁺]_o-dependent.

Introduction

The inward rectifier potassium current (I_{K1}) is an important regulator of the cardiac action potential, serving to stabilize the resting membrane potential (Sakmann and Trube 1984; Tourneur 1986), and contributing to late repolarization (Kass et al. 1990; Ibarra et al. 1991). The molecular basis of cardiac I_{K1} is attributed to the Kir2.x subfamily of inward rectifier potassium channel proteins (Dhamoon and Jalife 2005), which are strongly regulated by extracellular potassium concentration ([K⁺]_o). For

instance, hypokalemia is known to shift the reversal potential for I_{K1} to a more negative potential and reduce the slope conductance of the inward current (resulting in a decreased peak density of I_{K1}), as well as hyperpolarize the resting membrane potential (Scamps and Carmeliet 1989; Shimoni et al. 1992; Hirota et al. 2000), which together alters sodium channel availability and cardiac excitability. Studies have suggested that I_{K1} plays a critical role in modulating cardiac excitability and the incidence of arrhythmias including congenital atrial fibrillation (Deo et al. 2013), catecholaminergic polymorphic

ventricular tachycardia (Barajas-Martinez et al. 2011), ventricular fibrillation (Warren et al. 2003), and arrhythmias associated with Andersen-Tawil syndrome type I and short QT syndrome 3 (see Anumonwo and Lopatin (2010) for review). Furthermore, hypokalemia has been suggested to exacerbate conduction abnormalities, with reports of an increased risk of ventricular arrhythmias in Brugada patients (Araki et al. 2003; Notarstefano et al. 2005). Similarly, during hypokalemia patients with Andersen-Tawil syndrome type 1 have more pronounced ECG changes (Zhang et al. 2005), a greater burden of premature ventricular contractions (Tawil et al. 1994; Nichols et al. 1996), and an increased occurrence of ventricular arrhythmias (Tawil et al. 1994; Tristani-Firouzi et al. 2002). Lastly, in heart failure, which is associated with a loss of I_{K1} function (Kaab et al. 1996), both the complex pathologic state and common therapies can lead to electrolyte disturbances including hypokalemia (Leier et al. 1994). Thus, regulation of I_{K1} and potassium homeostasis has significant clinical implications for cardiac conduction and arrhythmogenesis.

Despite several decades of recognizing the importance of I_{K1} for cardiac function, the lack of specific and efficacious agonists/antagonists for Kir2.x channels has slowed progress toward understanding the physiologic and pathophysiologic roles of I_{K1} in the heart. Pharmacologic compounds targeting I_{K1} generally lack specificity for Kir2.x channels, or have toxic side effects that prohibit their use in vivo (de Boer et al. 2010; Bhoelan et al. 2014). Recently, seven analogs of the diamine antiprotozoal drug pentamidine were shown to inhibit I_{K1} at nanomolar concentrations. The sixth analog (PA-6) was shown to have high efficiency and specificity for inhibition of the Kir2.x-mediated current (i.e., I_{K1}) (Takanari et al. 2013). In isolated cardiac myocytes, PA-6 was previously shown to increase action potential duration (APD) (Takanari et al. 2013). Additionally, 200 nm PA-6 prolonged APD in ventricular myocardium of isolated rat hearts (Skarsfeldt et al. 2016).

Previously, we demonstrated that partially inhibiting I_{K1} with BaCl_2 prolongs ventricular APD and increases conduction velocity (CV) in ventricular myocardium of isolated guinea pig hearts (Poelzing and Veeraghavan 2007; Veeraghavan and Poelzing 2008), whose action potential morphology more closely mimics human action potentials than those of rats and smaller rodents. However, barium is known to have multiple off-target effects, which could confound these findings (Lesage et al. 1995).

The aim of this study was to investigate the effects of the selective I_{K1} inhibitor, PA-6, on action potential repolarization and conduction in an intact guinea pig heart preparation during normo- and hypokalemia. In this study, we demonstrate that in Langendorff-perfused adult guinea pig

hearts, inhibiting I_{K1} alone prolongs ventricular repolarization but does not substantially alter conduction. However, under hypokalemic conditions, which itself prolongs APD and decreases CV, treatment with PA-6 resulted in further APD prolongation and increased CV.

Materials and Methods

Animals

Animal care and experimental procedures were approved by the Institutional Animal Care and Use Committee at Virginia Polytechnic Institute and State University and conducted in compliance with the European convention for the protection of vertebrate animals used for experimental and other scientific purposes (Council of Europe No 123, Strasbourg 1985).

Male retired breeder, albino Hartley guinea pigs (Hilltop Lab Animals, Scottdale, PA; $n = 29$, approximately 900–1200 g, 13–20 months old) were placed in an induction chamber and anesthetized with 5% isoflurane mixed with 100% oxygen at 3 L/min. After losing consciousness, the animal was removed from the induction chamber and masked with 3–5% isoflurane mixed with 100% oxygen at 4 L/min. Once in a surgical plane of anesthesia, a thoracotomy was performed, the heart was excised and rinsed in Tyrode's solution (see below for details on perfusate composition).

Langendorff perfusion

After the heart was excised, the aorta was cannulated and perfused retrogradely with a modified Tyrode's solution containing (in mmol/L): NaCl 140, KCl 4.56, CaCl_2 1.25, dextrose 5.5, MgCl_2 0.7, and HEPES 10; pH was adjusted to 7.40–7.42 at 37°C using NaOH. The Tyrode's solution was bubbled with 100% oxygen and perfused at a constant flow to maintain a perfusion pressure of 40–55 mmHg. The atria were excised to reduce competitive stimulation, and the heart was placed in a custom-made tissue bath where it was immersed in the perfusate and maintained at 37°C. Hearts were stimulated with a unipolar silver chloride wire positioned on the epicardium of the anterior left ventricle (LV) and paced at a basic cycle length (BCL) of 300 msec using a pulse width of 5 msec and current strength at $1.5\times$ the diastolic threshold.

Optical mapping

Cardiac motion was suppressed by adding the electromechanical uncoupler 2,3-butanedione monoxime (BDM, 7.5 mmol/L) to the perfusate. The voltage sensitive dye di-4 ANEPPS (7.5 $\mu\text{mol/L}$; Biotium, Hayward, CA) was perfused

into the heart for 10 min followed by a 10 min washout period before the start of the experimental protocol. Di-4-ANEPPS was excited by illuminating the anterior surface of the heart with a 150 W halogen light source (MHAB-150 W, Moritex Corporation) and quartz fiber light guide (Moritex Corporation, Saitama, Japan) fitted with an excitation filter (center wavelength of 510/10 nm; Semrock, Rochester, NY). Fluoresced light was collected by a tandem lens assembly (0.63 \times magnification), transmitted through a 610 nm long pass filter (610FG01-50(T257), Andover Corporation, Salem, NH), and detected by a 100 \times 100 pixel CMOS camera (MiCAM Ultima-L, SciMedia, Costa Mesa, CA) with a field of view of 15.9 \times 15.9 mm² (0.159 mm interpixel resolution). Fluorescence was recorded at a sampling rate of 1000 frames/sec.

Reagents

Di-4-ANEPPS (5 mg) was dissolved in 1.3 mL of ethanol to create an 8 mmol/L stock solution. The pentamidine analog PA-6 (C₃₁H₃₂N₄O₂) was synthesized by Syngene, Bangalore, India. PA-6 (M.W. = 492.62) was dissolved in dimethyl sulfoxide (DMSO) and prepared as a 5 mmol/L stock solution. PA-6 is known to interact with the cytoplasmic domain of Kir2.1 (Takanari et al. 2013), and data from rat hearts suggest that maximal, stable effects of PA-6 occur between 45 and 90 min of exposure (Skarsfeldt et al. 2016). To maintain PA-6 in solution and facilitate intracellular uptake over a sustained period, at the time of the experiment, 40 μ L of the PA-6 stock solution was mixed with 40 μ L of Pluronic solution (1 g of Pluronic F-127, Sigma-Aldrich, dissolved in 5 mL of DMSO) and added to 1 L of Tyrode's solution for a final PA-6 concentration of 200 nmol/L. For vehicle control (Veh) studies, 40 μ L of DMSO and 40 μ L of Pluronic solution were added to 1 L of Tyrode's solution.

Experimental protocol

After the Di-4-ANEPPS washout period, steady-state optical action potentials were recorded at baseline, and after 30 and 60 min of no treatment (time control, TC, $n = 3$) or treatment with either PA-6 (200 nmol/L, $n = 8$) or Veh (DMSO + Pluronic, $n = 5$). In separate studies (PA-6: $n = 9$, vehicle: $n = 4$), hearts were perfused with Tyrode's solution containing 2 mmol/L [K⁺] (vs. 4.56 mmol/L) and the experimental protocol was performed as described above.

Data analysis

Fluorescence signals were binned 2 \times 2, yielding an effective spatial resolution of 0.318 mm. Action potential

activation times and 90% repolarization times were quantified as previously described (Girouard et al. 1996; Tian et al. 2004). APD at 90% repolarization (APD₉₀) was calculated as the difference between the 90% repolarization and activation times. CV was quantified from contour maps of activation times transverse (CV_T) and longitudinal (CV_L) to fiber orientation as previously described (Entz et al. 2016). All data are presented as mean \pm standard error of the mean. Two-tailed, paired Student's *t*-tests were used to compare mean APD₉₀, CV_T, and CV_L after treatment to corresponding baseline values within the same heart. Two-tailed, unpaired Student's *t*-tests were used to compare means between groups at a given time point. Differences were considered to be statistically significant for $P < 0.05$.

Results

Action potential prolongation with PA-6 inhibition of I_{K1}

To investigate the effects of I_{K1} inhibition on ventricular repolarization, Langendorff-perfused guinea pig hearts were treated with 200 nmol/L PA-6 for 60 min and APD₉₀ was measured from optical action potentials recorded at 0 (pretreatment), 30, and 60 min. The results from PA-6 treated hearts ($n = 8$) were compared to time (TC, $n = 3$) and vehicle control hearts (Veh, $n = 5$) to assess the effects of preparation stability and the vehicle solvent (DMSO + Pluronic) on APD₉₀. Shown in Figure 1A are superimposed action potentials from the same recording pixel at baseline (0 min) and 60 min for each treatment group: time control (TC), vehicle (Veh), and PA-6. These traces demonstrate that over 60 min, the action potentials recorded in the time control and vehicle hearts were relatively unchanged. In contrast, there were marked changes in the action potential obtained after 60 min of treatment with PA-6 (Fig. 1A, *bottom*). Specifically, PA-6 appeared to affect late repolarization (phase 3 of the action potential), resulting in a prolonged action potential duration (Δ APD₉₀ = 34 msec). Over all experiments ($n = 8$), PA-6 significantly prolonged APD₉₀ (Fig. 1B) at both 30 (Δ 14.2 msec, 108.7% of baseline, $P < 0.05$) and 60 min (Δ 22.3 msec, 113.6% of baseline, $P < 0.05$) of treatment. Over the course of 60 min, a small (Δ 5.7 msec, 103.4% of baseline) but statistically significant APD₉₀ prolongation was observed in untreated hearts (TC). At 60 min, APD₉₀ in PA-6 treated hearts was significantly longer than both Veh and TC hearts. There were no significant differences in APD₉₀ between Veh and TC at any of the time points. These data demonstrate that PA-6 increases APD₉₀ independent of any effects due to time or exposure to vehicle.




Figure 1. PA-6 prolongs APD₉₀ in normokalemic hearts. (A) Superimposed action potentials at 0 min (pretreatment) and after 60 min of time control (TC), or treatment with vehicle (Veh) or 200 nmol/L PA-6 at $[K^+]_o = 4.56$ mmol/L. The difference in APD₉₀ values are shown as Δ APD in the inset. (B) Summary data of mean APD₉₀ (top) and the change in APD₉₀ (as a percent of baseline, bottom) over 60 min in each of the three treatment groups. * $P < 0.05$ versus Veh, # $P < 0.05$ versus baseline.

Negligible effect of PA-6 on conduction velocity

We previously demonstrated that 10 μ mol/L BaCl₂ can increase cardiac conduction (Veeraraghavan and Poelzing 2008). In order to assess the effect of a more potent and specific inhibitor of I_{K1} on transverse and longitudinal conduction, CV_T and CV_L were measured before and after treatment with 200 nmol/L PA-6 during normokalemia ($[K^+]_o = 4.56$ mmol/L). Representative maps of activation isochrones are presented in Figure 2A. These maps demonstrate that after 60 min, there was very little change in the activation patterns due to time (TC) or treatment with Veh or PA-6. Summary data of CV (CV_T and CV_L) revealed a very modest effect of inhibiting I_{K1} on conduction (Fig. 2B). Specifically, treatment with 200 nmol/L PA-6 modestly, but significantly, increased CV_T by paired comparison from 21.9 ± 1.2 to 23.4 ± 1.1 cm/sec at 30 min ($P < 0.05$) and 23.4 ± 1.2 cm/sec at 60 min ($P = 0.05$). However, PA-6 did not increase conduction relative to TC or Veh, as determined by unpaired statistical comparisons. Hence, neither CV_T nor CV_L were significantly different in hearts treated with PA-6 compared to TC and Veh-treated hearts at any time point.

I_{K1} inhibition during hypokalemia

Lowering $[K^+]_o$ leads to a decrease in I_{K1} peak current density (Scamps and Carmeliet 1989), as well as changes

in resting membrane potential and excitability (Shimoni et al. 1992). Previously, we demonstrated that hypokalemia differentially modulates ventricular myocardial electrophysiology during partial I_{K1} blockade with 10 μ mol/L BaCl₂ (Poelzing and Veeraraghavan 2007). Therefore, we sought to test the effect of PA-6 on repolarization and conduction during conditions of hypokalemia. Isolated hearts were equilibrated in Tyrode's solution containing 2 mmol/L $[K^+]_o$, and the experimental protocol was repeated as before with 200 nmol/L PA-6 ($n = 9$) and Veh ($n = 4$). At baseline, hypokalemia by itself prolonged APD₉₀ (202.4 ± 5.3 compared to 164.1 ± 1.9 msec at 4.56 mmol/L $[K^+]_o$, $P < 0.05$) and decreased CV_T (18.0 ± 1.8 compared to 23.1 ± 0.7 cm/sec at 4.56 mmol/L $[K^+]_o$, $P < 0.05$) and CV_L (37.0 ± 2.7 compared to 54.0 ± 1.4 cm/sec at 4.56 mmol/L $[K^+]_o$, $P < 0.05$).

Representative action potentials in Figure 3A demonstrate the action potential changes induced by PA-6 during hypokalemia. While the action potential at baseline (0 min) is already longer than the corresponding action potentials in Figure 1A due to reduced $[K^+]_o$, subsequent treatment with 200 nmol/L PA-6 resulted in further prolongation of APD₉₀. In contrast, the action potentials from Veh-treated hearts were nearly superimposable, with no discernable effect on repolarization. The time-dependent effects of PA-6 on APD₉₀ during hypokalemia are summarized in Figure 3B. Within 30 min, PA-6 significantly increased APD₉₀ ($\Delta 13.9$ msec, 109.3% of baseline, $P < 0.05$) and the prolongation




Figure 2. Effect of PA-6 on CV in normokalemic hearts. (A) Representative contour maps of action potential activation times at 0 min (pretreatment) and 60 min of time control (TC), or treatment with vehicle (Veh) or 200 nmol/L PA-6 at $[K^+]_o = 4.56$ mmol/L. Each isochrone represents a 3 msec change in activation time. The pacing symbol at the center of each map indicates the site of stimulus delivery. (B) Summary data of mean CV_T (top) and CV_L (bottom) over 60 min in each of the three treatment groups. * $P < 0.05$ versus Veh, # $P < 0.05$ versus baseline.

persisted up to 60 min ($\Delta 8.9$ msec, 106.9% of baseline, $P < 0.05$). This APD prolongation (as a percent change from baseline) was significantly less than that observed after 60 min of PA-6 treatment with 4.56 mmol/L $[K^+]_o$ (6.9% vs. 13.6%, $P < 0.05$). The reduced effect of PA-6 on APD prolongation during hypokalemia could possibly be due to partial inhibition of I_{K1} by the lowered $[K^+]_o$ prior to application of PA-6. There were no significant differences in APD_{90} in Veh-treated hearts at either 30 or 60 min.

Comparing the representative baseline (0 min) activation maps at 2 mmol/L $[K^+]_o$ (Fig. 4A) to those at 4.56 mmol/L $[K^+]_o$ (Fig. 2A), it is apparent that there is crowding of isochrones during hypokalemia representing slower conduction due to the decrease in $[K^+]_o$. The pattern of activation and number of isochrones was relatively unchanged by 60 min of treatment with Veh. In contrast,

the activation map after 60 min of PA-6 revealed a marked effect on conduction, with far fewer isochrones (total activation time = 34 msec vs. 42 msec at baseline) and a corresponding increase in CV. In fact, the summary data presented in Figure 4B shows that for all experiments PA-6 reversed the decrease in CV_T that was observed with hypokalemia, and restored CV_T to values observed under normokalemia (23.6 ± 2.4 cm/sec vs. 23.1 ± 0.7 cm/sec). Mean CV_T was greater with PA-6 than Veh at both 30 min (23.3 ± 1.9 cm/sec vs. 14.1 ± 2.7 cm/sec, $P < 0.05$) and 60 min (23.6 ± 2.4 cm/sec vs. 12.6 ± 3.4 cm/sec, $P < 0.05$). A similar trend was observed for CV_L , where mean CV_L after 30 minutes of treatment with PA-6 was 45.4 ± 3.3 cm/sec (compared to Veh: 34.4 ± 4.6 , $P = 0.08$) and after 60 min CV_L was 44.3 ± 3.6 cm/sec (compared to Veh: 29.8 ± 7.9 , $P = 0.09$).




Figure 3. PA-6 prolongs APD₉₀ in hypokalemic hearts. (A) Superimposed action potentials at 0 min (pretreatment) and after 60 min of treatment with vehicle (Veh) or 200 nmol/L PA-6 at $[K^+]_o = 2$ mmol/L. The difference in APD₉₀ values are shown as Δ APD in the inset. (B) Summary data of mean APD₉₀ (left) and the change in APD₉₀ (as a percent of baseline, right) over 60 min during treatment with PA-6 or Veh. * $P < 0.05$ versus Veh, # $P < 0.05$ versus baseline.




Figure 4. PA-6 increases CV in hypokalemic hearts. (A) Representative contour maps of action potential activation times at 0 min (pretreatment) and 60 min of treatment with vehicle (Veh) or 200 nmol/L PA-6 at $[K^+]_o = 2$ mmol/L. Each isochrone represents a 3 msec change in activation time. The pacing symbol at the center of each map indicates the site of stimulus delivery. (B) Summary data of mean CV_T (top) and CV_L (bottom) over 60 min in each of the three treatment groups. * $P < 0.05$ versus Veh, # $P < 0.05$ versus baseline.

Discussion

Recently, selective inhibition of I_{K1} with the pentamidine analog PA-6 was shown to influence action potential repolarization and refractoriness in isolated rat hearts (Skarsfeldt et al. 2016). Prior to the development of

PA-6, our laboratory demonstrated that inhibiting I_{K1} with BaCl₂ led to APD prolongation (Poelzing and Veeraraghavan 2007) and enhanced conduction (Veeraraghavan and Poelzing 2008) in Langendorff-perfused guinea pig hearts. In this study, using the same experimental model, we found that 60 min of treatment with 200 nmol/L PA-

6: (1) significantly prolonged ventricular APD₉₀; (2) had minimal to no effect on CV_T or CV_L during normokalemia; (3) prior to administration of PA-6, hypokalemia (2 mmol/L [K⁺]_o) significantly prolonged APD₉₀ and decreased CV_T and CV_L; (4) during hypokalemia, PA-6 prolonged APD₉₀ further; and (5) in contrast to normokalemia, treatment with PA-6 during hypokalemia significantly increased CV_T and demonstrated a similar trend in CV_L.

A role for I_{K1} in cardiac repolarization has been demonstrated by studies conducted in a number of species, using a variety of pharmacologic agents to inhibit Kir2.x [e.g. BaCl₂ (Poelzing and Veeraghavan 2007; Wu et al. 1999; Baiardi et al. 2003), MS-551 (Nakaya et al. 1993; Sen et al. 1998) RP58866 (Rees and Curtis 1993), RP62719 (Williams et al. 1999; Biliczki et al. 2002), chloroquine (Noujaim et al. 2010), tamoxifen (He et al. 2003), carbon monoxide (Liang et al. 2014), and cesium (Morita et al. 2007)]. Related studies have consistently found that inhibiting I_{K1} prolongs the QT interval and APD, although the severity of the effect varies widely (Rees and Curtis 1993; Biliczki et al. 2002; He et al. 2003). Among these studies, those reporting the largest effect on APD predominantly utilized nonselective compounds or doses of inhibitors that are known to be nonselective for I_{K1} . Thus, it is possible that nonspecific effects influenced these findings, and that they represent an overestimation of the role of I_{K1} in normal cardiac repolarization.

In Langendorff-perfused guinea pig hearts, we found that 60 min of treatment with 200 nmol/L PA-6, a dose which has been shown to inhibit I_{K1} by 77–100% while having no effect on I_{Na} , I_{Ca} , I_{to} , I_{Kr} , or I_{Ks} (Takanari et al. 2013), prolonged APD₉₀ by 14%. Previously, 200 nmol/L PA-6 was shown to increase APD₉₀ by 74% in Langendorff-perfused rat hearts (Skarsfeldt et al. 2016). However, it is difficult to directly compare APD prolongation from studies in rat hearts to those of humans and other larger mammals with more pronounced action potential plateaus, as the composition and time course of their respective repolarizing currents are clearly distinct from one another. As a result, APD₉₀ in ventricular guinea pig myocardium is fourfold longer than in rat at baseline conditions [164 msec vs. 42 msec (Skarsfeldt et al. 2016)]. Interestingly, the absolute change in mean APD₉₀ induced by PA-6 was similar in the two species (22.3 msec in guinea pig, 30.8 msec in rat). However, since late repolarization (where I_{K1} is active) constitutes a greater percentage of total APD in rat than guinea pig, the resulting percent difference in APD prolongation for a given absolute change is much larger in rat myocardium. A review of the literature supports the assertion that APD prolongation due to I_{K1} inhibition is enhanced in species with no action potential plateau (e.g., rat and mouse) relative to species

with an action potential plateau (e.g., guinea pig, rabbit, canine, primate, and human) (Rees and Curtis 1993; Williams et al. 1999; Baiardi et al. 2003; Noujaim et al. 2010; Nagy et al. 2013). Accordingly, in isolated canine adult-ventricular cardiomyocytes, 200 nmol/L PA-6 increased APD₉₀ by 34% (Takanari et al. 2013), which is similar to our findings in guinea pig. Using arguably the most commonly used approach for inhibiting I_{K1} (i.e., 10 μ mol/L BaCl₂), we previously observed a 15–25% prolongation of APD in isolated guinea pig hearts, similar to the effect we are currently reporting with PA-6. Thus, in larger mammals with action potential morphologies similar to those in human, the effect of PA-6 on cardiac repolarization appears to be less than observed in smaller rodent species which lack a prominent plateau.

Given that I_{K1} plays a critical role in determining the resting membrane potential, it has been postulated that I_{K1} could oppose the depolarizing current through voltage-gated sodium channels (i.e. I_{Na}) during the early phase of AP activation. Consequently, inhibiting I_{K1} would lead to an increase in cardiac CV. Alternatively, decreased I_{K1} could potentially raise the resting membrane potential, resulting in sodium channel inactivation, thereby leading to a decrease in CV. However, the effect on the ventricular resting membrane potential following I_{K1} blockade has in most studies has been found to be either minor or undetectable, suggesting a resting membrane potential reserve of other potassium currents (see van der Heyden and Jespersen (2016) for review). Despite a well-recognized role for I_{K1} modulation of cardiac excitability, surprisingly few studies have directly tested the prevailing theories of the effect of I_{K1} inhibition on ventricular conduction. Escande et al. (1992) saw no change in CV with RP62719, while Noujaim et al. (2010) reported a 35% decrease in CV with chloroquine. However, both of these I_{K1} inhibitors have been demonstrated to block other potassium currents at the doses tested, as well as sodium and calcium currents in the case of chloroquine (Jurkiewicz et al. 1996; Yang et al. 1999; Fujita and Kurachi 2000). In support of the theory that I_{K1} opposes I_{Na} depolarization, we have previously reported that 10 μ mol/L BaCl₂ increased CV_T by approximately 25% (+6 cm/sec) (Veeraghavan and Poelzing 2008). The corresponding increase in CV_T in this study with 200 nmol/L PA-6 was 7% (+1.5 cm/sec). This is consistent with a lack of an effect of 200 nmol/L PA-6 on resting membrane potential in isolated canine cardiomyocytes (Takanari et al. 2013). Altogether, this suggests that less specific I_{K1} inhibitors, such as BaCl₂, may alter CV due to off-target effects, and/or on its own, selective inhibition of I_{K1} can significantly impact repolarization, but alone may not be sufficient to change cardiac excitability or appreciably alter CV.

To further compare our findings with PA-6 to those with BaCl₂, we repeated the experimental protocol under conditions of low [K⁺]_o. Consistent with previous results (Poelzing and Veeraraghavan 2007), lowering [K⁺]_o to 2 mmol/L prolonged APD by approximately 25%. Treatment with 200 nmol/L PA-6 prolonged APD by a further 4–7%. Importantly, hypokalemia alone significantly decreased CV_T by 22%, whereas subsequent treatment with PA-6 increased CV_T by 27%, effectively reversing the conduction loss due to hypokalemia and restoring CV_T to normokalemic values (23.6 cm/sec vs. 23.1 cm/sec). To our knowledge, this is the first study to investigate the effects of I_{K1} inhibition on cardiac conduction under conditions of hypokalemia. Therefore, these data suggest that PA-6 may actually rescue conduction slowing induced by hypokalemia. Lowering [K⁺]_o will hyperpolarize the resting membrane potential, leading to a delay in sodium channel activation and therefore a slowing of conduction. Subsequent I_{K1} blockade could potentially depolarize the membrane and restore normal resting membrane potential, thereby alleviating the conduction abnormality induced by hypokalemia. Which leads to the question—why does not PA-6 not increase CV at normokalemia? Perhaps cardiomyocyte excitability is more sensitive to small shifts (a few mV) in membrane potential at more hyperpolarized potentials than at normal resting membrane potential. Alternatively, perhaps I_{K1} inhibition alone is insufficient to significantly affect resting membrane potential or CV, and a further perturbation, such as hypokalemia, is required before an effect is observed. Data from Takanari et al. (2013) would support the latter, given that no change in resting membrane potential was observed in isolated canine cardiomyocytes treated with 200 nmol/L PA-6. Lastly, it is possible that PA-6 inhibition of I_{K1} is [K⁺]-dependent. It has been demonstrated that permeant ions (in this case K⁺) can influence ligand interactions with ion channels (Zhorov and Tikhonov 2013), and perhaps the greater effect of PA-6 on CV at low [K⁺]_o could be explained by such an interaction. These provocative hypotheses warrant further testing, particularly as new small molecule I_{K1} inhibitors such as ML133 are being developed for use in vivo (Wang et al. 2011).

Limitations

Due to the degradation of di-4-ANEPPS fluorescence signals over time, we limited our experimental protocol to 60 min of PA-6 treatment. While PA-6 has been demonstrated to have a stable effect on APD₉₀ within 45 min of treatment in Langendorff-perfused rat hearts, the ventricular effective refractory period increased up to 90 min (Skarsfeldt et al. 2016). Therefore, it is possible that longer exposure times could reveal more pronounced

effects of I_{K1} inhibition than observed in this study. While the specificity of PA-6 for I_{K1} has been rigorously tested in heterologous systems expressing cardiac ion channels from human and mouse, as well as in isolated canine cardiomyocytes (Takanari et al. 2013), specificity has not been tested in guinea pig cardiomyocytes.

Conclusion

Under normokalemic conditions, the I_{K1} inhibitor PA-6 significantly prolonged APD₉₀ without substantially affecting CV. During hypokalemia, PA-6 prolonged APD₉₀, although to a lesser degree, and significantly increased CV. Thus, in isolated guinea pig hearts, the electrophysiologic effects of PA-6 are [K⁺]_o-dependent. Furthermore, these results highlight the importance of using a selective inhibitor to investigate the role of I_{K1} in cardiac repolarization and conduction, as well as validating these results in a species with action potential morphologies similar to those in human. Given its superior selectivity for I_{K1} and advantageous safety profile, PA-6 will serve as an important tool for advancing our understanding of the physiologic and pathophysiologic roles of I_{K1} in vivo.

Conflict of Interest

None declared.

References

- Anumonwo, J. M., and A. N. Lopatin. 2010. Cardiac strong inward rectifier potassium channels. *J. Mol. Cell. Cardiol.* 48:45–54.
- Araki, T., T. Konno, H. Itoh, H. Ino, and M. Shimizu. 2003. Brugada syndrome with ventricular tachycardia and fibrillation related to hypokalemia. *Circ. J.* 67:93–95.
- Baiardi, G., A. P. Zumino, and E. R. Petrich. 2003. Effects of barium and 5-hydroxydecanoate on the electrophysiologic response to acute regional ischemia and reperfusion in rat hearts. *Mol. Cell. Biochem.* 254:185–191.
- Barajas-Martinez, H., D. Hu, G. Ontiveros, G. Caceres, M. Desai, E. Burashnikov, et al. 2011. Biophysical and molecular characterization of a novel de novo KCNJ2 mutation associated with Andersen-Tawil syndrome and catecholaminergic polymorphic ventricular tachycardia mimicry. *Circ. Cardiovasc. Genet.* 4:51–57.
- Bhoelan, B. S., C. H. Stevering, A. T. van der Boog, and M. A. van der Heyden. 2014. Barium toxicity and the role of the potassium inward rectifier current. *Clin. Toxicol. (Phila.)* 52:584–593.
- Biliczki, P., L. Virag, N. Iost, J. G. Papp, and A. Varro. 2002. Interaction of different potassium channels in cardiac repolarization in dog ventricular preparations: role of repolarization reserve. *Br. J. Pharmacol.* 137:361–368.

- de Boer, T. P., M. J. Houtman, M. Compier, and M. A. van der Heyden. 2010. The mammalian K(IR)2.x inward rectifier ion channel family: expression pattern and pathophysiology. *Acta Physiol. (Oxf)* 199:243–256.
- Deo, M., Y. Ruan, S. V. Pandit, K. Shah, O. Berenfeld, A. Blaurock, et al. 2013. KCNJ2 mutation in short QT syndrome 3 results in atrial fibrillation and ventricular proarrhythmia. *Proc. Natl. Acad. Sci. U. S. A.* 110:4291–4296.
- Dhamoon, A. S., and J. Jalife. 2005. The inward rectifier current (I_{K1}) controls cardiac excitability and is involved in arrhythmogenesis. *Heart Rhythm* 2:316–324.
- Entz, M. 2nd, S. A. George, M. J. Zeitz, T. Raisch, J. W. Smyth, and S. Poelzing. 2016. Heart rate and extracellular sodium and potassium modulation of gap junction mediated conduction in Guinea Pigs. *Front. Physiol.* 7:16.
- Escande, D., M. Mestre, I. Caverro, J. Brugada, and C. Kirchhof. 1992. RP 58866 and its active enantiomer RP 62719 (terikalant): blockers of the inward rectifier K⁺ current acting as pure class III antiarrhythmic agents. *J. Cardiovasc. Pharmacol.* 20(Suppl. 2):S106–S113.
- Fujita, A., and Y. Kurachi. 2000. Molecular aspects of ATP-sensitive K⁺ channels in the cardiovascular system and K⁺ channel openers. *Pharmacol. Ther.* 85:39–53.
- Girouard, S. D., K. R. Laurita, and D. S. Rosenbaum. 1996. Unique properties of cardiac action potentials recorded with voltage-sensitive dyes. *J. Cardiovasc. Electrophysiol.* 7:1024–1038.
- He, J., M. E. Kargacin, G. J. Kargacin, and C. A. Ward. 2003. Tamoxifen inhibits Na⁺ and K⁺ currents in rat ventricular myocytes. *Am. J. Physiol. Heart Circ. Physiol.* 285:H661–H668.
- van der Heyden, M. A., and T. Jespersen. 2016. Pharmacological exploration of the resting membrane potential reserve: impact on atrial fibrillation. *Eur. J. Pharmacol.* 771:56–64.
- Hirota, M., H. Ohtani, E. Hanada, H. Sato, H. Kotaki, H. Uemura, et al. 2000. Influence of extracellular K⁺ concentrations on quinidine-induced K⁺ current inhibition in rat ventricular myocytes. *J. Pharm. Pharmacol.* 52:99–105.
- Ibarra, J., G. E. Morley, and M. Delmar. 1991. Dynamics of the inward rectifier K⁺ current during the action potential of guinea pig ventricular myocytes. *Biophys. J.* 60:1534–1539.
- Jurkiewicz, N. K., J. Wang, B. Fermini, M. C. Sanguinetti, and J. J. Salata. 1996. Mechanism of action potential prolongation by RP 58866 and its active enantiomer, terikalant. Block of the rapidly activating delayed rectifier K⁺ current, I_{Kr} . *Circulation* 94:2938–2946.
- Kaab, S., H. B. Nuss, N. Chiamvimonvat, B. O'Rourke, P. H. Pak, D. A. Kass, et al. 1996. Ionic mechanism of action potential prolongation in ventricular myocytes from dogs with pacing-induced heart failure. *Circ. Res.* 78:262–273.
- Kass, R. S., J. P. Arena, and K. B. Walsh. 1990. Measurement and block of potassium channel currents in the heart - importance of channel type. *Drug Dev. Res.* 19:115–127.
- Leier, C. V., L. Dei Cas, and M. Metra. 1994. Clinical relevance and management of the major electrolyte abnormalities in congestive heart failure: hyponatremia, hypokalemia, and hypomagnesemia. *Am. Heart J.* 128:564–574.
- Lesage, F., E. Guillemare, M. Fink, F. Duprat, C. Heurteaux, M. Fosset, et al. 1995. Molecular properties of neuronal G-protein-activated inwardly rectifying K⁺ channels. *J. Biol. Chem.* 270:28660–28667.
- Liang, S., Q. Wang, W. Zhang, H. Zhang, S. Tan, A. Ahmed, et al. 2014. Carbon monoxide inhibits inward rectifier potassium channels in cardiomyocytes. *Nat. Commun.* 5:4676.
- Morita, H., D. P. Zipes, S. T. Morita, and J. Wu. 2007. Mechanism of U wave and polymorphic ventricular tachycardia in a canine tissue model of Andersen-Tawil syndrome. *Cardiovasc. Res.* 75:510–518.
- Nagy, N., K. Acsai, A. Kormos, Z. Sebok, A. S. Farkas, N. Jost, et al. 2013. [Ca(2)(+)] i-induced augmentation of the inward rectifier potassium current (I_{K1}) in canine and human ventricular myocardium. *Pflugers Arch.* 465:1621–1635.
- Nakaya, H., N. Tohse, Y. Takeda, and M. Kanno. 1993. Effects of MS-551, a new class III antiarrhythmic drug, on action potential and membrane currents in rabbit ventricular myocytes. *Br. J. Pharmacol.* 109:157–163.
- Nichols, C. G., E. N. Makhina, W. L. Pearson, Q. Sha, and A. N. Lopatin. 1996. Inward rectification and implications for cardiac excitability. *Circ. Res.* 78:1–7.
- Notarstefano, P., C. Pratola, T. Toselli, and R. Ferrari. 2005. Atrial fibrillation and recurrent ventricular fibrillation during hypokalemia in Brugada syndrome. *Pacing Clin. Electrophysiol.* 28:1350–1353.
- Noujaim, S. F., J. A. Stuckey, D. Ponce-Balbuena, T. Ferrer-Villada, A. Lopez-Izquierdo, S. Pandit, et al. 2010. Specific residues of the cytoplasmic domains of cardiac inward rectifier potassium channels are effective antifibrillatory targets. *FASEB J.* 24:4302–4312.
- Poelzing, S., and R. Veeraraghavan. 2007. Heterogeneous ventricular chamber response to hypokalemia and inward rectifier potassium channel blockade underlies bifurcated T wave in guinea pig. *Am. J. Physiol. Heart Circ. Physiol.* 292: H3043–H3051.
- Rees, S. A., and M. J. Curtis. 1993. Specific I_{K1} blockade: a new antiarrhythmic mechanism? Effect of RP58866 on ventricular arrhythmias in rat, rabbit, and primate. *Circulation* 87:1979–1989.
- Sakmann, B., and G. Trube. 1984. Conductance properties of single inwardly rectifying potassium channels in ventricular cells from guinea-pig heart. *J. Physiol.* 347:641–657.
- Scamps, F., and E. Carmeliet. 1989. Effect of external K⁺ on the delayed K⁺ current in single rabbit Purkinje cells. *Pflugers Arch.* 414(Suppl. 1):S169–S170.
- Sen, L., G. Cui, Y. Sakaguchi, and B. N. Singh. 1998. Electrophysiological effects of MS-551, a new class III agent:

- comparison with dl-sotalol in dogs. *J. Pharmacol. Exp. Ther.* 285:687–694.
- Shimoni, Y., R. B. Clark, and W. R. Giles. 1992. Role of an inwardly rectifying potassium current in rabbit ventricular action potential. *J. Physiol.* 448:709–727.
- Skarsfeldt, M. A., H. Carstensen, L. Skibsbye, C. Tang, R. Buhl, B. H. Bentzen, et al. 2016. Pharmacological inhibition of I_{K1} by PA-6 in isolated rat hearts affects ventricular repolarization and refractoriness. *Physiol. Rep.*, 4, 1–11.
- Takanari, H., L. Nalos, A. Stry-Weinzinger, K. C. de Git, R. Varkevisser, T. Linder, et al. 2013. Efficient and specific cardiac IK(1) inhibition by a new pentamidine analogue. *Cardiovasc. Res.* 99:203–214.
- Tawil, R., L. J. Ptacek, S. G. Pavlakis, D. C. DeVivo, A. S. Penn, C. Ozdemir, et al. 1994. Andersen's syndrome: potassium-sensitive periodic paralysis, ventricular ectopy, and dysmorphic features. *Ann. Neurol.* 35:326–330.
- Tian, X. L., S. L. Yong, X. Wan, L. Wu, M. K. Chung, P. J. Tchou, et al. 2004. Mechanisms by which SCN5A mutation N1325S causes cardiac arrhythmias and sudden death in vivo. *Cardiovasc. Res.* 61:256–267.
- Tourneur, Y. 1986. Action potential-like responses due to the inward rectifying potassium channel. *J. Membr. Biol.* 90:115–122.
- Tristani-Firouzi, M., J. L. Jensen, M. R. Donaldson, V. Sansone, G. Meola, A. Hahn, et al. 2002. Functional and clinical characterization of KCNJ2 mutations associated with LQT7 (Andersen syndrome). *J. Clin. Invest.* 110:381–388.
- Veeraraghavan, R., and S. Poelzing. 2008. Mechanisms underlying increased right ventricular conduction sensitivity to flecainide challenge. *Cardiovasc. Res.* 77:749–756.
- Wang, H. R., M. Wu, H. Yu, S. Long, A. Stevens, D. W. Engers, et al. 2011. Selective inhibition of the K(ir)2 family of inward rectifier potassium channels by a small molecule probe: the discovery, SAR, and pharmacological characterization of ML133. *ACS Chem. Biol.* 6:845–856.
- Warren, M., P. K. Guha, O. Berenfeld, A. Zaitsev, J. M. Anumonwo, A. S. Dhamoon, et al. 2003. Blockade of the inward rectifying potassium current terminates ventricular fibrillation in the guinea pig heart. *J. Cardiovasc. Electrophysiol.* 14:621–631.
- Williams, B. A., D. R. Dickenson, and G. N. Beatch. 1999. Kinetics of rate-dependent shortening of action potential duration in guinea-pig ventricle; effects of I_{K1} and I_{Kr} blockade. *Br. J. Pharmacol.* 126:1426–1436.
- Wu, M. H., M. J. Su, and S. S. Sun. 1999. Electrophysiological profile after inward rectifier K channel blockade by barium in isolated rabbit hearts. Altered repolarization and unmasked decremental conduction property. *Europace* 1:85–95.
- Yang, B. F., G. R. Li, C. Q. Xu, and S. Nattel. 1999. Effects of RP58866 on transmembrane K⁺ currents in mammalian ventricular myocytes. *Zhongguo Yao Li Xue Bao* 20:961–969.
- Zhang, L., D. W. Benson, M. Tristani-Firouzi, L. J. Ptacek, R. Tawil, P. J. Schwartz, et al. 2005. Electrocardiographic features in Andersen-Tawil syndrome patients with KCNJ2 mutations: characteristic T-U-wave patterns predict the KCNJ2 genotype. *Circulation* 111:2720–2726.
- Zhorov, B. S., and D. B. Tikhonov. 2013. Ligand action on sodium, potassium, and calcium channels: role of permeant ions. *Trends Pharmacol. Sci.* 34:154–161.