



Perturbation of Auxin Homeostasis and Signaling by PINOID Overexpression Induces Stress Responses in Arabidopsis

Saini, Kumud; AbdElgawad, Hamada; Markakis, Marios N.; Schoenaers, Sebastjen; Asard, Han; Prinsen, Els; Beemster, Gerrit T. S.; Vissenberg, Kris

Published in:
Frontiers in Plant Science

DOI:
[10.3389/fpls.2017.01308](https://doi.org/10.3389/fpls.2017.01308)

Publication date:
2017

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](#)

Citation for published version (APA):
Saini, K., AbdElgawad, H., Markakis, M. N., Schoenaers, S., Asard, H., Prinsen, E., ... Vissenberg, K. (2017). Perturbation of Auxin Homeostasis and Signaling by *PINOID* Overexpression Induces Stress Responses in Arabidopsis. *Frontiers in Plant Science*, 8, [1308]. <https://doi.org/10.3389/fpls.2017.01308>



Perturbation of Auxin Homeostasis and Signaling by *PINOID* Overexpression Induces Stress Responses in Arabidopsis

Kumud Saini¹, Hamada AbdElgawad^{1,2}, Marios N. Markakis^{1,3}, Sébastien Schoenaers¹, Han Asard¹, Els Prinsen¹, Gerrit T. S. Beemster¹ and Kris Vissenberg^{1,4*}

¹ Integrated Molecular Plant Physiology Research, University of Antwerp, Antwerp, Belgium, ² Department of Botany, Faculty of Science, Beni-Suef University, Beni Suef, Egypt, ³ Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark, ⁴ Plant and Biochemistry and Biotechnology Lab, Department of Agriculture, School of Agriculture, Food and Nutrition, Technological Educational Institute of Crete: University of Applied Sciences, Heraklion, Greece

OPEN ACCESS

Edited by:

Vicent Arbona,
Jaume I University, Spain

Reviewed by:

Luisa M. Sandalio,
Consejo Superior de Investigaciones
Científicas (CSIC), Spain
Christine Helen Foyer,
University of Leeds, United Kingdom

*Correspondence:

Kris Vissenberg
kris.vissenberg@uantwerpen.be

Specialty section:

This article was submitted to
Plant Abiotic Stress,
a section of the journal
Frontiers in Plant Science

Received: 07 April 2017

Accepted: 12 July 2017

Published: 02 August 2017

Citation:

Saini K, AbdElgawad H,
Markakis MN, Schoenaers S,
Asard H, Prinsen E, Beemster GTS
and Vissenberg K (2017) Perturbation
of Auxin Homeostasis and Signaling
by *PINOID* Overexpression Induces
Stress Responses in Arabidopsis.
Front. Plant Sci. 8:1308.
doi: 10.3389/fpls.2017.01308

Under normal and stress conditions plant growth require a complex interplay between phytohormones and reactive oxygen species (ROS). However, details of the nature of this crosstalk remain elusive. Here, we demonstrate that *PINOID* (*PID*), a serine threonine kinase of the AGC kinase family, perturbs auxin homeostasis, which in turn modulates rosette growth and induces stress responses in Arabidopsis plants. Arabidopsis mutants and transgenic plants with altered *PID* expression were used to study the effect on auxin levels and stress-related responses. In the leaves of plants with ectopic *PID* expression an accumulation of auxin, oxidative burst and disruption of hormonal balance was apparent. Furthermore, *PID* overexpression led to the accumulation of antioxidant metabolites, while *pid* knockout mutants showed only moderate changes in stress-related metabolites. These physiological changes in the plants overexpressing *PID* modulated their response toward external drought and osmotic stress treatments when compared to the wild type. Based on the morphological, transcriptome, and metabolite results, we propose that perturbations in the auxin hormone levels caused by *PID* overexpression, along with other hormones and ROS downstream, cause antioxidant accumulation and modify growth and stress responses in Arabidopsis. Our data provide further proof for a strong correlation between auxin and stress biology.

Keywords: auxin, *PINOID* (*PID*), reactive oxygen species (ROS), flavonoids, drought stress, osmotic stress

INTRODUCTION

The growth regulator auxin plays a role in many aspects of growth and development in plants (Teale et al., 2006; Benjamins and Scheres, 2008; Enders and Strader, 2015). There is increasing evidence for the involvement of auxin metabolism, transport and signaling in stress responses as well (Shibasaki et al., 2009; Shen et al., 2010; Zhang et al., 2012; Kazan, 2013; Remy et al., 2013). Morphogenesis and stress adaptive responses are closely linked to cellular hormonal homeostasis, including auxin. On one hand, mutants overproducing auxin or with altered auxin distribution show developmental defects such as inhibition of shoot growth, cotyledon and leaf epinasty, a longer hypocotyl and excessive lateral root growth (Boerjan et al., 1995; King et al., 1995; Delarue et al., 1998; Zhao et al., 2001). On the other hand, several stress conditions are also known to modify

auxin homeostasis and response. Auxin homeostasis could be perturbed by stress-induced changes in the concentration of phenolics compounds such as quercetin and kaempferol (auxin transport inhibitor affecting cycling of auxin efflux carriers), and changes in apoplastic pH affecting auxin uptake and distribution (Potters et al., 2007, 2009). Cadmium treated, phosphorus-, or sulfur-starved plants showed altered auxin metabolism (Potters et al., 2007). Water deficit conditions upregulated Gretchen Hagen 3 (*GH3*) genes, suppressed *YUC* genes and lowered indole-3-acetic acid (IAA) levels in rice (Zhang et al., 2009; Du et al., 2013). Hyper-osmolarity promoted PIN FORMED1 (PIN1) and PIN2 internalization while hypo-osmolarity showed opposite effect in Arabidopsis roots (Zwiewka et al., 2015). Salt stress is also shown to affect free and conjugated IAA levels in the developing xylems in poplar, resulting in negative effects on the expansion of xylem vessels (Junghans et al., 2006). Studies on rice and sorghum suggest that auxin response factors (*ARFs*) and several other early auxin-responsive genes also function in abiotic stress responses (Jain and Khurana, 2009; Wang et al., 2010). Alternatively, stress also induced changes in H₂O₂ levels that are co-factors for peroxidases to catalyze the oxidative degradation of IAA (Gazarian et al., 1998; Tognetti et al., 2012). Apoplastic reactive oxygen species (ROS) transiently decreased auxin signaling, modified auxin homeostasis and altered leaf morphology in ozone (O₃) treated plants (Blomster et al., 2011).

Interestingly, many studies have reported an increase in abiotic stress tolerance in transgenic plants, and in conditions with altered auxin levels (Shi et al., 2014; Naser and Shani, 2016). For instance, elevated auxin levels in transgenic lines overexpressing *YUCCA7* (a gene involved in the auxin biosynthesis pathway) increased resistance to drought stress in Arabidopsis (Lee et al., 2012). Similarly, overexpression of *YUCCA6* led to enhanced resistance to water stress in potato (Kim et al., 2013). Furthermore, ectopic expression of H₂O₂-inducible *UGT74E2* led to an increase in free indole-3-butyric acid (IBA, another form of active auxin) and IBA-glucose, and also increased the tolerance to drought stress (Tognetti et al., 2010). Interestingly, Park et al. (2007a), on the other hand, showed that overexpression of the *GH3* gene lowered IAA levels, reduced growth and enhanced the resistance to abiotic stresses. These studies clearly demonstrate a link between auxin physiology and response to environmental stress factors. Clearly, the relation between auxin and stress adaptations is complex. The specific role of auxin in the induction of stress responses also remains elusive.

The kinase PINOID (*PID*) regulates PIN localizations on the cellular membranes and thus regulates polar auxin transport (PAT; Friml et al., 2004; Kleine-Vehn et al., 2009). *PID* is an early auxin inducible gene and a member of the AGCVIII protein kinase family (Benjamins et al., 2001). AGC kinases are named after three classes of animal proteins involved in a receptor-mediated growth factor signal transduction: protein kinase A (PKA), cyclic GMP-dependent protein kinases (PKG) and protein kinase C (PKC). There are 37 such kinases in Arabidopsis, 23 of which belong to the AGCVIII group (Galván-Ampudia and Offringa, 2007). AGC kinases regulate cell growth, morphogenesis and cell immunity in animals, where mutations

and malfunctioning kinases can lead to diseases such as cancer (Pearce et al., 2010). In plants as well, apart from supporting normal development, AGC kinases also participate in stress signaling and in regulation of plant immunity (Devarenne et al., 2006; Hirt et al., 2011; Garcia et al., 2012). For example, the AGC kinase, oxidative signal-inducible 1 (OXI1), is essential for H₂O₂-mediated oxidative stress signaling, defense against the oomycete pathogen *Hyaloperonospora parasitica*, and immunity against *Pseudomonas syringae* (Rentel et al., 2004; Petersen et al., 2009).

We have performed a detailed growth analysis on two *pid* knockouts (*pid-14* and SALK_009478) and two *PID* overexpression lines (*PID^{OE}*), 35S::PID10 (addressed here after as P10) and 35S::PID21 (here after P21), and discussed their contrasting effect on auxin homeostasis and rosette phenotypes (Saini et al., 2017). *PID^{OE}* lines showed severely reduced rosette growth and high auxin levels in the first pair of leaves. RNA sequencing of the leaves of *PID^{OE}* lines pointed toward induction of stress-related responses. Therefore, we here investigate the effect of altered *PID* expression on the perturbation of auxin homeostasis and on the induction of stress responses in whole rosettes in Arabidopsis. We also investigate the effect of such perturbations on the modulation of whole plant responses toward osmotic and drought stress conditions.

MATERIALS AND METHODS

Plant Material and Growth Conditions of Arabidopsis

Arabidopsis thaliana Col-0 ecotype seeds were grown in half strength Murashige and Skoog (MS) medium including vitamins (Duchefa, Netherlands), at pH 5.8 containing 1% sugar, 0.5 g/L MES buffer and 0.7% agar. Seeds were sterilized briefly with 70% ethanol followed by 6% bleach for 5 to 10 min and finally rinsed in water. Plants were kept at 21°C in 16/8 h light period under a light intensity of 70–90 μmol m⁻²s⁻¹. The knockout T-DNA insertion line SALK_009478 was obtained from NASC. The *pid-14* mutant is SALK_049736 as reported by Haga et al. (2014). The two *PID^{OE}* lines, P10 and P21, were developed by Benjamins et al. (2001) by cloning the *PID* cDNA in sense orientation behind the strong Cauliflower Mosaic Virus 35S promoter (35S::PID) and introducing this construct into *Arabidopsis thaliana* ecotype Columbia (Col).

Drought Stress Application

Seeds (stratified in cold and dark) were sown in separate pots containing an equal amount of soil (Tref substrate). Pots were well watered initially to assist germination. To avoid heterogeneity, trays with pots were rotated 90° on a daily basis. Watering was subsequently withheld 4–5 days after germination. Three conditions were generated, where soil relative water content (RWC) was maintained by weighing pots daily and adjusted to desired values by controlled watering: control pots with 70% RWC, mild stress pots with 45% RWC and severe stress pots at 40% RWC. The experiment continued until 25 days after stratification (DAS). Rosette pictures were taken for

measurements with a Canon EOS 40D digital camera equipped with a Canon EF-S 60mm f/2.8 USM macro lens.

For *in vitro* osmotic stress, seeds were sown on half strength MS medium containing different concentrations of mannitol (Duchefa, Netherlands) and Sorbitol (Duchefa, Netherlands) and allowed to grow until 25 DAS. Digital rosette pictures were analyzed using ImageJ¹.

Hormone Measurements

To measure concentrations of IAA, abscisic acid (ABA), gibberellins, salicylic acid, cytokinins and jasmonic acid, whole rosettes were harvested at various time points between 7 to 22 DAS. The samples for three replicates were obtained from multiple plates in each experiment. Dissected samples were collected, frozen in liquid nitrogen, and ground using a MagNA Lyser (Roche) with 2 mm glass beads. The detailed hormone extraction procedures and measurements can be found in the Supplementary Material and Methods section.

qPCR Analysis

Quantitative PCR using SyBr green was performed at 9 and 22 DAS using the first pair of leaves. RNA was isolated using Purelink Plant RNA reagent (Ambion Life Technologies) and quantified with a nanodrop NZ 1000 (Thermo Scientific). An average of 1 µg of RNA was used for first strand cDNA synthesis using RQ1 RNase-Free DNase treatment and the GoScriptTM Reverse Transcription System (Promega). Actin 8 primers and gene-specific primers spanning the intron region were used at 55°C as annealing temperature (Supplementary Table 1). A SyBr green assay for qPCR was accomplished according to the developer's protocol using ROX SYBR Mastermix blue dTTP (Takyon) and a Step one plus Thermocycler (Life technologies). This experiment was done in three independent biological and three technical repetitions. The results were analyzed as $\Delta\Delta CT$ comparison with the StepOnePlusTM Real-Time PCR System (Life TechnologiesTM) software with a confidence level set at 95%.

RNA Sequencing

18 RNA samples (three genotypes, three replicates, two time points), originating from the first pair of leaves at 9 DAS and 16 DAS from wild type (WT) and both 35S::PID lines were sequenced using an IlluminaTM platform. Prior to library preparation the RNA quality and integrity was assessed according to IlluminaTM guidelines. Library preparation was done using the TruSeq[®] Stranded mRNA sample preparation 96-reaction kit (IlluminaTM) following the low sample protocol according to manufacturer's recommendations. In brief, approximately 2.5 µg of total RNA was diluted and purified using RNA purification beads targeting the poly-A tail of the mRNA and this was subsequently fragmented by means of the enzymes provided in the kit. After the cDNA synthesis adenylation of 3' ends and ligation of the adaptors were performed. Adaptors were ligated in 12-plex formations, allowing the pooling of 12 samples. Subsequently, the library was quantified using PicoGreen[®] dye (Life TechnologiesTM) as described in the manufacturer's

protocol. In order to accurately quantify the concentration in nM of the sample, the Kapa SYBR[®] FAST universal qPCR kit (Kapa BiosystemsTM) for IlluminaTM sequencing was used to quantify the number of the amplifiable molecules in the sample and the Bioanalyzer[®] machine (Agilent TechnologiesTM) to determine the average fragment size of the sample. These measurements allowed optimizing the flow cell clustering and proceeding with the Run. The sample was 50 bp pair-end sequenced in one lane of an IlluminaTM HiSeq1500 sequencer.

RNA Sequencing Data Analysis

Resulting sequence data was preliminary analyzed by CLC Genomics Workbench v.6 using the *Arabidopsis thaliana* (Col-0 TAIR10) sequence database² as reference genome. The RNA-Seq analysis was carried out for sequence reads obtained from the three genotypes. Throughout the analysis with CLC default settings were used. Briefly, after the trimming of the sequences they were mapped against the reference genome with the default settings. The expression values were calculated based on "reads per kilo base of exon model per million mapped reads" (RPKM) values (Mortazavi et al., 2008). The RNA-seq data was grouped accordingly and two group comparisons (unpaired) were performed. The expression values were normalized by scaling to the default setting of 10 million reads. Moderated t statistics for pairwise contrasts were calculated using the Baggerly's test (Baggerly et al., 2003). Genes with no counts in all three replicates for at least one of the genotype/time combinations were discarded as not detectable above the background. Statistical analysis for the effect of genotype and time was conducted by a two-way ANOVA on the MeV software (Multi Experiment Viewer 4.9.0). The Baggerly's p values were corrected for multiple testing for each contrast separately by means of false discovery rate (FDR; Benjamini and Hochberg, 1995) for significant genes based on ANOVA. FDR corrected p-value < 0.05 and log₂ of fold change > 0.75 was used as a cutoff. All significantly induced or repressed genes with known functions were classified into groups based on gene ontology information obtained from the TAIR Database³ using MapMan (Thimm et al., 2004) and overrepresented functions and gene enrichment studies were carried out in Cytoscape using the BiNGO plugin (Maere et al., 2005).

Metabolite Analyses

Malondialdehyde (MDA) and H₂O₂ were determined by the thiobarbituric acid-MDA (TBA-MDA) assay (Hodges et al., 1999) and xynol orange-based FOX1 method (Jiang et al., 1990), respectively, from a 100 mg frozen sample. Antioxidants were extracted by homogenizing 100 mg sample in 1.4 ml 80% ethanol (v/v). Total antioxidant capacity was measured after mixing the tissue extract with ferric reducing/antioxidant power (FRAP) assay reagent (0.3 M acetate buffer, pH 3.6, 0.01 mM TPTZ in 0.04 mM Hal, 0.02 M FeCl₃·6H₂O) at 600 nm using a microplate reader (Synergy Mx, Biotek Instruments Inc., Winooski, VT,

¹<http://rsbweb.nih.gov/ij/>

²<http://www.arabidopsis.org/>

³www.arabidopsis.org

United States) (Benzie and Strain, 1996). Trolox was used as standard.

Enzyme Assays

Glycolate oxidase was measured according to Feierabend and Beevers (1972). GO was measured by the formation of a glyoxylate complex with phenylhydrazine ($\epsilon_{324} = 17 \text{ mM}^{-1} \text{ cm}^{-1}$). NADPH oxidase (oxidative stress-related enzyme) was assayed according to Sarath et al. (2007) measuring NADPH-dependent superoxide generation as the reduction rate of NBT into monoformazan ($\epsilon_{530} = 12.8 \text{ mM}^{-1} \text{ cm}^{-1}$). Enzymes were measured in extracts obtained from 100 mg of frozen tissue, in 1 mL of extraction buffer: 50 mM MES/KOH (pH 6.0) containing 0.04 M KCl, 2 mM CaCl_2 and homogenized with a MagNA Lyser (Roche, Vilvorde, Belgium). Catalase activity was assayed by observing the H_2O_2 decomposition rate ($\epsilon_{240} = 39.4 \text{ M}^{-1} \text{ cm}^{-1}$) (Aebi, 1984). SOD activity was determined according to Dhindsa et al. (1982) by measuring the inhibition of NBT reduction at 560 nm. SOD and CAT measurements were scaled down for semi-high throughput measurement using a micro-plate reader (Synergy Mx, Biotek Instruments Inc., Winooski, VT, United States). Glutathione S-transferase (GST) activity was determined according to Habig et al. (1974). Enzyme activity was calculated by measuring the conjugation of GSH with excess 1-chloro-2,4-dinitrobenzene (CDNB) in 50 mM potassium phosphate buffer at 340 nm. The activity of glutaredoxin enzymes was measured according to Holmgren and Aslund (1995). The assay contains tris-HCl buffer (100 mM, pH 8.0), 100 $\mu\text{g}/\text{mL}$ bovine serum albumin, 1 mM GSH, 6 $\mu\text{g}/\text{mL}$ yeast glutathione reductase, 0.1 M Tris-HCl, 2 mM EDTA, 0.4 mM NADPH and 0.7 mM of 2-hydroxy-ethyl-disulfide (HED). The peroxiredoxin activity was performed by measuring the decrease in H_2O_2 concentration in the reaction mixture contained 100 mM K_2HPO_4 buffer (pH 7.0), 0.3 to 3 mM Prx, 10 mM DTT, and 100 μM H_2O_2 (Horling et al., 2003). The reaction was stopped with 12.5% trichloroacetic acid (TCA). To quantify the reduction in H_2O_2 content, 10 mM $\text{Fe}(\text{NH}_4)_2(\text{SO}_4)_2$ and 2.5 mM KSCN were added and the absorbance was measured at 480 nm. Ferredoxin enzyme activity measured in a 50 mM HEPES-KOH (pH 8.0) contains 5 mM MgCl_2 0.3 mM NADP^+ , 3 mM glucose 6-phosphate, 1 unit/mL glucose 6-phosphate dehydrogenase and 1 mM potassium ferricyanide. The decrease in NADP^+ was measured at 420 nm.

Superoxide Detection in Leaves

Superoxide radical detection was performed using the nitroblue tetrazolium (NBT) staining method (Ramel et al., 2009). NBT reacts with O_2^- to form a dark blue insoluble formazan compound. Arabidopsis plants were infiltrated in 0.1% (w/v) NBT solution in PBS buffer (0.2 g KCl + 0.2 g KH_2PO_4 + 1.15 g NaH_2PO_4 + 8 g NaCl in 1000 ml of H_2O ; pH 7.1) containing 10 mM of sodium azide (NaN_3), for 30 min under vacuum. After 4 h of incubation they were cleared and pictured under a Nikon AZ-100 macroscope equipped with a Nikon DS-Ri1 digital camera.

Statistical Analysis

Rosette and auxin measurements were analyzed by *t*-test using the R statistical package⁴. Conditions of normality of distribution and homogeneity of variance were checked and met. Metabolite and enzyme data were analyzed using SPSS (SPSS Inc, Chicago, IL, United States). Conditions of normality of distribution and homogeneity of variance were not examined due to the small sample size. One-way analyses of variance (ANOVA) were performed on the original data to evaluate the differences between genotypes and time (days). Significant differences between means were determined with the Duncan test ($P < 0.05$).

Accession Number

The Arabidopsis Genome Initiative locus identifier for the *PID* gene is AT2G34650. Raw RNA sequencing data were deposited in NCBI's Gene Expression Omnibus as GEO accession no. GSE82086.

RESULTS


Overexpression of *PID* Leads to Smaller Rosettes and Elevated Auxin Pools

In an attempt to deepen our knowledge on auxin-regulated leaf growth and development, we chose to analyze two *PID*^{OE} lines, P10 and P21, which clearly had smaller rosettes compared to WT at 22 DAS (days after stratification; **Figure 1A** and Saini et al., 2017). To quantify the *PID* transcript levels between WT and *PID*^{OE} lines and to understand the differences in the phenotype between P10 and P21, we performed qPCR. Relative quantification of *PID* transcript levels revealed that P21 had a two-fold higher *PID* overexpression level than P10 at 9 DAS, whereas at 22 DAS this was inverted and *PID* overexpression in P21 was four-fold lower compared to P10 (**Figure 1B**). The severity of the phenotype and *PID* transcript abundance differences between P10 and P21 suggests that *PID* affects rosette growth in a dose-dependent manner (**Figures 1A,B**). Since *PID* affects auxin transport and thus its distribution (Christensen et al., 2000; Raftopoulou, 2004; Kleine-Vehn et al., 2009), we measured the total pool of IAA (free IAA and IAA-conjugates). IAA levels were strongly elevated in the rosettes of both *PID*^{OE} lines compared to the WT from 7 to 22 DAS (**Figure 1C**). Significantly higher IAA levels in P10 and P21 from 12 DAS onward reflect the difference in expression levels and that a certain threshold of *PID* abundance is required to sustain these high IAA levels (**Figures 1B,C**).

Transcriptome Data Point to the Induction of Stress in *PID* Overexpression Lines

To gain further insight into *PID* function, we checked the expression levels of the AT2G34650 gene (*PID*) under various conditions in Arabidopsis using perturbation as condition search tool in Genevestigator. Publicly available microarray data

⁴<https://www.r-project.org/>



revealed that *PID* is differentially expressed in response to several abiotic stress conditions (Figure 2A). We performed RNA sequencing on the first pair of leaves of *PID*^{OE} lines and WT at 9 and 16 DAS to understand the small-rosette phenotype (Supplementary Data 1). In total there were 3805 genes that expressed differentially in at least one genotype and one time point (FDR < 0.05 and log₂ fold change > 0.75; Supplementary Data 2). Consistent with the link to abiotic stress suggested by Genevestigator, gene enrichment analysis of the 3805 differentially expressed genes (DEGs), using BiNGO (Maere et al., 2005), showed a significant overrepresentation of ABA signaling, abiotic and biotic stress-related genes, stimulus to chemical and stress, including response to water, water deprivation and osmotic stress (Figure 2B and Supplementary Data 3). Moreover, a MapMan (Thimm et al., 2004) plot of the DEGs also showed that many stress and redox-related genes were among induced or repressed genes in P10 and P21 (Figure 2C). The MapMan analysis also revealed that similar categories were changed in P10 and P21, but at a more pronounced level (more genes and higher fold changes) in P10. This is consistent with the *PID* expression level of P10 being higher than in P21 at 16 DAS when the transcriptome analysis was performed (Figure 1B).

Multiple Hormonal Responses Are Modulated in *PID* Overexpression Plants

Physiological and molecular mechanisms underlying plant growth and stress responses are complex and likely involve feedback mechanism and crosstalk with other hormones. Consistently, several auxin metabolism, transport and signaling-related genes were affected by *PID* overexpression that point at a perturbed auxin homeostasis and signaling. These include *GH3*, IAA-leucine conjugate hydrolase (*ILR1*), indole-3-butyric acid response (*IBR 1* and *2*), PINs, auxin/indole-3-acetic acid (Aux/IAAs), small auxin up RNAs (SAURs), Lateral Organ Boundaries-Domain (LBDs), and auxin responsive factors (ARFs) (Supplementary Data 2). Auxin is known to modulate the metabolism of other hormones (Nemhauser et al., 2006). In our data, besides auxin, other hormone signaling components were also modulated in the overexpression lines (Supplementary Figure 1). The expression level of signature genes of several hormones (Nemhauser et al., 2006) were significantly affected in at least one of the *PID*^{OE} lines on one time point: ABA-induced carotenoid dioxygenase (*NCED3*) and abscisic acid responsive elements-binding factor (*AREB1/ABF2*, *AREB2/ABF4*, and *ABF3*), gibberellic acid (GA) inducible, *RGL1*,





FIGURE 2 | Effect of abiotic stress on *PID* expression levels and of *PID* overexpression on stress-related genes. Genevestigator analysis shows up- (yellow) and downregulation (blue) of *PID* under various abiotic stress conditions (A). Gene enrichment analysis of differentially expressed genes in *PID* overexpression lines vs. WT, visualized via the Cytoscape plugin BINGO. The inset shows a representation of all 3805 genes, while the major overrepresented genes (orange part of the inset) are magnified (B). MapMan analysis of up- (blue squares) and downregulated (red squares) genes in P10 and P21 at 16 DAS compared to WT, shows the upregulation of biotic and abiotic stress-related genes and changes in redox status (C).




and jasmonic acid induced, lipoxygenases *LOX1* and *LOX2* were all upregulated, whereas cytokinin-induced response regulators, *ARR4* and *UGT76C2*, were downregulated (see Supplementary Table 2 for full list). Thus, these data suggest that the accumulation of IAA in the *PID*^{OE} lines leads to changes in signaling components of other hormones as well. Consistently, there was an overrepresentation of stress-related genes that are reported to function in ABA-dependent stress responses. This includes ABA insensitive (*ABI1* and *ABI2*), *AREB1/ABF2* and *AREB2/ABF4* and *ABF3* (Murata et al., 2001; Abe et al., 2003; Fujita et al., 2005; Yoshida et al., 2010). Multiple genes involved in stress regulation that are auxin inducible, such as *RAB18*, *RD22*, *RD29A*, *RD29B* (Shi et al., 2014; Supplementary Table 2), were also differentially expressed in our data, suggesting upregulation of hormone-related stress responses in *PID*^{OE} lines.

To study whether indeed the levels of other hormones were affected in the *PID*^{OE} lines, we quantified ABA, GA, cytokinins, salicylic acid, and jasmonic acid in the rosettes (Figure 3). Both *PID*^{OE} lines showed elevated ABA levels at 9 and 16 DAS compared to the WT, while P10 also maintained significantly high ABA levels at 22 DAS (Figure 3A). Thus ABA concentration varied in parallel with *PID* expression levels and the auxin accumulation. *PID*^{OE} lines had slightly increased GA4 levels at 16 DAS, while there was a clear increase in cytokinins levels

at 16 and 22 DAS. Salicylic acid and GA19 levels were not significantly changed between the genotypes (Figures 3B–E), whereas other GAs were either absent or below the detection limit. Also jasmonic acid concentrations were below the detection limit (1.41 pmol/g).

PID Overexpression Lines Show Changes in the Activity of Stress Markers

To verify the transcriptome data and study the relation between stress and *PID*, we tested the effect of altered *PID* expression on cellular stress markers using both *pid* knockouts and overexpression mutants. Being a common response to most adverse abiotic conditions, we looked at the parameters related to oxidative stress (Polle and Rennenberg, 1993). Photorespiration, NADPH oxidase, apoplastic peroxidases, acyl-CoA oxidase, and mitochondrial electron transport are among important ROS sources and some have also been involved in responses related to IAA (Elstner, 1982; Wingerl et al., 2000; Gill and Tuteja, 2010; Sandalio et al., 2016). We measured the parameters that were also differentially expressed in our transcriptome data (Supplementary Table 3). Our global transcript data suggest an upregulation of photorespiration (Supplementary Table 3), and both *PID*^{OE} lines showed elevated GO activities at 16 DAS and P10 at 22 DAS (Figure 4A). The elevation in GO activity was



closely linked to *PID* expression levels (Figure 1B). Compared to the WT, NADPH oxidase activity was increased in both the *PID*^{OE} lines at 16 and 22 DAS (Figure 4B). Unlike in the *PID*^{OE} lines, the activities of both enzymes were largely unchanged in *pid* mutants relative to the WT. Oxidative damage to the cell membranes can be measured in terms of MDA levels. Our results show increased MDA levels in the overexpression lines, but these were unchanged in *pid* mutants (Figure 4C), clearly suggesting lipid peroxidation in the membranes of *PID*^{OE} lines, unlike in the membranes of the WT and *pid* knockouts. H₂O₂, quantified using the xylenol orange-based FOX1 method (Jiang et al., 1990), showed a gradual increase from 16 to 22 DAS to six-fold higher levels in P10 at 22 DAS, while the mutants and P21 showed no differences compared to the WT (Figure 4D). Rosettes and leaves from all the three genotypes were stained for superoxides and both P10 and P21 showed high staining based on the reaction between NBT and O²⁻, leading to the formation of the blue colored formazan compound. The staining was most prominent at 9 DAS and particularly enriched in the top-half of the leaves of *PID*^{OE} lines, while


it was spread across the leaf blade in the WT suggesting a relation between auxin accumulation (i.e., top leaf blade in *PID*^{OE} lines) and superoxide radicals (Saini et al., 2017; Supplementary Figure 2).

PID Overexpression Lines Have High Cellular Antioxidant Levels

Given the dual role of ROS as signaling molecules and molecules causing cellular damage, plants strictly regulate ROS levels in various cellular compartments by means of various antioxidant systems (Noctor and Foyer, 1998; Arora et al., 2002; Neill et al., 2002; Foreman et al., 2003; Foyer and Noctor, 2005). Therefore, we determined the concentrations of some chemical antioxidants and enzymatic antioxidants in *pid* mutants, *PID*^{OE} lines and the WT. The transcripts related to phenylpropanoid and flavonoid biosynthesis pathways showed higher abundance in *PID*^{OE} lines compared to the WT in the transcriptome data (Supplementary Figure 3 and Data 2). Since the results could indicate higher concentrations of the end products, we measured the concentrations of flavonoids and anthocyanins along with the total antioxidant capacity in all the genotypes. Indeed, the total antioxidant capacity, reflecting overall changes in the radical scavenging capacity, was higher in P21 lines at 16 DAS and in both lines at 22 DAS compared to the WT (Figure 5A). In addition to the enhanced total antioxidant capacity, also the levels of one or several groups of antioxidant molecules increased gradually (over time, from 16 to 22 DAS) and considerably in at least one genotype and time point in *PID*^{OE} lines. This is particularly clear for flavonoids, anthocyanins, and polyphenols, more so in P10 than P21 (Figures 5B–D), which is again in congruence with comparatively higher auxin levels, *PID* transcript and ROS accumulation in P10 (Figures 1, 4D). Contrary to *PID*^{OE} lines, antioxidants levels were unchanged in *pid* knockout mutants compared to the WT, with exception of increased flavonoids in *pid-14* at 16 DAS. At the level of enzymatic antioxidants, with the exception of catalase (CAT), a large number of enzymes showed moderate changes in *pid* knockouts and overexpression lines [superoxide dismutase (SOD), peroxidases (POX), glutaredoxins, ferredoxins, and peroxiredoxins), while levels of GSTs, enzymes involved in detoxification of xenobiotic compounds (Marrs, 1996), were unchanged in all the genotypes (Supplementary Figure 4). These results demonstrate an altered antioxidant status as a consequence of altered *PID* expression.

PID Overexpression Lines Show Modulated Response to External Stresses

Many reports showed that modulation of auxin and antioxidant levels confer resistance or tolerance to drought or osmotic stress (Tognetti et al., 2010; Lee et al., 2012; Espinoza et al., 2013; Kim et al., 2013; Cha et al., 2014; Nakabayashi et al., 2014; Shi et al., 2014; Avramova et al., 2015, 2017; Islam et al., 2015). To investigate if the *PID*-mediated auxin responses in stress regulation affected whole plant responses to abiotic stresses, we



subjected *PID*^{OE} lines to osmotic and drought stress. Mannitol and sorbitol were used to lower the water potential in the media (Verslues et al., 2006; Claeys et al., 2014). Rosette area measurements were made at 25 DAS when the first pair of leaves have reached maturity (Beemster et al., 2005). Increasing concentrations of sorbitol or mannitol reduced rosette growth in WT (Figures 6A–D). All the lines showed growth reduction although both *PID*^{OE} lines, especially P10, showed less impact of the water deficit.

To further confirm the *in vitro* responses to low water potentials, plants were subjected to different water availabilities in soil by altered watering regimes. WT, P10, and P21 showed a growth reduction under mild (45% RWC) and severe (40% RWC) drought stress. Growth reductions in P10 were similar to WT and stronger in P21 (Figures 6E,F). After 4 weeks, WT plants had a 100% survival rate (i.e., number of plants surviving at 25 DAS), while the survival in P10 and P21 was reduced (90 and 80% in mild stress; 55 and 35% in severe stress, respectively).


DISCUSSION

PID Overexpression Perturbs the Homeostasis of Auxin and Other Hormones, Inducing Stress Responses in Arabidopsis Rosettes

Previously, we elaborated on the fact that *PID* overexpression causes reduced growth in Arabidopsis leaves due to elevated auxin levels. On the other hand, *pid* knockouts displayed similar rosette growth and total auxin levels as the WT (Bennett et al., 1995; Christensen et al., 2000; Saini et al., 2017). In order to get a better understanding of the molecular changes induced by the altered *PID* expression levels we performed RNA sequencing on WT and *PID*^{OE} lines. Our transcriptome data suggested differential expression of genes related to auxin metabolism and signaling in *PID*^{OE} lines (Supplementary Figure 1 and Data 2). Since *PID* is a regulator of *PAT* (Benjamins et al., 2001) and controls the subcellular localization of *PIN* proteins (Friml et al., 2004), the most likely explanation for the increased auxin levels is due to defective transport from the young leaves, which are generally considered as a site of auxin production (Ljung et al., 2001; Saini et al., 2017).

Several studies using genetic mutants with modified auxin metabolism or signaling have demonstrated a role for auxin in abiotic stress. Such studies have, so far, mostly assessed the plant's tolerance toward abiotic stresses. Here, we uniquely show that *PID*, a regulator of auxin transport, when overexpressed in Arabidopsis perturbs auxin homeostasis, which consequently leads to induction of stress responses in the rosettes (measured in terms of severity of phenotype, ROS, antioxidants accumulation and upregulation of genes known to be involved in stress and redox signaling in plants; Supplementary Data 2, 3). It is evident from public transcriptome data that *PID* is differentially expressed in stress conditions, although up to four-fold only (Figure 2), suggesting that *PID* might play a role in stress responses. However, a role for *PID* as a regulator of *PIN* polarity and thus, auxin transport, is well established and we believe that it is the auxin accumulation in the leaves that causes the reduction in rosette growth, induction of stress responses and modification of plant response toward drought treatment. Our study does strongly point to a relationship between *PID*-modulated auxin homeostasis and signaling, and the observed stress responses.


PID^{OE} plants also showed changes in levels of other hormones and their responsive genes, as shown by PageMan, hormone quantification and expression changes in our transcriptome data (Figure 3, Supplementary Figure 1, and Data 2). The involvement of multiple hormones suggests involvement of complex and interlinked hormonal regulatory pathways in growth and stress responses in Arabidopsis leaves, that is also frequently evidenced across literature (Mouchel et al., 2006; Nemhauser et al., 2006; Chapman and Estelle, 2009; Harrison, 2012; Lanza et al., 2012; Šimášková et al., 2015). Several reports indeed suggest that auxin conjugates could be involved in stress responses (Tognetti et al., 2010; Ludwig-Müller, 2011; Kinoshita et al., 2012). In fact a stress responsive *GH3* gene, *WES1*, is known to modulate the cross talk between auxin-SA and auxin-ABA (Park et al., 2007a,b;



Zhang et al., 2007). Upregulation of *MYB96*, a molecular link in the ABA-auxin crosstalk in stress conditions, along with other genes involved in the *MYB96* regulated pathways such as *RD-22* (drought stress inducing gene) and *GH3* suggests the involvement of auxin-ABA interactive responses in the *PID*^{OE} lines (Seo et al., 2009). However, since *PID* is only known to directly affect auxin, we primarily focus on auxin and speculate that other hormones act downstream of auxin or in concert with auxin to induce stress responses.

Like hormones, ROS are signaling molecules that co-regulate growth and development in plants and coordinate responses to environment cues (Neill et al., 2002; Mittler et al., 2004; Jaspers and Kangasjärvi, 2010). Plants respond to environmental stresses by adopting various developmental modulations that include altered growth and development (collectively known as stress

induced morphogenic responses; SIMR), reduced metabolism and increased antioxidant accumulation. In fact, plant growth is influenced by a controlled balance between ROS and hormones where auxin and ROS are seen as the key players in stress adaptive responses (Potters et al., 2007, 2009; Tognetti et al., 2012). Evidences showing frequent crosstalk between ROS signaling and hormonal networks are not uncommon. Interplay between ROS and phytohormones is evidenced in abiotic stress adaptation (Tognetti et al., 2010), induction of plant defense responses (Mühlenbock et al., 2008), programmed cell death (Kuriyama and Fukuda, 2002; Gechev et al., 2004), growth and developmental aspects such as cell cycle and cell elongation (Hirt, 2000; D'Haese et al., 2003; Teale et al., 2006) and regulation of stomatal aperture (Neill et al., 2002). Auxin induced changes in ROS levels are shown to facilitate root gravitropism (Joo et al., 2001), stomatal



opening (Song et al., 2006) and cell elongation (Schopfer, 2001). Several sources of ROS, i.e., NADPH oxidases, photo respiratory enzymes, apoplastic peroxidases, acyl-CoA oxidase, and mitochondrial electron transport, are also involved in IAA regulatory networks (Sandalio et al., 2016). *PID^{OE}* lines had elevated ROS levels that can possibly be explained by increased photorespiration, but also by increased NADPH oxidase activity (Peer et al., 2013). Auxin overproducing/accumulating mutants are known to have reduced rosette growth and epinastic leaves (Boerjan et al., 1995; Delarue et al., 1998; Zhao et al., 2001). Leaf epinasty is primarily controlled by auxin and involves ROS-auxin interplay as well (Sandalio et al., 2016). *PID^{OE}* lines also show enhanced auxin and ROS and downward leaf curling (Christensen et al., 2000; this study). We believe that the high auxin levels in *PID^{OE}* lines could be causing high ROS production, resulting in growth retardation and distinct morphological phenotypes in the leaves (Figure 7).

Apart from growth reduction, plants adapt to abiotic stress by mechanisms such as stomatal closure, accumulation of osmolytes, antioxidants, and the induction of LEA proteins (Verslues et al., 2006). Similarly, in response to induced stress conditions due to *PID^{OE}*, various antioxidants appeared to buffer ROS accumulation in plants. The increase in overall antioxidant capacity is probably caused by increase in flavonoids, anthocyanins, and other polyphenols. Consistently, most of the genes related to flavonoids/anthocyanin biosynthesis were also upregulated in the *PID^{OE}* lines (Supplementary Figure 3 and Data 2). Flavonoids and anthocyanins are plant secondary metabolites produced via the phenylpropanoid pathway that

affect several developmental processes including protection against UV, ROS, etc. (Taylor and Grotewold, 2005; Grotewold, 2006). Flavonoids are known to alter auxin transport and promote localized auxin accumulation in a tissue-specific manner (Peer et al., 2004; Peer and Murphy, 2006, 2007; Kuhn et al., 2011; Buer et al., 2013). Kuhn et al. (2017) showed that flavonol (a subgroup of flavonoids) accumulation in the *rol1-2* mutant, which is deficient in rhamnose synthase, affected shoot development and altered export of naphthalene-1-acetic (NAA), but not of IAA. Many reports suggest that flavonoids affect auxin transport directly or indirectly by modifying vesicular trafficking and PIN cycling, other auxin efflux proteins like ATP-binding cassette transporter superfamily (ABCB), or by modifying activities of PAT regulators, or protein phosphatase 2A (PP2A) and its antagonist PID (Peer et al., 2004; Terasaka et al., 2005; Geisler and Murphy, 2006; Kuhn et al., 2017). Clearly flavonoids are modulators of PAT, however, anthocyanins seem to have little or no effect on auxin transport (reviewed in Peer and Murphy, 2007). Interestingly, auxin accumulation is shown to promote flavonoid accumulation, presumably to quench ROS signal generated during auxin catabolism (Peer and Murphy, 2007; Peer et al., 2011). Similarly, enhanced flavonoid synthesis in auxin accumulating *PID^{OE}* lines could also help to scavenge elevated ROS molecules. Another interesting assumption is that gradual accumulation of flavonoids in the leaves of *PID^{OE}* lines further promotes localized auxin accumulation in the leaves, which is evident in both P10 and P21 (Figures 1C, 5B, 7).

In addition to increases in molecular antioxidants, also the activity of a relatively large range of antioxidant enzymes is, at

least moderately, modified. The decreased activity of enzymatic antioxidants such as peroxidases, peroxiredoxins could also be seen as contributory factor in ROS accumulation in *PID^{OE}* lines. The comparison of changes at the level of transcripts and enzyme activity is difficult to establish or rather inconclusive as most of redox-related genes were downregulated (Supplementary Table 3). The observation that a relatively large number of antioxidant enzymes and molecules change (mostly increase), suggest that a wide range of cellular redox processes are affected in the *PID^{OE}* lines, rather than any specific process. The fact that changes in specific metabolites (anthocyanins) and enzymes (GO) are closely related to *PID* overexpression levels, indicates a positive correlation between them. We believe that the differences between two overexpression lines itself are because of the differences in *PID* expression levels and thus auxin levels and the fact that even transient changes in auxin can hugely impact cellular and molecular events in the plant (Nemhauser et al., 2006; Paponov et al., 2008; Saini et al., 2017). These data clearly indicate the presence of enhanced (oxidative) stress responses, in *PID^{OE}* lines compared to *pid* knockouts, as shown by measurements of ROS, ROS producing and detoxifying enzymes, MDA and antioxidant molecules. To summarize, it is well established that environmental stresses impact cellular ROS levels, antioxidant concentration and their redox state (Mittler, 2002; Apel and Hirt, 2004; Hong-bo et al., 2008). Here, we demonstrate that in the absence of external stress, ROS and antioxidants levels change in response to cellular auxin perturbations, and modulate growth and stress adaptive responses.

***PID* Overexpression Does Not Confer Tolerance to Water Stress**

There are many reports providing links between abiotic stress, antioxidant abundance and auxin (Jung and Park, 2011; Hasanuzzaman et al., 2013; Rahman, 2013; Avramova et al., 2015, 2017). For example, activation of *YUCCA7* elevates IAA levels and enhances resistance to drought in *Arabidopsis* (Lee et al., 2012). However, as a result of the overexpression of this gene, growth was reduced and plants had narrow and curled leaves. Similarly, overexpression of *OsPIN3t* increases water tolerance in rice (Zhang et al., 2012). *Phot1* (Wan et al., 2012), a close homolog of PINOID belonging to the same AGC kinase family (Galván-Ampudia and Offringa, 2007), is also shown to improve drought tolerance in *Arabidopsis* seedlings (Galen et al., 2007). Similarly, overaccumulation of flavonoids is shown to enhance tolerance to drought and osmotic stress (Nakabayashi et al., 2014), and anthocyanins are also suggested to have an osmoprotectant role among others, such as ROS scavenging (Hughes et al., 2010, 2013).

In our results, all three genotypes responded to osmotic stress by reducing their rosette growth (Figure 6). The differences in growth reduction between *PID^{OE}* and WT could be due to already smaller size of non-stressed *PID^{OE}* rosettes that allowed a lower degree of growth reduction when subjected to osmotic stress conditions. Both *PID^{OE}* lines showed lower survival than the WT during drought stress assays in the soil. We believe that

the increased IAA levels, due to *PID* overexpression, already induce stress conditions in the rosettes and that the water deficit treatment is additive, thereby leading to higher lethality. Another explanation could be that *PID^{OE}* lines have much shorter roots and reduced numbers of lateral roots (Benjamins et al., 2001), making their quest for water in the soil very difficult, which could have resulted in poor performance under drought stress. Evidently, the shoot/root ratio and stress responses are closely related (Woo et al., 2007; Werner et al., 2010). This suggests that *PID*-induced auxin alterations induce the observed stress responses and clearly do not confer advantage in water deficit conditions.

To conclude, various reports show that plant AGC kinases participate in the response to biotic and abiotic stresses. We demonstrated that ectopic expression of *PID* causes changes in auxin levels and its response, and consequentially ROS is generated and other hormones are affected which results in the generation of growth inhibitory stress responses. Additionally, in contrast to several previous reports, our study uniquely shows that despite their high auxin and antioxidant levels, plants are impaired in their tolerance to osmotic and drought stress and show a low survival under severe drought conditions. For future work it will be interesting to know how direct the relation between *PID* and stress responses is and whether auxin accumulation in leaves is necessary or sufficient to cause the observed stress responses.

AUTHOR CONTRIBUTIONS

KS planned and performed most of the experiments, analyzed the data and wrote the article with contribution from HAS, EP, GB, and KV; HAb and SS provided assistance with data analysis; MM performed next generation sequencing; EP, GB, and KV conceived the project and KV supervised the research.

FUNDING

This work was supported by a concerted research activity (GOA) research grant, 'A Systems Biology Approach of Leaf Morphogenesis' from the research council of the University of Antwerp. This work was supported by the Research Foundation Flanders (FWO-Vlaanderen; G.0656.13N, 1.5.091.11.N.00).

ACKNOWLEDGMENTS

The 35S::*PID10* and 35S::*PID21* mutants were kindly provided by Eva Benkova (IST Austria). We sincerely thank Sevgi Öden and Tim Willems (Biology Department, UA) for the hormone measurements.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fpls.2017.01308/full#supplementary-material>

REFERENCES

- Aebi, H. (1984). Oxygen radicals in biological systems. *Methods Enzymol.* 105, 121–126. doi: 10.1016/S0076-6879(84)05016-3
- Abe, H., Urao, T., Ito, T., Seki, M., and Shinozaki, K. (2003). Transcriptional activators in abscisic acid signaling. *Society* 15, 63–78. doi: 10.1105/tpc.006130.salt
- Apel, K., and Hirt, H. (2004). Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* 55, 373–399. doi: 10.1146/annurev.arplant.55.031903.141701
- Arora, A., Sairam, R. K., and Srivastava, G. C. (2002). Oxidative stress and oxidative system in plants. *Curr. Sci.* 82, 1227–1238.
- Avramova, V., AbdElgawad, H., Vasileva, I., Petrova, A. S., Holek, A. S., Mariën, J., et al. (2017). High antioxidant activity facilitates maintenance of cell division in leaves of drought tolerant maize hybrids. *Front. Plant Sci.* 8:84. doi: 10.3389/fpls.2017.00084
- Avramova, V., AbdElgawad, H., Zhang, Z., Fotschki, B., Casadevall, R., Vergauwen, L., et al. (2015). Drought induces distinct growth response, protection, and recovery mechanisms in the maize leaf growth zone. *Plant Physiol.* 169, 1382–1396. doi: 10.1104/pp.15.00276
- Baggerly, K. A., Deng, L., Morris, J. S., and Aldaz, C. M. (2003). Differential expression in SAGE: accounting for normal between-library variation. *Bioinformatics* 19, 1477–1483. doi: 10.1093/bioinformatics/btg173
- Beemster, G. T. S., De Veylder, L., Vercruyse, S., West, G., Rombaut, D., Van Hummelen, P., et al. (2005). Genome-wide analysis of gene expression profiles associated with cell cycle transitions in growing organs of *Arabidopsis*. *Plant Physiol.* 138, 734–743. doi: 10.1104/pp.104.053884
- Benjamini, Y., and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc.* 57, 289–300. doi: 10.2307/2346101
- Benjamins, R., Quint, A., Weijers, D., Hooykaas, P., and Offringa, R. (2001). The PINOID protein kinase regulates organ development in *Arabidopsis* by enhancing polar auxin transport. *Development* 128, 4057–4067.
- Benjamins, R., and Scheres, B. (2008). Auxin: the looping star in plant development. *Annu. Rev. Plant Biol.* 59, 443–465. doi: 10.1146/annurev.arplant.58.032806.103805
- Bennett, R. M. S., Alvarez, J., Gerd, B., and Smyth, D. R. (1995). Morphogenesis in pinoid mutants of *Arabidopsis thaliana*. *Plant J.* 8, 505–520.
- Benzie, I. F. F., and Strain, J. J. (1996). The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: the FRAP assay. *Anal. Biochem.* 239, 70–76. doi: 10.1006/abio.1996.0292
- Blomster, T., Salojärvi, J., Sipari, N., Brosche, M., Ahlfors, R., Keinänen, M., et al. (2011). Apoplastic reactive oxygen species transiently decrease auxin signaling and cause stress-induced morphogenic. *Plant Physiol.* 157, 1866–1883. doi: 10.1104/pp.111.181883
- Boerjan, W., Cervera, M. T., Delarue, M., Beeckman, T., Dewitte, W., Bellini, C., et al. (1995). Superroot, a recessive mutation in *Arabidopsis*, confers auxin overproduction. *Plant Cell* 7, 1405–1419. doi: 10.1105/tpc.7.9.1405
- Buer, C. S., Kordbacheh, F., Truong, T. T., Hocart, C. H., and Djordjevic, M. A. (2013). Alteration of flavonoid accumulation patterns in transparent testa mutants disturbs auxin transport, gravity responses, and imparts long-term effects on root and shoot architecture. *Planta* 238, 171–189. doi: 10.1007/s00425-013-1883-3
- Cha, J. Y., Kim, J. Y., Jung, I. J., Kim, M. R., Melencion, A., Alam, S. S., et al. (2014). NADPH-dependent thioredoxin reductase A (NTRA) confers elevated tolerance to oxidative stress and drought. *Plant Physiol. Biochem.* 80, 184–191. doi: 10.1016/j.plaphy.2014.04.008
- Chapman, E. J., and Estelle, M. (2009). Cytokinin and auxin intersection in root meristems. *Genome Biol.* 10:210. doi: 10.1186/gb-2009-10-2-210
- Christensen, S. K., Dagenais, N., Chory, J., and Weigel, D. (2000). Regulation of auxin response by the protein kinase PINOID. *Cell* 100, 469–478.
- Clayes, H., Van Landeghem, S., Dubois, M., Maleux, K., and Inzé, D. (2014). What is stress? Dose-response effects in commonly used in vitro stress assays. *Plant Physiol.* 165, 519–527. doi: 10.1104/pp.113.234641
- Delarue, M., Prinsen, E., Van Onckelen, H., Caboche, M., and Bellini, C. (1998). Sur2 mutations of *Arabidopsis thaliana* define a new locus involved in the control of auxin homeostasis. *Plant J.* 14, 603–611. doi: 10.1046/j.1365-313X.1998.00163.x
- Devarenne, T. P., Ekengren, S. K., Pedley, K. F., and Martin, G. B. (2006). Adi3 is a Pdk1-interacting AGC kinase that negatively regulates plant cell death. *EMBO J.* 25, 255–265. doi: 10.1038/sj.emboj.7600910
- D’Haeze, W., De Rycke, R., Mathis, R., Goormachtig, S., Pagnotta, S., Verplancke, C., et al. (2003). Reactive oxygen species and ethylene play a positive role in lateral root base nodulation of a semiaquatic legume. *Proc. Natl. Acad. Sci. U.S.A.* 100, 11789–11794. doi: 10.1073/pnas.1333899100
- Dhindsa, R. S., Plumb-Dhindsa, P. L., and Reid, D. M. (1982). Leaf senescence and lipid peroxidation: effects of some phytohormones, and scavengers of free radicals and singlet oxygen. *Physiol. Plant.* 56, 453–457. doi: 10.1111/j.1399-3054.1982.tb04539.x
- Du, H., Liu, H., and Xiong, L. (2013). Endogenous auxin and jasmonic acid levels are differentially modulated by abiotic stresses in rice. *Front. Plant Sci.* 4:397. doi: 10.3389/fpls.2013.00397
- Elstner, E. F. (1982). Oxygen activation and oxygen toxicity. *Annu. Rev. Plant Physiol.* 33, 73–96.
- Enders, T. A., and Strader, L. C. (2015). Auxin activity: past, present, and future. *Am. J. Bot.* 102, 180–196. doi: 10.3732/ajb.1400285
- Espinoza, A., San Martín, A., López-Climent, M., Ruiz-Lara, S., Gómez-Cadenas, A., and Casaretto, J. A. (2013). Engineered drought-induced biosynthesis of α -tocopherol alleviates stress-induced leaf damage in tobacco. *J. Plant Physiol.* 170, 1285–1294. doi: 10.1016/j.jplph.2013.04.004
- Feierabend, J., and Beevers, H. (1972). Developmental studies on microbodies in wheat leaves. *Plant Physiol.* 49, 28–32.
- Foreman, J., Demidchik, V., Bothwell, J. H. F., Mylona, P., Miedema, H., Torres, M. A., et al. (2003). Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. *Nature* 422, 442–446. doi: 10.1038/nature01485
- Foyer, C. H., and Noctor, G. (2005). Oxidant and antioxidant signalling in plants: a re-evaluation of the concept of oxidative stress in a physiological context. *Plant Cell Environ.* 28, 1056–1071. doi: 10.1111/j.1365-3040.2005.01327.x
- Friml, J., Yang, X., Michniewicz, M., Weijers, D., Quint, A., Tietz, O., et al. (2004). A PINOID-dependent binary switch in apical-basal PIN polar targeting directs auxin efflux. *Science* 306, 862–865. doi: 10.1126/science.1100618
- Fujita, Y., Fujita, M., Satoh, R., Maruyama, K., Parvez, M. M., Seki, M., et al. (2005). AREB1 is a transcription activator of novel ABRE-dependent ABA signaling that enhances drought stress tolerance in *Arabidopsis*. *Plant Cell* 17, 3470–3488. doi: 10.1105/tpc.105.035659
- Galen, C., Rabenold, J. J., and Liscum, E. (2007). Functional ecology of a blue light photoreceptor: effects of phototropin-1 on root growth enhance drought tolerance in *Arabidopsis thaliana*. *New Phytol.* 173, 91–99. doi: 10.1111/j.1469-8137.2006.01893.x
- Galván-Ampudia, C. S., and Offringa, R. (2007). Plant evolution: AGC kinases tell the auxin tale. *Trends Plant Sci.* 12, 541–547. doi: 10.1016/j.tplants.2007.10.004
- García, A. V., Al-Yousif, M., and Hirt, H. (2012). Role of AGC kinases in plant growth and stress responses. *Cell. Mol. Life Sci.* 69, 3259–3267. doi: 10.1007/s00018-012-1093-3
- Gazarian, I. G., Lagrimini, L. M., Mellon, F. A., Naldrett, M. J., Ashby, G. A., and Thorneley, R. N. F. (1998). Identification of skatolyl hydroperoxide and its role in the peroxidase-catalysed oxidation of indol-3-yl acetic acid. *Biochem. J.* 333, 223–232.
- Gechev, T. S., Gadjev, I. Z., and Hille, J. (2004). An extensive microarray analysis of AAL-toxin-induced cell death in *Arabidopsis thaliana* brings new insights into the complexity of programmed cell death in plants. *Cell. Mol. Life Sci.* 61, 1185–1197. doi: 10.1007/s00018-004-4067-2
- Geisler, M., and Murphy, A. S. (2006). The ABC of auxin transport: the role of p-glycoproteins in plant development. *FEBS Lett.* 580, 1094–1102. doi: 10.1016/j.febslet.2005.11.054
- Gill, S. S., and Tuteja, N. (2010). Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol. Biochem.* 48, 909–930. doi: 10.1016/j.plaphy.2010.08.016
- Grotewold, E. (ed.). (2006). *The Science of Flavonoids*. Berlin: Springer. doi: 10.1007/978-0-387-28822-2
- Habig, W. H., Pabst, M. J., and Jakoby, W. B. (1974). Glutathione S-Transferases. *J. Biol. Chem.* 249, 7130–7139.
- Haga, K., Hayashi, K.-I., and Sakai, T. (2014). PINOID AGC kinases are necessary for phytochrome-mediated enhancement of hypocotyl phototropism in *Arabidopsis*. *Plant Physiol.* 166, 1535–1545. doi: 10.1104/pp.114.244434

- Harrison, M. A. (2012). "Cross-talk between phytohormone signaling pathways under both optimal and stressful environmental conditions," in *Phytohormones and Abiotic Stress Tolerance in Plants*, eds N. A. Khan, R. Nazar, N. Iqbal, and N. A. Anjum (Berlin: Springer Science & Business Media), 1–49. doi: 10.1007/978-3-642-25829-9_2
- Hasanuzzaman, M., Nahar, K., Gill, S. S., and Fujita, M. (2013). "Drought stress responses in plants, oxidative stress, and antioxidant defense," in *Climate Change and Plant Abiotic Stress Tolerance*, eds N. Tuteja and S. S. Gill (Weinheim: Wiley-VCH Verlag GmbH & Co), 209–250. doi: 10.1002/9783527675265.ch09
- Hirt, H. (2000). Connecting oxidative stress, auxin, and cell cycle regulation through a plant mitogen-activated protein kinase pathway. *Proc. Natl. Acad. Sci. U.S.A.* 97, 2405–2407. doi: 10.1073/pnas.97.6.2405
- Hirt, H., Garcia, A. V., and Oelmüller, R. (2011). AGC kinases in plant development and defense. *Plant Signal. Behav.* 6, 1030–1033. doi: 10.4161/psb.6.7.15580
- Hodges, D. M., DeLong, J. M., Forney, C. F., and Prange, R. K. (1999). Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. *Planta* 207, 604–611. doi: 10.1007/s004250050524
- Holmgren, A., and Aslund, F. (1995). *Biothiols Part B: Glutathione and Thioredoxin: Thiols in Signal Transduction and Gene Regulation*. Amsterdam: Elsevier. doi: 10.1016/0076-6879(95)52031-7
- Hong-bo, S., Li-ye, C., Ming-an, S., Abdul, J. C., and Hong-mei, M. (2008). Higher plant antioxidants and redox signaling under environmental stresses. *C. R. Biol.* 331, 433–441. doi: 10.1016/j.crvi.2008.03.011
- Horling, F., Lamkemeyer, P., König, J., Finkemeier, I., Kandlbinder, A., Baier, M., et al. (2003). Dependent regulation of expression of the peroxiredoxin gene family in Arabidopsis. *Plant Physiol.* 131, 317–325. doi: 10.1104/pp.010017
- Hughes, N. M., Carpenter, K. L., and Cannon, J. G. (2013). Estimating contribution of anthocyanin pigments to osmotic adjustment during winter leaf reddening. *J. Plant Physiol.* 170, 230–233. doi: 10.1016/j.jplph.2012.09.006
- Hughes, N. M., Reinhardt, K., Feild, T. S., Gerardi, A. R., and Smith, W. K. (2010). Association between winter anthocyanin production and drought stress in angiosperm evergreen species. *J. Exp. Bot.* 61, 1699–1709. doi: 10.1093/jxb/erq042
- Islam, T., Manna, M., and Reddy, M. K. (2015). Glutathione peroxidase of *Pennisetum glaucum* (PpGPx) is a functional Cd²⁺ dependent peroxiredoxin that enhances tolerance against salinity and drought stress. *PLoS ONE* 10:e0143344. doi: 10.1371/journal.pone.0143344
- Jain, M., and Khurana, J. P. (2009). Transcript profiling reveals diverse roles of auxin-responsive genes during reproductive development and abiotic stress in rice. *FEBS J.* 276, 3148–3162. doi: 10.1111/j.1742-4658.2009.07033.x
- Jaspers, P., and Kangasjärvi, J. (2010). Reactive oxygen species in abiotic stress signaling. *Physiol. Plant.* 138, 405–413. doi: 10.1111/j.1399-3054.2009.01321.x
- Jiang, Z. Y., Woollard, A. C., and Wolff, S. P. (1990). Hydrogen peroxide production during experimental protein glycation. *FEBS Lett.* 268, 69–71. doi: 10.1016/0014-5793(90)80974-N
- Joo, J. H., Bae, Y. S., and Lee, J. S. (2001). Role of auxin-induced reactive oxygen species in root gravitropism. *Plant Physiol.* 126, 1055–1056.
- Jung, J.-H., and Park, C.-M. (2011). Auxin modulation of salt stress signaling in Arabidopsis seed germination. *Plant Signal. Behav.* 6, 1198–1200. doi: 10.4161/psb.6.8.15792
- Junghans, U., Polle, A., DÜchting, P., Weiler, E., Kuhlman, B., Gruber, F., et al. (2006). Adaptation to high salinity in poplar involves changes in xylem anatomy and auxin physiology. *Plant Cell Environ.* 29, 1519–1531. doi: 10.1111/j.1365-3040.2006.01529.x
- Kazan, K. (2013). Auxin and the integration of environmental signals into plant root development. *Ann. Bot.* 112, 1655–1665. doi: 10.1093/aob/mct229
- Kim, J. I., Baek, D., Park, H. C., Chun, H. J., Oh, D. H., Lee, M. K., et al. (2013). Overexpression of arabidopsis YUCCA6 in potato results in high-auxin developmental phenotypes and enhance. *Mol. Plant* 6, 337–349. doi: 10.1093/mp/sss100
- King, J. J., Stimart, D. P., Fisher, R. H., and Bleeker, A. B. (1995). A mutation altering auxin homeostasis and plant morphology in Arabidopsis. *Plant Cell* 7, 2023–2037.
- Kinoshita, N., Wang, H., Kasahara, H., Liu, J., MacPherson, C., Machida, Y., et al. (2012). IAA-Ala resistant3, an evolutionarily conserved target of miR167, mediates Arabidopsis root architecture changes during high osmotic stress. *Plant Cell* 24, 3590–3602. doi: 10.1105/tpc.112.097006
- Kleine-Vehn, J., Huang, F., Naramoto, S., Zhang, J., Michniewicz, M., Offringa, R., et al. (2009). PIN auxin efflux carrier polarity is regulated by PINOID kinase-mediated recruitment into GNOM-independently trafficking in Arabidopsis. *Plant Cell* 21, 3839–3849. doi: 10.1105/tpc.109.071639
- Kuhn, B. M., Geisler, M., Bigler, L., and Ringli, C. (2011). Flavonols accumulate asymmetrically and affect auxin transport in Arabidopsis. *Plant Physiol.* 156, 585–595. doi: 10.1104/pp.111.175976
- Kuhn, B. M., Nodzyński, T., Errafi, S., Bucher, R., Gupta, S., Aryal, B., et al. (2017). Flavonol-induced changes in PIN2 polarity and auxin transport in the Arabidopsis thaliana roll-2 mutant require phosphatase activity. *Sci. Rep.* 7:41906. doi: 10.1038/srep41906
- Kuriyama, H., and Fukuda, H. (2002). Developmental programmed cell death in plants. *Curr. Opin. Plant Biol.* 5, 568–573. doi: 10.1016/S1369-5266(02)00305-9
- Lanza, M., Garcia-Ponce, B., Castrillo, G., Catarecha, P., Sauer, M., Rodriguez-Serrano, M., et al. (2012). Role of actin cytoskeleton in brassinosteroid signaling and in its integration with the auxin response in plants. *Dev. Cell* 22, 1275–1285. doi: 10.1016/j.devcel.2012.04.008
- Lee, M., Jung, J.-H., Han, D.-Y., Seo, P. J., Park, W. J., and Park, C.-M. (2012). Activation of a flavin monooxygenase gene YUCCA7 enhances drought resistance in Arabidopsis. *Planta* 235, 923–938. doi: 10.1007/s00425-011-1552-3
- Ljung, K., Bhalerao, R. P., and Sandberg, G. (2001). Sites and homeostatic control of auxin biosynthesis in Arabidopsis during vegetative growth. *Plant J.* 28, 465–474. doi: 10.1046/j.1365-313X.2001.01173.x
- Ludwig-Müller, J. (2011). Auxin conjugates: their role for plant development and in the evolution of land plants. *J. Exp. Bot.* 62, 1757–1773. doi: 10.1093/jxb/erq412
- Maere, S., Heymans, K., and Kuiper, M. (2005). BiNGO: a Cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks. *Bioinformatics* 21, 3448–3449. doi: 10.1093/bioinformatics/bti551
- Marrs, K. A. (1996). The functions and regulation of glutathione S-transferases in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 47, 127–158.
- Mittler, R. (2002). Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.* 7, 405–410. doi: 10.1016/S1360-1385(02)02312-9
- Mittler, R., Vanderauwera, S., Gollery, M., and Van Breusegem, F. (2004). Reactive oxygen gene network of plants. *Trends Plant Sci.* 9, 490–498. doi: 10.1016/j.tplants.2004.08.009
- Mortazavi, A., Williams, B. A., McCue, K., Schaeffer, L., and Wold, B. (2008). Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat. Methods* 5, 621–628. doi: 10.1038/nmeth.1226
- Mouchel, C. F., Osmont, K. S., and Hardtke, C. S. (2006). BRX mediates feedback between brassinosteroid levels and auxin signalling in root growth. *Nature* 443, 458–461. doi: 10.1038/nature05130
- Mühlenbock, P., Szechynska-Hebda, M., Plaszczyca, M., Baudo, M., Mateo, A., Mullineaux, P. M., et al. (2008). Chloroplast signaling and lesion simulating disease1 regulate crosstalk between light acclimation and immunity in Arabidopsis. *Plant Cell* 20, 2339–2356. doi: 10.1105/tpc.108.059618
- Murata, Y., Pei, Z., Mori, I. C., and Schroeder, J. (2001). Abscisic acid activation of plasma membrane Ca²⁺ channels in guard cells requires cytosolic NAD(P)H and is differentially disrupted upstream and downstream of reactive oxygen species production in abi1-1 and abi2-1 protein phosphatase 2C mutants. *Plant Cell* 13, 2513–2523. doi: 10.1105/tpc.010210.2514
- Nakabayashi, R., Yonekura-Sakakibara, K., Urano, K., Suzuki, M., Yamada, Y., Nishizawa, T., et al. (2014). Enhancement of oxidative and drought tolerance in Arabidopsis by overaccumulation of antioxidant flavonoids. *Plant J.* 77, 367–379. doi: 10.1111/tpj.12388
- Naser, V., and Shani, E. (2016). Auxin response under osmotic stress. *Plant Mol. Biol.* 91, 661–672. doi: 10.1007/s11103-016-0476-5
- Neill, S., Desikan, R., and Hancock, J. (2002). Hydrogen peroxide signalling. *Curr. Opin. Plant Biol.* 5, 388–395. doi: 10.1016/S1369-5266(02)00282-0
- Nemhauser, J. L., Hong, F., and Chory, J. (2006). Different plant hormones regulate similar processes through largely nonoverlapping transcriptional responses. *Cell* 126, 467–475. doi: 10.1016/j.cell.2006.05.050
- Noctor, G., and Foyer, C. H. (1998). Ascorbate and glutathione: keeping active oxygen under control. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 49, 249–279. doi: 10.1146/annurev.arplant.49.1.249

- Paponov, I. A., Paponov, M., Teale, W., Menges, M., Chakrabortee, S., Murray, J. A. H., et al. (2008). Comprehensive transcriptome analysis of auxin responses in *Arabidopsis*. *Mol. Plant* 1, 321–337. doi: 10.1093/mp/ssm021
- Park, J.-E., Park, J.-Y., Kim, Y.-S., Staswick, P. E., Jeon, J., Yun, J., et al. (2007a). GH3-mediated Auxin homeostasis links growth regulation with stress adaptation response in *Arabidopsis*. *J. Biol. Chem.* 282, 10036–10046. doi: 10.1074/jbc.M610524200
- Park, J. E., Seo, P. J., Lee, A. K., Jung, J. H., Kim, Y. S., and Park, C. M. (2007b). An *Arabidopsis* GH3 gene, encoding an auxin-conjugating enzyme, mediates phytochrome B-regulated light signals in hypocotyl growth. *Plant Cell Physiol.* 48, 1236–1241. doi: 10.1093/pcp/pcm086
- Pearce, L. R., Komander, D., and Alessi, D. R. (2010). The nuts and bolts of AGC protein kinases. *Nat. Rev. Mol. Cell Biol.* 11, 9–22. doi: 10.1038/nrm2822
- Peer, W. A., Bandyopadhyay, A., Blakeslee, J. J., Makam, S. N., Chen, R. J., Masson, P. H., et al. (2004). Variation in expression and protein localization of the PIN family of auxin efflux facilitator proteins in flavonoid mutants with altered auxin transport in *Arabidopsis thaliana*. *Plant Cell* 16, 1898–1911. doi: 10.1105/tpc.021501
- Peer, W. A., Blakeslee, J. J., Yang, H., and Murphy, A. S. (2011). Seven things we think we know about auxin transport. *Mol. Plant* 4, 487–504. doi: 10.1093/mp/ssr034
- Peer, W. A., Cheng, Y., and Murphy, A. S. (2013). Evidence of oxidative attenuation of auxin signalling. *J. Exp. Bot.* 64, 2629–2639. doi: 10.1093/jxb/ert152
- Peer, W. A., and Murphy, A. S. (2006). “Flavonoids as signal molecules,” in *The Science of Flavonoids*, ed. E. Grotewold (Berlin: Springer), 239–268.
- Peer, W. A., and Murphy, A. S. (2007). Flavonoids and auxin transport: modulators or regulators? *Trends Plant Sci.* 12, 556–563. doi: 10.1016/j.tplants.2007.10.003
- Petersen, L. N., Ingle, R. A., Knight, M. R., and Denby, K. J. (2009). OXII protein kinase is required for plant immunity against *Pseudomonas syringae* in *Arabidopsis*. *J. Exp. Bot.* 60, 3727–3735. doi: 10.1093/jxb/erp219
- Polle, A., and Rennenberg, H. (1993). “Significance of antioxidants in plant adaptation to environmental stress,” in *Plant Adaptations to Environmental Stresses*, eds L. Fowden, T. Mansfield, and J. Stoddart (London: Chapman & Hall), 263–273.
- Potters, G., Pasternak, T. P., Guisez, Y., and Jansen, M. A. K. (2009). Different stresses, similar morphogenic responses: integrating a plethora of pathways. *Plant Cell Environ* 32, 158–169. doi: 10.1111/j.1365-3040.2008.01908.x
- Potters, G., Pasternak, T. P., Guisez, Y., Palme, K. J., and Jansen, M. A. K. (2007). Stress-induced morphogenic responses: growing out of trouble? *Trends Plant Sci.* 12, 98–105. doi: 10.1016/j.tplants.2007.01.004
- Raftopoulou, M. (2004). PINOID pinpoints auxin. *Nat. Cell Biol.* 6, 1149–1149. doi: 10.1038/ncb1204-1149
- Rahman, A. (2013). Auxin: a regulator of cold stress response. *Physiol. Plant* 147, 28–35. doi: 10.1111/j.1399-3054.2012.01617.x
- Ramel, F., Sulmon, C., Bogard, M., Coué, I., and Gouesbet, G. (2009). Differential patterns of reactive oxygen species and antioxidative mechanisms during atrazine injury and sucrose-induced tolerance in *Arabidopsis thaliana* plantlets. *BMC Plant Biol.* 9:28. doi: 10.1186/1471-2229-9-28
- Remy, E., Cabrito, T. R., Baster, P., Batista, R. A., Teixeira, M. C., Friml, J., et al. (2013). A major facilitator superfamily transporter plays a dual role in polar auxin transport and drought stress tolerance in *Arabidopsis*. *Plant Cell* 25, 901–926. doi: 10.1105/tpc.113.110353
- Rentel, M. C., Lecourieux, D., Ouaked, F., Usher, S. L., Petersen, L., Okamoto, H., et al. (2004). OXII kinase is necessary for oxidative burst-mediated signalling in *Arabidopsis*. *Nature* 427, 858–861. doi: 10.1038/nature02353
- Saini, K., Markakis, M. N., Zdanio, M., Balcerowicz, D. M., Beekman, T., De Veylder, L., et al. (2017). Alteration in auxin homeostasis and signaling by overexpression of PINOID kinase causes leaf growth defects in *Arabidopsis thaliana*. *Front. Plant Sci.* 8:1009. doi: 10.3389/fpls.2017.01009
- Sandalio, L. M., Rodríguez-Serrano, M., and Romero-Puertas, M. C. (2016). Leaf epinasty and auxin: a biochemical and molecular overview. *Plant Sci.* 253, 187–193. doi: 10.1016/j.plantsci.2016.10.002
- Sarath, G., Hou, G., Baird, L. M., and Mitchell, R. B. (2007). Reactive oxygen species, ABA and nitric oxide interactions on the germination of warm-season C4-grasses. *Planta* 226, 697–708. doi: 10.1007/s00425-007-0517-z
- Schopfer, P. (2001). Hydroxyl radical-induced cell-wall loosening in vitro and in vivo: implications for the control of elongation growth. *Plant J.* 28, 679–688. doi: 10.1046/j.1365-313x.2001.01187.x
- Seo, P. J., Xiang, F., Qiao, M., Park, J.-Y., Lee, Y. N., Kim, S.-G., et al. (2009). The MYB96 transcription factor mediates abscisic acid signaling during drought stress response in *Arabidopsis*. *Plant Physiol.* 151, 275–289. doi: 10.1104/pp.109.144220
- Shen, C., Bai, Y., Wang, S., Zhang, S., Wu, Y., Chen, M., et al. (2010). Expression profile of PIN, AUX/LAX and PGP auxin transporter gene families in *Sorghum bicolor* under phytohormone and abiotic stress. *FEBS J.* 277, 2954–2969. doi: 10.1111/j.1742-4658.2010.07706.x
- Shi, H., Chen, L., Ye, T., Liu, X., Ding, K., and Chan, Z. (2014). Modulation of auxin content in *Arabidopsis* confers improved drought stress resistance. *Plant Physiol. Biochem.* 82, 209–217. doi: 10.1016/j.plaphy.2014.06.008
- Shibasaki, K., Uemura, M., Tsurumi, S., and Rahman, A. (2009). Auxin response in *Arabidopsis* under cold stress: underlying molecular mechanisms. *Plant Cell* 21, 3823–3838. doi: 10.1105/tpc.109.069906
- Šimášková, M., O’Brien, J. A., Khan, M., Van Noorden, G., Ötvös, K., Vieten, A., et al. (2015). Cytokinin response factors regulate PIN-FORMED auxin transporters. *Nat. Commun.* 6:8717. doi: 10.1038/ncomms9717
- Song, X.-G., She, X.-P., He, J.-M., Huang, C., and Song, T. (2006). Cytokinin- and auxin-induced stomatal opening involves a decrease in levels of hydrogen peroxide in guard cells of *Vicia faba*. *Funct. Plant Biol.* 33, 573. doi: 10.1071/FP05232
- Taylor, L. P., and Grotewold, E. (2005). Flavonoids as developmental regulators. *Curr. Opin. Plant Biol.* 8, 317–323. doi: 10.1016/j.pbi.2005.03.005
- Teale, W. D., Paponov, I. A., and Palme, K. (2006). Auxin in action: signalling, transport and the control of plant growth and development. *Nat. Rev. Mol. Cell Biol.* 7, 847–859. doi: 10.1038/nrm2020
- Terasaka, K., Blakeslee, J. J., Titapiwatanakun, B., Peer, W. A., Bandyopadhyay, A., Makam, S. N., et al. (2005). PGP4, an ATP binding cassette P-glycoprotein, catalyzes auxin transport in *Arabidopsis thaliana* roots. *Plant Cell* 17, 2922–2939. doi: 10.1105/tpc.105.035816
- Thimm, O., Bläsing, O., Gibon, Y., Nagel, A., Meyer, S., Krüger, P., et al. (2004). Mapman: a user-driven tool to display genomics data sets onto diagrams of metabolic pathways and other biological processes. *Plant J.* 37, 914–939. doi: 10.1111/j.1365-313X.2004.02016.x
- Tognetti, V. B., Mühlentock, P., and van Breusegem, F. (2012). Stress homeostasis - the redox and auxin perspective. *Plant Cell Environ.* 35, 321–333. doi: 10.1111/j.1365-3040.2011.02324.x
- Tognetti, V. B., Van Aken, O., Morreel, K., Vandenbroucke, K., van de Cotte, B., De Clercq, I., et al. (2010). Perturbation of indole-3-Butyric acid homeostasis by the UDP-glucosyltransferase UGT74E2 modulates *Arabidopsis* architecture and water stress tolerance. *Plant Cell* 22, 2660–2679. doi: 10.1105/tpc.109.071316
- Verslues, P. E., Agarwal, M., Katiyar-Agarwal, S., Zhu, J., and Zhu, J.-K. (2006). Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J.* 45, 523–539. doi: 10.1111/j.1365-313X.2005.02593.x
- Wan, Y., Jasik, J., Wang, L., Hao, H., Volkman, D., Menzel, D., et al. (2012). The signal transducer NPH3 integrates the phototropin1 photosensor with PIN2-based polar auxin transport in *Arabidopsis* root phototropism. *Plant Cell* 24, 551–565. doi: 10.1105/tpc.111.094284
- Wang, S., Bai, Y., Shen, C., Wu, Y., Zhang, S., Jiang, D., et al. (2010). Auxin-related gene families in abiotic stress response in *Sorghum bicolor*. *Funct. Integr. Genomics* 10, 533–546. doi: 10.1007/s10142-010-0174-3
- Werner, T., Nehnevajova, E., Kollmer, I., Novak, O., Strnad, M., Kramer, U., et al. (2010). Root-specific reduction of cytokinin causes enhanced root growth, drought tolerance, and leaf mineral enrichment in *Arabidopsis* and tobacco. *Plant Cell* 22, 3905–3920. doi: 10.1105/tpc.109.072694
- Wingler, A., Lea, P. J., Quick, W. P., and Leegood, R. C. (2000). Photorespiration: metabolic pathways and their role in stress protection. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 355, 1517–1529. doi: 10.1098/rstb.2000.0712
- Woo, Y. M., Park, H. J., Su’udi, M., Yang, J. I., Park, J. J., Back, K., et al. (2007). Constitutively wilted 1, a member of the rice YUCCA gene family, is required for maintaining water homeostasis and an appropriate root to shoot ratio. *Plant Mol. Biol.* 65, 125–136. doi: 10.1007/s11103-007-9203-6
- Yoshida, T., Fujita, Y., Sayama, H., Kidokoro, S., Maruyama, K., Mizoi, J., et al. (2010). AREB1, AREB2, and ABF3 are master transcription factors that cooperatively regulate ABRE-dependent ABA signaling involved in drought stress tolerance and require ABA for full activation. *Plant J.* 61, 672–685. doi: 10.1111/j.1365-313X.2009.04092.x

- Zhang, Q., Li, J., Zhang, W., Yan, S., Wang, R., Zhao, J., et al. (2012). The putative auxin efflux carrier OsPIN3t is involved in the drought stress response and drought tolerance. *Plant J.* 72, 805–816. doi: 10.1111/j.1365-313X.2012.05121.x
- Zhang, S.-W., Li, C.-H., Cao, J., Zhang, Y.-C., Zhang, S.-Q., Xia, Y.-F., et al. (2009). Altered architecture and enhanced drought tolerance in rice via the down-regulation of indole-3-acetic acid by TLD1/OsGH3.13 activation. *Plant Physiol.* 151, 1889–1901. doi: 10.1104/pp.109.146803
- Zhang, Z., Li, Q., Li, Z., Staswick, P. E., Wang, M., Zhu, Y., et al. (2007). Dual regulation role of GH3.5 in salicylic acid and auxin signaling during *Arabidopsis-Pseudomonas syringae* interaction. *Plant Physiol.* 145, 450–464. doi: 10.1104/pp.107.106021
- Zhao, Y., Christensen, S. K., Fankhauser, C., Cashman, J. R., Cohen, J. D., Weigel, D., et al. (2001). A role for flavin monooxygenase-like enzymes in auxin biosynthesis. *Science* 291, 306–309. doi: 10.1126/science.291.5502.306
- Zwiewka, M., Nodzyński, T., Robert, S., Vanneste, S., and Friml, J. (2015). Osmotic stress modulates the balance between exocytosis and clathrin-mediated endocytosis in *Arabidopsis thaliana*. *Mol. Plant* 8, 1175–1187. doi: 10.1016/j.molp.2015.03.007

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Saini, AbdElgawad, Markakis, Schoenaers, Asard, Prinsen, Beemster and Vissenberg. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.