



Draft Genome Sequence of *Bacillus* sp. Strain K2I17, Isolated from the Rhizosphere of *Deschampsia antarctica* Desv.

Pavón, Alequis; Orellana, Paz; Salazar, Lorena; Céspedes, Sandra; Muiño, Laura; Gutiérrez, Ana; Castillo Bermúdez, Daniel Elías; Corsini, Gino

Published in:
Genome Announcements

DOI:
[10.1128/genomeA.00786-17](https://doi.org/10.1128/genomeA.00786-17)

Publication date:
2017

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Pavón, A., Orellana, P., Salazar, L., Céspedes, S., Muiño, L., Gutiérrez, A., ... Corsini, G. (2017). Draft Genome Sequence of *Bacillus* sp. Strain K2I17, Isolated from the Rhizosphere of *Deschampsia antarctica* Desv. *Genome Announcements*, 5(33), [e00786-17]. <https://doi.org/10.1128/genomeA.00786-17>



Draft Genome Sequence of *Bacillus* sp. Strain K2I17, Isolated from the Rhizosphere of *Deschampsia antarctica* Desv.

Alequis Pavón,^a Paz Orellana,^a Lorena Salazar,^a Sandra Céspedes,^a
Laura Muiño,^a Ana Gutiérrez,^{a,b} Daniel Castillo,^c Gino Corsini^a

Facultad de Ciencias de la Salud, Instituto de Ciencias Biomédicas, Universidad Autónoma de Chile, Santiago, Chile^a; BIOREN-UFRO and Departamento de Producción Agropecuaria, Facultad de Ciencias Agropecuarias y Forestales, Universidad de la Frontera, Temuco, Chile^b; Marine Biological Section, University of Copenhagen, Helsingør, Denmark^c

ABSTRACT We present here the draft genome sequence of *Bacillus* sp. strain K2I17, which was isolated from the rhizosphere of *Deschampsia antarctica* Desv. The genomic sequence contained 6,113,341 bp. This genome provides insights into the possible new biomedical and biotechnical applications of this specific Antarctic bacterium.

The genus *Bacillus* is a phenotypically large, diverse collection of Gram-positive, endospore-forming, aerobic or facultatively anaerobic bacteria that have undergone considerable reclassification due to high phylogenetic heterogeneity (1). In addition, this genus is distributed widely in extreme environments and includes psychrophilic, thermophilic, acidophilic, alkalophilic, and halophilic bacteria that utilize a wide range of carbon sources for growth (2, 3).

Bacillus spp. comprise one of the most interesting groups of bacteria in biotechnology due to their resistance to pH and temperature changes, which are important parameters in industrial processes (4). To date, the presence of *Bacillus* spp. has been demonstrated in different environments, including the rhizosphere, the layer of soil influenced by plant-root metabolism. In comparison to root-free soil or bulk soil, the rhizosphere displays higher contents of nutrients, which could explain the activity of *Bacillus* spp. in these dynamic environments (5), and thus new approaches have begun to utilize the biotechnological applications of *Bacillus* and related species for agricultural and pharmaceutical products (6). Here, we announce the draft genome sequence of *Bacillus* sp. strain K2I17, which was isolated from the rhizosphere of *Deschampsia antarctica* Desv., at the Collins Glacier, Fildes Peninsula, King George Island, Antarctica (62°10'S, 58°55'W).

Bacillus sp. strain K2I17 was grown at 18°C in LB broth (catalog no. 12106-05; Mo Bio, Inc.). Genomic DNA was extracted using the Power Biofilm DNA isolation kit (catalog no. 24000-50; Mo Bio, Inc.) according to the manufacturer's protocol. A sequencing library was prepared using the Illumina HiSeq 2500 platform (Macrogen, Republic of Korea) with paired-end read sizes of 100 bp. A total of 15,092,729 paired-end reads were used for the *de novo* assembly in Geneious version 9.1.6 (7). Short and low-coverage contigs were filtered out, resulting in a set of 146 with an average coverage of 89× (N_{50} , 142,492 bp). Annotation was performed with the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (8). Additionally, the genomes were analyzed on the Rapid Annotation using Subsystems Technology (RAST) server (9). Acquired antibiotic resistance genes were identified using ResFinder version 2.1 (10), virulence factors using virulencefinder version 1.2 (11), and prophage-related sequences using PHASTER (12).

Received 25 June 2017 Accepted 28 June 2017 Published 17 August 2017

Citation Pavón A, Orellana P, Salazar L, Céspedes S, Muiño L, Gutiérrez A, Castillo D, Corsini G. 2017. Draft genome sequence of *Bacillus* sp. strain K2I17, isolated from the rhizosphere of *Deschampsia antarctica* Desv. *Genome Announc* 5:e00786-17. <https://doi.org/10.1128/genomeA.00786-17>.

Copyright © 2017 Pavón et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Daniel Castillo, daniel.castillo@bio.ku.dk, or Gino Corsini, gino.corsini@uautonoma.cl.

The final assembly for *Bacillus* sp. strain K2117 had a total length of 6,113,341 bp and a G+C content of 34.9%. Genome annotation resulted in 6,125 coding sequences (CDSs), 22 tRNAs, 326 pseudogenes, and 7 rRNAs; 59% of the CDSs were classified as hypothetical proteins. Five prophage-like elements between 19 and 36 kb were detected. Interestingly, one of them encodes a putative cell-adhesion protein. Genes resistant to the antibiotics tetracycline [*tet*(M) and *tet*(O)], penicillin (*bl*, *bla*, *ampS*, *pse2*, and *ybxI*), vancomycin (*vanA* and *vanH*), fosfomycin (*fosA*, *fosB*, and *fosX*), and fluoroquinolones (*parC* and *parE*) were detected. Also, secondary metabolism genes that encode thiazole/oxazole-modified microcin synthesis, auxin biosynthesis, pyridoxine (vitamin B₆) biosynthesis, and lanthionine biosynthesis were identified. In addition, 27 membrane-bound and secreted *Listeria* internalin-like proteins (LPXTG and GW motifs) were identified. Thus, this genome sequence can facilitate the understanding of the genetic diversity within the *Bacillus* genus, which could be useful for the development of new biotechnological applications.

Accession number(s). The draft genome sequence of *Bacillus* sp. strain K2117 has been deposited in GenBank under the accession no. [NJGF00000000](https://www.ncbi.nlm.nih.gov/nuclseq/NJGF00000000). The version described in this paper is the first version, NJGF01000000.

ACKNOWLEDGMENTS

This work was supported by grants DIP66-2016, DIP92-2016, and DIUA97-2017 from the Universidad Autónoma de Chile, and by grant DIUFRO-DI14-0045 from the Universidad de la Frontera.

REFERENCES

- Logan NA, De Vos P. 2015. *Bacillus*, p 1–164. In Whitman WB (ed), *Bergey's manual of systematics of archaea and bacteria*, 2nd ed, vol 1. John Wiley and Sons, Hoboken, NJ.
- Harrell LJ, Andersen GL, Wilson KH. 1995. Genetic variability of *Bacillus anthracis* and related species. *J Clin Microbiol* 33:1847–1850.
- Maughan H, Van der Auwera G. 2011. *Bacillus* taxonomy in the genomic era finds phenotypes to be essential though often misleading. *Infect Genet Evol* 11:789–797. <https://doi.org/10.1016/j.meegid.2011.02.001>.
- Schallmey M, Singh A, Ward OP. 2004. Developments in the use of *Bacillus* species for industrial production. *Can J Microbiol* 50:1–17. <https://doi.org/10.1139/w03-076>.
- Smalla K, Wieland G, Buchner A, Zock A, Parzy J, Kaiser S, Roskot N, Heuer H, Berg G. 2001. Bulk and rhizosphere soil bacterial communities studied by denaturing gradient gel electrophoresis: plant-dependent enrichment and seasonal shifts revealed. *Appl Environ Microbiol* 67:4742–4751. <https://doi.org/10.1128/AEM.67.10.4742-4751.2001>.
- Giddings LA, Newman DJ. 2015. Bioactive compounds from terrestrial extremophiles, p 1–75. In Tiquia-Arashi SM, Mormile M (ed), *Extremophilic bacteria*, 1st ed, vol 1. Springer, New York, NY.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the rapid annotation of microbial genomes using subsystems technology (RAST). *Nucleic Acids Res* 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
- Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
- Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM, Aarestrup FM. 2014. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J Clin Microbiol* 52:1501–1510. <https://doi.org/10.1128/JCM.03617-13>.
- Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.