



Elevated CO₂ leads to enhanced photosynthesis but decreased growth in early life stages of reef building coralline algae

Ordoñez, Alexandra; Wangpraseurt, Daniel; Lyndby, Niclas Heidelberg; Kühl, Michael; Diaz-Pulida, G.

Published in:
Frontiers in Marine Science

DOI:
[10.3389/fmars.2018.00495](https://doi.org/10.3389/fmars.2018.00495)

Publication date:
2019

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Ordoñez, A., Wangpraseurt, D., Lyndby, N. H., Kühl, M., & Diaz-Pulida, G. (2019). Elevated CO₂ leads to enhanced photosynthesis but decreased growth in early life stages of reef building coralline algae. *Frontiers in Marine Science*, 5, 1-11. [495]. <https://doi.org/10.3389/fmars.2018.00495>



Elevated CO₂ Leads to Enhanced Photosynthesis but Decreased Growth in Early Life Stages of Reef Building Coralline Algae

Alexandra Ordoñez¹, Daniel Wangpraseurt^{2,3,4}, Niclas Heidelberg Lyndby^{4,5}, Michael Kühl^{4,6} and Guillermo Diaz-Pulido^{1,7*}

¹ School of Environment and Science, Australian Rivers Institute, Griffith University, Brisbane, QLD, Australia, ² Department of Chemistry, University of Cambridge, Cambridge, United Kingdom, ³ Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA, United States, ⁴ Marine Biological Section, Department of Biology, University of Copenhagen, Helsingør, Denmark, ⁵ Laboratory for Biological Geochemistry, School of Architecture, Civil and Environmental Engineering, Ecole Polytechnique Fédérale de Lausanne, Lausanne, Switzerland, ⁶ Climate Change Cluster, University of Technology Sydney, Ultimo, NSW, Australia, ⁷ ARC Centre of Excellence for Coral Reef Studies, Townsville, QLD, Australia

OPEN ACCESS

Edited by:

Laurie Carol Hofmann,
Max Planck Institute for Marine
Microbiology (MPG), Germany

Reviewed by:

Maggy Nugues,
Université de Sciences Lettres
de Paris, France
Raphael Ritson-Williams,
California Academy of Sciences,
United States

*Correspondence:

Guillermo Diaz-Pulido
g.diaz-pulido@griffith.edu.au

Specialty section:

This article was submitted to
Marine Ecosystem Ecology,
a section of the journal
Frontiers in Marine Science

Received: 29 August 2018

Accepted: 11 December 2018

Published: 07 January 2019

Citation:

Ordoñez A, Wangpraseurt D,
Lyndby NH, Kühl M and
Diaz-Pulido G (2019) Elevated CO₂
Leads to Enhanced Photosynthesis
but Decreased Growth in Early Life
Stages of Reef Building Coralline
Algae. *Front. Mar. Sci.* 5:495.
doi: 10.3389/fmars.2018.00495

Crustose coralline algae (CCA) are key organisms in coral reef ecosystems, where they contribute to reef building and substrate stabilization. While ocean acidification due to increasing CO₂ can affect the biology, physiology and ecology of fully developed CCA, the impacts of elevated CO₂ on the early life stages of CCA are much less explored. We assessed the photosynthetic activity and growth of 10-day-old recruits of the reef-building crustose coralline alga *Porolithon* cf. *onkodes* exposed to ambient and enhanced CO₂ seawater concentration causing a downward shift in pH of ~0.3 units. Growth of the CCA was estimated using measurements of crust thickness and marginal expansion, while photosynthetic activity was studied with O₂ microsensors. We found that elevated seawater CO₂ enhanced gross photosynthesis and respiration, but significantly reduced vertical and marginal growth of the early life stages of *P.* cf. *onkodes*. Elevated CO₂ stimulated photosynthesis, particularly at high irradiance, likely due to increased availability of CO₂, but this increase did not translate into increased algal growth as expected, suggesting a decoupling of these two processes under ocean acidification scenarios. This study confirms the sensitivity of early stages of CCA to elevated CO₂ and identifies complexities in the physiological processes underlying the decreased growth and abundance in these important coral reef builders upon ocean acidification.

Keywords: ocean acidification, global warming, early stages, red algae, algal physiology, photosynthesis, respiration

INTRODUCTION

Crustose coralline algae (CCA) are abundant in tropical and temperate reef ecosystems, where they provide important ecosystem functions (Littler, 1972; Littler and Doty, 1975). CCA contribute to the stabilization of coral reef frameworks by consolidating and cementing loose rubble and by sealing porous dead corals skeleton against mechanical erosion (Littler, 1972; Adey, 1998). CCA also provide food and habitat for a range of reef organisms such as parrot fish and sea urchins

(Littler and Doty, 1975; Steneck, 1983) and are preferred settlement substrates for larvae of corals and other invertebrates (Heyward and Negri, 1999; Harrington et al., 2004). The presence of CCA is thus crucial for the health and recovery of coral reefs following disturbances.

The current and projected increase in atmospheric $p\text{CO}_2$ and the derived ocean acidification (OA) due to a decrease in seawater pH and calcium carbonate (CaCO_3) saturation state (Ω) is critical for the integrity and fitness of calcifying organisms (Anthony et al., 2008; Kroeker et al., 2010). CCA are among the marine organisms most sensitive to OA as their skeleton is predominantly formed by high-magnesium calcite, a highly soluble form of CaCO_3 (Morse et al., 2007). Rates of net calcification and growth of CCA are generally negatively affected by OA (Anthony et al., 2008; Kuffner et al., 2008; Martin and Gattuso, 2009) due to reduced availability of carbonate ions (CO_3^{2-}) for calcification (Feely et al., 2004) and increased skeletal dissolution of existing calcium carbonate (Diaz-Pulido et al., 2014). On the other hand, increased atmospheric CO_2 increases the concentration of dissolved CO_2 and bicarbonate ions (HCO_3^-) in seawater potentially alleviating inorganic carbon limitation (Larkum et al., 2003) and thus stimulating algal photosynthesis. However, the available experimental evidence shows that the responses of CCA photosynthesis to OA are highly variable, including positive (Semesi et al., 2009) or negative effects (Anthony et al., 2008; Gao and Zheng, 2010; Martin et al., 2013), as well as no significant response (Noisette et al., 2013; Johnson et al., 2014; Comeau et al., 2017). This variability in response to OA may be due to a number of reasons, including use of different experimental setups and time scales (e.g., 1 year experiment with *Lithophyllum cabiochae* vs. 3 months experiment with *Lithophyllum incrustans* Martin et al., 2013; Noisette et al., 2013), different life history stages (e.g., recruits vs. adults), and flexibility of different CCA species in their use of inorganic carbon uptake strategies, i.e., use of carbon concentrating mechanisms (CCM) vs. diffusive CO_2 uptake (Cornwall et al., 2012, 2015; Diaz-Pulido et al., 2016).

Increased availability of dissolved inorganic carbon (DIC) is expected to enhance algal photosynthesis and consequently algal growth rates (Roleda et al., 2012). However, growth responses to OA have been less explored in coralline algae compared to fleshy macroalgae (e.g., Kroeker et al., 2013), and in most studies changes in CCA growth rates have been based on changes in crust weight rather than on a direct measurement of algal growth (e.g., Anthony et al., 2008; Johnson and Carpenter, 2012). Changes in crust weight have the innate difficulty of not allowing distinction between new growth and skeletal dissolution (Lewis et al., 2017). The relationship between photosynthesis and growth rates has been previously studied in fleshy macroalgae (e.g., Koch et al., 2013), where elevated CO_2 enhanced photosynthetic activity and tissue growth in two species of red algae *Gracilaria* (Gao et al., 1993), but similar relations in CCA remain unexplored.

Existing knowledge on the physiological responses of elevated CO_2 on tropical CCA comes from studies on adult crusts

(Anthony et al., 2008; Johnson et al., 2014; Comeau et al., 2017). In our previous studies using the reef building CCA *Porolithon onkodes*, we explored the impacts of OA and warming on spore germination and germling growth and found these processes to be highly sensitive to elevated CO_2 (Ordoñez et al., 2017). Roleda et al. (2015) tested the effects of OA on growth of temperate coralline algae recruits and found a direct negative response of crust size upon exposure to OA conditions. Studies of the early life stages of tropical (Kuffner et al., 2008; Fabricius et al., 2015; Ordoñez et al., 2017) and temperate (Bradassi et al., 2013; Roleda et al., 2015; Guenther et al., 2018) CCA have demonstrated a high vulnerability to elevated CO_2 , as well as other environmental stressors (Santelices, 1990). However, very little is known about the effects of OA on photosynthesis and the coupling between photosynthesis and growth rates in the early life stages of CCA (but see Cornwall et al., 2013). In the present study, we explore the effect of elevated CO_2 conditions on the photosynthetic activity and growth rates of recruits of a dominant crustose coralline alga in the Great Barrier Reef, Australia. We used O_2 microsensors due to the small size of the CCA recruits (around 0.004–0.002 mm^2 surface area), and because microsensors can decouple light respiration from gross oxygen production, as compared to chamber based incubation measurements (Revsbech and Jørgensen, 1983; Larkum et al., 2003).

MATERIALS AND METHODS

Experimental Approach

The effects of ocean acidification on rates of growth and photosynthesis of CCA recruits were investigated by exposing *Porolithon* cf. *onkodes* spores to elevated and ambient CO_2 conditions in the laboratory during a 4 week experiment. Dynamics of O_2 concentration (measured using O_2 microsensors) and crust thickness (measured using scanning electron microscopy, SEM) of 10-day-old recruits were estimated at the end of the experiment. The experiment was conducted in an indoor laboratory at Heron Island Research Station (HIRS), Great Barrier Reef (GBR), Australia during the summer of 2014 (February–March 2014). The species *P. cf. onkodes* was used for this experiment due to its important role as a reef builder and cementer (Littler, 1972), and because it is a common alga in tropical reefs, especially in the Great Barrier Reef (Ringeltaube and Harvey, 2005; Dean et al., 2015). Identification of *P. cf. onkodes* (Heydrich) Foslie was obtained with field and laboratory observations. Microscope techniques such as Scanning Electron Microscopy and histology were used to examine morphological and anatomical characteristics. The encrusting coralline alga *P. cf. onkodes* used for this experiment is usually found on the reef crest (approx. 3–5 m depth), has a pink-orange color surface with granular appearance due to the presence of numerous tightly packed mega cells in horizontal fields (trichocytes fields), reproductive structures (conceptacles) are unipored, small, flush or slightly raised, cell fusions are present and thallus has a non-coaxial organization. DNA sequences of individuals of this species can be found

in GenBank (Accession No. MF979936, see Gabrielson et al., 2018).

Cultivation of Recruits and Experimental Setup

To obtain CCA spores for the experiment, spore release from reproductive *P. cf. onkodes* fragments was induced in the laboratory following a combination of methods described in Jones and Moorjani (1973); Ichiki et al. (2000), Roleda et al. (2012), and Ordoñez et al. (2017). Adult *P. cf. onkodes* fragments (3 cm × 3 cm) were collected from the reef crest (ca 5 m depth at highest tide) at Harry's Bommie (Heron Island, GBR, 23°27'631"S, 151°55'798"E) using hammer and chisel. After collection, fragments were cleaned from epiphytes with a soft brush and rinsed with filtered sterilized seawater. Seawater was filtered twice: first using a hand-made sand filter containing a cotton filter mat and then a house water filter with a 5 µm cartridge (house water filter OMNI OPAQUE), seawater was then sterilized using an aquarium ultra violet sterilizer (Pro Aqua UV sterilizer). Spore release was then induced by placing the fragments on a tray with no water and in a dark, cold (18°C) room for 30 min. Subsequently, sterilized ambient seawater was added to the tray, which was then placed under an artificial metal-halide lamp (Aqua Medic Ocean Light Plus equipped with one 150 W aqualine bulb and two 24 W T5 blue fluorescent bulbs) for 7 h. Immediately after the illumination period, adults commenced spore release and were carefully transferred to the experimental containers (1 L plastic containers) where spore release continued under experimental conditions. Three replicate containers ($n = 3$) were used for each CO₂ treatment (6 experimental containers in total, 3 for ambient CO₂ and 3 for high CO₂). One transparent polystyrene Petri dish (94 mm × 16 mm diameter) was placed on the bottom of each experimental container to provide substratum for spore settlement and growth. Three adult fragments were randomly assigned to each experimental container to obtain sufficient spores for analysis. However, adults were removed after 4 h, when the first spore attachment was detected, for the following two reasons: (1) To avoid having individuals at different growth stages and (2) to have sufficient number of spores to get a good estimate of their individual growth, but only enough spores to avoid those that had settled close to each other to coalesce. The margin of coalesced spores are difficult to distinguish making individual measurements hard to perform.

Spores were cultured for a period of 10 days in seawater under (i) ambient CO₂ (400.8–448.1 µatm) and pH 8.00–8.03 (National Bureau of Standards/National Institute of Standards and Technology) and (ii) high CO₂ (998.9–1070.6 µatm) and pH 7.67–7.70. The high CO₂ treatment corresponded to the representative concentration pathway 8.5 scenario (RCP 8.5), i.e., the worst case OA scenario predicted by the end of the century (year 2100) according to the Intergovernmental Panel on Climate Change (IPCC, 2014). To achieve the target pH for the high CO₂ treatment, analytical grade CO₂ gas (BOC Limited Australia) was injected into a 120 L mixing sump using an aquarium control system (Aquatronica, AEB

Technologies, Italy). Seawater pH was monitored in the mixing sump by temperature compensated pH electrodes (inPro4501VP, Mettler-Toledo, Switzerland). When seawater pH exceeded the desired threshold, the control system opened solenoid valves to inject CO₂ into the mixing sump as previously described in Diaz-Pulido et al. (2011). CO₂ conditioned seawater was pumped continuously into experimental containers at a rate of 500 mL min⁻¹. The same set up was used for ambient CO₂ treatment, but with untreated seawater in the mixing sump. Mixing sumps were constantly fed with seawater from the Heron Island reef flat. pH probes were calibrated daily to 0.01 pH units with three NIST-certified pH buffers (Mettler-Toledo, Switzerland). Temperatures in both sumps (ambient CO₂ and high CO₂) and in one container of each treatment was constantly monitored every 30 s and recorded by the aquarium control system. In addition, pH and temperature measurements were frequently taken on experimental containers with a portable pH and temperature meter (SG98-B-SevenGo Duo Pro, Mettler-Toledo, Switzerland) to ensure the pH and temperature were kept constant. Illumination was provided with metal-halide lamps (Aqua Medic Ocean Light Plus equipped with one 150 W aqualine bulb and two 24 W T5 blue fluorescent bulbs) over a 12 h light: 12 h dark photoperiod under an irradiance of 160 µmol photons m⁻² s⁻¹, as measured by cosine corrected quantum sensor connected to a light meter (LICOR, United States). Total alkalinity, pH and salinity values were used to estimate the concentration of dissolved inorganic carbon (pCO₂, HCO₃⁻ and CO₃²⁻) using Microsoft Excel CO₂SYS version 2.1 (Pierrot et al., 2006). The saturation state of seawater with respect to high-Mg-calcite was calculated for a 16.4 mol% MgCO₃, following a protocol described in (Diaz-Pulido et al., 2012). Carbonate chemistry parameters are shown in Table 1.

O₂ Microsensor Measurements

Microscale O₂ concentration measurements were done with Clark-type O₂ microsensors (Revsbech, 1989) connected to a pA-meter (PA2000, Unisense A/S, Aarhus, Denmark) interfaced to a PC via an A/D converter (Figure 1). The O₂ microsensors (OX25; Unisense, Denmark) had a measuring tip diameter of 25 µm, a t_{90} response time of <0.5 s, a stirring sensitivity of the measuring signal of <2–3%, and a detection limit of ~0.3 µM¹. The O₂ microsensors were linearly calibrated from signal readings in air saturated seawater and anoxic seawater (flushed with N₂). The O₂ concentration (µmol O₂ L⁻¹) in air saturated seawater at experimental temperature and salinity was determined using tabulated values of O₂ solubility in water at defined temperature and salinity¹.

The O₂ microsensors were mounted on a PC-controlled motorized micromanipulator for automatic profiling (Pyro Science GmbH, Germany) at an angle of 15° relative to the vertically incident collimated light from a fiber-optic tungsten-halogen lamp (KL-2500, Schott GmbH, Germany), equipped

¹<http://www.unisense.com>

TABLE 1 | Summary of water chemistry parameters for the different CO₂ levels.

Treatment	Temperature °C	TA μmol kg ⁻¹	pH	pCO ₂ μatm	HCO ₃ ⁻ μmol kg ⁻¹	CO ₃ ²⁻ μmol kg ⁻¹	Ω _{High Mg Calcite} 16.4 mol% MgCO ₃
Control CO ₂	26.1 (±0.3) (25.0–27.2)	2323.7 (±4.8) (2319.35– 2333.61)	8.01 (±0.005) (8.0–8.03)	425 (±6.4) (400.8– 448.1)	1801.76 (±10.1) (1763.1–1846.1)	213.34 (±2.8) (201.7–220.9)	1.14 (±0.02) (1.06–1.19)
High CO ₂	26.1 (±0.3) (25.0–27.2)	2347.4 (±8.9) (2311.55– 2379.98)	7.69 (±0.004) (7.67–7.7)	1026.67 (±9.4) (998.9– 1070.6)	2065.06 (±7.6) (2038–2096.9)	115.95 (±1.4) (110.1–120.6)	0.61 (±0.01) (0.58–0.65)

Values represent means (±SEM) (range) for n = 8 biological replicates.




FIGURE 1 | Setup for O₂ microsensors on crustose coralline algae. General overview of the set-up showing a flow-chamber with a mounted CCA sample (A,B), with an O₂ microsensor mounted in a manual micromanipulator, illumination via a fiber-optic halogen lamp, and observation of the sample via a dissection microscope (B). (C) Close up image (via the dissection microscope) of an O₂ microsensor performing O₂ measurements on CCA recruits (pink crust).

with a heat filter and a collimating lens. Positioning of the microsensor as well as data acquisition of microsensor signals was facilitated by customized software (Profix, Pyro-Science GmbH, Germany). Experiments were conducted with CCA recruits placed in a custom-made black acrylic flow chamber supplied with seawater at a flow velocity of $\sim 1\text{--}2\text{ cm s}^{-1}$ (Brodersen et al., 2014). We used the same seawater as in the different CO₂ treatments. The downwelling photon irradiance (PAR, 400–700 nm) was measured for defined lamp settings with a calibrated cosine corrected quantum sensor connected to a light meter (LI-250A, Li-COR). For the high CO₂ treatment, the seawater taken from the treatment sumps was exchanged every 20–30 min in order to account for changes in pH related to atmospheric CO₂ exchange. The pH in the water was monitored constantly with a portable pH meter (SG98-B-SevenGo Duo Pro, Mettler-Toledo, Switzerland) calibrated with NIST-certified pH buffers (Mettler-Toledo, Switzerland) to 0.01 pH units. Recruits from 3 ambient and 2 high CO₂ experimental tanks were used for O₂ microsensor measurements. Only two replicates could be

used from high CO₂ treatments due to unexpected experimental constraints. However, around 6–9 measurements were taken on different spots within a single recruit with the objective of including and understanding spatial variations within the recruit.

After measurements of steady-state O₂ concentrations at the surface of CCA samples under defined photon irradiance levels, the microsensor tip was positioned at the CCA surface, where the local volume-specific gross photosynthesis rate (in units of $\text{nmol O}_2\text{ cm}^{-3}\text{ s}^{-1}$) was measured from the immediate O₂ depletion rate during a brief 1–3 s darkening according to the microsensor light-dark shift technique (Revsbech and Jørgensen, 1983); for these measurements, microsensor signals were recorded on a fast-responding strip-chart recorder (BD25, Kipp & Zonen, Netherlands).

The diffusive flux of O₂ between CCA and the overlaying water across the diffusive boundary layer (DBL), *J*, was determined from steady state O₂ concentration profiles using




FIGURE 2 | Scanning electron microscopy images of surface view (**A,C**) and cross section (**B,D**) of 10-day-old *Porolithon cf. onkodes* recruits developed under ambient CO₂ (CO₂ between 400.8 and 448.1 μatm, pH 8.0–8.03) and high CO₂ (CO₂ between 998.9 and 1070.6 μatm, pH 7.67–7.70) conditions. Arrows in panels (**B,D**) indicate vertical growth direction. Short white arrows in panel (**A**) indicate non-collapsed epithallial/perithallial cells and in panel (**C**) collapsed epithallial/perithallial cells. White T in panels (**A,C**) indicate first appearance of mega cells (trichocytes).

Fick's first law of diffusion: $J = -D_0 \frac{dC}{dz}$, where dC/dz is the slope of the O₂ concentration profile in the DBL, and D_0 is the molecular diffusion coefficient of O₂ in seawater at experimental temperature and salinity ($2.1707 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$), as taken from tabulated values².

To examine the effects of elevated CO₂ on the photosynthetic performance of the CCA, we measured gross and net photosynthesis as a function of increasing photon irradiance (P-E curves). Each specimen was incubated for about 15 min at each irradiance level before microsensor measurements commenced to provide steady state O₂ conditions. Gross photosynthesis data was fitted to an exponential function (Webb et al., 1974): $P = P_{\max} * 1 - e^{-\alpha * \frac{E}{P_{\max}}}$, where P_{\max} is the maximum photosynthetic rate and α is the photosynthetic efficiency, i.e., the initial slope of the P-E curve. The saturation irradiance (E_k) was calculated as $E_k = P_{\max}/\alpha$, and describes the photon irradiance above which photosynthesis approaches saturation. Net photosynthesis data were fitted to the modified exponential function used in Roberts et al. (2002): $P = P_{\max} * 1 - e^{-\alpha * \frac{E}{P_{\max}}} + R$. This

function considers the term of respiration (R) and assumes absence of photoinhibition.

Vertical and Marginal Growth Measurements

Vertical growth of the juvenile CCA samples was estimated at the end of the experiment by measuring the crust thickness using a scanning electron microscope (JSM-6510 series, JEOL). To obtain cross sections, recruits attached to the petri dish were carefully detached from the substrate and sliced into sections using a razor blade. The resulting fragments were mounted on a metal stub with adhesive. Samples were carbon coated using a sputter coater (JFC-1600 auto fine coater, JEOL) for SEM analyses. Images were taken with a magnification of $\times 3000$ at a high voltage of 5 kV with a spot size of 50 nm and a working distance (WD) of 12–16 mm. We measured a total of 15 points along the crust from different fragments. To estimate thickness of the crust, the distance from the surface of the crust to the bottom was measured using the SEM software (JEOL Scanning electron microscope software) (Figures 2B,D). Growth rates in units of μm crust increment day⁻¹ were obtained by normalizing crust thickness to the duration of

²<http://www.unisense.com>

the experiment, i.e., 10 days. Marginal growth was estimated by measuring the change in surface area. Final and initial photographs of individual recruits were taken and images were analyzed using Image J software (University of Wisconsin-Madison).

Statistical Analyses

Photosynthetic parameters were compared between treatments using one-way ANOVA. Pearson correlation was used to test the relationships between vertical growth and gross photosynthesis. Normal distribution of data and homogeneity of variances were tested using Kolmogorov-Smirnov and Cochran's test and data was arc-sin transformed when needed. All statistical analyses were performed using SPSS (version 25).

RESULTS

Photosynthesis and Respiration


The measured P-E curves showed a significant difference between recruits growing under elevated CO₂ and recruits growing under ambient CO₂ conditions (**Figures 3A,B**). Maximum gross photosynthesis was 50% higher under elevated CO₂ [mean $P_{\max} = 75.36 \text{ nmol O}_2 \text{ cm}^{-3} \text{ s}^{-1}$ ($\pm 9.7 \text{ SE}$)] as compared to ambient CO₂ conditions [mean $P_{\max} = 37.36 \text{ nmol O}_2 \text{ cm}^{-3} \text{ s}^{-1}$ ($\pm 7.8 \text{ SE}$)], ANOVA $p = 0.044$). Photosynthetic efficiency and saturation irradiance were slightly higher under high CO₂ when compared to ambient CO₂, but the difference was not statistically significant (ANOVA, α : $p = 0.22$, E_k : $p = 0.12$). At the maximum photon irradiance used in the experiment ($340 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$), gross photosynthesis was 39% higher under high CO₂ than under ambient CO₂ conditions but the difference was not significant (ANOVA $p = 0.081$, **Figure 4A**).

The responses of net photosynthetic rates to the CO₂ treatments followed a similar trend as the gross photosynthesis, with increased P_{\max} , and slightly higher α and E_k values under elevated CO₂ than under ambient CO₂ conditions. However, the difference across CO₂ treatments was less pronounced and not statistically significant (ANOVA, $p = 0.12$, $p = 0.78$, $p = 0.35$, for P_{\max} , α and E_k , respectively, **Figure 3B**).

Dark respiration was significantly enhanced by elevated CO₂, with recruits showing a 48.7% increase in respiration rates compared to recruits from ambient CO₂ (ANOVA, $p = 0.0001$, **Figure 4B**). In addition, surface dynamics of O₂ concentration during experimental dark-light shifts showed higher O₂ production and faster response of CCA recruits under elevated CO₂ compared to measurements under ambient CO₂ (**Figure 5**).

Growth Rates


Vertical and marginal growth rates were significantly affected by high CO₂ (ANOVA, $p = 0.01$ and 0.03 , respectively). *P. cf. onkodes* recruits grown under ambient CO₂ conditions showed thicker and larger crusts as compared to recruits exposed to high CO₂ conditions (**Figures 6A,B**). Crust vertical growth rates were reduced from $2.8 \pm 0.24 \mu\text{m d}^{-1}$ under ambient CO₂ to $2.05 \mu\text{m d}^{-1}$ ($\pm 0.04 \text{ SE}$) under high CO₂, i.e., a 28.7% change. Marginal



growth rates were also reduced by 19%. Structural damage of the cells under high CO₂ was observed on the crust surface, where epithallial/perithallial cells seemed to have collapsed (**Figure 2C**) as opposed to observations of the crust surface of germlings growing under ambient conditions where cells seem to preserve their structure (**Figure 2A**). Furthermore, cross sections of CCA recruits showed cells with irregular shapes and thinner cell walls in samples exposed to elevated CO₂ (**Figure 2D**) compared to those under ambient conditions (**Figure 2B**).

Growth vs. Photosynthesis


We found a negative correlation between gross photosynthesis and crust thickness of *P. cf. onkodes* recruits at an incident



irradiance of 340 μmol photons m⁻² s⁻¹ (Pearson correlation coefficient 0.89, *p* = 0.01, *n* = 5, **Figure 7**). The highest gross photosynthetic rate (55.16 nmol O₂ cm⁻³ s⁻¹) was found on thinnest recruits (1.93 μm d⁻¹) growing under elevated CO₂, whereas the lowest gross photosynthetic rate (3.29 nmol O₂ cm⁻³ s⁻¹) was measured on the thickest recruits (3.35 μm d⁻¹) from ambient CO₂ condition.

DISCUSSION


The impacts of ocean acidification (OA) on adult coralline algae have been relatively well documented (Hofmann and Bischof, 2014), and there is a growing body of evidence supporting the sensitivity of the early life history stages of coralline algae to OA. In particular, recent studies have shown reduced spore germination (Bradassi et al., 2013; Ordoñez et al., 2017) and attachment (Guenther et al., 2018) and declined germling abundance [e.g., % cover, Kuffner et al. (2008); Ordoñez et al. (2014); Fabricius et al. (2015); Ordoñez et al. (2017)] and



growth (Roleda et al., 2015; Ordoñez et al., 2017) under elevated CO₂ conditions. However, whether metabolic processes such as photosynthesis and respiration of early life stages of CCA also show a negative response to elevated CO₂ has not been well documented. Our study tests for the first time the effects of elevated CO₂ on the photosynthesis and respiration of recruits of a major reef building coralline algae. We demonstrate that increased CO₂ can potentially enhance gross photosynthesis and dark respiration, but can reduce the growth of *P. cf. onkodes* recruits. Our findings also reveal a possible decoupling between photosynthesis and growth rates in CCA recruits, suggesting that the carbon fixed during the photosynthetic process does not necessarily translate into enhanced algal growth. Understanding the relationships between physiological and vital processes under OA, in particular on vulnerable early life stages, is essential to improve our understanding of OA impacts on coral reef organisms and the important ecosystem services they support.


Effects of Elevated CO₂ on Photosynthesis

Porolithon cf. onkodes recruits growing under high CO₂ conditions showed significantly higher maximum gross photosynthetic rates at the thallus surface compared to those under ambient CO₂ conditions, suggesting that photosynthesis in our experimental CCA was carbon limited under ambient CO₂ conditions, in line with, e.g., measurements on epilithic algal layers on coral tiles (Larkum et al., 2003). The photosynthetic responses of coralline algae to elevated CO₂ (reduced pH) documented in the literature are highly variable and depend on the nature of the experiments, the techniques employed to quantify O₂ dynamics, and the species of algae considered in the experiments. For example, two studies using adult *Porolithon onkodes* from French Polynesia found no effect of elevated CO₂ on gross photosynthesis and respiration (Johnson et al., 2014;



Comeau et al., 2017), although in the Johnson et al. (2014) study, high CO₂ decreased net photosynthesis, similar to results from the Great Barrier Reef using the same CCA species (Anthony et al., 2008). Studies using different taxa from temperate regions found gross photosynthesis to remain constant under elevated CO₂ [e.g., articulated algae *Corallina officinalis* (Hofmann et al., 2012), *Corallina elongata* (Noisette et al., 2013), CCA *Lithophyllum incrustans*, *Lithothamnion corallioides* (Noisette et al., 2013), *Lithothamnion glaciale* (Kamenos et al., 2013)]. Enhanced productivity in response to elevated CO₂ has been observed for some adult tropical corallines (Semese et al., 2009; Yildiz et al., 2014). Using O₂ microsensors on adult CCA, Hofmann et al. (2016) found elevated photosynthetic rates under reduced pH, similar to the findings in our study.

Enhanced photosynthesis in response to elevated CO₂ concentrations may indicate that photosynthesis in early life stages of *P. onkodes* is limited by intracellular inorganic carbon availability. There are two main mechanisms by which algae acquire CO₂ for carbon fixation within the cell: (1) the active



uptake of HCO₃⁻ via a number of carbon concentrating mechanisms (CCMs), and (2) the diffusive uptake of CO₂ (Giordano et al., 2005; Raven et al., 2005). These mechanisms could be altered by elevated seawater CO₂ concentrations. Coralline algae have CCMs that facilitate active transport of HCO₃⁻ across cellular membranes leading to higher concentrations of CO₂ at the site of RuBisCO (Comeau et al., 2013; Hofmann and Heesch, 2018) and given the high availability of HCO₃⁻ in seawater, enhanced CO₂ may thus not significantly enhance DIC transport to the site of photosynthesis. However, since CCM activity is energetically costly, it is likely that elevated seawater CO₂ alleviates the use of CCMs, effectively downregulating CCM activity upon enhanced and energetically less costly diffusive CO₂ uptake under OA conditions. Flexibility in the use of carbon acquisition strategies under OA has been suggested for a number of articulated coralline algae including *C. officinalis* (Cornwall et al., 2012) and *Arthrocardia corymbosa* (Cornwall et al., 2013). Our study cannot discern the mechanistic basis for the increased rates of photosynthesis observed in our recruits under OA, but could suggest that photosynthesis in our early life stages of CCA is carbon limited. The majority of studies examining the effects of elevated CO₂ on CCA photosynthesis have used incubation chambers with larger O₂ sensors and have shown negative or no response of OA on photosynthetic activity (Anthony et al., 2008; Gao and Zheng, 2010; Hofmann et al., 2012; Cornwall et al., 2013; Martin et al., 2013; Noisette et al., 2013; Johnson et al., 2014; Comeau et al., 2017). However, when microsensors have been used to test CO₂ effects on coralline algae photosynthesis, increased O₂ production has been detected (e.g., Hofmann et al., 2016, but see Cornwall et al., 2013). Microsensors measure O₂ dynamics directly at the surface of the CCA, with minimum interference from epiphytic algae (e.g., diatoms, Roleda et al., 2015) or bacteria, or endolithic algae (e.g., Anthony et al., 2008), presumably providing a more local and direct measurement of the effects of CO₂ on CCA physiology.

Declined Growth and Relationship With Photosynthesis

Photosynthesis is the major source of carbon for CCA algal tissue construction, therefore, it is generally expected that increased photosynthetic rates and consequently the amount of fixed carbon would be positively correlated with algal growth. As detailed earlier, photosynthetic rates of the experimental recruits increased with increased CO₂ concentration, however, this increase in CO₂ did not translate directly in enhanced algal growth. On the contrary, both vertical and marginal growth rates of the algal crusts were depressed under high CO₂ (a 28 and 19% decrease under high compared to ambient CO₂, **Figures 6A,B**), and we additionally observed a significant inverse relationship between algal photosynthesis and growth (Pearson correlation $R^2 = 0.89$, $p = 0.01$, **Figure 7**). The decreased growth of CCA under elevated CO₂ shown in our study agrees with findings from several studies using both adult and early life stages of CCA (Kuffner et al., 2008; Bradassi et al., 2013; Ordoñez et al., 2017). The decoupling between increased photosynthetic rates and algal growth rates may be explained by two key processes. First, and perhaps the most important process, high CO₂ enhances skeletal dissolution of the coralline algal crusts. In fact, SEM images showed distorted crust tissues, collapsed epithelial cells (**Figure 2C**), and apparently a thinning of calcified cell walls (**Figure 2D**), similar to observations on *Lithothamnion glaciale* from the Mediterranean (Ragazzola et al., 2012) and *Hydrolithon samoense* from Japan exposed to high CO₂ (Kato et al., 2014). Several studies have attributed such reduction in growth of CCA recruits (e.g., Bradassi et al., 2013; Roleda et al., 2015) and adults (Diaz-Pulido et al., 2014) to increased skeletal dissolution due to undersaturation of seawater with respect to the dominant form of calcium carbonate in the CCA, i.e., high-magnesium-calcite. **Table 1** shows a high Mg-calcite saturation state <1 in the high CO₂ treatment, indicating undersaturation of seawater calcium carbonate and hence conditions for skeletal dissolution. Undersaturation may also affect deposition of new calcium carbonate in the algal skeleton. Secondly, since algal growth is a function of photosynthesis minus algal respiration and excretion of photosynthates, this suggests that carbon losses due to respiratory and excretion/secretion processes are higher than carbon gains from the photosynthetic process (Falkowski and Raven, 2013). In this regard, respiration rates were indeed higher under high CO₂, which may explain the observed reduced growth, albeit net photosynthesis was also slightly higher under elevated CO₂. It is also likely that a significant fraction of photosynthetically fixed carbon is excreted/secreted as dissolved organic carbon (DOC) to the external environment, leaving less carbon readily available for algal growth (e.g., Iñiguez et al., 2016, 2017) and potentially for calcification. This situation may be particularly important when nutrients limit algal growth, as shown in symbiotic algae (Yellowlees et al., 2008) and polar algae (Iñiguez et al., 2016). Recent work shows that high CO₂ concentrations stimulated the release of DOC in a range of tropical reef algae, particularly red algae (Diaz-Pulido and Barron, personal observation). The mechanisms explaining the decoupling of algal photosynthesis and growth under OA conditions require further investigations.

Although our study did not directly quantify the calcification rate of the early life stage of *P. cf. onkodes*, we expect that the observed reduction in both vertical and marginal crust growth is associated with a lowering in the amount of calcium carbonate deposited by the algal crusts. On the other hand, a direct positive relationship between photosynthesis and calcification has also been postulated (Borowitzka, 1981; Hofmann et al., 2016), although decoupling between photosynthesis and calcification on adult coralline algae has also been shown (Semesi et al., 2009). Our finding of increased photosynthetic rate under elevated CO₂ does not seem to be coupled with elevated rates of calcification, given the reduced growth observed when exposed to high CO₂, and the considerable alterations in crust thickness and ultrastructure. Any positive effect of elevated DIC (both CO₂ and HCO₃⁻) on CCA photosynthesis and potentially on calcification may thus not be sufficient to compensate for the reduction in high-Mg-calcite saturation state and resulting skeletal dissolution, as discussed by Comeau et al. (2013) and Bradassi et al. (2013). The very thin crusts of the early life stages of *P. cf. onkodes* could contribute to some extent to the high vulnerability of these recruits to OA, thus CCA recruits may represent one of the most sensitive life stages to OA. A further understanding of the metabolic processes underpinning decreased growth and calcification is fundamental for understanding the underlying causes for decreased abundance in key reef building coralline algae.

DATA AVAILABILITY STATEMENT

Data are available at the Australian Rivers Institute – Coast and Estuaries repository system at Griffith University.

AUTHOR CONTRIBUTIONS

AO and GD-P designed the experiments, collected and analyzed the growth data, and drafted the manuscript. AO performed the experiments. DW and NL collected and analyzed the photosynthesis data. All authors contributed with manuscript writing and gave final approval for publication.

FUNDING

This project was supported by the Australian Research Council (DP-120101778), the Great Barrier Reef Foundation, a Sapere-Aude Advanced grant from the Independent Research Fund Denmark (MK), and the Carlsberg Foundation (DW). Samples for this study were collected under the permit G12/34877.1 granted by Great Barrier Reef Marine Park Authority.

ACKNOWLEDGMENTS

We thank P. Gartrell, C. Barron, B. Lewis, L. Gomez, E. Kennedy, and HIRS staff who assisted with the experiments.

REFERENCES

- Adey, W. H. (1998). Coral reefs: algal structured and mediated ecosystems in shallow, turbulent, alkaline waters. *J. Phycol.* 34, 393–406. doi: 10.1046/j.1529-8817.1998.340393.x
- Anthony, K. R. N., Kline, D. I., Diaz-Pulido, G., Dove, S., and Hoegh-Guldberg, O. (2008). Ocean acidification causes bleaching and productivity loss in coral reef builders. *Proc. Natl. Acad. Sci. U.S.A.* 105, 17442–17446. doi: 10.1073/pnas.0804478105
- Borowitzka, M. A. (1981). Photosynthesis and calcification in the articulated coralline red algae *Amphiroa anceps* and *A. foliacea*. *Mar. Biol.* 62, 17–23. doi: 10.1007/BF00396947
- Bradassi, F., Cumani, F., Bressan, G., and Dupont, S. (2013). Early reproductive stages in the crustose coralline alga *Phymatolithon lenormandii* are strongly affected by mild ocean acidification. *Mar. Biol.* 160, 2261–2269. doi: 10.1007/s00227-013-2260-2
- Brodersen, K. E., Lichtenberg, M., Ralph, P. J., Kühl, M., and Wangpraseurt, D. (2014). Radiative energy budget reveals high photosynthetic efficiency in symbiont-bearing corals. *J. Royal Soc. Interface* 11:20130997. doi: 10.1098/rsif.2013.0997
- Comeau, S., Carpenter, R. C., and Edmunds, P. J. (2013). Coral reef calcifiers buffer their response to ocean acidification using both bicarbonate and carbonate. *Proc. R. Soc. B* 280:1753. doi: 10.1098/rspb.2012.2374
- Comeau, S., Carpenter, R. C., and Edmunds, P. J. (2017). Effects of pCO₂ on photosynthesis and respiration of tropical scleractinian corals and calcified algae. *ICES J. Mar. Sci.* 74, 1092–1102. doi: 10.1093/icesjms/fsv267
- Cornwall, C. E., Hepburn, C. D., McGraw, C. M., Currie, K. I., Pilditch, C. A., Hunter, K. A., et al. (2013). Diurnal fluctuations in seawater pH influence the response of a calcifying macroalga to ocean acidification. *Proc. R. Soc. B* 280:1772. doi: 10.1098/rspb.2013.2201
- Cornwall, C. E., Hepburn, C. D., Pritchard, D., Currie, K. I., McGraw, C. M., Hunter, K. A., et al. (2012). Carbon-use strategies in macroalgae: differential responses to lowered pH and implications for ocean acidification. *J. Phycol.* 48, 137–144. doi: 10.1111/j.1529-8817.2011.01085.x
- Cornwall, C. E., Revill, A. T., and Hurd, C. L. (2015). High prevalence of diffusive uptake of CO₂ by macroalgae in a temperate subtidal ecosystem. *Photosynth. Res.* 124, 181–190. doi: 10.1007/s11120-015-0114-0
- Dean, A. J., Steneck, R. S., Tager, D., and Pandolfi, J. M. (2015). Distribution, abundance and diversity of crustose coralline algae on the Great Barrier Reef. *Coral Reefs* 34, 581–594. doi: 10.1007/s00338-015-1263-5
- Diaz-Pulido, G., Anthony, K. R. N., Kline, D. I., Dove, S., and Hoegh-Guldberg, O. (2012). Interactions between ocean acidification and warming on the mortality and dissolution of coralline algae. *J. Phycol.* 48, 32–39. doi: 10.1111/j.1529-8817.2011.01084.x
- Diaz-Pulido, G., Cornwall, C., Gartrell, P., Hurd, C., and Tran, D. V. (2016). Strategies of dissolved inorganic carbon use in macroalgae across a gradient of terrestrial influence: implications for the Great Barrier Reef in the context of ocean acidification. *Coral Reefs* 35, 1327–1341. doi: 10.1007/s00338-016-1481-5
- Diaz-Pulido, G., Gouezo, M., Tilbrook, B., Dove, S., and Anthony, K. R. N. (2011). High CO₂ enhances the competitive strength of seaweeds over corals. *Ecol. Lett.* 14, 156–162. doi: 10.1111/j.1461-0248.2010.01565.x
- Diaz-Pulido, G., Nash, M. C., Anthony, K. R. N., Bender, D., Opdyke, B. N., Reyes-Nivia, C., et al. (2014). Greenhouse conditions induce mineralogical changes and dolomite accumulation in coralline algae on tropical reefs. *Nat. Commun.* 5:3310. doi: 10.1038/ncomms4310
- Fabricius, K. E., Kluibenschedl, A., Harrington, L., Noonan, S., and De'ath, G. (2015). In situ changes of tropical crustose coralline algae along carbon dioxide gradients. *Sci. Rep.* 5:9537. doi: 10.1038/srep09537
- Falkowski, P. G., and Raven, J. A. (2013). *Aquatic Photosynthesis*. Princeton, NJ: Princeton University Press. doi: 10.1515/9781400849727
- Feely, R. A., Sabine, C. L., Lee, K., Berelson, W., Kleypas, J., Fabry, V. J., et al. (2004). Impact of anthropogenic CO₂ on the CaCO₃ system in the oceans. *Science* 305, 362–366. doi: 10.1126/science.1097329
- Gabrielson, P. W., Hughey, J. R., and Diaz-Pulido, G. (2018). Genomics reveals abundant speciation in the coral reef building alga *Porolithon onkodes* (Corallinales, Rhodophyta). *J. Phycol.* 54, 429–434. doi: 10.1111/jpy.12761
- Gao, K., Aruga, Y., Asada, K., and Kiyohara, M. (1993). Influence of enhanced CO₂ on growth and photosynthesis of the red algae *Gracilaria* sp. and *G. chilensis*. *J. Appl. Phycol.* 5, 563–571. doi: 10.1007/BF02184635
- Gao, K., and Zheng, Y. (2010). Combined effects of ocean acidification and solar UV radiation on photosynthesis, growth, pigmentation and calcification of the coralline alga *Corallina sessilis* (Rhodophyta). *Glob. Change Biol.* 16, 2388–2398. doi: 10.1111/j.1365-2486.2009.02113.x
- Giordano, M., Beardall, J., and Raven, J. A. (2005). CO₂ concentrating mechanisms in algae: mechanisms, environmental modulation, and evolution. *Annu. Rev. Plant Biol.* 56, 99–131. doi: 10.1146/annurev.arplant.56.032604.144052
- Guenther, R., Miklasz, K., Carrington, E., and Martone, P. T. (2018). Macroalgal spore dysfunction: ocean acidification delays and weakens adhesion. *J. Phycol.* 54, 153–158. doi: 10.1111/jpy.12614
- Harrington, L., Fabricius, K., De'Ath, G., and Negri, A. (2004). Recognition and selection of settlement substrata determine post-settlement survival in corals. *Ecology* 85, 3428–3437. doi: 10.1890/04-0298
- Heyward, A. J., and Negri, A. P. (1999). Natural inducers for coral larval metamorphosis. *Coral Reefs* 18, 273–279. doi: 10.1007/s00338005193
- Hofmann, L. C., and Bischof, K. (2014). Ocean acidification effects on calcifying macroalgae. *Aquat. Biol.* 22, 261–279. doi: 10.3354/ab00581
- Hofmann, L. C., and Heesch, S. (2018). Latitudinal trends in stable isotope signatures and carbon concentrating mechanisms of northeast Atlantic *rhodoliths*. *Biogeosciences* 15, 6139–6149. doi: 10.5194/bg-2017-399
- Hofmann, L. C., Koch, M., and de Beer, D. (2016). Biotic control of surface pH and evidence of light-induced H⁺ pumping and Ca²⁺-H⁺ exchange in a tropical crustose coralline alga. *PLoS One* 11:e0159057. doi: 10.1371/journal.pone.0159057
- Hofmann, L. C., Yildiz, G., Hanelt, D., and Bischof, K. (2012). Physiological responses of the calcifying rhodophyte, *Corallina officinalis* (L.), to future CO₂ levels. *Mar. Biol.* 159, 783–792. doi: 10.1007/s00227-011-1854-9
- Ichiki, S., Mizuta, H., and Yamamoto, H. (2000). Effects of irradiance, water temperature and nutrients on the growth of sporelings of the crustose coralline alga *Lithophyllum yessoense* Foslie (Corallinales, Rhodophyceae). *Phycol. Res.* 48, 115–120. doi: 10.1046/j.1440-1835.2000.00193.x
- Iñiguez, C., Carmona, R., Lorenzo, M. R., Niell, F. X., Wiencke, C., and Gordillo, F. J. L. (2016). Increased CO₂ modifies the carbon balance and the photosynthetic yield of two common Arctic brown seaweeds: *Desmarestia aculeata* and *Alaria esculenta*. *Polar Biol.* 39, 1979–1991. doi: 10.1007/s00300-015-1724-x
- Iñiguez, C., Heinrich, S., Harms, L., and Gordillo, F. J. L. (2017). Increased temperature and CO₂ alleviate photoinhibition in *Desmarestia anceps*: from transcriptomics to carbon utilization. *J. Exp. Bot.* 68, 3971–3984. doi: 10.1093/jxb/erx164
- IPCC (2014). “Climate change 2013: the physical science basis,” in *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung et al. (Cambridge: Cambridge University Press).
- Johnson, M. D., and Carpenter, R. C. (2012). Ocean acidification and warming decrease calcification in the crustose coralline alga *Hydrolithon onkodes* and increase susceptibility to grazing. *J. Exp. Mar. Biol. Ecol.* 43, 94–101. doi: 10.1016/j.jembe.2012.08.005
- Johnson, M. D., Moriarty, V. W., and Carpenter, R. C. (2014). Acclimatization of the crustose coralline alga *Porolithon onkodes* to variable pCO₂. *PLoS One* 9:e87678. doi: 10.1371/journal.pone.0087678
- Jones, W. E., and Moorjani, A. S. (1973). *Attachment and Early Development of the Tetraspores of Some Coralline Red Algae*. Cochín: MBI Special Publication dedicated to Dr. N K Panikkar 1, 293–304.
- Kamenos, N. A., Burdett, H. L., Aloisio, E., Findlay, H. S., Martin, S., Longbone, C., et al. (2013). Coralline algal structure is more sensitive to rate, rather than the magnitude, of ocean acidification. *Glob. Change Biol.* 19, 3621–3628. doi: 10.1111/gcb.12351
- Kato, A., Hikami, M., Kumagai, N. H., Suzuki, A., Nojiri, Y., and Sakai, K. (2014). Negative effects of ocean acidification on two crustose coralline species using genetically homogeneous samples. *Mar. Environ. Res.* 94, 1–6. doi: 10.1016/j.marenvres.2013.10.010

- Koch, M., Bowes, G., Cliff, R., and Xing-Hai, Z. (2013). Climate change and ocean acidification effects on seagrasses and marine macroalgae. *Glob. Change Biol.* 19, 103–132. doi: 10.1111/j.1365-2486.2012.02791.x
- Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., et al. (2013). Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Glob. Change Biol.* 19, 1884–1896. doi: 10.1111/gcb.12179
- Kroeker, K. J., Kordas, R. L., Crim, R. N., and Singh, G. G. (2010). Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. *Ecol. Lett.* 13, 1419–1434. doi: 10.1111/j.1461-0248.2010.01518.x
- Kuffner, I. B., Andersson, A. J., Jokiel, P. L., Rodgers, K. U. S., and Mackenzie, F. T. (2008). Decreased abundance of crustose coralline algae due to ocean acidification. *Nat. Geosci.* 1, 114–117. doi: 10.1038/ngeo100
- Larkum, A. W. D., Koch, E.-M. W., and Kühl, M. (2003). Diffusive boundary layers and photosynthesis of the epilithic algal community of coral reefs. *Mar. Biol.* 142, 1073–1082. doi: 10.1007/s00227-003-1022-y
- Lewis, B., Kennedy, E. V., and Diaz-Pulido, G. (2017). Seasonal growth and calcification of a reef-building crustose coralline alga on the Great Barrier Reef. *Mar. Ecol. Prog. Ser.* 568, 73–86. doi: 10.3354/meps12074
- Littler, M. M. (1972). The crustose corallinaceae. *Oceanogr. Mar. Biol. Annu. Rev.* 10, 103–120.
- Littler, M. M., and Doty, M. S. (1975). Ecological components structuring the seaward edges of tropical pacific reefs: the distribution, communities and productivity of *Porolithon*. *J. Ecol.* 63, 117–129. doi: 10.2307/2258846
- Martin, S., Coahu, S., Vignot, C., Zimmerman, G., and Gattuso, J. P. (2013). One-year experiment on the physiological response of the Mediterranean crustose coralline alga, *Lithophyllum cabiochae*, to elevated pCO₂ and temperature. *Ecol. Evol.* 3, 676–693. doi: 10.1002/ece3.475
- Martin, S., and Gattuso, J. P. (2009). Response of Mediterranean coralline algae to ocean acidification and elevated temperature. *Glob. Change Biol.* 15, 2089–2100. doi: 10.1111/j.1365-2486.2009.01874.x
- Morse, J. W., Arvidson, R. S., and Lüttge, A. (2007). Calcium carbonate formation and dissolution. *Chem. Rev.* 107, 342–381. doi: 10.1021/cr050358j
- Noiset, F., Egilsdottir, H., Davoult, D., and Martin, S. (2013). Physiological responses of three temperate coralline algae from contrasting habitats to near-future ocean acidification. *J. Exp. Mar. Biol. Ecol.* 448, 179–187. doi: 10.1016/j.jembe.2013.07.006
- Ordoñez, A., Doropoulos, C., and Diaz-Pulido, G. (2014). Effects of ocean acidification on population dynamics and community structure of crustose coralline algae. *Biol. Bull.* 226, 255–268. doi: 10.1086/BBLv226n3p255
- Ordoñez, A., Kennedy, E. V., and Diaz-Pulido, G. (2017). Reduced spore germination explains sensitivity of reef-building algae to climate change stressors. *PLoS One* 12:e0189122. doi: 10.1371/journal.pone.0189122
- Pierrot, D., Lewis, E., and Wallace, D. W. R. (2006). *MS Excel Program Developed for CO₂ System Calculations*. ORNL/CDIAC-105a. Oak Ridge, TN: Carbon Dioxide Information Analysis Center. doi: 10.3334/CDIAC/otg.CO2SYS_XLS_CDIA105a
- Ragazzola, F., Foster, L. C., Form, A., Anderson, P. S. L., Hansteen, T. H., and Fietzke, J. (2012). Ocean acidification weakens the structural integrity of coralline algae. *Glob. Change Biol.* 18, 2804–2812. doi: 10.1111/j.1365-2486.2012.02756.x
- Raven, J. A., Ball, L. A., Beardall, J., Giordano, M., and Maberly, S. C. (2005). Algae lacking carbon-concentrating mechanisms. *Can. J. Bot.* 83, 879–890. doi: 10.1139/b05-074
- Revsbech, N. P. (1989). An oxygen microsensor with a guard cathode. *Limnol. Oceanogr.* 34, 474–478. doi: 10.4319/lo.1989.34.2.0474
- Revsbech, N. P., and Jørgensen, B. B. (1983). Photosynthesis of benthic microflora measured with high spatial resolution by the oxygen microprofile method: capabilities and limitations of the method. *Limnol. Oceanogr.* 28, 749–756. doi: 10.4319/lo.1983.28.4.0749
- Ringeltaube, P., and Harvey, A. (2005). Non-geniculate coralline algae (Corallinales, Rhodophyta) on heron reef, Great Barrier Reef (Australia). *Bot. Mar.* 43, 409–498.
- Roberts, R. D., Kühl, M., Glud, R. N., and Rysgaard, S. (2002). Primary production of crustose coralline red algae in a high arctic fjord. *J. Phycol.* 38, 273–283. doi: 10.1046/j.1529-8817.2002.01104.x
- Roleda, M. Y., Cornwall, C. E., Feng, Y., McGraw, C. M., Smith, A. M., and Hurd, C. L. (2015). Effect of ocean acidification and pH fluctuations on the growth and development of coralline algal recruits, and an associated benthic algal assemblage. *PLoS One* 10:e0140394. doi: 10.1371/journal.pone.0140394
- Roleda, M. Y., Morris, J. N., McGraw, C. M., and Hurd, C. L. (2012). Ocean acidification and seaweed reproduction: increased CO₂ ameliorates the negative effect of lowered pH on meiospore germination in the giant kelp *Macrocystis pyrifera* (Laminariales, Phaeophyceae). *Glob. Change Biol.* 18, 854–864. doi: 10.1111/j.1365-2486.2011.02594.x
- Santelices, B. (1990). Patterns of reproduction dispersal and recruitment in seaweeds. *Oceanogr. Mar. Biol. Annu. Rev.* 28, 177–276.
- Semesi, I. S., Kangwe, J., and Björk, M. (2009). Alterations in seawater pH and CO₂ affect calcification and photosynthesis in the tropical coralline alga, *Hydrolithon* sp. (Rhodophyta). *Estuar. Coast. Shelf Sci.* 84, 337–341. doi: 10.1016/j.ecss.2009.03.038
- Steneck, R. S. (1983). Escalating herbivory and resulting adaptive trends in calcareous algal crusts. *Paleobiology* 9, 44–61. doi: 10.2307/2400629
- Webb, W. L., Newton, M., and Starr, D. (1974). Carbon dioxide exchange of *Alnus rubra*. *Oecologia* 17, 281–291. doi: 10.1007/bf00345747
- Yellowlees, D., Rees, T. A. V., and Leggat, W. (2008). Metabolic interactions between algal symbionts and invertebrate hosts. *Plant Cell Environ.* 31, 679–694. doi: 10.1111/j.1365-3040.2008.01802.x
- Yildiz, G., Hofmann, L. C., Bischof, K., and Dere, S. (2014). *Ultraviolet Radiation Modulates the Physiological Responses of the Calcified Rhodophyte Corallina officinalis to Elevated CO₂*. Washington, DC: USDA. doi: 10.1594/PANGAEA.831729

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Ordoñez, Wangpraseurt, Lyndby, Kühl and Diaz-Pulido. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.