



## Hallmarks of human small antral follicle development

### Implications for regulation of ovarian steroidogenesis and selection of the dominant follicle

Kristensen, Stine G.; Mamsen, Linn S.; Jeppesen, Janni V.; Bøtkjær, Jane Alrø; Pors, Susanne E.; Borgbo, Tanni; Ernst, Erik; Macklon, Kirsten T.; Andersen, Claus Yding

*Published in:*  
Frontiers in Endocrinology

*DOI:*  
[10.3389/fendo.2017.00376](https://doi.org/10.3389/fendo.2017.00376)

*Publication date:*  
2018

*Document version*  
Publisher's PDF, also known as Version of record

*Document license:*  
[CC BY](https://creativecommons.org/licenses/by/4.0/)

*Citation for published version (APA):*  
Kristensen, S. G., Mamsen, L. S., Jeppesen, J. V., Bøtkjær, J. A., Pors, S. E., Borgbo, T., ... Andersen, C. Y. (2018). Hallmarks of human small antral follicle development: Implications for regulation of ovarian steroidogenesis and selection of the dominant follicle. *Frontiers in Endocrinology*, 8(JAN), [376]. <https://doi.org/10.3389/fendo.2017.00376>



# Hallmarks of Human Small Antral Follicle Development: Implications for Regulation of Ovarian Steroidogenesis and Selection of the Dominant Follicle

Stine G. Kristensen<sup>1</sup>, Linn S. Mamsen<sup>1</sup>, Janni V. Jeppesen<sup>1</sup>, Jane Alrø Bøtkjær<sup>1</sup>, Susanne E. Pors<sup>1</sup>, Tanni Borgbo<sup>1</sup>, Erik Ernst<sup>2</sup>, Kirsten T. Macklon<sup>3</sup> and Claus Yding Andersen<sup>1\*</sup>

<sup>1</sup> Faculty of Health and Medical Sciences, Laboratory of Reproductive Biology, The Juliane Marie Centre for Women, Children and Reproduction, Copenhagen University Hospital, University of Copenhagen, Copenhagen, Denmark, <sup>2</sup> The Fertility Clinic, Aarhus University Hospital, Aarhus, Denmark, <sup>3</sup> The Fertility Clinic, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark

## OPEN ACCESS

### Edited by:

Katja Teerds,  
Wageningen University, Netherlands

### Reviewed by:

Livio Casarini,  
University of Modena and Reggio  
Emilia, Italy  
Johan Smitz,  
University Hospital Brussels, Belgium

### \*Correspondence:

Claus Yding Andersen  
yding@rh.dk

### Specialty section:

This article was submitted  
to Reproduction,  
a section of the journal  
Frontiers in Endocrinology

**Received:** 08 June 2017

**Accepted:** 20 December 2017

**Published:** 12 January 2018

### Citation:

Kristensen SG, Mamsen LS, Jeppesen JV, Bøtkjær JA, Pors SE, Borgbo T, Ernst E, Macklon KT and Andersen CY (2018) Hallmarks of Human Small Antral Follicle Development: Implications for Regulation of Ovarian Steroidogenesis and Selection of the Dominant Follicle. *Front. Endocrinol.* 8:376. doi: 10.3389/fendo.2017.00376

Regulation of human ovarian steroidogenesis differs from other species and precise knowledge on how human small antral follicles (hSAF) develop and acquire competence for continued growth and steroid output is still incomplete. The present study has characterized almost 1,000 normal hSAF collected in connection with cryopreservation of ovarian tissue for fertility preservation. The antral follicles (ranging from 3 to 13 mm) were generally aspirated from one ovary surgically removed during the natural cycle, and the follicular fluid (FF) and the granulosa cells (GC) were isolated and snap-frozen. In FF, the following hormones were measured: inhibin-B, inhibin-A, AMH, follistatin, PAPP-A, estradiol, progesterone, testosterone, and androstenedione. In GC, mRNA gene expressions using q-PCR were measured for the following genes: *FSHR*, *AMH*, *CYP19*, and *AR*. All samples in which one of the abovementioned parameters was measured were included, but typically multiple parameters were measured. Highly significant differences in concentration and follicular content in relation to follicular diameter were found for all measured hormones despite massive variability in-between follicles for any given diameter. The results demonstrate that profound changes take place in the hormonal microenvironment around follicular diameters of 8–11 mm corresponding to when follicular selection occurs. At this point, inhibin-B and inhibin-A showed distinct peaks concomitant with a significant reduction in both AMH protein and mRNA expression. Concentrations of inhibins, androgens, *FSHR*, and *AR* were intimately associated, and it is suggested that inhibin-B in combination with PAPP-A and thereby IGF2 activity exerts important paracrine signaling at follicular selection. At the same time upregulation of estradiol synthesis and *CYP19* mRNA expression increased steroid output profoundly. Furthermore, the highly significant association between *FSHR* and *AR* mRNA gene expression enforces important functions of androgens in follicular development. Collectively, these data reintroduce the understanding of the follicular phase as two parted in which regulation of steroidogenesis differs. The profound changes taking place around follicular selection highlight important paracrine actions of TGF- $\beta$  family members and IGFs for securing dominance of the selected follicle.

**Keywords:** human ovarian steroidogenesis, human small antral follicles, follicle-stimulating hormone, luteinizing hormone, granulosa cells, follicular fluid, inhibin-B, paracrine signaling

## INTRODUCTION

Follicles are the functional units of the ovaries. Each follicle consists of a single oocyte surrounded by somatic granulosa cells (GC) covering the internal surface of the basal membrane that encompasses follicles and prevents vessels from entering. Early in development, follicles recruit theca cells immediately adjacent to the basal membrane encircling the follicle. The theca cells become highly vascularized and secure transport of hormones including gonadotrophins to and from both the follicle and the theca cells. When human follicles reach a diameter of around 200–300  $\mu\text{m}$  a fluid-filled antral cavity starts to form, which contains the follicular fluid (FF). The FF consists of filtrations or transudates from circulation that penetrate the basal membrane and of GC secretions including a number of steroid and peptide hormones (1). The composition of hormones in FF is very dynamic in relation to developmental stage and very different from conditions found in circulation (2–4). For instance, steroidogenesis is very prominent throughout most of follicular development with synthesis of androgens in the theca cells by stimulation with luteinizing hormone (LH), which to a high degree is converted to estrogens in GC *via* stimulation with follicle-stimulating hormone (FSH), illustrating the two-cell, two-gonadotropin concept. As the follicle matures, the concentration of steroids such as estradiol and progesterone increases and reaches levels 1–10,000-fold higher than in circulation. In addition, several peptide hormones are specifically synthesized by the GC including AMH, inhibin-A, and inhibin-B. They also accumulate in FF often with concentrations more than 1,000 times higher than in circulation (4, 5). Growth and development of human small antral follicles (hSAF) are affected by these and other hormones, and a number of both autocrine and paracrine effects are dependent on the very different and local concentrations present in FF and in theca cells (6–8). It has been suggested that inhibin-B in addition to attenuate pituitary FSH secretion also plays an important paracrine effect in augmenting theca cell androgen production in synergy with LH (9–12). These fine-tuned local mechanisms are likely to exert an important function in the process of selecting the dominant follicle that takes place in each individual menstrual cycle (5). Almost four decades ago, it was realized that each individual follicle possesses a unique hormonal fingerprint and that the microenvironment in the follicle to a large extent determines its further developmental capacity (13). Furthermore, the dynamic changes in FF in relation to the developmental stage of the human follicles are massive, probably reflecting the variable hormonal and nutritional requirement of the developing follicle and the enclosed oocyte (14). Although FF from large preovulatory follicles, especially as collected in connection with IVF treatment, has received a lot of attention scientifically, normal hSAF have only to a limited degree been studied mainly because of unavailability. However, this has changed in recent years because women in connection with fertility preservation often have one entire ovary excised for cryopreservation (15). The hSAF do not sustain freezing and are normally discarded. During the last 10 years, we have accumulated a large number of hSAF including FF and corresponding GC.

The aim of the present study was to characterize the hallmarks of hormonal and genetic markers of normal hSAF, spanning in follicular diameter from 3 to 13 mm. This cohort of follicles thereby constitutes the follicular starting point for ovarian stimulation with exogenous hormones as used in connection with infertility treatment.

## MATERIALS AND METHODS

All follicles were collected from surplus ovarian tissue from women undergoing fertility preservation, having ovarian tissue cryopreserved at the Laboratory of Reproductive Biology, Rigshospitalet, Denmark. The fertility preservation procedure normally involved excision of one entire ovary from which individual visible antral follicles were aspirated with a 23G needle attached to a syringe. The fluid was centrifuged, the FF aspirated and immediately stored at  $-80^{\circ}\text{C}$  or snap frozen in liquid nitrogen. The diameter of the follicles was calculated based on the aspirated volume. In some cases, the GC was also collected and snap frozen in liquid nitrogen.


Collection of follicles has taken place during the last 15 years, and parts of the present material of both FF and GC have been used in several individual studies (16–26). Included in the present study is any follicle on which at least one of the reported parameters has been measured. Excluded from the analysis are girls below the age of 16 years and those who had a diagnosis for fertility preservation that included a disease of the ovary (e.g., ovarian cancer). The diagnoses of the women included were as follows: cancer mammae: 133; Md. Hodgkin: 41; non-Hodgkin and other lymphoma: 20; Ewing sarcoma and other sarcoma: 14; cervix: 13; leukemia: 13; cancer cerebrum: 12; colon cancer: 7; autoimmune diseases: 5; others: 28. The collection of FF had no effect on the fertility preservation procedure. In all cases, the ovary had overall gross normal appearance.

In total, 958 follicles were collected from 286 women, with a mean age of 28 years (ranging from 16 to 43 years). The number of follicles collected from each woman ranged from 1 to 13 (mean 3.3 follicles/patient; median 3.0). The number of women in relation to the number of follicles aspirated is given in **Figure 1A**. Follicle diameters ranged from 3 to 13 mm, as calculated from the aspirated volume, assuming a spherical structure of the follicle. The frequency of follicles aspirated with a specific diameter (e.g., 5 mm follicles comprise follicles with 4.5–5.5 mm in diameter) from the cohort of follicles is depicted in **Figure 1B**.

The hormone milieu in hSAF has been shown to be independent of circulating levels of gonadotropins and has no major impact on the serum hormone levels, thus serum hormone levels were not included in this study (22, 26, 27). The follicles were collected at various times during the menstrual cycle, as the hormone levels of hSAF remain constant throughout the menstrual cycle (23, 27, 28).

## Hormone Measurements

The concentrations of AMH, inhibin-A, and inhibin-B were measured using commercially available ELISA assays. AMH was measured using the Ultra-Sensitive AMH/MIS ELISA kit (AI-105-i, Ansh Labs, Webster, TX, USA), Inhibin-A ELISA kit



(AL-123-i, Ansh Labs, Webster, TX, USA), and Inhibin-B ELISA kit (AL-107-i Ansh Labs, Webster, TX, USA) according to the manufacturer's protocol, with an appropriate dilution of the FF samples using the supplied assay buffer. Estradiol, progesterone, androstenedione, and testosterone were initially measured using commercially available RIA kits (DSL-43100, DSL-3400, DSL-3800, and DSL-4000; DSL, Webster, TX, USA). However, during the course of sample collection, the RIA assays became unavailable, and it was necessary to switch to ELISA assays (NovaTec Immundiagnostica, Dietzenbach, Germany, DNOV 002, 003, 006, and 008, respectively). Due to limited sample material, it was not possible to reanalyze the FF with ELISA kits. Based on highly significant linear correlations between the results from the two types of assays (correlation coefficients: estradiol  $r = 0.99$ , progesterone  $r = 0.93$ , testosterone  $r = 0.91$ , and androstenedione  $r = 0.91$ ), mathematical conversion of values to the ELISA standard was performed as previously published (21). The remaining follicles were analyzed using NovaTec ELISA assays, according to the manufacturer's protocol, using in-house prepared steroid-free serum for FF dilution.

## RNA Purification

From individual GC samples, total RNA was purified using a combination of Tri Reagent (Sigma-Aldrich, St. Louis, MO, USA) and the RNeasy Mini Kit (Qiagen, Hilden, Germany). For each sample, the quality of the purified RNA was analyzed using an Agilent 2100 Bioanalyzer and an RNA 6000 Pico LabChip (RNA 6000 Pico assay kit, Agilent Technologies, Waldbronn, Germany). The RNA integrity number should be higher than the internal threshold value in order to qualify samples for the study. Total RNA quantity in each sample was measured using a Beckman Coulter Du730 life science UV/vis spectrophotometer.

## cDNA Synthesis and qPCR Analysis

First-strand cDNA was synthesized using the High Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Carlsbad, CA, USA). Briefly, a master mix containing 2.0  $\mu$ l 10 $\times$  RT Buffer, 0.8  $\mu$ l 25 $\times$  dNTP Mix (100 mM), 2.0  $\mu$ l 10 $\times$  RT Random Primers, 1.0  $\mu$ l MultiScribe<sup>TM</sup> Reverse Transcriptase (50 U/ $\mu$ l), 0.5  $\mu$ l RNase inhibitor, and 4.7  $\mu$ l nuclease-free (DEPC) water was prepared for each 20  $\mu$ l reaction. 11  $\mu$ l master mix was added to 9  $\mu$ l of total

RNA. Samples were centrifuged briefly at 12,000  $\times g$  and then incubated at room temperature for 10 min, followed by 37°C for 2 h and finally, 85°C for 5 s.

Gene expression levels were evaluated by relative quantification-PCR analysis using TaqMan<sup>®</sup> technology (Applied Biosystems) and the LightCycler 480 qPCR instrument (Roche, Copenhagen, Denmark). Predesigned AMH, FSHR, AR, and CYP19a1 TaqMan<sup>®</sup> Gene Expression Assays, as well as the Endogenous Control Assays for human  $\beta$ -actin and glyceraldehyde 3-phosphatdehydrogenase (GAPDH) were purchased from Applied Biosystems [Assay id no.:  $\beta$ -actin: #4326315E, GAPDH: #4333764E, AMH: #Hs00174915\_m1, FSH-R: #Hs00174865\_m1, AR: #Hs00171172\_m1, and Cyp19a1 (aromatase): #Hs00903413]. Data were quantified according to the comparative CT method (LightCycler480 Software, Roche); expression levels are presented as  $2^{-\Delta\Delta CT}$  normalized to GAPDH (20).

## Statistical Analysis

Statistical analysis was performed using GraphPad Prism 6.07 program (GraphPad Software, Inc., CA, USA). One-way analysis of variance (ANOVA) was used to determine the significant differences in hormone levels and follicular content in relation to the follicular diameter of hSAFs, and mRNA gene expression of GC-specific substances in hSAF in relation to follicular diameter. Spearman's rank correlation was used to evaluate correlations between various hormones in FF and between genes expressed in GC. A level of  $P < 0.05$  was considered statistically significant.

## RESULTS

The number of follicles aspirated per woman varied from 1 to 13, with the vast majority having one or two follicles aspirated (Figure 1A). The diameter of follicles aspirated from women above the age of 15 years is depicted in Figure 1B. Follicles with diameters from 4 to 7 mm account for more than 80% of the all analyzed follicles.

## Follicular Content and Concentrations of Peptide and Steroid Hormones

The intrafollicular concentration of inhibin-B showed a highly significant variation with follicular diameter showing peak

concentrations around diameters of 9–11 mm. The follicular content of inhibin-B also showed a highly significant variation with follicular diameter with a peak at 11 mm and high flanking values in 10 and 12 mm follicles (**Figure 2**; Tables S1 and S2 in Supplementary Material).

Inhibin-A exhibited a peak at around 9–10 mm similar to inhibin-B, but inhibin-A was present in concentrations comprising around 8–20% of that of inhibin-B with a relative concentration of inhibin-A increasing as the follicular diameter increased (3–11 mm). When the diameter exceeded 13 mm, the picture reversed and the concentration and content of inhibin-A exceeded that of inhibin-B (**Figure 2**; Tables S1 and S2 in Supplementary Material).

Both the concentration and the content of AMH showed a highly significant association with the follicular diameter with peak values at around a diameter of 8 mm (**Figure 2**; Tables S1 and S2 in Supplementary Material). Noteworthy is the follicular content of inhibin-B, inhibin-A, and AMH (calculated in nanograms), which was in a similar order of magnitude comprising around 100–200 ng at peak levels with a little less inhibin-A. Despite the relative smooth curves based on the mean levels and mean content of the follicles, the variability in follicles with a given diameter was very high (Tables S1 and S2 in Supplementary Material). There was a highly significant negative association between the FF concentrations of AMH and inhibin-B irrespective of follicular diameter (Spearman rank sum test:  $N = 538$ ;  $R = 0.46$ ;  $P < 0.00001$ ).

The intrafollicular concentrations of the four evaluated steroids showed different associations with the follicular diameter. Androstenedione and testosterone were both relatively constant, but testosterone showed a peak at a follicular diameter of 10 mm (**Figure 2**). In contrast, estradiol starts to increase at around 8–10 mm (**Figure 2**), coinciding with a reduction in intrafollicular AMH concentrations and GC expression (**Figures 2 and 3**). Progesterone remained relatively constant but ended with a steep increase in concentration at a follicular diameter of 11–12 mm (**Figure 2**). The association between the follicular diameter and the intrafollicular concentrations of follistatin and PAPP-A is also shown in **Figure 2**. PAPP-A showed a steep increase in the concentration starting with a follicular diameter of 8 mm, whereas follistatin was relatively constant with a drop in concentration at around 8–10 mm.

## Gene Expression Profiles of Selected Genes

The mRNA expression profiles of GC showed that *FSHR* expression is highly dependent on the follicular diameter and decreases with increasing diameter (**Figure 3**). This contrasts *CYP19* mRNA expression that in parallel to the concentration of estradiol starts to increase at diameters of 8–10 mm (**Figure 3**). The *AR* mRNA expression in relation to follicular diameter showed a decreasing expression as the follicular diameter increase in close association with the *FSHR* expression (**Figure 3**). There was a highly significant association between mRNA expression of *AR* and *FSHR* (Spearman rank sum test:  $N = 310$ ;  $R = 0.77$ ;  $P < 0.00001$ ). The *AMH* expression decreased with increasing

follicular diameter as expected (**Figure 3**). As for the substances present in FF, the variability of the gene expression was massive in between individual follicles with similar diameter (Table S3 in Supplementary Material).

## DISCUSSION

Including data from almost 1,000 normal hSAF, the present study documented profound changes taking place in the average concentrations of multiple key FF hormones and GC-expressed genes in relation to follicular development. It is especially noticeable that for all the secreted hormones, major alterations in the intrafollicular concentrations emerge with diameters from around 8 to 11 mm (**Figure 4**). This corresponds to the follicular stage in which selection of the dominant follicle takes place in each individual menstrual cycle and demonstrates the dramatic changes that occur during the transition of follicles from the recruitable stage to the selected follicle destined to ovulate with a pronounced shift in steroidogenesis resulting in a massive synthesis of estradiol/*CYP19* (aromatase) and progesterone. Overall, this enforces the notion introduced by Yen and Jaffe (29) almost half a century ago, hormonal regulation of the follicular phase of the menstrual cycle differs in two periods during a first half lasting to around cycle day 7 and a second half starting from the day of follicle selection and until ovulation.

Furthermore, the current data not only corroborate and extend that regulation of ovarian steroidogenesis during the follicular phase of the menstrual cycle is governed by FSH and LH but also highlight that local TGF- $\beta$  growth factors exert important paracrine and autocrine functions that act to secure the development of one competent follicle selected for ovulation per cycle.

The present study demonstrates that intrafollicular concentrations of estradiol start to rise rapidly in the window of follicular selection (i.e., 8–11 mm) paralleled by concomitant increase in *CYP19* mRNA expression. Several mechanisms are probably involved in this strong upregulation. First, inhibins become synthesized as a result of FSH stimulation. Thus, the follicle that secretes high amounts of inhibins probably also possesses high *FSHR* expression. So not only does FSH secure substrate for aromatization *via* inhibin and subsequent androgen synthesis but does also stimulate aromatase and LHR expression, which are essential for estradiol production during the second half of the follicular phase (5). Despite the high levels of FSH in first half of the follicular phase aromatase expression and estradiol production remain relatively low (30) as the FF data of the present study also suggest. This appears to be an effect of AMH which in the very high concentrations present in hSAF exert a negative effect on estradiol production (18, 22, 31). At follicular selection, the dominant follicle acquires strong aromatase expression and upregulate estradiol production, which to a large extent appears to be driven by decreasing levels of AMH (attenuating) and increasing levels of inhibins (augmenting). In this scenario, AMH and inhibins could be expected to be inversely associated, which indeed is the case in the present study and confirms an earlier study (22). This suggests that both AMH and inhibin-B are intimately associated and both involved in the regulation of follicular estradiol synthesis.







FIGURE 2 | Continued

**FIGURE 2** | Intrafollicular content and concentrations of peptide and steroid hormones. Graphs show the follicular fluid concentrations of estradiol, progesterone, androstenedione, testosterone, inhibin B, inhibin A, AMH, PAPP-A, and follistatin, and follicular content in nanograms (bottom panel) of inhibin B, inhibin A, and AMH according to follicular diameter. Values are presented as mean  $\pm$  SEM. Statistically significant differences were observed in intrafollicular hormone levels of estradiol [analysis of variance (ANOVA);  $N = 264$ ;  $P < 0.001$ ], androstenedione (ANOVA;  $N = 276$ ;  $P < 0.03$ ), progesterone (ANOVA;  $N = 267$ ;  $P < 0.003$ ), PAPP-A (ANOVA;  $N = 96$ ;  $P < 0.03$ ), inhibin-B (ANOVA;  $N = 627$ ;  $P < 0.002$ ), inhibin-A (ANOVA;  $N = 258$ ;  $P < 0.001$ ), AMH (ANOVA;  $N = 655$ ;  $P < 0.001$ ), but not testosterone (ANOVA;  $N = 566$ ;  $P > 0.10$ ) and follistatin (ANOVA;  $N = 148$ ;  $P > 0.10$ ). Regarding follicular content (bottom panel), the intrafollicular contents of inhibin-B (ANOVA;  $N = 611$ ;  $P < 0.001$ ), inhibin-A (ANOVA;  $N = 258$ ;  $P < 0.001$ ), and AMH (ANOVA;  $N = 654$ ;  $P < 0.001$ ) were observed to be significantly different in relation to follicular diameter. Specific numbers for the analyzed follicle groups divided according to follicular diameter are shown in Tables S1 and S2 in Supplementary Material.




**FIGURE 3** | Granulosa cell expression of selected genes in human small antral follicles. Graphs show relative expression levels of *FSHR*, *CYP19a1*, *AR*, and *AMH* in relation to follicular diameter. Values are presented as mean  $\pm$  SEM. Statistically significant differences were observed in the mRNA expression levels of *FSHR* [analysis of variance (ANOVA);  $N = 315$ ;  $P < 0.002$ ] and *AR* (ANOVA;  $N = 316$ ;  $P < 0.03$ ), but not for *CYP19a1* (ANOVA;  $N = 252$ ;  $P > 0.10$ ) and *AMH* (ANOVA;  $N = 277$ ;  $P > 0.10$ ). Specific numbers for the analyzed follicle groups divided according to follicular diameter are shown in Table S3 in Supplementary Material.

In addition, IGF2 is also likely to play an important role in the regulation of estradiol synthesis by the selected follicle. The present study showed that the intrafollicular concentration of PAPP-A increased almost threefold in follicles with diameters increasing from 8 to 12 mm. Again, assuming that PAPP-A expression reflects IGF bioactivity (32–36), it is noteworthy that PAPP-A expression moves from the theca cells to the GC as the follicles become dominant (24, 34–36). The increased IGF activity may serve two purposes: first, the exceptional high proliferation of GC during the second half of the follicular phase probably requires a strong local signal especially since the *FSHR* expression becomes downregulated (21). Second, IGF2 on its own stimulates human GC estradiol synthesis (37–40), which has recently been shown with highly significant associations between PAPP-A activity in FF and estradiol concentrations in FF from different-sized follicles (24, 34).

It is surprising that *FSHR* expression per cell declines with follicular diameter and is significantly reduced at follicular selection

during the natural menstrual cycle, confirming an earlier study (21). When GC from preovulatory follicles prior to ovulation induction were challenged by similar concentrations of FSH and LH, aromatase *in vitro* induction was clearly biggest after stimulation with LH, and in connection with ovarian stimulation for infertility treatment it has been shown that just a modest addition of LH-like activity results in a pronounced augmentation of estradiol. Collectively suggesting that LH probably also plays a prominent role in stimulation of aromatase expression. Furthermore, it is likely that there is a redundancy between FSH and LH in induction of aromatase expression and that ovarian stimulation with exogenous FSH preparations overcomes a reduction in *FSHR* expression. Furthermore, as the present results are expressed as *FSHR* expression per cell, the collective follicular *FSHR* number may still increase because of the hectic proliferation of GC as the follicle enlarges.

The inhibitory effect of inhibins in concentrations of picograms per milliliter on pituitary FSH release is well recognized, but the



importance of the simultaneous paracrine effect of inhibins in synergy with LH on theca cell androgen synthesis continues to be unrevealed in human ovaries. The intrafollicular concentration of inhibin-B plus inhibin-A remained relatively constant around 100–120 ng/ml in follicles with diameters of 3–7 mm but doubled to 200–220 ng/ml and showed a peak in follicles with diameters of 9–10 mm. Previous *in vitro* studies of human theca cells showed that inhibins act in synergy with LH to upregulate synthesis of androstenedione (8–11, 41–45). The maximal concentration of inhibins tested in these studies was 100 ng/ml. Indeed, the collective intrafollicular concentrations reach at least twice this concentration and make it conceivable that the theca cells actually are exposed to such high concentrations *in vivo*. Obviously, theca produced androstenedione requires further metabolism to testosterone and to diffuse to the follicular compartment to exert a true androgenic effect. In this context, it is noteworthy that intrafollicular concentrations of testosterone in parallel to intrafollicular inhibins also peak in follicles with diameters of around 9–11 mm in line with recent results showing strong

significant associations between inhibin-B and androstenedione and testosterone in hSAFs (5). The intrafollicular testosterone concentrations exert a stimulatory effect on the GC expression of *FSHR* (19, 46, 47) and subsequently expression of *LHR*. Androgens act through the AR, and the notion is enforced by the highly significant association between *FSHR* and *AR* mRNA levels in GC from hSAFs of the present study (19). Thereby, the selected follicle on one side secures not only a sufficient substrate for a high estrogen production but also a sufficient *FSHR* expression, which in light of declining FSH levels is required to secure sufficient *LHR* expression to sustain ovulation. In fact, the shape of the *FSHR* and *AR* mRNA expression profile in relation to follicular diameter is almost superimposable and suggests a close interrelationship.

It has also been reported that in humans especially IGF2 augment theca cell-derived androgen production in synergy with LH [human (48–50), primate (51), and rats (11, 52, 53)]. It has previously been difficult to account for the biological activity of IGFs because there are several different binding proteins (i.e., IGFBP 1–6) inactivating IGF activity, proteases that specifically cleave different IGFBPs to release bound IGFs (i.e., PAPP-A and PAPP-A2), and furthermore specific proteins that inactivate the proteolytic activity of PAPP-A (i.e., stanniocalcin 1 and 2) (32, 34). On top of this complicated system for regulating IGF bioactivity, it appears that the presence of PAPP-A covalently linked to the cell membrane secures a local microenvironment where IGFs released after specific cleavage by PAPP-A have a possibility to activate IGF receptors and cause signal transduction (32). Collectively, however, several studies have now shown that concentrations of PAPP-A appear to be a good surrogate marker for the bioactivity of IGFs (32, 34). It has been shown that PAPP-A becomes expressed in the theca cells of hSAF moving to the GC as the follicle develops further (24). These results suggest a local production of IGFs that corroborate *in vitro* studies (11, 48–52) that showed a synergistic effect of LH and IGFs on theca cell androgen production.

Collectively, these new data support the original hypothesis proposed by Hillier and coworkers in their landmark studies on human tissue in the early 1990s and place especially inhibin-B and IGF2 in a physiological context as important paracrine regulators augmenting steroid output by especially the selected follicle in the second half of the human follicular phase.

Both inhibins and IGFs themselves can independently stimulate androgen production in synergy with LH, but it appears that they utilize separate mechanisms. Inhibins are capable of enhancing the combined maximal stimulatory effect of LH and IGF1 with an up to 10-fold higher androgen output in human theca cells *in vitro* (12, 43).

While inhibins augment theca cell androgen production, activins, in contrast, have been shown to exert the opposite effect and reduce androgen output *via* downregulating expression of CYP17 and 3 $\beta$ -hydroxy steroid dehydrogenase [human (12, 54), rat (9), bovine (55, 56), sheep (42, 44, 57)]. No specific receptors for inhibins have been identified and no corresponding second messenger signaling pathway system identified. The affinity of activin for the activin type II receptor is around 10-fold higher than that of inhibins (58), but by binding to the membrane bound  $\beta$ -glycan,



inhibins augment interaction with the activin type II receptor and enhance its effect as an activin antagonist (6). In this context, it is interesting to notice that FF concentrations of inhibin-A start to surpass those of inhibin-B when follicles exceed a diameter of around 13–14 mm. In circulation, this is only observed close to ovulation when the effect of massive inhibin-B production in connection with selection is being reduced and suggests that theca cell androgen production is maintained at a high level until ovulation.

Taken together, regulation of androgen production in hSAF is likely to involve an upregulation of LHR number *via* locally expressed PAPP-A, which leads to enhanced IGF2 activity that acts on the IGFR being expressed on theca cells (59). Independent of the IGF2 action, inhibins in FF of hSAF being present in high concentrations reduce a negative impact of activins and enhance the 17-hydroxylase/C<sub>17-20</sub> lyase activity that further augments the androgen producing capacity.

In this context, it is noteworthy that LHR numbers have been shown to be a major determinant for hCG-induced steroidogenic output from cells during culture highlighting that LHR density is of importance for LH action (60). Therefore, the follicle where the surrounding theca cells respond with highest production of androgen is most likely to become the selected follicle destined to ovulate as recently described (5). The present data emphasize the exquisite sensitivity of theca cells to LH stimulation in combination with inhibins, IGFs, and potentially other locally produced growth factors in the regulation of ovarian steroidogenesis and function.

Collectively, a local amplification of thecal androgen production by inhibins and IGF2 suggests a mechanism through which follicular dominance is acquired and maintained in human reproductive cycles.

A major limitation of the present study is the inability to distinguish between healthy follicles and follicles which have started to undergo atresia. A fraction of the follicles in this study have already become atretic to some degree (61), but will be saved by administration of exogenous FSH preparations. It is well-known that oocytes from a cohort of follicles as a result of exogenous FSH stimulation may result in several children pointing at a clear clinical interest. The degree of atresia is likely to affect steroidogenesis and to what extent this affects the outcome measures as reported in the present study is unknown. This may reflect that the FF concentrations as reported in the present study show a massive range, often more than three orders of magnitude from one follicle to the other, which, however, may not be linked to atresia but rather reflect other factors. Obviously, there is a whole range of methods to detect the rate of atresia, but the often very limited number of GC from each individual follicle does not allow for detailed analysis.

The present study has only addressed a limited number of albeit prominent TGF- $\beta$  growth factors and their presence in FF from hSAF and their potential effect on ovarian steroidogenesis. Indeed, other TGF- $\beta$  growth factors have been reported to affect ovarian steroidogenesis, and it is likely that regulation involves a number of other factors such as BMP 4, 6, and 7 (42, 54, 62, 63).

It is interesting to notice that a number of the mechanisms proposed to regulate ovarian steroidogenesis involve release of biological active substances in the vicinity of the cell membrane. Covalently coupled PAPP-A appears to cleave IGFBP to release

IGFs close to the membrane bound receptor (33), and locally high concentrations of inhibins can attenuate even locally produced activin from interaction with its receptor are examples that may improve our understanding of how the fine-tuned hormonal regulation of ovarian steroidogenesis takes place *in vivo*.

In conclusion, the present study suggests that regulation of human ovarian steroidogenesis during the follicular phase differs between the first part of the follicular phase in which recruitment takes place and the second half where development of the dominant follicle concomitant with a massive output of estradiol. Especially with follicular diameters of around 8–11 mm, regulation of steroid production change profoundly with significant impact from paracrine and autocrine regulation of TGF- $\beta$  growth factors especially inhibins and activins that together with IGF2 *via* PAPP-A modulate and fine-tune the effects of FSH and LH in order to secure the development of one dominant follicle in each menstrual cycle. The pronounced effects of inhibin-B are noticeable acting both to reduce pituitary output of FSH, but also as a follicle derived signal to augment androgen production locally by the theca cells, which in turn upregulate *FSHR* mRNA expression that enables the dominant follicle to respond to the decreasing levels of FSH by having aromatase and *LHR* mRNA expression upregulated.

## ETHICS STATEMENT

This study was carried out in accordance with the recommendations of the ethical committee of the municipalities of Copenhagen and Frederiksberg with written informed consent from all subjects. All subjects gave written informed consent in accordance with the Declaration of Helsinki. The protocol was approved by the ethical committee of the municipalities of Copenhagen and Frederiksberg (journal number; H-2-2011-044).

## AUTHOR CONTRIBUTIONS

CA and SK designed the study and wrote the manuscript. KM and EE enrolled patients for the study. SK, JJ, and SP collected follicular fluids and GCs for the study. TB, JB, and JJ performed the gene expression analysis and ELISA/RIA analysis.

## ACKNOWLEDGMENTS

The University Hospital of Copenhagen, the EU interregional project ReproUnion, and Vera and Carl Johan Michaelsens Legat are gratefully acknowledged for financial support. They had no role in the study design, collection and analysis of data, data interpretation, or in writing the report. Follicular fluid samples used in this study were obtained from women undergoing fertility preservation by having one ovary frozen at our institution. All personnel including the clinical activities in fertility preservation are thanked for their passionate work.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at <http://www.frontiersin.org/articles/10.3389/fendo.2017.00376/full#supplementary-material>.

## REFERENCES

- Yding Andersen C. Characteristics of human follicular fluid associated with successful conception after in vitro fertilization. *J Clin Endocrinol Metab* (1993) 77:1227–34. doi:10.1210/jcem.77.5.7521343
- McNatty KP, Baird DT, Bolton A, Chambers P, Corker CS, McLean H. Concentration of oestrogens and androgens in human ovarian venous plasma and follicular fluid throughout the menstrual cycle. *J Endocrinol* (1976) 71:77–85. doi:10.1677/joe.0.0710077
- Yding Andersen C. Possible new mechanism of cortisol action in female reproductive organs: physiological implications of the free hormone hypothesis. *J Endocrinol* (2002) 173:211–7. doi:10.1677/joe.0.1730211
- Yding Andersen C, Schmidt KT, Gry Kristensen S, Rosendahl M, Byskov AG, Ernst E. Concentrations of AMH and Inhibin-B in relation to follicular diameter in normal human small antral follicles. *Hum Reprod* (2010) 25:1282–7. doi:10.1093/humrep/deq019
- Yding Andersen C. Inhibin-B secretion and FSH isoform distribution may play an integral part of follicular selection in the natural menstrual cycle. *Mol Hum Reprod* (2017) 23:16–24. doi:10.1093/molehr/gaw070
- Knight PG, Glistler C. Local roles of TGF-beta superfamily members in the control of ovarian follicle development. *Anim Reprod Sci* (2003) 78:165–83. doi:10.1016/S0378-4320(03)00089-7
- Kranc W, Budna J, Kahan R, Chachula A, Bryja A, Ciesiolka S, et al. Molecular basis of growth, proliferation, and differentiation of mammalian follicular granulosa cells. *J Biol Regul Homeost Agents* (2017) 31:1–8.
- Juengel JL, McNatty KP. The role of proteins of the transforming growth factor-beta superfamily in the intraovarian regulation of follicular development. *Hum Reprod Update* (2005) 11:143–60. doi:10.1093/humupd/dmh061
- Hsueh AJ, Dahl KD, Vaughan J, Tucker E, Rivier J, Bardin CW, et al. Heterodimers and homodimers of inhibin subunits have different paracrine action in the modulation of luteinizing hormone-stimulated androgen biosynthesis. *Proc Natl Acad Sci USA* (1987) 84:5082–6. doi:10.1073/pnas.84.14.5082
- Campbell BK, Baird DT, Webb R. Effects of dose of LH on androgen production and luteinization of ovine theca cells cultured in a serum-free system. *J Reprod Fertil* (1998) 112:69–77. doi:10.1530/jrf.0.1120069
- Cara JF, Rosenfield RL. Insulin-like growth factor I and insulin potentiate luteinizing hormone-induced androgen synthesis by rat ovarian thecal-interstitial cells. *Endocrinology* (1988) 123:733–9. doi:10.1210/endo-123-2-733
- Hillier SG, Yong EL, Illingworth PJ, Baird DT, Schwall RH, Mason AJ. Effect of recombinant inhibin on androgen synthesis in cultured human thecal cells. *Mol Cell Endocrinol* (1991) 75(2):R1–6. doi:10.1016/0303-7207(91)90234-J
- McNatty KP, Smith DM, Makris A, Osathanondh R, Ryan KJ. The microenvironment of the human antral follicle: interrelationships among the steroid levels in antral fluid, the population of granulosa cells, and the status of the oocyte in vivo and in vitro. *J Clin Endocrinol Metab* (1979) 49(12):851–60. doi:10.1210/jcem-49-6-851
- Humaidan P, Westergaard LG, Mikkelsen AL, Fukuda M, Yding Andersen C. Levels of the EGF-like peptide amphiregulin in follicular fluid reflect the mode of triggering ovulation: a comparison between GnRHa and hCG. *Fertil Steril* (2011) 95:2034–8. doi:10.1016/j.fertnstert.2011.02.013
- Rosendahl M, Schmidt KT, Ernst E, Rasmussen PE, Loft A, Byskov AG, et al. Cryopreservation of ovarian tissue for a decade in Denmark – an overview of the technique. *Reprod Med Online* (2011) 22:162–71. doi:10.1016/j.rbmo.2010.10.015
- Yding Andersen C, Byskov AG. Is oestradiol an important regulator for secretion of anti-Müllerian hormone, inhibin-A and inhibin-B: analysis of fluid from small antral and preovulatory human follicles. *J Clin Endocrinol Metab* (2006) 91:4064–9. doi:10.1210/jc.2006-1066
- Yding Andersen C, Rosendahl M, Byskov AG. Concentration of anti-Müllerian hormone and inhibin-B in relation to steroids and age in follicular fluid from small antral human follicles. *J Clin Endocrinol Metab* (2008) 93:2344–9. doi:10.1210/jc.2007-2342
- Eilsø Nielsen M, Rasmussen IA, Fukuda M, Westergaard LG, Yding Andersen C. Concentrations of anti-Müllerian hormone in fluid from small human antral follicles shows a negative correlation with CYP19 mRNA expression in the corresponding granulosa cells. *Mol Hum Reprod* (2010) 16:637–43. doi:10.1093/molehr/gaq001
- Eilsø Nielsen M, Rasmussen IA, Kristensen SG, Christensen ST, Møllgård K, Wreford Andersen E, et al. Expression of androgen-receptor mRNA in granulosa cells from human small antral follicles and the corresponding follicular fluid concentrations of androgens are positively correlated to granulosa cell FSH receptor mRNA expression. *Mol Hum Reprod* (2011) 17:63–70. doi:10.1093/molehr/gaq073
- Jeppesen JV, Nielsen ME, Kristensen SG, Yding Andersen C. Concentration of activin A and follistatin in follicular fluid from human small antral follicles associated to gene expression of the corresponding granulosa cells. *Mol Cell Endocrinol* (2012) 356:48–54. doi:10.1016/j.mce.2011.07.051
- Jeppesen JV, Kristensen SG, Nielsen ME, Humaidan P, Dal Canto M, Fadini R, et al. LH-receptor gene expression in human granulosa and cumulus cells from antral and preovulatory follicles. *J Clin Endocrinol Metab* (2012) 97:1524–31. doi:10.1210/jc.2012-1427
- Jeppesen JV, Anderson RA, Kelsey TW, Christiansen SL, Kristensen SG, Jayaprakasan K, et al. Which follicles make the most anti-Müllerian hormone in humans? Evidence for an abrupt decline in AMH production at the time of follicle selection. *Mol Hum Reprod* (2013) 19:519–27. doi:10.1093/molehr/gat024
- Borgbo T, Jeppesen JV, Lindgren I, Lundberg Giwerzman Y, Hansen LL, Yding Andersen C. Effect of the FSH receptor single nucleotide polymorphisms (FSHR 307/680) on the follicular fluid hormone profile and the granulosa cell gene expression in human small antral follicles. *Mol Hum Reprod* (2015) 21:255–61. doi:10.1093/molehr/gau106
- Botkjaer JA, Jeppesen JV, Wissing ML, Kløverpris S, Oxvig C, Mason JI, et al. Pregnancy-associated plasma protein A in human ovarian follicles and its association with intrafollicular hormone levels. *Fertil Steril* (2015) 104:1294–301. doi:10.1016/j.fertnstert.2015.07.1152
- Borgbo T, Macek M Sr, Chrudimska J, Jeppesen JV, Hansen LL, Andersen CY. Size matters: associations between the androgen receptor CAG repeat length and the intrafollicular hormone milieu. *Mol Cell Endocrinol* (2016) 419:12–7. doi:10.1016/j.mce.2015.09.015
- Botkjaer JA, Borgbo T, Kløverpris S, Noer PR, Oxvig C, Andersen CY. Effect of pregnancy-associated plasma protein-A (PAPP-A) single-nucleotide polymorphisms on the level and activity of PAPP-A and the hormone profile in fluid from normal human small antral follicles. *Fertil Steril* (2016) 106:1778–86. doi:10.1016/j.fertnstert.2016.09.008
- Westergaard L, McNatty KP, Christensen IJ. Steroid concentrations in fluid from human ovarian antral follicles during pregnancy. *J Endocrinol* (1985) 107:133–6. doi:10.1677/joe.0.1070133
- Westergaard L, Christensen IJ, McNatty KP. Steroid levels in ovarian follicular fluid related to follicle size and health status during the normal menstrual cycle in women. *Hum Reprod* (1986) 1:227–32. doi:10.1093/oxfordjournals.humrep.a136390
- Yen SSC. The human menstrual cycle (integrative function of the hypothalamic-pituitary-ovarian-endometrial axis). In: Yen SSC, Jaffe RB, editors. *Reproductive Endocrinology*, Philadelphia: WB Saunders Co. (1978) 126 p.
- Greame NP, Illingworth PJ, O'Brien M, Pai R, Rodger FE, Mather JB, et al. Measurement of dimeric inhibin-B throughout the human menstrual cycle. *J Clin Endocrinol Metab* (1996) 81:1401–5. doi:10.1210/jcem.81.4.8636341
- Dewailly D, Andersen CY, Balen A, Broekmans F, Dilaver N, Fanchin R, et al. The physiology and clinical utility of anti-Müllerian hormone in women. *Hum Reprod Update* (2014) 20:370–85. doi:10.1093/humupd/dmt062
- Oxvig C. The role of PAPP-A in the IGF system: location, location, location. *J Cell Commun Signal* (2015) 9:177–87. doi:10.1007/s12079-015-0259-9
- Monget P, Oxvig C. PAPP-A and the IGF system. *Ann Endocrinol (Paris)* (2016) 77:90–6. doi:10.1016/j.ando.2016.04.015
- Jepsen MR, Kløverpris S, Botkjaer JA, Wissing ML, Andersen CY, Oxvig C. The proteolytic activity of pregnancy-associated plasma protein-A is potentially regulated by stanniocalcin-1 and -2 during human ovarian follicle development. *Hum Reprod* (2016) 31:866–74. doi:10.1093/humrep/dew013
- Fortune JE, Rivera GM, Yang MY. Follicular development: the role of the follicular microenvironment in selection of the dominant follicle. *Anim Reprod Sci* (2004) 8(2–83):109–26. doi:10.1016/j.anireprosci.2004.04.031
- Conover CA, Faessen GF, Ilg KE, Chandrasekher YA, Christiansen M, Overgaard MT, et al. Pregnancy-associated plasma protein-a is the insulin-like growth factor binding protein-4 protease secreted by human ovarian granulosa cells and is a marker of dominant follicle selection and the corpus luteum. *Endocrinology* (2001) 142(5):2155. doi:10.1210/endo.142.5.8286

37. Giudice LC. Insulin-like growth factor family in Graafian follicle development and function. *J Soc Gynecol Invest* (2001) 8:26–9. doi:10.1016/S1071-5576(00)00102-7
38. Thierry van Dessel HJ, Chandrasekhar Y, Yap OW, Lee PD, Hintz RL, Faessen GH, et al. Serum and follicular fluid levels of insulin-like growth factor I (IGF-I), IGF-II, and IGF-binding protein-1 and -3 during the normal menstrual cycle. *J Clin Endocrinol Metab* (1996) 81(3):1224–31. doi:10.1210/jcem.81.3.8772603
39. Silva JR, Figueiredo JR, van den Hurk R. Involvement of growth hormone (GH) and insulin-like growth factor (IGF) system in ovarian folliculogenesis. *Theriogenology* (2009) 71(8):1193–208. doi:10.1016/j.theriogenology.2008.12.015
40. Spicer LJ, Aad PY. Insulin-like growth factor (IGF) 2 stimulates steroidogenesis and mitosis of bovine granulosa cells through the IGF1 receptor: role of follicle-stimulating hormone and IGF2 receptor. *Biol Reprod* (2007) 77(1):18–27. doi:10.1095/biolreprod.106.058230
41. Duleba AJ, Spaczynski RZ, Olive DL, Behrman HR. Divergent mechanisms regulate proliferation/survival and steroidogenesis of theca-interstitial cells. *Mol Hum Reprod* (1999) 5:193–8. doi:10.1093/molehr/5.3.193
42. Hoang YD, McTavish KJ, Chang RJ, Shimasaki S. Paracrine regulation of theca androgen production by granulosa cells in the ovary. *Fertil Steril* (2013) 100:561–7. doi:10.1016/j.fertnstert.2013.04.016
43. Nahum R, Thong KJ, Hillier SG. Metabolic regulation of androgen production by human thecal cells in vitro. *Hum Reprod* (1995) 10:75–81. doi:10.1093/humrep/10.1.75
44. Young JM, McNeilly AS. Theca: the forgotten cell of the ovarian follicle. *Reproduction* (2010) 140(4):489–504. doi:10.1530/REP-10-0094
45. Young JM, McNeilly AS. Inhibin removes the inhibitory effects of activin on steroid enzyme expression and androgen production by normal ovarian thecal cells. *J Mol Endocrinol* (2012) 48(1):49–60. doi:10.1530/JME-11-0134
46. Weil S, Vendola K, Zhou J, Bondy CA. Androgen and follicle-stimulating hormone interactions in primate ovarian follicle development. *J Clin Endocrinol Metab* (1999) 84:2951–6. doi:10.1210/jcem.84.8.5929
47. Weil SJ, Vendola K, Zhou J, Adesanya OO, Wang J, Okafor J, et al. Androgen receptor gene expression in the primate ovary: cellular localization, regulation, and functional correlations. *J Clin Endocrinol Metab* (1998) 83:2479–85. doi:10.1210/jcem.83.7.4917
48. Barbieri RL, Makris A, Randall RW, Daniels G, Kistner RW, Ryan KJ. Insulin stimulates androgen accumulation in incubations of ovarian stroma obtained from women with hyperandrogenism. *J Clin Endocrinol Metab* (1986) 62(5):904–10. doi:10.1210/jcem-62-5-904
49. Yong EL, Baird DT, Yates R, Reichert LE Jr, Hillier SG. Hormonal regulation of the growth and steroidogenic function of human granulosa cells. *J Clin Endocrinol Metab* (1992) 74:842–9. doi:10.1210/jcem.74.4.1548349
50. Bergh C, Carlsson B, Olsson JH, Selleskog U, Hillensjö T. Regulation of androgen production in cultured human thecal cells by insulin-like growth factor I and insulin. *Fertil Steril* (1993) 59:323–31. doi:10.1016/S0015-0282(16)55675-1
51. Vendola K, Zhou J, Wang J, Bondy CA. Androgens promote insulin-like growth factor-I and insulin-like growth factor-I receptor gene expression in the primate ovary. *Hum Reprod* (1999) 14:2328–32. doi:10.1093/humrep/14.9.2328
52. Cara JF, Fan J, Azzarello J, Rosenfield RL. Insulin-like growth factor-I enhances luteinizing hormone binding to rat ovarian theca-interstitial cells. *J Clin Invest* (1990) 86(2):560–5. doi:10.1172/JCI114745
53. Adashi EY, Resnick CE, Hurwitz A, Ricciarelli E, Hernandez ER, Roberts CT, et al. Insulin-like growth factors: the ovarian connection. *Hum Reprod* (1991) 6:1213–9. doi:10.1093/oxfordjournals.humrep.a137514
54. Dooley CA, Attia GR, Rainey WE, Moore DR, Carr BR. Bone morphogenetic protein inhibits ovarian androgen production. *J Clin Endocrinol Metab* (2000) 85:3331–7. doi:10.1210/jcem.85.9.6835
55. Wrathall JH, Knight PG. Effects of inhibin-related peptides and oestradiol on androstenedione and progesterone secretion by bovine theca cells in vitro. *J Endocrinol* (1995) 145:491–500. doi:10.1677/joe.0.1450491
56. Shukovski L, Dyson M, Findlay JK. The effects of follistatin, activin and inhibin on steroidogenesis by bovine thecal cells. *Mol Cell Endocrinol* (1993) 97:19–27. doi:10.1016/0303-7207(93)90207-Z
57. Young JM, Henderson S, Souza C, Ludlow H, Groome N, McNeilly AS. Activin B is produced early in antral follicular development and suppresses thecal androgen production. *Reproduction* (2012) 143:637–50. doi:10.1530/REP-11-0327
58. Knight PG, Satchell L, Glistler C. Intra-ovarian roles of activins and inhibins. *Mol Cell Endocrinol* (2012) 359(1–2):53–65. doi:10.1016/j.mce.2011.04.024
59. Poretsky L, Bhargava G, Levitan E. Type I insulin-like growth factor receptors in human ovarian stroma. *Horm Res* (1990) 33:22–6. doi:10.1159/000181441
60. Hirakawa T, Galet C, Ascoli M. MA-10 cells transfected with the human lutropin/choriogonadotropin receptor (hLHR): a novel experimental paradigm to study the functional properties of the hLHR. *Endocrinology* (2002) 143:1026–35. doi:10.1210/endo.143.3.8702
61. Gougeon A. Regulation of ovarian follicular development in primates: facts and hypotheses. *Endocr Rev* (1996) 7(2):121–55. doi:10.1210/edrv-17-2-121
62. Glistler C, Richards SL, Knight PG. Bone morphogenetic proteins (BMP)-4, -6, and -7 potentially suppress basal and luteinizing hormone-induced androgen production by bovine theca interna cells in primary culture: could ovarian hyperandrogenic dysfunction be caused by a defect in thecal BMP signaling? *Endocrinology* (2005) 146:1883–92. doi:10.1210/en.2004-1303
63. Glistler C, Satchell L, Knight PG. Changes in expression of bonemorphogenetic proteins (BMPs), their receptors and inhibin co-receptor betaglycan during bovine antral follicle development: inhibin can antagonize the suppressive effect of BMPs on thecal androgen production. *Reproduction* (2010) 140:699–712. doi:10.1530/REP-10-0216

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Kristensen, Mamsen, Jeppesen, Bøtkjær, Pors, Borgbo, Ernst, Macklon and Andersen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.