



## **Night work and postpartum depression a national register-based cohort study**

Hammer, Paula; Hageman, Ida; Garde, Anne; Begtrup, Luise; Flachs, Esben; Hansen, Johnni; Hansen, Åse Marie; Hougaard, Karin; Kolstad, Henrik; Larsen, Ann; Pinborg, Anja; Specht, Ina; Bonde, Jens Peter

*Published in:*  
Scandinavian Journal of Work, Environment & Health

*DOI:*  
[10.5271/sjweh.3831](https://doi.org/10.5271/sjweh.3831)

*Publication date:*  
2019

*Document version*  
Publisher's PDF, also known as Version of record

*Document license:*  
[CC BY](https://creativecommons.org/licenses/by/4.0/)

*Citation for published version (APA):*  
Hammer, P., Hageman, I., Garde, A., Begtrup, L., Flachs, E., Hansen, J., ... Bonde, J. P. (2019). Night work and postpartum depression: a national register-based cohort study. *Scandinavian Journal of Work, Environment & Health*, 45(6), 577-587. <https://doi.org/10.5271/sjweh.3831>



## Original article

Scand J Work Environ Health 2019;45(6):577-587

doi:10.5271/sjweh.3831

### Night work and postpartum depression: a national register-based cohort study

by [Hammer P](#), [Hageman I](#), [Garde A](#), [Begtrup L](#), [Flachs E](#), [Hansen J](#), [Hansen Å](#), [Hougaard K](#), [Kolstad H](#), [Larsen A](#), [Pinborg A](#), [Specht I](#), [Bonde JP](#)

The novel aspects of our study are the use of clinically diagnosed severe postpartum depression, objective information on pregnancy period-specific exposure and within-night work comparisons. The results suggest that changes of working schedules might mask a potential risk of postpartum depression among women who work at night during pregnancy.

**Affiliation:** Department of Occupational and Environmental Medicine, Bispebjerg and Frederiksberg University Hospital, Bispebjerg Bakke 23, indgang 20F - 2400 Copenhagen, Denmark. paulahammer@dadlnet.dk

Refers to the following texts of the Journal: [2011;37\(5\):402-410](#) [2015;41\(3\):268-279](#) [2018;44\(4\):403-413](#) [2018;44\(5\):512-520](#)

**Key terms:** [cohort study](#); [depression](#); [depressive disorder](#); [doctor](#); [healthy worker effect](#); [night work](#); [nurse](#); [postpartum depression](#); [pregnancy](#); [register-based cohort study](#); [shift work](#); [shift worker](#)

This article in PubMed: [www.ncbi.nlm.nih.gov/pubmed/31125110](http://www.ncbi.nlm.nih.gov/pubmed/31125110)

### Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).



This work is licensed under a [Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/).

## Night work and postpartum depression: a national register-based cohort study

by Paula Hammer, MD, PhD,<sup>1</sup> Ida Hageman, MD, PhD,<sup>2</sup> Anne Garde, PhD,<sup>3,4</sup> Luise Begtrup, MD, PhD,<sup>1</sup> Esben Flachs, MSc, PhD,<sup>1</sup> Johnni Hansen, MSc, PhD,<sup>5</sup> Åse Hansen, DMSc, PhD,<sup>3,4</sup> Karin Hougaard, MSc, PhD,<sup>3,4</sup> Henrik Kolstad, MD, PhD,<sup>6</sup> Ann Larsen, MSc, PhD,<sup>3</sup> Anja Pinborg, MD, PhD,<sup>7</sup> Ina Specht, MSc, PhD,<sup>8</sup> Jens Peter Bonde, MD, PhD<sup>1,4</sup>

Hammer P, Hageman I, Garde A, Begtrup L, Flachs E, Hansen J, Hougaard K, Kolstad H, Larsen A, Pinborg A, Specht I, Bonde JP. Night work and postpartum depression: a national register-based cohort study. *Scand J Work Environ Health*. 2019;45(6):577–587. doi:10.5271/sjweh.3831

**Objective** We aimed to investigate the association of night work during pregnancy with the risk of severe postpartum depression (PPD).

**Methods** We performed a nationwide register-based cohort study of workers in all Danish public hospitals. Daily information on working hours was retrieved from the Danish Working Hour Database from January 2007 to December 2015. Pregnancies, covariates and outcome were identified from national registries for births and hospital contacts. We performed logistic regression of the risk of severe PPD in relation to the number and duration of night shifts, spells of consecutive night shifts, and short shift intervals during the first 32 pregnancy weeks. Analyses were adjusted for age, body mass index, socioeconomic status, parity, sickness absence three months prior to pregnancy, and prior diagnosis of severe depression.

**Results** The study cohort comprised 25 009 singleton pregnancies from 19 382 workers. The majority were nurses or physicians. Overall, we did not observe an increased risk of PPD for any of the dimensions of night work analyzed. We found, however, an increased risk of PPD (adjusted odds ratio 2.08, 95% confidence interval 1.09–4.00) among women who stopped working night shifts after the first pregnancy trimester (N=3094).

**Conclusion** Overall, our results do not support night work during pregnancy as a risk factor for severe PPD among hospital employees. However, we observed a 2-fold increased risk of PPD among women who stopped working night shifts after the first pregnancy trimester. This may reflect the influence of the healthy worker survivor effect and warrants further attention.

**Key terms** depressive disorder; doctor; healthy worker effect; nurse; pregnancy; shift work; shift worker.

The prevalence of postpartum depression (PPD) is estimated to be 10–15%, but considerably higher rates have been reported across countries (1, 2). PPD imposes potential severe consequences for the mother, child, family and society and is one of the primary causes of maternal mortality in many countries (3, 4). It is therefore considered a global public health problem (5).

The pathogenesis of PPD is complex and multifactorial and involves endocrine, genetic and environmental factors (1, 6). The strongest known risk factors for PPD

are prior own or family history of major depressive disorder (1, 6, 7). Other factors shown to be related to the development of PPD include adverse life events, inadequate social support, and low socioeconomic status (SES) (8–10). Identification of potentially modifiable risk factors for PPD is crucial for prevention.

A recent systematic review of 11 longitudinal studies on night work and the risk of depression performed a meta-analysis of the 5 studies with observation periods of  $\geq 2$  years revealing some indication of an increased

<sup>1</sup> Department of Occupational and Environmental Medicine, Bispebjerg and Frederiksberg University Hospital, Copenhagen, Denmark.

<sup>2</sup> Mental Health Services, Capital Region of Denmark, Copenhagen University Hospital, Copenhagen, Denmark.

<sup>3</sup> National Research Centre for the Working Environment, Copenhagen, Denmark.

<sup>4</sup> University of Copenhagen, Department of Public Health, Copenhagen, Denmark.

<sup>5</sup> Danish Cancer Society Research Centre, Copenhagen, Denmark.

<sup>6</sup> Department of Occupational Medicine, Aarhus University Hospital, Aarhus, Denmark.

<sup>7</sup> Fertility Clinic, Rigshospitalet – Copenhagen University Hospital, Copenhagen, Denmark.

<sup>8</sup> The Parker Institute, Bispebjerg and Frederiksberg University Hospital, Copenhagen, Denmark.

Correspondence to: Paula Edeusa Cristina Hammer, Department of Occupational and Environmental Medicine, Bispebjerg and Frederiksberg University Hospital, Bispebjerg Bakke 23, indgang 20F - 2400 Copenhagen, Denmark. [E-mail: paulahammer@dadlnet.dk]

risk of depression among night workers [risk estimate 1.42, 95% confidence interval (CI) 0.92–2.19] (11). Another similar systematic review also including 11 studies used different inclusion criteria. Therefore, only 3 studies contributed to both reviews, and the 8 remaining studies were of cross-sectional design. This review found that night work was associated with an increased risk of depression (relative risk 1.43, 95% CI 1.24–1.64) across sex, night work duration, type of occupation, continent, and type of publication (12). These reviews did not inform whether PPD was included in any of the studies. As pointed out by Angerer and colleagues (11), prior studies on depression in relation to night work are challenged by cross-sectional design, self-reported exposure and outcome, and the possibility of the healthy worker survivor effect, ie, the selection of individuals with poorer health status out of night work (13, 14).

Night work is common among women in the reproductive age. In 2016, around 14% of the female European workers aged <50 years engaged in night work (15). The mechanisms linking night work to the risk of PPD are suggested to involve hormonal dysfunction, inflammation, and sleep disorders (16–25). Sleep disorders are particularly important because they add to the physiological sleep disturbances imposed by pregnancy (25).

We aimed to investigate the association of different dimensions of night work, expressed by frequency and duration of night shifts throughout pregnancy, with the risk of severe PPD. The novel aspect of our study was the use of clinically diagnosed outcome and objective information on pregnancy period-specific exposure.

## Methods

### Design

Similar to our previously published study (26), we conducted a register-based cohort study with information from the following Danish national registries linked on individual level through the civil registration number: (i) the Danish Working Hour Database (DWHD), a national payroll database covering >250 000 employees in the Danish administrative regions including all public hospital employees from January 2007 to December 2015 (27); (ii) the Danish Medical Birth Registry, which contains information from all births in Denmark since 1973 (28); (iii) the Danish National Patient Registry, which contains data on hospital admissions since 1977 and on outpatients since 1994 (29); and (iv) the Danish Psychiatric Central Research Registry, providing information on psychiatric inpatient treatment since 1969 and on all outpatient treatment since 1995 (30).

### Cohort

The source population were women from the DWHD who gave birth  $\geq 1$  between 2007 and 2015 (N=43 833 women with 70 306 births). We excluded women  $\leq 18$  and  $\geq 50$  years (N=16 pregnancies); multiple pregnancies (N=3243); pregnancies resulting in still births (N=254); pregnancies conceived in 2006 (N=6383) because they lacked payroll data from conception to January 2007; and pregnancies without  $\geq 1$  day or night shift during the first 32 pregnancy weeks (N=4954). To ensure that payroll data was available throughout the first 32 pregnancy weeks, we excluded pregnancies with employment in one of the administrative regions of Denmark, ie, registration in the DWHD, starting after conception or ending before 32 pregnancy weeks (N=25 330). Finally, we excluded pregnancies with other working schedules than fixed day shifts or  $\geq 1$  night shift during the first 32 pregnancy weeks (N=5117), for example fixed evening shifts, leaving 25 009 singleton pregnancies from 19 382 women eligible for analysis (figure 1).


### Exposure

The DWHD provided daily information on the exact time of start and end of all workdays and all types of paid and unpaid leave, job title and place of employment. Shifts during the first 32 pregnancy weeks, including on-call shifts, lasting  $\geq 3$  hours were defined as day (start time after 06:00 and end time before 21:00 hours) or night shifts (any start and end time including any duration of working hours 23:00–06:00 hours). They corresponded to 84% of all registered shifts, the rest being early morning or evening shifts. We applied the cut-off of 32 pregnancy weeks because workers in the Danish administrative regions are entitled to pregnancy leave eight weeks prior to their due date. The different dimensions of night work were expressed by the number and duration of night shifts, consecutive night shifts and quick returns as defined below.

*Number of night shifts.* The cumulated number of night shifts during the first 32 pregnancy weeks was categorized as 0 (fixed day work), 1–8, or  $\geq 9$  night shifts (roughly corresponding to  $\geq 1$  night shift per pregnancy month on average).

*Duration of night shifts.* The duration of night shifts was defined as  $\leq 8$  hours, 9–12 hours or  $>12$  hours (long night shifts). The categories used for analysis were either working only night shifts  $\leq 8$  hours or  $\geq 1$  night shift of  $>12$  hours during the first 32 pregnancy weeks.

*Consecutive night shifts.* Categories of consecutive night shifts were 1 (only single night shifts), 2–3 ( $\geq 1$  spell of



2–3 consecutive night shifts and no spells of  $\geq 4$  consecutive night shifts), or  $\geq 4$  ( $\geq 1$  spell of  $\geq 4$  consecutive night shifts) during the first 32 pregnancy weeks.

**Quick returns.** We defined quick returns as intervals of  $< 11$  hours between any type of shifts according to the European Union's Working Time Directive (31) or of  $< 28$  hours after a night shift, as suggested by a prior study using payroll data (32). The cumulated number of quick returns during the first 32 pregnancy weeks was categorized as 0, 1–8 or  $\geq 9$  quick returns (roughly corresponded to  $\geq 1$  quick return per pregnancy month on average).

**Reference group.** In all analyses, we used two reference groups, fixed day work for comparisons of night versus day work, and the lowest category of exposure for within night work comparisons.

## Outcome

The outcome severe PPD was identified by ICD-8 (296.0 – involuntal melancholia – and 300.4 – depressive neurosis) or ICD-10 (F32 – depressive episode – and F33 – recurrent depressive disorder) codes registered as the primary diagnose at somatic or psychiatric hospital departments either as in- or outpatient treatment. As

most of the women with perinatal psychiatric disorders in Denmark are treated in primary care (33), the outcome of our study was severe depression requiring in- and/or outpatient treatment at a hospital. The dates of registration were classified as prior (from 1969 to conception date) or postpartum (from birth to one year postpartum) in relation to each pregnancy.

## Covariates

Covariates were chosen a priori based on data availability and previous literature (1, 8, 34, 35).

Age ( $< 30$ , 30–35,  $> 35$  years), body mass index (BMI  $< 25$ , 25.0–29.9,  $\geq 30$  kg/m<sup>2</sup>) and parity (1, 2,  $\geq 3$ ) registered by the midwife or family doctor at the first antenatal visit derived from the Danish Medical Birth Registry. The classification of socioeconomic status (SES) into high, low or medium derived from Statistics Denmark and was based on the two Danish versions of the International Standard Classification of Occupations from 2007–2009 and 2010–2015 (ISCO-88 and ISCO-08), respectively (36, 37). Sickness absence three months prior to pregnancy (0, 1–9 or  $\geq 10$  days) expressed the sum of all days registered with  $\geq 3$  hours of sickness absence in DWHD during this period. Prior diagnosed severe depression (yes/no) derived from the Danish National Patient Registry and the Danish Psychiatric Central Research Registry as outlined above.

## Statistical analysis

We performed logistic regression of the risk of PPD by different dimensions of night work during the first 32 pregnancy weeks. We applied generalized estimating equations to account for repeated pregnancies within participants. We presented crude and adjusted estimates as odds ratio (OR) with 95% CI. Adjusted analyses included categories of age, BMI, SES, parity, sickness absence three months prior to pregnancy and prior diagnosed severe depression.

We performed the following sensitivity analyses: (i) restricted to nulliparous women; (ii) restricted to pregnancies without prior diagnosed severe depression; (iii) restricted to nurses, who represented most of the cohort; (iv) restricted to exposure in the first trimester; and (v) within-night work analysis with restriction to night work during the first or first and second trimester compared to night work throughout pregnancy. The reason for analyses (iv) and (v) was to evaluate the potential presence of healthy worker survivor effect in

the main analysis. In another words, we accounted for the possibility of women, who are somehow susceptible to PPD, stopping night shift work during pregnancy and, therefore, not contributing to the group of workers with high cumulated exposure to night work.

We used a significance level of 0.05. The analyses were done with SAS 9.4 software (SAS Institute, Cary, NC, USA).

## Results

The study cohort comprised 25 009 singleton pregnancies from 19 382 women. Their personal and working time characteristics are presented in table 1. In around 61% of pregnancies, the woman worked  $\geq 1$  night shift during pregnancy. Among those, 64% were nurses and 17% were physicians, while most of the day workers were medical secretaries (18%), nurses (15%), and physiotherapists (14%). In only 84 (0.3%) pregnan-

**Table 1.** Characteristics of 25 009 singleton pregnancies among 19 382 public hospital employees in Denmark, 2007–2015. [SD=standard deviation.]

	Day work <sup>a</sup> (N = 9642)				Night work <sup>b</sup> (N = 15 367)			
	N	%	Mean	SD	N	%	Mean	SD
Age (years)			32.3	4.0			31.2	3.8
Body mass index (kg/m <sup>2</sup> )			23.9	6.6			23.9	7.6
Parity								
1	3128	32.9			6204	40.8		
2	4402	46.2			6169	40.5		
$\geq 3$	1990	20.9			2841	18.7		
Socioeconomic status								
High	2393	25.1			3541	23.1		
Medium	5390	56.6			10 818	70.7		
Low	1741	18.3			948	6.2		
Most frequent occupations								
Laboratory technician	709	7.4			395	2.6		
Medical secretary	1785	18.5			92	0.6		
Nurse	1447	15.0			9842	64.0		
Nurse assistant	279	2.9			801	5.2		
Physician	744	7.7			2654	17.3		
Physiotherapist	1358	14.1			44	0.3		
Psychologist	683	7.1			4	0.03		
Type and number of shifts <sup>c</sup>								
Day			107.4	30.0			60.6	26.7
Early morning			0	0			0.04	0.9
Evening			0	0			13.5	13.9
Night			0	0			14.5	12.5
$\leq 8$ hours			0	0			5.7	10.3
9–12 hours			0	0			5.0	9.4
$> 12$ hours			0	0			3.8	7.0
Spells of 2–3 night shifts <sup>d</sup>			0	0			5.3	4.5
Spells of $\geq 4$ night shifts <sup>e</sup>			0	0			17.7	11.4
Quick returns			0	0			3.2	4.1
Quick returns after a night shift			0	0			7.9	8.5
Weekly working hours <sup>e</sup>			24.1	7.2			23.5	7.1
Sickness absence three months prior to pregnancy, days <sup>f</sup>			2.9	7.0			2.6	5.4

<sup>a</sup> At least one day shift and no night, evening or early morning shift during the first 32 pregnancy weeks.

<sup>b</sup> At least one night shift during the first 32 pregnancy weeks.

<sup>c</sup> During the first 32 pregnancy weeks.

<sup>d</sup> At least one spell of 2–3 and no spells of  $\geq 4$  consecutive night shifts during the first 32 pregnancy weeks.

<sup>e</sup> At least one spell of  $\geq 4$  consecutive night shifts.

<sup>f</sup> Paid and unpaid leave excluded.

cies the working schedule was fixed night shifts. Night workers had on average 14.5 night shifts during the first 32 pregnancy weeks. Around 39% of night shifts lasted  $\leq 8$  hours, 32% lasted 9–12 hours and 29%  $> 12$  hours. Women who worked  $\geq 1$  spell of  $\geq 4$  consecutive night shifts had considerably higher number of consecutive night shifts in total (mean 17.7) compared to those with spells of 2–3 and no spells of  $\geq 4$  consecutive night shifts (mean 5.3). These categories represented therefore both length of spells and total number of consecutive night shifts. Even though quick returns after a night shift (shift intervals  $< 28$  hours) accounted for the compulsory day off following a night shift, they were more frequent than quick returns between any type of shifts ( $< 11$  hours) – see table 1.

We identified 80 cases of PPD (0.3%) and 427 cases of prior diagnosed severe depression (1.7%). Approximately 20% of the women who developed PPD had a prior diagnose of severe depression, compared to 2% among women who did not develop PPD. Among women with prior diagnosed severe depression 4% developed PPD compared to 0.3% among those without a prior diagnosed severe depression.

Table 2 shows the results for the analysis of number of night shifts. Adjusted OR for PPD among women

who worked  $\geq 9$  night shifts during the first 32 pregnancy weeks was 0.59 (95% CI 0.33–1.06) compared to day work and 0.66 (95% CI 0.36–1.23) compared to having 1–8 night shifts.

The results for the analysis of duration of night shifts are presented in table 3. Adjusted OR for PPD among women who worked long night shifts ( $> 12$  hours) was 0.71 (95% CI 0.38–1.36) compared to day work and 0.76 (95% CI 0.29–1.97) compared to working only night shifts lasting  $\leq 8$  hours.

Table 4 presents the results for number of consecutive night shifts. Adjusted OR for PPD among women who worked  $\geq 4$  consecutive night shifts weeks was 0.54 (95% CI 0.22–1.30) compared to day work and 0.69 (95% CI 0.23–2.08) compared to working only single night shifts.

Table 5 presents the results for quick returns. Adjusted OR for PPD among women who worked with  $\geq 9$  quick returns was 0.75 (95% CI 0.26–2.18) compared to day work, and 0.76 (95% CI 0.24–2.38) compared to night work without quick return. Adjusted OR for PPD among pregnancies with  $\geq 9$  quick returns after a night shift was 0.57 (95% CI 0.27–1.19) compared to day work, and 0.29 (95% CI 0.11–0.72) compared to night work without quick return after a night shift.

**Table 2.** Odds ratios (OR) of postpartum depression by number of night shifts during the first 32 pregnancy weeks among 19 382 public hospital employees in Denmark, 2007–2015. [CI=confidence interval]

Number of night shifts	Pregnancies		Cases		Crude analysis			Adjusted analysis <sup>a</sup>		
	N	%	N	%	OR	95% CI	P for trend	OR	95% CI	P for trend
All pregnancies (N=25 009)										
Day work <sup>b</sup>	9642	38.5	36	0.4	1.00	Referent	0.06	1.00	Referent	0.08
1–8	5827	23.3	23	0.4	1.06	0.62–1.81		0.89	0.50–1.58	
$\geq 9$	9540	38.2	21	0.2	0.59	0.34–1.02		0.59	0.33–1.06	
Pregnancies with $\geq 1$ night shift during the first 32 pregnancy weeks (N=15 367)										
1–8	5827	37.9	23	0.4	1.00	Referent		1.00	Referent	
$\geq 9$	9540	62.1	21	0.2	0.56	0.31–1.01		0.66	0.36–1.23	

<sup>a</sup> Adjusted for age, body mass index, socioeconomic status, parity, sickness absence three months prior to pregnancy and prior diagnosed severe depression.

<sup>b</sup> At least one day shift and no night, evening or early morning shift during the first 32 pregnancy weeks.

**Table 3.** Odds ratios (OR) of postpartum depression by duration of night shifts during the first 32 pregnancy weeks among 19 382 public hospital employees in Denmark, 2007–2015. [CI=confidence interval].

Duration of night shifts	Pregnancies		Cases		Crude analysis			Adjusted analysis <sup>a</sup>		
	N	%	N	%	OR	95% CI	P for trend	OR	95% CI	P for trend
All pregnancies, N=19 125										
Day work <sup>b</sup>	9642	50.4	36	0.4	1.00	Referent	0.31	1.00	Referent	0.26
$\leq 8$ hours <sup>c</sup>	3427	17.9	10	0.3	0.78	0.38–1.59		0.74	0.35–1.55	
$> 12$ hours <sup>d</sup>	6056	31.7	17	0.3	0.75	0.42–1.35		0.71	0.38–1.36	
Pregnancies with $\geq 1$ night shift during the first 32 pregnancy weeks, N=9483										
$\leq 8$ hours	3427	36.1	10	0.3	1.00	Referent		1.00	Referent	
$> 12$ hours	6056	63.9	17	0.3	0.96	0.44–2.10		0.76	0.29–1.97	

<sup>a</sup> Adjusted for age, body mass index, socioeconomic status, parity, sickness absence three months prior to pregnancy and prior diagnosed severe depression.

<sup>b</sup> At least one day shift and no night, evening or early morning shift during the first 32 pregnancy weeks.

<sup>c</sup> Only night shifts lasting  $\leq 8$  hours.

<sup>d</sup> At least one night shift lasting  $> 12$  hours.

**Table 4.** Odds ratios (OR) of postpartum depression by consecutive night shifts during the first 32 pregnancy weeks among 19 382 public hospital employees in Denmark, 2007–2015. [CI=confidence interval]

Consecutive night shifts	Pregnancies		Cases		Crude analysis			Adjusted analysis <sup>a</sup>		
	N	%	N	%	OR	95% CI	P for trend	OR	95% CI	P for trend
All pregnancies (N=25 009)										
Day work <sup>b</sup>	9642	38.5	36	0.4	1.00	Referent	0.12	1.00	Referent	0.08
1 <sup>c</sup>	5172	20.7	18	0.4	0.93	0.52–1.66		0.93	0.49–1.77	
2–3 <sup>d</sup>	7409	29.7	20	0.3	0.72	0.41–1.26		0.63	0.35–1.16	
≥4 <sup>e</sup>	2786	11.1	6	0.2	0.58	0.24–1.38		0.54	0.22–1.30	
Pregnancies with ≥ 1 night shift during the first 32 pregnancy weeks (N=15 367)										
1	5172	33.7	18	0.4	1.00	Referent	0.26	1.00	Referent	0.50
2–3	7409	48.2	20	0.3	0.78	0.41–1.47		0.79	0.35–1.80	
≥4	2786	18.1	6	0.2	0.62	0.25–1.56		0.69	0.23–2.08	

<sup>a</sup> Adjusted age, body mass index, socioeconomic status, parity, sickness absence three months prior to pregnancy and prior diagnosed severe depression.

<sup>b</sup> At least one day shift and no night, evening or early morning shift during the first 32 pregnancy weeks.

<sup>c</sup> Only single night shifts.

<sup>d</sup> At least one spell of 2–3 consecutive night shifts and no spells of ≥ 4 consecutive night shifts.

<sup>e</sup> At least one spell of ≥ 4 consecutive night shifts.

**Table 5.** Odds ratios (OR) of postpartum depression by number of quick returns<sup>a</sup> and quick returns after a night shift<sup>b</sup> during the first 32 pregnancy weeks among 19 382 public hospital employees in Denmark, 2007–2015. [CI=confidence interval]

	Pregnancies		Cases		Crude analysis			Adjusted analysis <sup>c</sup>		
	N	%	N	%	OR	95% CI	P for trend	OR	95% CI	P for trend
Quick returns, all pregnancies (N=25 009)										
Day work <sup>d</sup>	9642	38.5	36	0.4	1.00	Referent	0.09	1.00	Referent	0.07
0 <sup>e</sup>	4605	18.4	19	0.4	1.11	0.63–1.95		1.05	0.57–1.93	
1–8	9274	37.1	21	0.2	0.61	0.35–1.05		0.54	0.30–0.96	
≥9	1488	6.0	4	0.3	0.72	0.25–2.03		0.75	0.26–2.18	
Pregnancies with ≥ 1 night shift during the first 32 pregnancy weeks (N=15 367)										
0	4605	30.0	19	0.4	1.00	Referent	0.16	1.00	Referent	0.30
1–8	9274	60.3	21	0.2	0.55	0.29–1.02		0.54	0.27–1.08	
≥9	1488	9.7	4	0.3	0.65	0.22–1.91		0.76	0.24–2.38	
Quick returns after night shift, all pregnancies (N=25 009)										
Day work	9642	38.5	36	0.4	1.00	Referent	0.06	1.00	Referent	0.04
0	1078	4.3	8	0.7	2.00	0.92–4.33		1.99	0.89–4.48	
1–8	9159	36.6	25	0.3	0.73	0.43–1.23		0.63	0.36–1.09	
≥9	5130	20.5	11	0.2	0.57	0.29–1.14		0.57	0.27–1.19	
Pregnancies with ≥ 1 night shift during the first 32 pregnancy weeks (N=15 367)										
0	1078	7.0	8	0.7	1.00	Referent	0.06	1.00	Referent	0.10
1–8	9159	59.6	25	0.3	0.37	0.16–0.81		0.29	0.13–0.66	
≥9	5130	33.4	11	0.2	0.29	0.12–0.72		0.29	0.11–0.72	

<sup>a</sup> Less than 11 hours between any type of shifts

<sup>b</sup> Less than 28 hours between a night shift and the next shift

<sup>c</sup> Adjusted for age, body mass index, socioeconomic status, parity, sickness absence three months prior to pregnancy and prior diagnosed severe depression

<sup>d</sup> At least one day shift and no night, evening or early morning shift during the first 32 pregnancy weeks

<sup>e</sup> At least one night shift and no quick returns

## Sensitivity analyses


*Analysis restricted to nulliparous women* (N=9332) did not substantially change the results from the main analyses (results not shown).

*Analysis restricted to pregnancies without prior diagnosed severe depression* (N=24 582) revealed adjusted OR 0.60 (95% CI 0.33–1.11) for women who worked ≥9 night shifts; OR 0.61 (95% CI 0.30–1.25) for women who worked long night shifts; OR 0.57 (95% CI 0.22–1.47) for women who worked ≥4 consecutive night shifts;

OR 0.42 (95% CI 0.10–1.75) for women who worked ≥9 quick returns; and OR 0.59 (95% CI 0.28–1.28) for women who worked ≥9 quick returns after a night shift.

*Analysis restricted to nurses* (N=11 298) tended to an increased risk of PPD among women in the lowest categories of exposure compared with day work (adjusted OR 1.49, 95% CI 0.44–5.04 for 1–8 night shifts; OR 1.12, 95% CI 0.31–4.05 for night shifts of ≤8 hours; OR 1.27, 95% CI 0.31–5.10 for night work without consecutive night shifts; OR 1.56, 95% CI 0.42–5.82





**Figure 2.** Odds ratios of postpartum depression by night work<sup>a</sup> during the first and second pregnancy trimester compared to night work throughout<sup>b</sup> pregnancy among 19 382 public hospital employees in Denmark, 2007–2015.

<sup>a</sup> At least one night shift.

<sup>b</sup> At least one night shift in each pregnancy trimester up to 32 pregnancy weeks.

<sup>c</sup> Adjusted for age, body mass index, socioeconomic status, parity, sickness absence three months prior to pregnancy and previously diagnosed severe depression.

for night work without quick returns; and OR 1.25, 95% CI 0.22–6.97 for night work without quick returns after a night shift).

*Analysis restricted to exposure during the first trimester* revealed overall slightly increased estimates, especially in comparisons of night versus day work – see supplementary tables ([www.sjweh.fi/show\\_abstract.php?abstract\\_id=3831](http://www.sjweh.fi/show_abstract.php?abstract_id=3831)).

*Within-night work sensitivity analysis* revealed that women who stopped working night shifts after the first trimester (N=3094) presented with an increased risk of PPD (adjusted OR 2.08 95% CI 1.09–4.00) compared to women who worked night shifts throughout pregnancy – see figure 2. The risk of PPD was not increased (adjusted OR 0.79, 95% CI 0.35–1.80) among women who stopped working night shifts after the second trimester (N=5325). Women who stopped working night shifts after the first trimester had lower proportion of high SES (14% versus 25%) but had otherwise similar demographic characteristics as women with night work throughout pregnancy. Both groups had also similar prevalence of prior diagnosed severe depression (1.8 and 1.5% respectively).

## Discussion

Overall, we did not find an increased risk of severe PPD in relation to night work during pregnancy in this nationwide cohort of Danish public hospital employees. We found, however, a 2-fold increased risk of PPD among women who stopped working night shifts after the first trimester.

To our knowledge, this is the first study investigating the association between different dimensions of night work during pregnancy and the risk of severe PPD. Our results are in line with prior longitudinal studies among hospital employees, which all found no overall association between night work and depression (38–45). Direct comparisons are not possible though as these studies applied self-reported information on depressive symptoms, not clinically diagnosed depression, and crude assessed exposure. The one study which accounted for changes of working schedules found that selection out of night work masked an effect of night work on the risk of common mental disorders (OR 1.25, 95% CI 1.03–1.52) (43). Similarly, the lack of association between PPD and the different dimensions of night work observed in our study can, at least partially, be explained by potential selection out of night work of

susceptible individuals, ie, the healthy worker survivor effect. This is supported by findings of increased risk of severe PPD among women who stopped working night shifts after the first trimester (figure 2), and slightly increased estimates in analysis restricted to exposure during the first trimester (supplementary tables). The latter analysis excluded though a potential effect of night work during the second and trimester on the risk of PPD. The different categories of night work in our study were expressed by the cumulated exposure during the first 32 pregnancy weeks. Accordingly, the women with the highest exposure were those who endured night work throughout pregnancy. These women had, presumably, a relatively higher health status. As pointed out by Nabe-Nielsen and colleagues (46), their study surprisingly shows that shift workers had better mental health than day workers “choosing night work might be a deliberate choice based on personal needs and preferences” (46). It is however important to acknowledge that we did not have information on the reason for changing working schedules among women who stopped working night shifts after the first trimester. We can only hypothesize that these women were somehow at higher risk of developing PPD. It is possible that women in our cohort, as healthcare professionals, were more aware of the occurrence of depressive symptoms than other occupational groups. Nevertheless, a study of >12 000 workers from several sectors and trades in The Netherlands has shown that workers with depressive symptoms more often changed from shift to day work (relative risk 1.98, 95% CI 1.13–3.47) and from shift work to sick leave (relative risk 2.96, 95% CI 2.00–4.29) (45). Another study using the same cohort investigated changes in mental health as predictors of changes in working time and found that workers with prolonged fatigue, need for recovery or psychological distress tended to change from shift to day work (OR 3.44, 95% CI 1.42–8.38; OR 1.36, 0.34–5.45; and OR 2.26, 0.84–6.04, respectively) (47). Finally, a Finish study of >30 000 workers from different occupations (“from semi-skilled cleaners to physicians and mayors”) found that night workers who developed common mental disorders had increased odds of moving to non-night work (OR 1.68, 95% CI 1.30–2.17) (43). One possible explanation for the association between depression and changes of working schedules is that worsening of depressive symptoms in relation to night work may be more easily recognizable than other disorders, such as hormonal dysfunction. Furthermore, adequate sleep is one of the non-pharmacological therapies in the treatment of depression (48).

It is important to identify women who work at night and who might be at risk of developing PPD, especially those who do not adapt to night work. As pregnant workers are not generally restrained from night work in Denmark, the identification of potential negative

health effects of night work during pregnancy is based on individual clinical evaluation usually by a general practitioner, an obstetrician or an occupational physician. It has been shown that women with predisposition for depression may be more susceptible to hormonal and pro-inflammatory changes during pregnancy and postpartum period (17, 18, 20, 21). It is therefore possible that these women might also be more susceptible to hormonal changes and sleep disorders (22, 23, 25) induced by night work.

It is possible that the gradient of higher exposures corresponding to higher health status is also present within night work. In our study, this hypothesis is supported by findings of an increased risk of PPD among night workers without quick returns after a night shift (table 5). Quick returns are frequently not part of planned schedules (49–51), but they might occur in periods of increased workload at the workplace. In this scenario, workers with poorer health status might not take extra shifts, avoiding quick returns. Also, analysis restricted to nurses, who represent a more homogeneous group regarding organization of night shifts, workload and job tasks, suggested elevated risks of PPD in the lowest categories of exposure to night work.

Further investigation of the healthy worker survivor effect among night workers requires specific epidemiological and statistical methods (14, 52–55). This would require information on the time of onset of depressive symptoms along with information on the reason for changing work schedules. Assessment of pregnancy period-specific exposure is also necessary as changes of working schedules occur generally during the end of the first trimester, when many women, at least in Denmark, inform their workplace about their pregnancy. Methods accounting for change of working schedules after the occurrence of disease, such as performed by Beltagy and colleagues (43), are suitable for accounting for changes of working schedule in relation to the occurrence of depressive symptoms during pregnancy. However, they are not applicable to PPD as it occurs per definition after birth, and therefore most probably when the woman is on maternity leave.

### Strengths and limitations

The primary strengths of our study include nationwide sample size, prospective design, detailed and register-based exposure and outcome assessment, and within-night work comparisons.

Even though our cohort is nationwide, our results are based on relatively few cases as only the most severe cases of PPD are treated in hospitals in Denmark. As shown by Munk-Olsen and colleagues (33), for each birth resulting in inpatient psychiatric treatment of postpartum disorder in Denmark, 2.5 births

resulted in outpatient treatment and 12 births resulted in general practitioner-provided pharmacological treatment. Therefore, even though the prevalence of severe PPD in our cohort (0.3%) corresponds to prior Danish population-based studies (33), low power is a limitation of our study. Misclassification of the mild cases of PPD might be an important source of bias towards the null in our cohort. The extent of this bias depends on whether night work is a risk factor specifically for mild cases of PPD. This is yet to be elucidated. Furthermore, it is possible that the low prevalence of severe PPD we observed among nurses and physicians might also reflect the possibility of receiving informal care from the work environment when having depressive symptoms. Nevertheless, including only severe cases of PPD increased the specificity of the outcome.

We lacked information on how well the participants adapted to night work. Psychological resilience (hardiness) and coping seem to be strongly related to psychological well-being (56). Further, we lacked information on chronotype. The evening chronotype, which is normally more frequent among night than day workers, has been independently associated with the risk and severity of depression (57). This might have biased the results towards the null if workers with evening chronotype, rather than other chronotypes, tended to change working schedules to non-night work due to depressive symptoms.

### Concluding remarks

Overall, our results do not suggest that night work during pregnancy is a risk factor for severe PPD. However, we observed a 2-fold increased risk of PPD among women who stopped working night shifts after the first pregnancy trimester. This may reflect the influence of the healthy worker survivor effect, but the interpretation of this finding requires caution and warrants further attention.

Future research should include cases of PPD not requiring hospital treatment, use pregnancy period-specific exposure assessment, and apply epidemiological and statistical methods that account for the healthy worker survivor effect.

### Acknowledgements

The Danish administrative regions are acknowledged for the participation and willingness to provide data to the Danish Working Hour Database. Jens Worm Begtrup, Lisbeth Nielsen and Anders Ørberg are thanked for valuable work with data management. Statistics Denmark is acknowledged for ensuring appropriate anonymization

of the data in accordance to the current Danish Data Protection legislation.

### Funding

This work is financially supported by the Danish Working Environment Research Fund grant 31-2015-03 2015001705. The establishment of the Danish Working Hour Database has been financed by research grants from the Danish Working Environment Research Fund (grant 23-2012-09), the Nordic Program of Health and Welfare – Nordforsk (grant 74809), and the National Research Center for the Working Environment. The Danish administrative regions have partially financed the transfer of the data to the cohort.

### References

1. Meltzer-Brody S, Howard LM, Bergink V, Vigod S, Jones I, Munk-Olsen T et al. Postpartum psychiatric disorders. *Nat Rev Dis Primers* 2018 Apr;4:18022. <https://doi.org/10.1038/nrdp.2018.22>.
2. Becker M, Weinberger T, Chandy A, Schumker S. Depression During Pregnancy and Postpartum. *Curr Psychiatry Rep* 2016 Mar;18(3):32. <https://doi.org/10.1007/s11920-016-0664-7>.
3. Orsolini L, Valchera A, Vecchiotti R, Tomasetti C, Iasevoli F, Fornaro M et al. Suicide during Perinatal Period: Epidemiology, Risk Factors, and Clinical Correlates. *Front Psychiatry* 2016 Aug;7:138. <https://doi.org/10.3389/fpsy.2016.00138>.
4. Khalifeh H, Hunt IM, Appleby L, Howard LM. Suicide in perinatal and non-perinatal women in contact with psychiatric services: 15 year findings from a UK national inquiry. *Lancet Psychiatry* 2016 Mar;3(3):233–42. [https://doi.org/10.1016/S2215-0366\(16\)00003-1](https://doi.org/10.1016/S2215-0366(16)00003-1).
5. Almond P. Postnatal depression: a global public health perspective. *Perspect Public Health* 2009 Sep;129(5):221–7. <https://doi.org/10.1177/1757913909343882>.
6. Meltzer-Brody S. New insights into perinatal depression: pathogenesis and treatment during pregnancy and postpartum. *Dialogues Clin Neurosci* 2011;13(1):89–100.
7. Bauer AE, Maegbaek ML, Liu X, Wray NR, Sullivan PF, Miller WC et al. Familiarity of Psychiatric Disorders and Risk of Postpartum Psychiatric Episodes: A Population-Based Cohort Study. *Am J Psychiatry* 2018 Aug;175(8):783–91. <https://doi.org/10.1176/appi.ajp.2018.17111184>.
8. Guintivano J, Sullivan PF, Stuebe AM, Penders T, Thorp J, Rubinow DR et al. Adverse life events, psychiatric history, and biological predictors of postpartum depression in an ethnically diverse sample of postpartum women. *Psychol Med* 2018 May;48(7):1190–200. <https://doi.org/10.1017/S0033291717002641>.

9. Meltzer-Brody S, Larsen JT, Petersen L, Guintivano J, Florio AD, Miller WC et al. Adverse life events increase risk for postpartum psychiatric episodes: A population-based epidemiologic study. *Depress Anxiety* 2018 Feb;35(2):160–7. <https://doi.org/10.1002/da.22697>.
10. Clare CA, Yeh J. Postpartum depression in special populations: a review. *Obstet Gynecol Surv* 2012 May;67(5):313–23. <https://doi.org/10.1097/OGX.0b013e318259cb52>.
11. Angerer P, Schmoock R, Elfantel I, Li J. Night Work and the Risk of Depression. *Dtsch Arztebl Int* 2017 Jul;114(24):404–11.
12. Lee A, Myung SK, Cho JJ, Jung YJ, Yoon JL, Kim MY. Night Shift Work and Risk of Depression: Meta-analysis of Observational Studies. *J Korean Med Sci* 2017 Jul;32(7):1091–6. <https://doi.org/10.3346/jkms.2017.32.7.1091>.
13. Pearce N, Checkoway H, Kriebel D. Bias in occupational epidemiology studies. *Occup Environ Med* 2007 Aug;64(8):562–8. <https://doi.org/10.1136/oem.2006.026690>.
14. Chowdhury R, Shah D, Payal AR. Healthy Worker Effect Phenomenon: Revisited with Emphasis on Statistical Methods - A Review. *Indian J Occup Environ Med* 2017 Jan-Apr;21(1):2–8. [https://doi.org/10.4103/ijoem.IJOEM\\_53\\_16](https://doi.org/10.4103/ijoem.IJOEM_53_16).
15. Parent-Thirion A, Biletta I, Cabrita J, Vargas O, Vermeulen G, Wilczynska A et al. Eurofond (2016), Sixth European Working Conditions Survey - Overview report. Luxembourg; 2016.
16. Dmitrzak-Weglarz M, Reszka E. Pathophysiology of Depression: Molecular Regulation of Melatonin Homeostasis - Current Status. *Neuropsychobiology* 2017;76(3):117–29. <https://doi.org/10.1159/000489470>.
17. Seth S, Lewis AJ, Galbally M. Perinatal maternal depression and cortisol function in pregnancy and the postpartum period: a systematic literature review. *BMC Pregnancy Childbirth* 2016 May;16(1):124. <https://doi.org/10.1186/s12884-016-0915-y>.
18. Brummelte S, Galea LA. Depression during pregnancy and postpartum: contribution of stress and ovarian hormones. *Prog Neuropsychopharmacol Biol Psychiatry* 2010 Jun;34(5):766–76. <https://doi.org/10.1016/j.pnpbp.2009.09.006>.
19. Parry BL, Meliska CJ, Sorenson DL, Lopez AM, Martinez LF, Nowakowski S et al. Plasma melatonin circadian rhythm disturbances during pregnancy and postpartum in depressed women and women with personal or family histories of depression. *Am J Psychiatry* 2008 Dec;165(12):1551–8. <https://doi.org/10.1176/appi.ajp.2008.08050709>.
20. Osborne LM, Monk C. Perinatal depression--the fourth inflammatory morbidity of pregnancy?: theory and literature review. *Psychoneuroendocrinology* 2013 Oct;38(10):1929–52. <https://doi.org/10.1016/j.psyneuen.2013.03.019>.
21. Chang JJ, Pien GW, Duntley SP, Macones GA. Sleep deprivation during pregnancy and maternal and fetal outcomes: is there a relationship? *Sleep Med Rev* 2010 Apr;14(2):107–14. <https://doi.org/10.1016/j.smr.2009.05.001>.
22. Okun ML. Disturbed Sleep and Postpartum Depression. *Curr Psychiatry Rep* 2016 Jul;18(7):66. <https://doi.org/10.1007/s11920-016-0705-2>.
23. Lawson A, Murphy KE, Sloan E, Uleryk E, Dalfen A. The relationship between sleep and postpartum mental disorders: A systematic review. *J Affect Disord* 2015 May;176:65–77. <https://doi.org/10.1016/j.jad.2015.01.017>.
24. Bhati S, Richards K. A systematic review of the relationship between postpartum sleep disturbance and postpartum depression. *J Obstet Gynecol Neonatal Nurs* 2015 May-Jun;44(3):350–7. <https://doi.org/10.1111/1552-6909.12562>.
25. Bei B, Coo S, Trinder J. Sleep and Mood During Pregnancy and the Postpartum Period. *Sleep Med Clin* 2015 Mar;10(1):25–33. <https://doi.org/10.1016/j.jsmc.2014.11.011>.
26. Hammer P, Flachs E, Specht I, Pinborg A, Petersen S, Larsen A et al. Night work and hypertensive disorders of pregnancy: a national register-based cohort study. *Scand J Work Environ Health* 2018 Jul;44(4):403–13. <https://doi.org/10.5271/sjweh.3728>.
27. Garde AH, Hansen J, Kolstad HA, Larsen AD, Pedersen J, Petersen JD et al. Payroll data based description of working hours in the Danish regions. *Chronobiol Int* 2018 Jun;35(6):795–800. <https://doi.org/10.1080/07420528.2018.1466797>.
28. Knudsen LB, Olsen J. The Danish Medical Birth Registry. *Dan Med Bull* 1998 Jun;45(3):320–3.
29. Schmidt M, Schmidt SA, Sandegaard JL, Ehrenstein V, Pedersen L, Sørensen HT. The Danish National Patient Registry: a review of content, data quality, and research potential. *Clin Epidemiol* 2015 Nov;7:449–90. <https://doi.org/10.2147/CLEP.S91125>.
30. Mors O, Perto GP, Mortensen PB. The Danish Psychiatric Central Research Register. *Scand J Public Health* 2011 Jul;39(7 Suppl):54–7. <https://doi.org/10.1177/1403494810395825>.
31. European Union's Working Time Directive (2003/88/EC) of the European Parliament and Council. 2003.
32. Härmä M, Ropponen A, Hakola T, Koskinen A, Vanttola P, Puttonen S et al. Developing register-based measures for assessment of working time patterns for epidemiologic studies. *Scand J Work Environ Health* 2015 May;41(3):268–79. <https://doi.org/10.5271/sjweh.3492>.
33. Munk-Olsen T, Maegbaek ML, Johannsen BM, Liu X, Howard LM, di Florio A et al. Perinatal psychiatric episodes: a population-based study on treatment incidence and prevalence. *Transl Psychiatry* 2016 Oct;6(10):e1919. <https://doi.org/10.1038/tp.2016.190>.
34. Jung SJ, Woo HT, Cho S, Park K, Jeong S, Lee YJ et al. Association between body size, weight change and depression: systematic review and meta-analysis. *Br J Psychiatry* 2017 Jul;211(1):14–21. <https://doi.org/10.1192/bjp.bp.116.186726>.

35. Jantaratnotai N, Mosikanon K, Lee Y, McIntyre RS. The interface of depression and obesity. *Obes Res Clin Pract* 2017 Jan - Feb;11(1):1–10. <https://doi.org/10.1016/j.orcp.2016.07.003>.
36. Statistics Denmark DISCO-88 - Danmarks Statistiks Fagklassifikation. 1996.
37. Statistics Denmark DISCO-08 - Danmark Statistiks Fagklassifikation. 2011.
38. Bohle P, Tilley AJ. The impact of night work on psychological well-being. *Ergonomics* 1989 Sep;32(9):1089–99. <https://doi.org/10.1080/00140138908966876>.
39. Lin PC, Chen CH, Pan SM, Pan CH, Chen CJ, Chen YM et al. Atypical work schedules are associated with poor sleep quality and mental health in Taiwan female nurses. *Int Arch Occup Environ Health* 2012 Nov;85(8):877–84. <https://doi.org/10.1007/s00420-011-0730-8>.
40. Thun E, Bjorvatn B, Torsheim T, Moen BE, Mageroy N, Pallesen S. Night work and symptoms of anxiety and depression among nurses: a longitudinal study. *Work Stress* 2014;28:376–86. <https://doi.org/10.1080/02678373.2014.969362>.
41. Flo E, Pallesen S, Moen BE, Waage S, Bjorvatn B. Short rest periods between work shifts predict sleep and health problems in nurses at 1-year follow-up. *Occup Environ Med* 2014 Aug;71(8):555–61. <https://doi.org/10.1136/oemed-2013-102007>.
42. Berthelsen M, Pallesen S, Mageroy N, Tyssen R, Bjorvatn B, Moen BE et al. Effects of Psychological and Social Factors in Shiftwork on Symptoms of Anxiety and Depression in Nurses: A 1-Year Follow-Up. *J Occup Environ Med* 2015 Oct;57(10):1127–37. <https://doi.org/10.1097/JOM.0000000000000532>.
43. Beltagy MS, Pentti J, Vahtera J, Kivimäki M. Night work and risk of common mental disorders: analyzing observational data as a non-randomized pseudo trial. *Scand J Work Environ Health* 2018 Sep;44(5):512–20. <https://doi.org/10.5271/sjweh.3733>.
44. Driesen K, Jansen NW, Kant I, Mohren DC, van Amelsvoort LG. Depressed mood in the working population: associations with work schedules and working hours. *Chronobiol Int* 2010 Jul;27(5):1062–79. <https://doi.org/10.3109/07420528.2010.489877>.
45. Driesen K, Jansen NW, van Amelsvoort LG, Kant I. The mutual relationship between shift work and depressive complaints--a prospective cohort study. *Scand J Work Environ Health* 2011 Sep;37(5):402–10. <https://doi.org/10.5271/sjweh.3158>.
46. Nabe-Nielsen K, Garde AH, Albertsen K, Diderichsen F. The moderating effect of work-time influence on the effect of shift work: a prospective cohort study. *Int Arch Occup Environ Health* 2011 Jun;84(5):551–9. <https://doi.org/10.1007/s00420-010-0592-5>.
47. De Raeve L, Kant I, Jansen NW, Vasse RM, van den Brandt PA. Changes in mental health as a predictor of changes in working time arrangements and occupational mobility: results from a prospective cohort study. *J Psychosom Res* 2009 Feb;66(2):137–45. <https://doi.org/10.1016/j.jpsychores.2008.05.007>.
48. Fang H, Tu S, Sheng J, Shao A. Depression in sleep disturbance: A review on a bidirectional relationship, mechanisms and treatment. *J Cell Mol Med* 2019 Apr;23(4):2324–32. <https://doi.org/10.1111/jcmm.14170>.
49. Overenskomst for underordnede læger [Work Union Agreement for subordinate physicians] 2015 [Available from: <https://www.laeger.dk/sygehusoverenskomsten>].
50. Overenskomst for overlæger [Work Union Agreement for chief physicians]. 2015.
51. Overenskomst for ikke-ledende personale på sundhedskartellets område [Work Union Agreement for subordinate health care professionals]. 2016.
52. Li CY, Sung FC. A review of the healthy worker effect in occupational epidemiology. *Occup Med (Lond)* 1999 May;49(4):225–9. <https://doi.org/10.1093/occmed/49.4.225>.
53. Naimi AI, Richardson DB, Cole SR. Causal inference in occupational epidemiology: accounting for the healthy worker effect by using structural nested models. *Am J Epidemiol* 2013 Dec;178(12):1681–6. <https://doi.org/10.1093/aje/kwt215>.
54. Buckley JP, Keil AP, McGrath LJ, Edwards JK. Evolving methods for inference in the presence of healthy worker survivor bias. *Epidemiology* 2015 Mar;26(2):204–12. <https://doi.org/10.1097/EDE.0000000000000217>.
55. Joffe MM. Structural nested models, g-estimation, and the healthy worker effect: the promise (mostly unrealized) and the pitfalls. *Epidemiology* 2012 Mar;23(2):220–2. <https://doi.org/10.1097/EDE.0b013e318245f798>.
56. Tahghighi M, Rees CS, Brown JA, Breen LJ, Hegney D. What is the impact of shift work on the psychological functioning and resilience of nurses? An integrative review. *J Adv Nurs* 2017 Sep;73(9):2065–83. <https://doi.org/10.1111/jan.13283>.
57. Taylor BJ, Hasler BP. Chronotype and Mental Health: recent Advances. *Curr Psychiatry Rep* 2018 Jul;20(8):59. <https://doi.org/10.1007/s11920-018-0925-8>.

Received for publication: 8 November 2018