



Biotic and abiotic factors affecting phorophyte preference in epiphytic orchids: a pilot study from southern China

Pecoraro, Lorenzo; Rasmussen, Hanne Nina; Rasmussen, Finn N.; Gomes, Sofia I.F.;
Merckx, Vincent S.F.T.; Cai, Lei

Publication date:
2019

Document version
Publisher's PDF, also known as Version of record

Document license:
[Unspecified](#)

Citation for published version (APA):
Pecoraro, L., Rasmussen, H. N., Rasmussen, F. N., Gomes, S. I. F., Merckx, V. S. F. T., & Cai, L. (2019). *Biotic and abiotic factors affecting phorophyte preference in epiphytic orchids: a pilot study from southern China*. 114. Abstract from International Orchid Conservation Congress, London, United Kingdom.

Saturday 1 June, 14:30 - 14:45 (parallel session)

Biotic and abiotic factors affecting phorophyte preference in epiphytic orchids: a pilot study from southern China.

Lorenzo Pecoraro^{1,2}, Hanne N. Rasmussen³, Finn N. Rasmussen⁴, Sofia I.F. Gomes⁵, Vincent S.F.T. Merckx⁵ & Lei Cai²

¹School of Pharmaceutical Science and Technology, Health Sciences Platform, Tianjin University, Tianjin, China ²State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China ³Institute for Geoscience and Nature Management, University of Copenhagen, Denmark ⁴Natural History Museum of Denmark, University of Copenhagen, Denmark ⁵Naturalis Biodiversity Center, Leiden, The Netherlands

Epiphytic orchids exhibit varying degrees of phorophyte tree specificity. Some orchid species show a strong phorophyte preference, while other species associate with a broad range of tree hosts. Although trees that carry epiphytic orchids play a crucial role in orchid life cycle, little is known about this phorophyte-epiphyte relationship. We performed a pilot study to investigate why epiphytic orchids prefer or avoid certain phorophyte trees. We selected two orchid species, *Panisea uniflora* and *Bulbophyllum odoratissimum* co-occurring in a forest habitat in southern China. *Panisea uniflora* was found to grow on *Quercus yiwuensis*, but avoided *Pistacia weinmannifolia* trees. The latter phorophyte host carried *B. odoratissimum* that was never found to colonize *Q. yiwuensis*. We also selected *Beilschmiedia percoriacea* as a neutral tree, carrying several epiphytes, but neither *B. odoratissimum* nor *P. uniflora*. We analysed a number of biotic and abiotic environmental factors potentially influencing the relationship between studied orchids and trees, including bark and orchid root fungal diversity. We hypothesised that the tree host specialization of the studied orchids was influenced by the presence of orchid mycorrhizal fungi, which were in turn biased toward particular tree species. Both morphological and molecular culture-based methods, combined with metagenomic analyses, were used to assess fungal communities associated with studied orchids and trees. A total of 232 fungal species in 72 genera were isolated from bark samples. Ascomycetous fungi belonging to *Cladosporium*, *Cyphellophora*, *Fusicolla*, *Penicillium*, *Pestalotiopsis* and *Trichoderma* were dominant. The highest fungal diversity was observed on neutral trees with 71 taxa, followed by *P. weinmannifolia* trees with 47, and *Q. yiwuensis* with 31. Metagenomic analysis confirmed the presence of significantly different fungal communities on the three investigated tree species. Different fungi were also found associated with the two analysed orchid species. Fungal diversity, influenced by bark features, may affect tree selection by epiphytic orchids.

Keywords: epiphytic orchids, phorophyte trees, fungal communities, orchid mycorrhizal fungi, bark.