



Advances in legume research in the genomics era

Egan, Ashley N.; Vatanparast, Mohammad

Published in:
Australian Systematic Botany

DOI:
[10.1071/SB19019](https://doi.org/10.1071/SB19019)

Publication date:
2019

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY-NC-ND](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Citation for published version (APA):
Egan, A. N., & Vatanparast, M. (2019). Advances in legume research in the genomics era. *Australian Systematic Botany*, 32(5-6), 459-483. <https://doi.org/10.1071/SB19019>

Advances in legume research in the genomics era

Ashley N. Egan ^{A,D} and Mohammad Vatanparast^{B,C}

^ADepartment of Bioscience, Aarhus University, Ny Munkegade 116, DK-8000 Aarhus C, Denmark.

^BDepartment of Geosciences and Natural Resource Management, University of Copenhagen, DK- 1958 Frederiksberg C, Denmark.

^CDepartment of Botany, National Museum of Natural History, Smithsonian Institution, Washington, DC 20560, USA.

^DPresent address, Department of Biology, Utah Valley University, 800 W University Parkway, Orem, UT 84058, USA. Corresponding author. Email: aegan@uvu.edu

Abstract. Next-generation sequencing (NGS) technologies and applications have enabled numerous critical advances in legume biology, from marker discovery to whole-genome sequencing, and will provide many new avenues for legume research in the future. The past 6 years in particular have seen revolutionary advances in legume science because of the use of high-throughput sequencing, including the development of numerous types of markers and data useful for evolutionary studies above and below the species level that have enabled resolution of relationships that were previously unattainable. Such resolution, in turn, affords opportunities for hypothesis testing and inference to improve our understanding of legume biodiversity and the patterns and processes that have created one of the most diverse plant families on earth. In addition, the genomics era has seen significant advances in our understanding of the ecology of legumes, including their role as nitrogen fixers in global ecosystems. The accumulation of genetic and genomic data in the form of sequenced genomes and gene-expression profiles made possible through NGS platforms has also vastly affected plant-breeding and conservation efforts. Here, we summarise the knowledge gains enabled by NGS methods in legume biology from the perspectives of evolution, ecology, and development of genetic and genomic resources.

Additional keywords: crop genomes, Fabaceae, genome-wide research, Leguminosae, next-generation sequencing, phylogenomics, RADseq, sequence capture, target enrichment.

Received 4 March 2019, accepted 16 July 2019, published online 30 September 2019

Introduction

In the late 20th century, Sanger sequencing (Sanger *et al.* 1977) transformed biology and medicine, enabling many genetic advances, the greatest being completion of the human genome project (International Human Genome Sequencing Consortium 2001). Three decades later, second- or next-generation sequencing (NGS) ushered in the genomics era, producing massive amounts of sequence data at a fraction of the cost and time. For comparison, the cost of sequencing a human genome in September of 2001 by Sanger sequencing was ~US\$95 million; today sequencing the same genome will cost less than US\$1500 (The Human Genome Research Institute; see <https://www.genome.gov/sequencingcosts/>, accessed 28 August 2019).

Numerous NGS methods have been introduced, each with strengths and weaknesses (for review, see Egan *et al.* 2012; Soltis *et al.* 2013; Reuter *et al.* 2015). NGS technologies provide unprecedented opportunities in fields such as crop genomics, molecular systematics, evolutionary genomics or plant breeding, prompting scientific understanding across the tree of life. In particular, plant biology has blossomed through NGS

applications, including transcriptomics (e.g. Matasci *et al.* 2014; Wen *et al.* 2015), phylogenomics (Ruhfel *et al.* 2014; Wickett *et al.* 2014; Soltis *et al.* 2018), genome-wide single-nucleotide polymorphism (SNP) sequencing by genome-reduction techniques (Andrews *et al.* 2016; Jiang *et al.* 2016) and whole-genome sequencing (Zhang *et al.* 2011; Koboldt *et al.* 2013).

Leguminosae (Fabaceae) is the third-largest family of flowering plants after Orchidaceae and Asteraceae (Lewis *et al.* 2005), comprising ~770 genera and ~19500 species (Legume Phylogeny Working Group 2013a, 2017). Our understanding of the classification and evolutionary relationships within the family has been transformed in recent years, with the impact of NGS methods as summarised by Doyle (2013). The family was recently reclassified from the classical three subfamilies (Caesalpinioideae, Mimosoideae, Papilionoideae) into six subfamilies, corresponding to the six main clades, on the basis of an international effort involving ~100 scientists, using molecular systematics (Fig. 1; Legume Phylogeny Working Group 2013b, 2017). Fabaceae is




Fig. 1. Numbers of species and genera partitioned across the six Leguminosae subfamilies. Subfamily Duparquetioideae consists of a single monotypic genus and the single species of *Duparquetia* is not distinguishable in the species circle.

distributed in all of the world's vegetation types (biomes) except polar ice. Legumes are second only to grasses in economic importance, with uses common to nearly all facets of life, including food, medicine, oils, timber, fibres, industry, fodder, soil stabilisation and soil enrichment (Graham and Vance 2003). Legume research has benefited from NGS innovations, with studies employing genomic sequencing techniques to address questions in evolution, ecology, conservation and plant breeding. The sequencing of legume genomes, both nuclear and plastid, along with transcriptomic and other genomic profiles, has greatly improved genetic and genomic data resources, providing foundations on which researchers can build. This review is wide in scope, both in terms of genomic methods, as well as their impacts on legume subdisciplines, from systematics to plant breeding. This review expands on that of Doyle (2013), focusing especially on work published during the intervening 6 years. We summarise recent insights across selected topics and review ongoing research regarding NGS technologies and how they are contributing to our understanding of legumes.

Evolutionary aspects

Given the incredible diversity and vast ecological and economic importance of legumes, understanding the patterns and processes underlying the evolution of the legume family is an important endeavour. Robust inferences of phylogenetic trees are fundamental to any subsequent analyses, above or below the species level. In this section, we summarise knowledge gains in legume evolutionary biology afforded by genomic advances.

Phylogenetics v. Phylogenomics

Investigation of evolutionary relationships among plants by using molecular data began in the late 1980s. Pioneering legume studies included size polymorphism of chloroplast DNA in *Pisum* L. (Palmer *et al.* 1985) and nuclear rDNA (rDNA) repeat-length and restriction-enzyme site locations in soybean and relatives (Doyle and Beachy 1985). Sanger sequencing revolutionised phylogenetics through ease of use

and reproducibility (Sanger *et al.* 1977). The chloroplast gene *matK* is the most comprehensively sequenced phylogenetic marker in legumes. Wojciechowski *et al.* (2004) produced one of the earliest generic-level phylogenies of legumes by using 330 *matK* gene sequences, demonstrating monophyly of Papilionoideae and resolving multiple papilionoid subclades. Many subsequent studies have used *matK* (e.g. Bruneau *et al.* 2008; Simon *et al.* 2009; Stefanovic *et al.* 2009; Cardoso *et al.* 2012; de Queiroz *et al.* 2015; Egan *et al.* 2016; Snak *et al.* 2016), culminating in the Legume Phylogeny Working Group (2017) phylogeny that included 3842 *matK* sequences representing 3696 species (~20% of the family) and 698 of the 765 genera that had been recognised. Whereas *matK* robustly supported each of the newly recognised six subfamilies, basal nodes key to understanding subfamilial relationships and early legume evolution remained unresolved, which was likely because of the lack of resolving power available from a single marker.

The use of nuclear markers for legume phylogenetics has lagged behind chloroplast markers, largely because of the biparental inheritance and the high incidence of gene duplications in the nuclear genome, which make orthology assessment and primer design difficult (Zimmer and Wen 2013). The advent of genomics has transformed and facilitated nuclear marker discovery (Zimmer and Wen 2015). For example, Scherson *et al.* (2005) screened several nuclear loci from the *Medicago truncatula* Gaertn. genome for phylogeny reconstruction in the hyper-diverse legume genus *Astragalus* L. Similarly, Choi *et al.* (2006) tested 274 putative single-copy genes garnered from comparative analysis of 15 legume genomes from six species and identified 129 single-copy loci that were tested across 95 legume species.

As helpful as NGS methods are for improving nuclear-marker discovery, nuclear loci that are putatively single-copy in one lineage may not be so in others, a phenomenon that makes finding the 'silver bullet' of nuclear markers difficult in plants (e.g. Manzanilla and Bruneau 2012), and which argues for having many different markers to mitigate issues with a few. This is where the true utility of NGS technologies comes in. The move from single amplicon-based Sanger sequencing to NGS-based, simultaneous sequencing of numerous markers is rapidly transforming our understanding of legume evolution (Doyle 2013), not least through development of a variety of new approaches for NGS marker data for phylogenetic and population-genetic studies. These include microsatellites, RNAseq or transcriptomics (Wang *et al.* 2009; Wen *et al.* 2015), restriction site-associated DNA tags (RADseq; Miller *et al.* 2007), genotyping-by-sequencing (GBS; Elshire *et al.* 2011), genome skimming (Straub *et al.* 2012), targeted enrichment (also known as sequence capture or hybrid enrichment; Gnirke *et al.* 2009), simultaneous amplicon sequencing (Bybee *et al.* 2011), and whole-genome sequencing (e.g. Stein *et al.* 2018; Grover *et al.* 2019).

Studies have capitalised on massively parallel sequencing for discovery and optimisation of microsatellite markers for population studies, for example, using transcriptome sequencing, including in legumes (e.g. Chapman 2015; Vatanparast *et al.* 2016; Sathyanarayana *et al.* 2017; Haynsen *et al.* 2018). Other popular NGS methods for population genetics include RADseq and GBS, which are also employed in

phylogenetics below the genus level. For instance, Wong *et al.* (2015) used GBS across 60 accessions of seven species of *Lens* and produced a phylogeny in which all seven species were reciprocally monophyletic. Grillo *et al.* (2016) employed RADseq across 191 accessions of *Medicago truncatula* in Europe to investigate population structure and screen for candidate symbiosis genes. They found evidence that suggests that one gene, *DM11*, is under adaptive selection. Their work detailed five distinct genetic clusters and aided in correct identification of species. That said, work by others has found that different parameter perturbations produced significant differences in phylogenetic networks of species relationships within *Medicago* by using RADseq (Blanco-Pastor *et al.* 2018), suggesting caution when using population-level markers above species level.

RNAseq is often used as the first line for phylogenetic-marker development, whether for microsatellite development, amplicon sequencing (e.g. Chapman 2015) or targeted enrichment. For example, Vatanparast *et al.* (2018) used 30 transcriptomes across the legume family to select over 500 nuclear markers for targeted enrichment, which were tested across 25 legume taxa. This same target set proved useful as far out as Rosales (M. Vatanparast and A.N. Egan, unpubl. data), showing that targeted enrichment can be useful for both lower- and higher-level phylogenomics, as has been found by others (Kadlec *et al.* 2017; Chau *et al.* 2018).

The first targeted enrichment-gene set in legumes selected 50 nuclear loci from the *Medicago* genome to resolve relationships among six *Medicago* species and other genera of tribe Trifolieae (de Sousa *et al.* 2014). Targeted enrichment may be especially useful for resolving relationships among rapidly radiated, species-rich groups by virtue of the large amount of data produced. For example, Nicholls *et al.* (2015) applied a targeted enrichment method using transcriptomes from three species to isolate 264 nuclear loci for sequencing *Inga* Mill., a genus of ~300 neotropical rainforest trees that diversified rapidly during the late Miocene (2–10 million years ago). Of these loci, 194 were used for phylogeny reconstruction across 22 *Inga* species, resulting in a highly resolved phylogenetic tree. Similarly, Ojeda *et al.* (2019) used 289 nuclear loci identified from transcriptomes across Detarioideae to construct a phylogenomic hypothesis of relationships within the *Anthonotha* clade, discovering an overall general trend towards petal reduction in this florally diverse group. In addition to the target-gene sets outlined here, several other, as yet unpublished, gene sets have been generated for legumes.

As legume researchers adopt NGS for phylogenomics, it could be worth considering selection and adoption of a core set of nuclear target genes for use across legumes, along lines similar to the 353 nuclear-gene set for targeted enrichment in angiosperms (Johnson *et al.* 2019). Doing so would also facilitate barcoding efforts (Hollingsworth *et al.* 2016). With the recent addition of the first genome sequences for mimosoids and cercidoids (Table 1, Fig. 2), plus a significant number of new transcriptomes, it would now be possible to design a legume-wide, legume-specific bait set more efficiently. Such a design might include genes specifically selected for phylogenetics, as well as genes that are related to particular legume functional traits such as nodulation, compound-leaf development or floral symmetry, as has been done for Caryophyllales (Moore *et al.*

2018). Whereas the number of target gene sets used in legumes will undoubtedly increase, including a core subset of genes in every target-gene set would enable published sequences from different studies to be combined for wider analyses. Doing so would foster wide collaboration among legume systematists, while still enabling project-specific objectives to be met. That said, issues with the conflation of orthology and paralogy need to be dealt with when using universal gene sets because of whole-genome duplications (WGDs) and differential gene birth and death events, issues that are particularly troubling if targets are used for DNA barcoding.

Species diversity

Understanding the dynamics of diversification within the third-largest species-rich plant family is complex and only a few attempts have been made to estimate species diversification and speciation–extinction rates within legumes (e.g. Sanderson and Wojciechowski 1996; Richardson *et al.* 2001; Scherson *et al.* 2008). Koenen *et al.* (2013) estimated species diversification rates and tested for rate shifts by using species-level, time-calibrated phylogenies of the following four legume clades: *Calliandra* Benth., Indigofereae, *Lupinus* L. and *Mimosa* L., finding evidence for significant among-lineage diversification-rate variation. A major challenge for diversification studies is having a well resolved phylogeny, a non-trivial task when dealing with rapid, recent radiations (Hughes *et al.* 2015). The most species-rich legume tribe, Galegeae, includes *Astragalus*, the largest genus of any biological group, with nearly 3000 species (Kazempour Osaloo *et al.* 2003; Podlech *et al.* 2014), as well as *Oxytropis* DC., its sister genus with between 310 and 450 species (Malyshev 2008). Recent molecular studies of these groups exemplify the difficulties of resolving phylogenies for rapidly radiating lineages (e.g. Azani *et al.* 2017; Bagheri *et al.* 2017). Where phylogenies based on data from single or a few genes lack resolution, genomic-scale data may provide a solution. Application of anchored enrichment of 527 gene regions (i.e. targeted enrichment) in *Oxytropis* produced a robust phylogeny compared to relying on conventional markers (Shahi Shavvon *et al.* 2017).

Next-generation sequencing data provide more than just phylogenetic resolution and their application to elucidating adaptive radiations and diversification is just beginning. This is exemplified in ground-breaking studies on *Lupinus*, a genus of ~280 species, with a series of nested and parallel rapid radiations in North and South America (Hughes and Eastwood 2006) where the genus exhibits startling diversity in growth form and habitat in the Andes, a radiation that is likely to have been spurred by Pleistocene glacial cycles (Nevado *et al.* 2018). Furthermore, lineage-specific diversification rates were detected across the phylogeny, but the ability to ascertain the processes underpinning rapid species radiations remained elusive. Today, investigating adaptive radiations from a genomic perspective is shedding light on how speciation and trait diversification occur. Contreras-Ortiz *et al.* (2018) used RADseq to investigate species diversification in Andean lupines and found evidence for both adaptive, ecological and non-adaptive, geographical drivers influencing their radiation.

Table 1. Legume genomes published as of 11 May 2019
 Some genomes were sequenced by multiple groups and not all are summarised here, in which case the genome summarised has the first author underlined. Gb, gigabases; Mb, megabases; bp, base pairs; nr, not reported; SNP, single-nucleotide polymorphism

Organism	Relevance	Genome size	Length assembled	Number of genes	Method	Sequence produced	Assembly status	References
Reference quality								
<i>Glycine max</i> (soybean) Williams 82 ^A	Protein and oil crop	1115 Mb	955 Mb	56044	Sanger shotgun & BAC sequencing	Iterative	8.04× coverage; 17191 contigs of L50 182.8 kb, 1190 scaffolds of L50 47.8 Mb; 20 chromosomes produced with BAC-end sequences	<u>Schmutz et al. 2010</u> ; Shen et al. 2018
<i>Glycine soja</i> (wild soybean)	Wild relative	1115 Mb	915.4 Mb	nr	Illumina (genome resequencing); 454 (variant validation)	48.8 Gb	52× coverage; 75195 contigs of N50 ~250 bp; re-sequenced reference: this genome was assembled based on soybean reference	<u>Kim et al. 2010</u> ; Xie et al. 2019
Reference draft quality								
<i>Arachis duranensis</i>	A genome diploid wild ancestor	1.25 Gb	1211 Mb	36734	Illumina	325.73 Gb	154× coverage; 765406 contigs of N50 22.3 kb; 635392 scaffolds of N50 948 kb; assembled to 10 pseudomolecules using BAC sequences and GBS-derived genetic map	<u>Bertioli et al. 2016</u> ; Chen et al. 2016
<i>Arachis ipaensis</i>	B genome diploid wild ancestor	1.56 Gb	1512 Mb	41840	Illumina	416.59 Gb	163× coverage; 869435 contigs of N50 23.5 kb; 759499 scaffolds of N50 5.34 Mb; assembled to 10 pseudomolecules using BAC sequences and GBS-derived genetic map	<u>Bertioli et al. 2016</u> ; Lu et al. 2018
<i>Arachis hypogaea</i> cv. Tifrunner (cultivated peanut)	Crop and oil plant	~2.7 Gb	2.54 Gb	90519	PacBio (sequencing), Illumina (polishing), Hi-C (scaffolding)	173.6 Gb	76.74× coverage; 17.75 Mreads of average length 9784 bp; 4.037 contigs of N50 1.5 Mb; 384 scaffolds of N50 134.0 Mb; assembled to 20 chromosomes using Hi-C	<u>Bertioli et al. 2019</u> ; Chen et al. 2019
<i>Cajanus cajan</i> (Pigeon pea) var. <i>Asha</i>	Crop plant	833.07 Mb	605.78 Mb	48680	Illumina, Sanger BAC-end libraries	237.2 Gb	~163.4× coverage; 173708 contigs of N50 21.95 kb; 137 542 scaffolds of N50 536 kb; anchored to 11 chromosomes using BAC-ends and genetic map	Varshney et al. 2012
<i>Cajanus cajan</i> (Pigeon pea) var. <i>Asha</i> ^A	Crop plant	858 Mb	648.2 Mb	56888	454, Illumina	195.4 Gb	360028 contigs of N50 5341 bp	Mahato et al. 2018
							>10× coverage; 59681 scaffolds of N50 13.9 kb; anchored to 11 chromosomes using linkage map of 347 SNPs	Singh et al. 2012

<i>Cicer arietinum</i> (chickpea)	Crop plant	738 Mb	544.73 Mb	28 269	Illumina	153 Gb	207.32× coverage; 62 619 contigs of N50 23.54 kb; 7163 scaffolds of N50 40 Mb; eight pseudomolecules produced with help of BAC sequences and genetic map	Varshney <i>et al.</i> 2013
<i>Cicer arietinum</i> L. (chickpea)	Crop plant	740 Mb	519.8 Mb	27 571	454, Illumina	57 Gb	15 × 454 coverage; 181 462 scaffolds of N50 77.3 Kb; 8 pseudomolecules produced with help of BAC sequences and genetic map	Jain <i>et al.</i> 2013
<i>Lotus japonicus</i> (bird's-foot trefoil) ^A	Model legume	472 Mb	315.1 Mb	39 734	Sanger shotgun and BAC sequencing; Illumina added later	Iterative	Sanger: 2.4 × coverage; 109 986 contigs; 110 940 scaffolds; six chromosomes produced with BAC and TAC sequences. Illumina 40 × coverage; 23 572 contigs with N50 (anchored) of 118 kb; further hypothetical chromosome 0 produced	Sato <i>et al.</i> 2008
<i>Lupinus angustifolius</i> (narrow-leaved lupin)	Health food	924 Mb	609 Mb	33 076	Illumina	150.4 Gb	162.8 × coverage; 1 068 669 contigs of N50 4246 bp; 14 379 scaffolds of N50 (?); 20 pseudomolecules derived from genetic map	Hane <i>et al.</i> 2017
<i>Medicago truncatula</i> (barrel medic) ^A	Model legume	465 Mb	384.5 Mb	50 894	Sanger, BAC end, Optical Mapping, 454, Illumina	Iterative	All data types (v.4.0): contig N50 102 kb; scaffold N50 4.24 Mb; eight pseudomolecules produced from optical maps and GBS-based linkage map	Young <i>et al.</i> 2011; Tang <i>et al.</i> 2014
<i>Phaseolus vulgaris</i> (common bean) ^A	Model bean and protein crop	587 Mb	537.2 Mb	36 995	Sanger, 454, and Illumina; PacBio added later	Iterative	All data types (v.2.1): 1044 contigs of N50 1.9 Mb; 478 scaffolds of N50 49.7 Mb; 11 chromosomal pseudomolecules produced with BACs and linkage map	Schmutz <i>et al.</i> 2014
<i>Trifolium pratense</i> (red clover)	Forage legume	420 Mb	309 Mb	40 868	Illumina, Sanger BAC-end sequencing	nr	Hybrid assembly: 30 × coverage; 39 904 scaffolds of N50 223 kb; seven pseudomolecules produced with BACs, a physical, and two genetic maps	De Vega <i>et al.</i> 2015
<i>Trifolium subterraneum</i> ^A	Annual wild relative of forage crop	540 Mb	512 Mb	32 333	Illumina, 454, Bionano	157 Gb (Illumina + 454)	290.7 × coverage; 27 257 super-scaffolds of N50 410.5 kb; eight pseudomolecules produced by genetic map of >35 000 SNPs and a Bionano physical map	Hirakawa <i>et al.</i> 2016; Kaur <i>et al.</i> 2017

Table 1. (continued)

Organism	Relevance	Genome size	Length assembled	Number of genes	Method	Sequence produced	Assembly status	References
<i>Trifolium subterraneum</i> ^A	Annual wild relative of forage crop	540 Mb	403.4 Mb	nr	Illumina, 454, Hi-C	nr	48× coverage (Hi-C); 46 453 contigs of N50 22 377; 5285 scaffolds of N50 56.3 Mb; eight chromosome-length scaffolds with lengths ranging from 49.5 to 65.2 Mb produced from Hi-C mapping	Hirakawa <i>et al.</i> 2016; Dudchenko <i>et al.</i> 2018
<i>Vigna angularis</i> (adzuki bean)	Protein crop	612 Mb	443 Mb	26 857	Illumina, 454	172 Gb	291.2× coverage; 36 516 contigs of N50 21.9 kb; 3883 scaffolds of N50 703 kb; 11 pseudomolecules produced with GBS-based genetic map	Kang <i>et al.</i> 2015
<i>Vigna radiata</i> var. <i>radiata</i> (mungbean)	Protein crop	579 Mb	421 Mb	22 427	Illumina, 454		25 922 contigs of N50 41.8 kb; 2748 scaffolds of N50 1516 kb; 11 pseudomolecules produced by GBS-based genetic map	Kang <i>et al.</i> 2014
<i>Vigna unguiculata</i> (cowpea)	Protein crop	620 Mb	519.4 Mb	29 773	PacBio (genome), Illumina (gene prediction)	56.8 Gb	~6 million PacBio reads of N50 14.5 kb for 91× coverage; 11 pseudomolecules produced using two optical maps and 10 genetic maps comprising >44 000 SNPs	Lonardi <i>et al.</i> 2019
Draft quality								
<i>Ammopiptanthus nanus</i>	Endangered desert shrub	889 Mb	823.74 Mb	37 188	PacBio (genome), Illumina (gene prediction)	64.72 Gb (PacBio); 55.97 Gb (Illumina)	72.59× coverage; 7.92 million reads of N50 12.79 kb; 1099 contigs of N50 2.76 Mb	Gao <i>et al.</i> 2018
<i>Cercis canadensis</i> (redbud)	Horticultural tree	301 Mb	330 Mb	34 023	Illumina	335.07 Gb	220.74× coverage; gap-filled scaffolding N50 of 12 883 bp; gap-filled scaffold N50 of 421 kb	Griesmann <i>et al.</i> (2018)
<i>Chamaecrista fasciculata</i> (partridge pea)	Annual legume	550 Mb	429 Mb	32 832	Illumina	381.97 Gb	492.29× coverage; gap-filled scaffolding N50 of 14 934 bp; gap-filled scaffold N50 of 96.6 kb	Griesmann <i>et al.</i> (2018)
<i>Dipteryx oleifera</i>	Tropical timber species	1.89 Gb	1.16 Gb	nr	Illumina, PacBio	50.78 Gb	44× Illumina coverage, 5× PacBio coverage. Combined assembly: 381 857 contigs of N50 8194 bp	Jimenez-Madrizal 2018
<i>Faidherbia albida</i> (ring acacia)	Agroforest crop	661 Mb	653.7 Mb	28 979	Illumina	nr	nr	Chang <i>et al.</i> 2019
<i>Glycine latifolia</i>	Wild relative	1.13 Gb	939 Mb	54 475	10X Genomics, Chromium linked reads on Illumina	78.39 Gb	66 825 contigs of N50 (?); 42 539 scaffolds of N50 853.5 kb; 20 pseudomolecules produced using two genetic maps and the G.max genome	Liu <i>et al.</i> 2018

<i>Glycyrrhiza uralensis</i> (chinese licorice)	Medicinal legume	400 Mb	379 Mb	34445	PacBio, Illumina	387.29 Gb	817× (Illumina) and 16.86× (PacBio) coverage. Hybrid assembly: 72 148 contigs of N50 7324 bp; 12 528 scaffolds of N50 109.3 kb	Mochida <i>et al.</i> 2017
<i>Lablab purpureus</i> (hyacinth bean)	Ancient crop	423 Mb	395.5 Mb	20946	Illumina	nr	nr	Chang <i>et al.</i> 2019
<i>Lupinus angustifolius</i> (narrow-leaved lupin)	Health food	1152 Mb	598 Mb	57807	Illumina	31 Gb	26.9× coverage; 457917 contigs of N50 5.8 kb; 234 534 scaffolds of N50 12.5 kb, RAD-seq produced 20 linkage groups but pseudomolecules not built	Yang <i>et al.</i> 2013
<i>Mimosa pudica</i> (sensitive plant)	Thigmotropic plant	896 Mb	557 Mb	33108	Illumina	370.07 Gb	287.65× coverage; gap-filled scaffig N50 of 11 069 bp; gap-filled scaffold N50 of 119.7 kb	Griesmann <i>et al.</i> (2018)
<i>Nissolia schottii</i> (Schott's yellowhood)	Desert perennial	471 Mb	466 Mb	36369	Illumina	144.06 Gb	149.48× coverage; gap-filled scaffig N50 of 20 655 bp; gap-filled scaffold N50 of 179.7 kb	Griesmann <i>et al.</i> (2018)
<i>Trifolium pratense</i> (red clover)	Forage legume	420 Mb	314.6 Mb	47398	Illumina	24.6 Gb	58.8× coverage; 236 989 contigs of N50 2397 bp; 176 760 scaffolds of N50 4750 bp	Ištvánek <i>et al.</i> 2014
<i>Vigna radiata</i> var. <i>sublobata</i>	Wild relative	501.7 Mb	423 Mb	22834	Illumina	nr	45 606 contigs of N50 16.4 kb; 8161 scaffolds of N50 214 kb	Kang <i>et al.</i> 2014
<i>Vigna reflexo-pilosa</i> var. <i>glabra</i>	Wild relative	967.8 Mb	791.6 Mb	41844	Illumina	nr	163 809 contigs of N50 4.6 kb; 29 166 scaffolds of N50 63 kb	Kang <i>et al.</i> 2014
<i>Vigna subterranea</i> (bambara groundnut)	Food crop	550 Mb	535 Mb	31707	Illumina	nr	nr	Chang <i>et al.</i> 2019
Unassembled								
<i>Vigna angularis</i> var. <i>nipponensis</i>	Wild relative	562 Mb	nr	nr	Illumina	43.9 Gb	72.04× coverage	Kang <i>et al.</i> 2015
<i>Vigna nakashimae</i>	Wild relative	756 Mb	431 Mb	23197	Illumina	36.2 Gb	47.91× coverage	Lestari <i>et al.</i> 2014
<i>Vigna nepalensis</i>	Wild relative	610 Mb	nr	nr	Illumina	45 Gb	80× coverage	Kang <i>et al.</i> 2015

^A Genome was improved by addition of next-generation sequence data after initial publication and statistics reported are of the improved version.




Fig. 2. Leguminosae genomes sequenced as of 11 May 2019. Taxa sequenced are listed, with their genera highlighted in blocks on the Legume Phylogeny Working Group (2017) *matK* best-scoring maximum-likelihood tree. Subfamilies are outlined using blocks in the inside of the phylogeny, with a scheme similar to that of Legume Phylogeny Working Group (2017). Numbers in the legend are numbers sampled/total number for genera and species respectively. *Vigna* species (*spp.*): *angularis* var. *angularis*, *angularis* var. *nipponensis*, *nakashimae*, *nepalense*, *radiata* var. *radiata*, *radiata* var. *sublobata*, *reflexa-pilosa*, *subterranean* and *unguiculata*.

Using transcriptomes from slowly and rapidly diversifying lupin lineages, Nevado *et al.* (2016) verified, for the first time in plants, the role of adaptive evolution in rapid radiations. Rapidly diversifying lineages had two to three times more positively selected genes than did slowly diversifying lineages, suggesting a genome-wide response to adaptation. Further, the rapidly diversifying Andean lineage exhibited a higher gene-expression divergence than did the slowly diversifying lineages, suggesting that underlying genomic shifts in expression happened during adaptive radiations. Also, shifts in gene-expression level were non-randomly clustered around significant evolutionary time points, including at the base of the Andean clade when lupins moved into novel, extreme montane environments, and near a branch signifying a shift from annual to perennial life-history. NGS has gone beyond simply detecting shifts in diversification rates, by enabling the determination of the how and why behind diversification in plants.

Polyploidy

Whole-genome duplication (WGD), or polyploidy, is a major evolutionary process underlying speciation in plants (Van de Peer *et al.* 2017), and legumes are no exception (Doyle 2012). Polyploid crop legumes, such as peanut, alfalfa and soybean are well known and, in soybean (*Glycine max* [L.] Merr.), studies have confirmed genomic evidence for both ancient and recent polyploidisation events just before the radiation of the genus *c.* 10–12 million years ago, and between 40 and 66 million years ago (Egan and Doyle 2010). Legume WGDs were further characterised by Cannon *et al.* (2015), by using transcriptomic and genomic data from 20 diverse legumes and 17 outgroups to determine that WGDs coincide with the origins of major legume lineages. A follow-up study incorporating genome data for the genus *Cercis* L. (Cercidoideae), suggested that *Cercis* may represent the only extant legume lineage lacking a polyploid

history, providing a plausible hypothesis of what the ancestral legume genome looked like (Stai *et al.* 2019). In contrast, Koenen *et al.* (2019) used thousands of nuclear genes and 72 protein-coding chloroplast genes to find evidence for WGDs at the stem of all legumes, as well as nested WGDs subtending radiation of subfamilies Papilionoideae and Detarioideae. Koenen *et al.* (2019) also described difficulties in resolving the initial divergence of the legume family tree and suggested that polyploidy may play a key role in the lack of support for deep-branching relationships in the family. Further improvements in long-read NGS platforms may solve such recalcitrant nodes and facilitate research into polyploidy and its ramifications.

Genome sequencing also enables us to detect and characterise more recent WGDs. Allopolyploidy within *Glycine* has been shown to be rampant and complex, but genomic data have helped unravel relationships within the genus by using transcriptomics (Bombarely *et al.* 2014) and GBS (Sherman-Broyles *et al.* 2017). Similarly, comparison of rDNA, genomic and fluorescence *in situ* hybridisation data and whole plastome sequences generated by Illumina-based sequencing of *Stylosanthes scabra* Vogel, an important forage legume, and its hypothesised genome donors *S. hamata* (L.) Taub. or *S. seabrana* B.L.Maass & 't Mannetje (A genome) and *S. viscosa* (L.) Sw. (B genome), have provided evidence for an allopolyploid origin of *S. scabra* and showed the genomic impacts of subsequent homogenisation following 'genomic shock' (Marques *et al.* 2018). Capitalising on a target-gene set derived from the *Medicago truncatula* genome (de Sousa *et al.* 2014), Eriksson *et al.* (2017) tested the hypothesis that *M. prostrata* is a homoploid hybrid, while accounting for the impact and signature of introgression from *M. sativa* as a contributor of genetic variation therein. Eriksson *et al.* (2018) characterised the allopolyploid origin of two *Medicago* species and showed the importance of allele phasing. Using NGS methods to investigate the mechanisms of polyploidisation can resolve duplicated regions by phasing and comparative analyses.

Ecological aspects

Quantification and integration of information on species, genetic, population and ecosystem diversity from NGS-based metagenomic methods can contribute to biodiversity and conservation assessment and understanding of evolutionary history, population processes, community assembly, and ecosystem equilibrium and services (Papadopoulou *et al.* 2015; de la Harpe *et al.* 2017). As primary mediators of nitrogen fixation, legumes are an integral source of fixed nitrogen within terrestrial communities from grasslands to forests. Here, we review some of the recent advances obtained through genomic studies within ecological arenas.

Forestry and range management

From providing essential ecosystem and natural resources to sustaining natural and human infrastructures, legumes are fundamental and abundant components of biomes across the globe, from desert sands to towering trees of the Amazon rainforests and temperate grasslands where legume

biodiversity is directly related to the overall biological diversity and community health. Metagenomic comparison of above- and below-ground plant-species richness in a grassland employed Sanger sequencing and Roche 454 pyrosequencing to illustrate that below-ground diversity was higher, demonstrating the power of NGS methods to detect dormant plant diversity (Hiiesalu *et al.* 2012). Assessments of the portion of biodiversity that legumes represent in grasslands, and perhaps other ecosystems, through metagenomic barcoding may help in maintaining and managing biodiversity in ecosystems, be they native or range managed, particularly as loss of legume biodiversity is directly linked to a decline in the nitrogen budget and the overall health and biodiversity of plant communities (Spehn *et al.* 2002).

Legumes are particularly prevalent in Neotropical forests and savannas, where plot data suggest that they make up more than 11% of species (Oliveira-Filho *et al.* 2013; Yahara *et al.* 2013). With increasing pressures on forest-ecosystem dynamics, including fragmentation, over-logging of particular species, and wholesale deforestation, understanding how to mitigate the effects of such events and their toll on ecosystem health, as well as the roles that legumes play therein, are urgent needs. Genomic methods are poised to help (Neale and Kremer 2011). Creation of genomic-marker sets is an important starting point. For example, RADseq has been used to generate 330 SNPs for population genetic analyses in *Robinia pseudoacacia* L., an economically important eastern North American tree widely cultivated and invasive in Europe and elsewhere (Verdu *et al.* 2016); employment of such markers may help forest managers capitalise on its use and limit its dynamic spread. Similarly, several studies have generated SNP marker sets for *Dipteryx* Schreb., an economically important genus of Neotropical canopy trees, various species of which are threatened. These include development of microsatellite markers (Soares *et al.* 2012), a nuclear and plastid SNP MassArray panel (Honorio Coronado *et al.* 2019), and a draft genome (Jimenez-Madrigal 2018), providing useful genetic data resources for management and conservation through detailed GD assessment. Gailing *et al.* (2017) used RADseq to produce a framework genetic-linkage map for *Gleditsia triacanthos* L., a common North American hardwood forest tree, providing an important genetic resource for future quantitative trait-locus mapping.

Landscape genomic data coupled with an assessment of GD can promote sound forest and range management, quantify the impacts of deforestation, fragmentation, climate change and habitat restoration. For example, *Acacia koa* A.Gray, a Hawaiian endemic and one of two dominant canopy hardwood tree species in Hawaiian forests, is under increasing pressures from logging and changing climate. Gugger *et al.* (2018) used GBS to assess GD across 311 *Acacia koa* trees sampled over its geographical, elevational and climatic range, and found evidence for genetic differentiation among islands and a strong association between genetic structure and the mean annual rainfall. These results suggest that changing rainfall patterns could cause a genetic offset between adaptation and extant populations, placing future survival of this species at risk. Such knowledge can be used for future management planning. Similarly, Grando (2015) assessed GD across native and restored populations of *Piptadenia gonoacantha* (Mart.) J.F.Macbr., a

species that is often used in reforestation across the Brazilian Atlantic Forest because of its rapid growth and regeneration. These data provided insights into which native populations should be used as seed sources for restoration efforts and determined that GD was similar across native and restored sites, which is evidence of successful capture and maintenance of biodiversity during restoration efforts. As NGS becomes more affordable for high sample numbers, landscape genomics and GD assessment using such data can ensure sound scientific foundations for natural-resource management and biodiversity conservation in the face of global change.

Conservation biology and genetics

As one of the largest plant families and key components of tropical and temperate forest ecosystems, legumes are the focus of global legume-diversity assessment (GLDA; Yahara *et al.* 2013) to quantify biodiversity and species loss stemming from rapid deforestation taking place across South-East Asia. Within the GLDA, biodiversity assessments of rosewoods (*Dalbergia* spp.; Vatanparast *et al.* 2013), *Bauhinia* L., *Mucuna* Adans. (Moura *et al.* 2016) and *Desmodium* Desv. have been prioritised. These ongoing studies are incorporating species-distribution modelling with biodiversity metrics. To assess conservation and biodiversity metrics correctly, a complete time-calibrated phylogenetic tree of a target group is required. However, for many lineages or communities, a complete phylogeny at the species level is not available. Combining biodiversity metrics with enhanced NGS-based phylogenies can enable greater understanding of the contribution of legumes to the overall biodiversity and aid in conservation efforts (for a review on biodiversity metrics, see Kellar *et al.* 2015). Similarly, Ahrends *et al.* (2016) used Illumina shotgun sequencing of rosoid species in a Nebraska grassland to isolate ~80 plastid genes for 45 species, 22 of which were legumes. This enabled the reconstruction of a plastid phylogeny for a complete community for divergence dating and estimation of conservation metrics, illustrating the potential of genomics methods for conservation-biology research.

Conservation often involves efforts to characterise genetic diversity within vulnerable or endangered species by using microsatellite markers. Genomic sequencing has revolutionised discovery of such markers (Zalapa *et al.* 2012), whether species-specific (Abdelkrim *et al.* 2009) or for use across a broader taxonomic group (Hodel *et al.* 2016), including for several legumes (e.g. Borges *et al.* 2015; Morris *et al.* 2016). For example, microsatellite markers developed from the *Lotus japonicus* (Regel) K.Larsen genome (Sato *et al.* 2008) were used through cross-species amplification in *Lotus sessilifolius* DC., a species endemic to Macaronesia (Yang *et al.* 2018). To determine which populations should be prioritised for conservation management, eight populations across four islands were assessed for their GD across 11 microsatellite markers, highlighting a population from Tejina-Milán as strongly distinct and of low genetic diversity relative to the others, suggesting this as one population to target. Next-generation sequencing platforms coupled with amplicon sequencing can now be used to obtain microsatellite data *en masse* (Zhan *et al.* 2017).

Although microsatellites are a time-tested and effective tool for assessing GD, issues of small sample sizes and low numbers

of markers limit their power for assessing population structure and dynamics. Reduced representation techniques such as GBS or RADseq, or targeted enrichment and amplicon-sequencing methods linked to RADseq and GBS (e.g. GTseq and Rapture), offer cost-effective methods for generating orders of magnitude more marker sites (thousands to millions of SNPs) than do microsatellites (Meek and Larson 2019) for assessing GD and establishing population and species relationships. For example, Harrison *et al.* (2019) garnered thousands of SNPs by GBS to compare genetic variation among infraspecific taxa of the *Astragalus lentiginosus* Hook. species complex including var. *piscinensis* Barneby, that inhabits just 8 km². They showed that, in spite of rarity, significant genetic diversity and population structure exists within and among varieties. Harrison *et al.* (2019) exemplified the power of NGS to produce large numbers of SNPs for population-genomic and conservation studies.

Invasion biology

The numerous markers generated by reduced-representation genomic methods can also resolve recent population divergences such as those arising from anthropogenic plant invasions (Chown *et al.* 2015) and help understand whether pre-adaptation to the novel environment or rapid adaptive changes account for invasions. For example, Helliwell *et al.* (2018) used 9658 SNPs genotyped across 446 accessions of *Medicago polymorpha* L. within its native European and introduced New World ranges. They showed that latitudinal variation in phenology that facilitated invasion resulted from rapid evolutionary adaptation across this clinal gradient following a single introduction and subsequent range expansion. Similarly, M. S. Haynsen and A. N. Egan (unpubl. data; A. N. Egan, pers. comm.) genotyped ~600 loci by using GBS over 600 individuals of kudzu (*Pueraria montana* (Lour.) Merr. var. *lobata* (Willd.) Maesen & Almeida ex Sanjappa & Pradeep), a notorious invasive vine that has now spread over half of the USA, to detail its introduction history.

Understanding the evolutionary mechanisms behind invasiveness and tracing patterns of introduction history are important for managing invasive species. Alternatively, investigating the reasons behind dieback within an invasive species inside its introduced range may also aid management. For example, Steinrücken (2017) used Illumina metagenomic sequencing of fungal and bacterial soil and plant communities within healthy and diseased populations in the native and introduced ranges of *Parkinsonia aculeata* L., an invasive caesalpinoid legume tree introduced from Venezuela to Australia, to determine that fungal endophytes, not bacterial ones, were likely to be responsible for dieback in the invasive range; this knowledge may prove useful for biological control of *P. aculeata* in Australia. Studies that integrate gene-expression profiling with population-level sampling will be able to truly determine how invasiveness arises, knowledge that we as yet lack at the genomic level.

Nitrogen fixation

Legumes provide numerous inputs to agricultural and natural ecosystems, with perhaps the most important being soil enrichment by nitrogen fixation. The majority of legume

species have the ability to form symbioses with rhizobial bacteria that transform atmospheric nitrogen to ammonia, making atmospheric nitrogen bioavailable in the soil as amino acids and other cellular constituents. Metagenomic approaches are commonly used to characterise soil microbiomes (Andújar *et al.* 2015), advancing understanding of the interactions of legumes and the environment through nodulation and nitrogen fixation (e.g. Afkhami and Stinchcombe 2016). Birnbaum *et al.* (2018) investigated symbionts of *Acacia rostellifera* Benth. across the natural-soil fertility gradient of the Jurien Bay Dune chronosequence. Using Illumina-based *nif* metabarcoding, they delineated which species of Rhizobiaceae inhabited nodules and how the composition of nodule symbionts changed with soil fertility. They noted that the older soils with the lowest soil phosphorus had more unclassified operational taxonomic units, suggesting a shift to a unique set of nitrogen-fixing bacteria more adapted to limited soil fertility. Associations between legume species and their root-nodule symbionts can be both generalist and specific, but our understanding of how specificity is controlled remains fragmentary. Keller *et al.* (2018) attempted to delineate the hows and whys of host specificity by meta-sequencing three *Lupinus* nodulomes, discovering differential compatibility between lupine and *Bradyrhizobium* and that different hormone, secondary metabolite and plant-defence mechanisms activated depended on host compatibility. Similarly, GBS was used to show lack of local adaptation to different prevailing soil species of *Ensifer* between northern and southern populations of *Medicago lupulina* L. in North America (Harrison *et al.* 2017). Knowledge of which nitrogen-fixing bacteria are optimal is of key importance to maximise crops yields. High-throughput sequencing of nodules or rhizospheres has determined the symbionts of legume crops, forage legumes and invasive species, including cowpea (Chidebe *et al.* 2018), lucerne (alfalfa; Wigley *et al.* 2017), rooibos tea (Le Roux *et al.* 2017) and silver wattle (Kamutando *et al.* 2017). Furthermore, characterisation and sequencing of bacterial genomes, including those that are key to nitrogen fixation, is now routine. For example, a new species, *Rhizobium hidalgonense*, was recently isolated and characterised from a *Phaseolus vulgaris* L. nodule growing in acidic soil (Yan *et al.* 2017), and draft genome sequences of *Bradyrhizobium* (Tian *et al.* 2015) and another new species, *Ensifer aridi* (Le Quéré *et al.* 2017), have been completed and characterised.

The evolutionary origin of nodulation has baffled researchers for many years, and particularly whether it evolved once or multiple times, whether some sort of cryptic precursor could have predisposed lineages in the nitrogen-fixing clade of angiosperms to evolve nodulation, and whether polyploidy may have been involved in its origination (Werner *et al.* 2014; Doyle, 2016). Griesmann *et al.* (2018) used a comparative genomics approach to address this question by sequencing genomes from across the nitrogen-fixing clade including several non-nodulating species, to look for genes known to be vital to nitrogen-fixing nodulation (NFN). They discovered that all NFN genes were conserved in all but one nodulating species and found evidence for multiple independent losses of the nodule-inception (*NIN*) gene in 10 of 13 non-nodulating species, attesting to the key role of this gene within the nodulation pathway, and suggesting multiple

evolutionary losses of nodulation (van Velzen *et al.* 2019). Comparative analysis of the legume *Medicago truncatula* and the non-legume *Parasponia andersonii* Planch. also supported the idea of a single origin and multiple losses of nodulation across the nitrogen-fixing angiosperm clade (van Velzen *et al.* 2018, 2019). However, others disagree, citing differences in transcriptome profiles as evidence for a two-step process in origination of nodulation (Battenberg *et al.* 2018). In addition to working out its origin, understanding the subsequent evolution of nodulation is also important. One unique example is the ability of some members of photosynthesising *Bradyrhizobium* strains to prompt nodulation and fix nitrogen in the absence of *nodABC* genes that are key to modulating nodule formation. Strains of this bacterium form symbioses with *Aeschynomene evenia* C. Wright, and transcriptomics has enabled the creation of a gene-map to delineate genes involved in this unique type of symbiosis (Chaintreuil *et al.* 2016).

Genetic and genomics resources

The genomics era and all genetic knowledge owes homage to the pea (*Pisum sativum* L.), the legume crop that captured Gregor Mendel's attention and led to his monumental discovery of genetic inheritance. The importance of nitrogen-fixing legumes as the most important protein and rotation crops has driven high-throughput sequencing and genomics to establish genetic and genomic data to underpin plant-breeding research. Here, we discuss some of these resources and subsequent discoveries.

Sequenced genomes

With soybean providing nearly 70% of the world's edible protein, the completion of the soybean genome (Schmutz *et al.* 2010) marked an important milestone in legume research. Even though the soybean genome was not sequenced using NGS methods, it remains a gold standard in plant-genome sequencing, providing a vital reference for other work, including the assembly and annotation of other legume genomes. Draft genome sequences of the model legumes *Lotus japonicus* (Sato *et al.* 2008) and *Medicago truncatula* (Young *et al.* 2011), and common bean, *Phaseolus vulgaris* (Schmutz *et al.* 2014), based on Sanger sequencing, have been improved to reference quality by using NGS data (e.g. Tang *et al.* 2014).

Plant genomes are complex relative to other eukaryotic genomes, owing to large genome sizes, higher repetitive fraction, prevalence of polyploidy and difficulty of obtaining high-molecular weight DNA caused by presence of the cell wall, polysaccharides and secondary metabolites that impair enzymes or damage DNA (Jiao and Schneeberger 2017). Nevertheless, the advent of NGS has enabled whole-genome sequencing of a rapidly growing number of species and multiple accessions within species (Fig. 2, Table 1), as costs have fallen (from the ~US\$20 million cost of the soybean genome; Marris 2008) and techniques have advanced such that a single laboratory can now rapidly produce multiple genomes. For example, Griesmann *et al.* (2018) sequenced draft genomes of four legume genera for comparative genomic analysis of nodulation genes, and Chang *et al.* (2019) sequenced three African orphan legume crop species to support crop breeding. Liu *et al.* (2019) set out to

barcode 761 vascular plants from the Riuli Botanical Garden by using NGS whole-genome sequencing, 71 of which are legumes (because identification and genome assembly remain incomplete, these genomes are not listed in Table 1).

Owing to plant genome size and complexity, NGS-based plant genomes often have poorer assembly statistics than do those of vertebrates and are commonly assembled only to draft status. However, a draft genome is still worthwhile! For example, genomic sequencing of multiple accessions of the recent domesticated, *Lupinus angustifolius* L., has allowed mapping of key disease resistance and domestication traits (Yang *et al.* 2013; Hane *et al.* 2017). Even without assembly to draft status, sequencing the full genomic content of an organism can yield important information. For example, Kang *et al.* (2015) sequenced the genomes of *Vigna angularis* var. *nipponensis* (Ohwi) Ohwi & H. Ohashi and *V. nepalensis* Tateishi & Maxted, wild relatives of cultivated adzuki bean, *V. angularis* var. *angularis* (Willd.) Ohwi & H. Ohashi. Even though the reads were not fully assembled, nor genes called *de novo*, variation in the two wild relatives called against the draft genome of the cultivated bean provided insights into the timing of domestication and variation across the genus.

Early NGS platforms provided vast amounts of data; however, limitations, such as short read length and limited read output, made genome assembly from such platforms alone challenging. As technologies have improved and new mapping methods have been devised, draft genome assemblies can now be improved on after the fact. For example, subterranean clover (*Trifolium subterraneum* L.), an annual relative of the forage legumes *T. repens* L. (white clover) and *T. pratense* L. (red clover), was sequenced using Illumina and 454 pyrosequencing as a reference in the genus *Trifolium* L. (Hirakawa *et al.* 2016), producing 27 228 scaffolds representing the draft genome, TSUd_r1.1. Subsequently, Kaur *et al.* (2017) applied a Bionano Genomics (San Diego, CA, USA) optical map and a transcriptome atlas to improve the TSUd_r1.1 assembly by anchoring unplaced contigs, correcting mis-assemblies, and improving gene annotation, resulting in the coalescence of 264 contigs into 97 super-scaffolds representing 43% of the genome and a 1.4-fold increase in the scaffold N50 to create the Tsub_RefV2.0 assembly. Similarly, also building on TSUd_r1.1, application of Hi-C contact mapping, a chromosome conformation technology, corrected misjoins, anchored and oriented scaffolds into eight chromosome-length pseudomolecules that included 95% of the sequenced bases in the input assembly, with the resulting TrSub3 assembly improving the initial scaffold N50 from 287 kb to 56 Mb (Dudchenko *et al.* 2018).

Whereas Illumina sequencing produces the most data for the lowest cost, long-read sequencing is increasingly affordable and can sequence through repetitive regions that often make plant genomes difficult to assemble. For example, Lonardi *et al.* (2019) used PacBio (Pacific Biosciences of California, Inc., Menlo Park, CA, USA) to sequence the genome of cowpea (*Vigna unguiculata* [L.] Walp) by using 56.8 Gb of sequence data with a read N50 of 14 595 bp, i.e. longer than many *de novo* assembled contigs and scaffolds. In addition, they employed two Bionano optical maps and a novel 'stitching' assembly method that combined eight different assemblies created using three

different programs. Long-read lengths coupled with iterative cross-checking methods to remove chimeric joins enabled a highly accurate and complete assembly that was able to push through the long repetitive regions that break other plant assemblies into numerous contigs. They also detected a 4.2-Mb chromosomal inversion that may confer resistance to a parasitic weed, *Striga gesnerioides* (Willd.) Vatke. Cross-platform genome-sequencing strategies that use both short- and long-read platforms coupled with physical mapping are, thus, good contemporary approaches for plant-genome sequencing, including enhancement of existing assemblies from draft to reference quality.

As the number of sequenced genomes has grown, genetic and genome-scope databases have been developed, including Phytozome (phytozome.jgi.doe.gov, accessed 30 May 2019) and the Legume Information System (legumeinfo.org, accessed 30 May 2019; Dash *et al.* 2016), to facilitate access to genome content and information. These databases can be integrated with programs and platforms such as CyVerse (cyverse.org, accessed 30 May 2019) to enable public access to datasets, management and integration of personal data and access to high-performance computing platforms, thus enabling collaborative and critical breakthroughs through the combined power of genetic and computational resource platforms. However, none of these databases integrates other types of information, such as morphological-trait variation or geographical-distribution data, prompting legume systematists to devise a legume portal that can integrate across platforms and data types (Bruneau *et al.* 2019). Development of integrated database systems across genomic, taxonomic, geographical, morphological, population and expression-profile data will enhance our ability to parse and use information from NGS data and plant genomes.

Organelle genomes

With the abundance of chloroplast DNA, compared to nuclear, within a cell, 10–20% of NGS reads are chloroplast, such that entire chloroplast genomes can be assembled from even low-coverage genome skimming. Furthermore, the shorter length and less-repetitive nature of the chloroplast makes assembly easier, even for degraded herbarium material (Bakker *et al.* 2016). Thus, the use and sequencing of chloroplast genomes has rocketed forward with the advent of NGS methods. Some have even suggested the use of whole chloroplast genomes as DNA barcodes for plants (Li *et al.* 2015).

Comparative phylogenetic analyses have shown that structural changes in the chloroplast genome are often synapomorphies for large legume clades (Wang *et al.* 2018). For example, the loss of one copy of the inverted repeat marks the inverted repeat lacking clade (IRLC), a large papilionoid clade including tribes Cicereae, Hedysareae, Trifolieae and Fabeae (Vicieae), among others (Wojciechowski *et al.* 2000). Recent sequencing of chloroplasts from eight Cercidoideae genera showed structural diversification characteristic of that subfamily (Wang *et al.* 2018). In contrast, comparative analyses of chloroplast genomes determined that Papilionoideae have reduced genome sizes and are more divergent from the ancestral angiosperm chloroplast-genome

organisation than are other subfamilies (Schwarz *et al.* 2015). Structural changes have also been identified at a generic level. *Trifolium subterraneum* L. was shown to have a highly unusual chloroplast genome greatly expanded by repetitive regions (Cai *et al.* 2008). To determine the evolutionary origin of this unique plastome type, Sveinsson and Cronk (2014) sequenced eight other *Trifolium* chloroplast genomes, showing that the expanded *T. subterraneum*-type plastome is shared by members of ‘core *Trifolium*’, providing a synapomorphy for what they called the ‘refractory clade’. The expansion of the inverted repeat region across the large Ingeae + *Acacia s.s.* clade of mimosoid legumes (Wang *et al.* 2017) has provided another synapomorphic structural plastid mutation apparently characterising that clade, and is associated with shifts in evolutionary rate or selection pressures on proximate chloroplast gene regions (Mensous *et al.* 2017).

Numerous chloroplast genomes have been sequenced in legumes; the NCBI archive of full-length chloroplast genomes includes 284 accessions (97.2% circularised, 2.8% linear; 66.4% complete, 35.6% partial), representing 72 genera and 202 species, ranging from 120289 bp in *Lathyrus odoratus* L. to 178887 bp in *Ebenopsis ebano* (Berlandier) Barneby & Grimes, including 118 NCBI designated reference sequences (Table S1, available as Supplementary material to this paper). This list is by no means exhaustive and some accessions may represent duplicate chloroplast assemblies at various stages; however, every effort to remove duplicates was taken. Even so, this list exemplifies the power of the chloroplast for evolutionary studies in legumes, including for phylogeny estimation of recalcitrant clades. For example, whole chloroplast alignments showed that 13 of 15 *Guibourtia* Benn. species were reciprocally monophyletic, with evidence of a single dispersal event from the Old to the New World c. 12 million years ago (Tosso *et al.* 2018). Of the nearly 300 legume accessions with sequenced chloroplasts, 95 are from *Acacia*, a lineage of >1000 species. Williams *et al.* (2016) used whole chloroplast sequences of 65 *Acacia* species to build a robust backbone constraint tree, adding amplicon

sequences of four chloroplast and two nuclear ribosomal loci for 508 other *Acacia* species, to produce a more robust phylogeny than from amplicon sequences alone. This suggests that combining legacy Sanger sequencing datasets with large, phylogenomic datasets will likely yield positive results.

Another byproduct of NGS is sequencing of the mitochondrion, which has significantly fewer genomes sequenced than does the chloroplast (Table 2). The first legume mitochondrion sequenced was that of *Vigna radiata* (L.) Wilczek (Alverson *et al.* 2011) and the Sanger-based sequence of the *Vicia faba* L. mitochondrion was not far behind (Negruk 2013). The first legume mitochondria sequenced by NGS methods were *Lotus japonicus* and *Milletia pinnata* (L.) Panigrahi, on the Illumina platform (Kazakoff *et al.* 2012). Soon after, the mitochondrial genomes of soybean (Chang *et al.* 2013) and *Medicago truncatula* (Bi *et al.* 2016) were sequenced on the 454 platform, and *Vigna angularis* was compiled using both 454 and Illumina sequence reads (Naito *et al.* 2013). The paucity of sequenced plant mitochondrial genomes is largely due to the highly recombinant nature of mitochondria, which makes assembly difficult. Shi *et al.* (2018) combined PacBio with Illumina reads to investigate the repetitive complement of the *Styphnolobium japonicum* (L.) Schott mitochondrial genome, and discovered that small repeats (<100 bp) had a disproportionate impact on the evolution of *Styphnolobium* mitochondria through mediation of recombination along intronic regions. The first non-papilionoid legume mitochondrion to be sequenced was *Leucaena trichandra* (Zucc.) Urb., completed using PacBio (Kovar *et al.* 2018) to obtain long-reads, enabling assessment of variable assemblies and investigation of mitochondrial genome-size variation in legumes. Further, overlaying transcriptomic data enabled comparative study of RNA editing among *Leucaena* species, providing knowledge that can yield useful information regarding close species relationships and hybrid origins (Kovar *et al.* 2018).

Despite these advances, the repetitive nature of the mitochondrion and its evolutionary implications are only

Table 2. List of mitochondrial genomes sequenced in Fabaceae from NCBI

All are complete and circularised. bp, base pairs; accession numbers preceded by NC_ are designated as reference sequences by NCBI. References for each genome can be found by querying the accession number in GenBank. Data were accessed 23 February 2019

Species	Length (bp)	Platform	Accession number	GI number	Reference
<i>Acacia ligulata</i>	698 138	Illumina	MH933866.1	1 552 055 398	Sanchez-Puerta <i>et al.</i> 2019
<i>Ammopiptanthus mongolicus</i>	475 396	Illumina	NC_039660.1	1 511 253 925	Yu <i>et al.</i> 2018
<i>Castanospermum australe</i>	542 079	Illumina	MK426679.1		Zhang <i>et al.</i> 2019
<i>Glycine max</i>	402 558	454	NC_020455.1	476 507 670	Chang <i>et al.</i> 2013
<i>Glycine soja</i>	402 545	Illumina	NC_039768.1	1 511 246 382	Asaf <i>et al.</i> 2018
<i>Leucaena trichandra</i>	722 009	PacBio	NC_039738.1	1 511 244 784	Kovar <i>et al.</i> 2018
<i>Lotus japonicus</i>	380 861	Illumina	NC_016743.2	387 866 040	Kazakoff <i>et al.</i> 2012
<i>Medicago truncatula</i>	271 618	454	NC_029641.1	1 003 725 997	Bi <i>et al.</i> 2016
<i>Milletia pinnata</i>	425 718	Illumina	NC_016742.1	372 450 249	Kazakoff <i>et al.</i> 2012
<i>Senna occidentalis</i>	447 106	Illumina	NC_038221.1	1 442 330 107	Kang <i>et al.</i> 2019a
<i>Senna tora</i>	566 589	Illumina	NC_038053.1	1 436 049 411	Kang <i>et al.</i> 2019b
<i>Styphnolobium japonicum</i>	484 916	Illumina, PacBio	NC_039596.1	1 509 839 239	Shi <i>et al.</i> 2018
<i>Vicia faba</i>	588 000	Sanger	KC189947.1	442 803 095	Negrak 2013
<i>Vigna angularis</i>	404 466	Illumina, 454	NC_021092.1	501 594 995	Naito <i>et al.</i> 2013
<i>Vigna radiata</i> var. <i>radiata</i>	401 262	Sanger	NC_015121.1	323 149 028	Alverson <i>et al.</i> 2011

beginning to be understood, particularly with regard to cross-species interactions. That said, Sanchez-Puerta *et al.* (2019) presented an elegant study of host–parasite horizontal gene transfer between the holoparasite *Lophophytum mirabile* Schott & Endl. and its host *Acacia ligulata* A.Cunn., showing that ~60% of the *L. mirabile* mitochondrion is derived from its host, including 34 of 43 protein-coding genes (also see Sanchez-Puerta *et al.* 2017; Kovar *et al.* 2018). In addition, ~26 of its native genes were replaced by host genes through homologous recombination, and with a large portion of intergenic regions also host-derived. This work provided incontrovertible support for mitochondrial-to-mitochondrial horizontal gene transfer, a phenomenon well documented in *Amborella trichopoda* Baill. (Rice *et al.* 2013). Furthermore, Kovar *et al.* (2018) also showed that non-coding mitochondrial DNA was horizontally transferred, suggesting capture of an entire mimosoid mitochondrial genome during the evolutionary history of the *Lophophytum* parasite. Knowledge such as this may shed light on the incredible length diversity of plant mitochondrial genomes, which range from 271 618 bp in *Medicago truncatula* to 698 138 bp in *Acacia ligulata*. At the time of writing, 15 legume mitochondria had been sequenced (Table 2). As this number expands, broader comparative analyses become possible; sequencing of the wild soybean, *Glycine soja* (Asaf *et al.* 2018), provided insights into a likely progenitor to the soybean mitochondrion, whereas that of *Ammopiptanthus mongolicus* (Kom.) S.H.Cheng provided a glimpse into a Tertiary relic (Yu *et al.* 2018). Comparative analysis of the most recently sequenced legume mitochondrion, namely that of *Castanospermum australe* A.Cunn. & C.Fraser, enabled phylogenetic analysis of 33 mitochondrial genes across legumes, producing a fully supported phylogeny (Zhang *et al.* 2019), suggesting that despite the fewer phylogenetically informative sites than in chloroplast genomes (Palmer and Herbon 1988), the mitochondrion has potential utility for molecular systematics of legumes.

Plant breeding

Simply put, the genomics era has revolutionised plant breeding in legumes. We cannot possibly summarise it all here; however, some studies are essential for discussion. Major goals of plant breeding are improvement of crop traits useful for humans and adaptation to environment such as increasing yield, freeze and drought tolerance, disease resistance, nutritional quality and seed size. A genome sequence provides the basis for genome-wide association studies (GWAS), functional genomics, quantitative trait-loci (QTL) analysis and linkage studies, SNP variant detection, and genetic modification by CRISPR, among many others. For example, sequencing of the licorice genome (*Glycyrrhiza uralensis* Fisch.) enabled assessment of genes involved in flavonoid and saponin synthesis, producing candidate genes for improving yield of glycyrrhizin, an active chemical component used in traditional Chinese medicine (Mochida *et al.* 2017). Of wider importance are several crop genomes, in addition to those already discussed. Chickpea (*Cicer arietinum* L.) is surpassed only by soybean, peanut and *Phaseolus* ssp. as a widely grown legume crop (Fig. 3, Table S2, available as Supplementary material to this paper).

Its sequenced genomes (Jain *et al.* 2013; Varshney *et al.* 2013), coupled with resequencing of numerous cultivars and wild accessions from 10 countries, enabled discovery of gene regions associated with key domestication, agronomic and disease-resistance traits (Varshney *et al.* 2013). Likewise, sequencing of *Cajanus cajan* [L.] Millsp., the pigeon pea (Singh *et al.* 2012; Varshney *et al.* 2012; Mahato *et al.* 2018), one of the most widely cultivated and consumed orphan legume crops in India, highlighted drought-tolerance genes important during its domestication (Varshney *et al.* 2012). These genomic resources have aided successful translational improvements of both chickpea and pigeon pea (for review, see Varshney 2016).

To illustrate the types of NGS applications and advances made in legume-crop breeding, we focus on *Vigna* Savi, a pantropical genus of over 100 species of which 10 species have been domesticated and are cultivated in both warm, humid and dry, seasonal climates (Table 3; Harouna *et al.* 2018). As such, *Vigna* species provide food for nearly half of the world's population. Ten species or varieties of *Vigna* have had their genomes sequenced (Tables 1, 3), including six crop species (Kang *et al.* 2014, 2015; Chang *et al.* 2019; Lonardi *et al.* 2019) and four wild relatives (Kang *et al.* 2014, 2015; Lestari *et al.* 2014). The impacts of these genome sequences are just beginning to be realised. For example, comparative searches of the *V. angularis* and *V. radiata* genomes for homologues of *ONSEN*-like sequences, a heat-activated retrotransposon isolated from *Arabidopsis thaliana* (L.) Heynh., produced several key hits in *V. angularis*, sequences that proved to be polymorphic in different accessions (Masuta *et al.* 2018). These retrotransposons were associated with accumulation of extrachromosomal DNA and were employed to successfully induce retrotransposition of *V. angularis* *ONSEN*-like elements in regenerated *V. angularis* callus tissue, providing a new tool for molecular breeding in *Vigna*.

Many plant-breeding efforts require a genetic map. RADseq was used to create a high-density genetic-linkage map by using 170 individuals to enable QTL detection of loci related to yield in cowpea (Pan *et al.* 2017), a species that includes two broad cultivar types, namely, bushy, short-podded grain grown predominantly in Africa, and long-podded climbing vegetables grown mostly in Asia. Genomic scans confirmed that pod length was selected for during domestication of the vegetable variety (Xu *et al.* 2017). Furthermore, GWAS for genomic regions controlling pod length between cultivars of these two types discovered 72 SNPs whose pod-length association was verified across 299 cowpea accessions. This knowledge, coupled with transcriptomic analysis, suggested the involvement of sugar, gibberellin and nutrition as key factors in pod-length regulation and that cell proliferation rather than cell elongation was key to pod length.

Applications of transcriptomics are often used to study stress responses such as low temperature-stress resistance in *V. subterranea*, providing gene modules for plant-breeding improvement (Bonthala *et al.* 2016), among many others. These studies often identify candidate genes useful for functional genomic characterisation. For instance, acidic soils are often characterised by accumulation of aluminium, with aluminium toxicity causing reduced yields in crop plants.




Fig. 3. Total production of cultivated legumes by year from 2008 to 2017. Data from FAOSTAT (www.fao.org/faostat/, accessed 25 April 2019). Numbers were summed across all countries, with China counted only once. nes, not elsewhere specified. Data are listed in Table S2.

Table 3. Summary of genetic and genomic resources available for species of *Vigna*

Including all cultivated species and those wild relatives with next-generation sequencing data. References for genomes can be had from Tables 1, 2 and S1. Nu, nuclear; Mito, mitochondrial; Cp, chloroplast; SRA, number of BIO-projects available in short-read archive (SRA) database of the NCBI as of 28 February 2019; ?, genome sequenced with NGS data in SRA, but not yet published

Common name	<i>Vigna</i> species	Cultivated area or distribution	Genome			
			Nu	Mito	Cp	SRA
Moth bean	<i>V. aconitifolia</i> (Jacq.) Marechal	Semi-arid India, South-East Asia				1
Adzuki bean	<i>V. angularis</i> (Willd.) Ohwi & H. Ohashi	Eastern Asia, Australia, New Zealand	X	X	X	7
wild relative	<i>V. angularis</i> var. <i>nipponensis</i> (Ohwi) Ohwi & H. Ohashi	Japan, Korea, China, Bhutan, Nepal, India (Himalaya)	X			
Black gram (urid bean)	<i>V. mungo</i> (L.) Hepper	Southern and South-East Asia				4
wild relative	<i>V. nakashimae</i> (Ohwi) Ohwi & H. Ohashi	Northern China, Korea, northern Japan	X			
wild relative	<i>V. nepalensis</i> Tateishi & Maxted	Bhutan, eastern Nepal, north-eastern India	X			
Mung bean	<i>V. radiata</i> var. <i>radiata</i> (L.) R. Wilczek	Southern, eastern and South-East Asia	X	X	X	11
wild relative	<i>V. radiata</i> var. <i>sublobata</i> (Roxb.) Verdc.	Asia, Africa, Australia	X			
Creole bean	<i>V. reflexo-pilosa</i> Hayata	India, Mauritius, Philippines, Vietnam	X			
Minni payaru	<i>V. stipulacea</i> Kuntze	India, Sri Lanka, Indonesia, New Guinea				
Bambara groundnut	<i>V. subterranea</i> (L.) Verdc.	Semi-arid Africa	X			2
Rice bean	<i>V. umbellata</i> (Thunb.) Ohwi & Ohashi	Asian tropics	X?			2
Cowpea	<i>V. unguiculata</i> (L.) Walp	Semi-arid regions of Africa, Asia, USA, Europe, Central and South America	X		X	12
Tuber cowpea	<i>V. vexillata</i> (L.) A. Rich.	Indonesia				

Rice bean (*V. umbellata* (Thunb.) Ohwi & H. Ohashi) is tolerant of soils with high aluminium accumulation. Studies of such resistance in other plants have suggested a role for the

abscisic acid (ABA) pathway in dealing with aluminium toxicity. Transcriptomic analysis and functional genomic studies in rice bean support this hypothesis (Fan *et al.* 2019),

but its involvement was shown to be dependent on *AB15*, a transcription factor that mediated changes in cell-wall modification and osmoregulation.

Another focus of plant-breeding efforts involve resistance to pests and pathogens. Micro-RNAs (miRNAs) are short (20–24 nucleotides) non-coding RNAs that act to regulate gene expression, particularly during response to stress, such as that inflicted on a plant by viral infection. The mungbean yellow mosaic India virus (MYMIV) significantly decreases yield across many South-East Asian countries where *Vigna* species are staple crops. Understanding the mechanisms of infection and host response are key to crop survival. Kundu *et al.* (2017) identified miRNAs involved in the stress response of *V. mungo* (L.) Hepper to MYMIV infection, by comparing gene-expression patterns and miRNAs across resistant and non-resistant cultivars, with putative target genes known to be involved in pathogen-stress response, such as *NB-LRR*, *ARF*, *SOD*, *SPB*, and Basic blue copper protein, linked to and validated as being regulated by miRNAs in stressed and non-stressed plants. Another proverbial pest problem plaguing legume crops is bruchid beetles (*Callosobruchus* Pic. spp.), which infest seeds in the field, then multiply and destroy seed during storage. Bruchid resistance has been found in wild mungbean, *V. radiata* var. *sublobata* (Roxb.) Verdc., and in one mungbean cultivar. Schafleitner *et al.* (2016) used GBS to map inbred recombinant lines for each of these resistant populations and discovered one QTL associated with bruchid resistance shared between both resistant entities. The markers associated with this QTL were validated as 100% predictive of bruchid resistance, providing an excellent screening tool for developing resistant cultivars. Similar genome-based research across the full spectrum of important legume crops is underway. Rapidly expanding knowledge of gene functional pathways from comparative genomics and the development of genome-based selection, is accelerating and revolutionising legume-crop breeding; these efforts will aid in future crop and food security.

This short summary would be incomplete without some discussion of the genomics of crop wild relatives, which can contribute key genes and diversity to crops to increase pest and disease resistance and extend environmental tolerances of important crop legumes. One of the first crop wild relatives to have its genome sequenced was *Glycine soja* Sieb. and Zucc. (Kim *et al.* 2010), one of the first genome re-sequencing projects in plants. In the study of Kim *et al.* (2010), *G. soja* was sequenced and its genome assembled against that of soybean, with a comparison between the two suggesting that *G. max* diverged from *G. soja* c. 0.27 million years ago, i.e. hundreds of thousands of years before domestication of soybean. The draft genome of *Glycine latifolia* (Benth.) C.A.Newell & Hymowitz, a perennial wild relative of soybean, was recently sequenced using only linked-reads from a single 10X Genomics (Pleasanton, CA, USA) Chromium library (Liu *et al.* 2018), presenting a valuable resource of alleles and genes for soybean improvement. Like soybean, the cultivated peanut (or groundnut; *Arachis hypogaea* L.) is of polyploid origin. The large allotetraploid peanut genome sequence (http://peanutbase.org/peanut_genome, accessed 30 May 2019; Chen *et al.* 2019; ~2.7 Gb) comprises the two recently diverged subgenomes of its diploid ancestors, *Arachis duranensis* Krapov. & W.C.Greg. and

A. ipaensis Krapov. & W.C.Greg, which were used to assist assembly of the domesticated-peanut genome and detect genetic recombination among peanut subgenomes, providing key information regarding the origin of the cultivated peanut (Bertioli *et al.* 2016). As climate changes and demands for better yield increase, wild relatives offer plant breeders sources of diverse and adapted traits to incorporate into cultivars.

Securing the future of legume germplasm diversity

Given the central roles that legumes play in agriculture, ecosystems and the global nitrogen cycle, securing the future of legume genetic resources is both urgent and of paramount importance. Application of genomics methods, from capturing the genome of a fading species to guiding restoration and management efforts and assisting and speeding up plant breeding, can enable advances in legume research that were previously unattainable that will help secure their future.

A good example is *Ammopiptanthus* S.H.Cheng, a genus of two evergreen broadleaf desert shrub species endemic to central Asia. This taxonomically isolated genus is hypothesised to be a relic from the Tertiary, having adapted to aridification from a moist and humid climate of the evergreen broadleaf forest characteristic of the Tethyan flora (Zhang *et al.* 2015). As a relic, *Ammopiptanthus* has evolved drought-, cold- and wind-resistance, among other stress-tolerant characteristics. Both species are considered threatened because of low seed set and increasing anthropogenic disturbance, with *A. nanus* (Popov) S. H.Cheng listed as *Critically Endangered* (www.iucnredlist.org, accessed 19 February 2019). *Ammopiptanthus* has been the focus of abiotic stress studies, using transcriptomics to understand its drought (Zhou *et al.* 2012) and cold (Pang *et al.* 2013) adaptations. These studies have enabled the cloning, characterisation and validation of candidate genes shown as beneficial in other model study systems, conferring salinity and heat tolerance to *Escherichia coli* by the *A. nanus* betaine aldehyde dehydrogenase gene (Yu *et al.* 2014), or cold tolerance by the *A. nanus* antifreeze gene *AnAFP* to both *E. coli* and tobacco (Deng *et al.* 2014), among many others. Recently, the *A. nanus* genome was sequenced on the PacBio platform (Table 1; Gao *et al.* 2018), providing an essential resource for functional genomics and improvement studies, with *Ammopiptanthus* fast becoming a model for understanding drought and cold tolerance.

Along with alfalfa (*Medicago sativa*) and several related species, clovers (*Trifolium* spp.) are important forage crops and key components of natural grazing systems. Red clover (*Trifolium pratense*) is an excellent short-term hay-rotation crop and pasture or field restoration plant. Its dual use as protein-rich fodder and soil-fertility enhancer led to its early adoption in crop-rotation schemes and is one reason why some tout it as a perfect plant for conservation agriculture, a movement aimed at sustainable intensification of food and forage for increasing world needs (McKenna *et al.* 2018). The recent sequencing of its genome (Ištváněk *et al.* 2014; De Vega *et al.* 2015) has provided another forage genome resource for comparative analyses to assist breeding (Annicchiarico *et al.* 2015). Furthermore, transcriptomic studies have enabled advances in seed set (Kovi *et al.* 2017), drought (Yates *et al.* 2014) and leaf

senescence (Chao *et al.* 2018a). For example, Chao *et al.* (2018b) used PacBio to sequence and analyse full-length transcripts, enabling the detection of over 30 000 novel isoforms and 5492 alternative splicing events, the majority of which involved intron retention. Using two-dimensional difference gel electrophoresis, Bertrand *et al.* (2016) characterised the proteome of red clover, particularly with reference to freezing tolerance; selection detailed the involvement of a small number of cold-regulated proteins, suggesting them as potential targets for breeding programs. These studies illustrated the incredible protein diversity arising from the complement of known genes and provided a third layer of information on top of genomic and expression-level knowledge.

The impacts of climate change will be particularly felt on food security, with changes in temperature and rainfall patterns affecting what, where and how crops will flourish. Genomics can help improve food security by providing fundamental molecular-data resources through genome sequencing, comparative genomics and rapid evaluation of genetic variation, to enable assessment of adaptation priority, determine import of specific traits for breeding programs, and discover adaptive genes and traits (Mousavi-Derazmahalleh *et al.* 2019), as previously discussed. Many have suggested that the potential of legumes to enhance food security and reduce meat consumption has not been realised nor explored adequately (Foyer *et al.* 2016). *Phaseolus*, with a sequenced genome and well-developed breeding programs, provides a good example. Key areas of inquiry remain unexplored. For instance, phenotyping of the world's germplasm collections is important to understand the diversity currently held and how that compares to wild populations (McClellan *et al.* 2011), and with the advent of genomics techniques, the time is now ripe for doing so *en masse*. Genotyping-by-sequencing and genome–environment associations within wild populations of common bean have discovered several genomic regions associated with drought adaptation, pinpointing genomic signatures potentially useful for marker-assisted selection (Cortés and Blair 2018). Yet, much still needs to be done. As wild relatives and related species often harbour greater GD than do cultivars, efforts to assess such genetic and phenotypic diversity and unlock the adaptive potential of these entities should be a priority for future food security (Porch *et al.* 2013). Efforts to identify gaps in germplasm collections for *Phaseolus* have begun (Ramírez-Villegas *et al.* 2010), yet, much is still needed (Dohle *et al.* 2019).

Conclusions

From evolution and ecology to classical and applied genetics, the genomics era has and will continue to contribute to our understanding of legume biology in many important ways, and this is set to expand and accelerate in coming years as sequencing costs continue to fall. Partial or whole genomes of hundreds to thousands of legume species are expected to be available in the near future, with the 10 000 genomes project already targeting over 300 legume genera (Cheng *et al.* 2018). While such an unprecedented accumulation of genomic data presents significant computational and analytical challenges, there will soon be unparalleled opportunities to address large-scale comparative genomic questions. Analyses of numerous

legume genomes will provide massive insights into synteny, micro- and macro-level gene and genome duplication, and chromosome structure coupled with gene expression analyses and recombination perspectives. At the same time, variant detection-panel screening across population-level sampling schemes will enable exceptional advances in understanding patterns and processes of evolution at micro- and macro-scales. Such knowledge will help secure the future of legumes as vital components of ecological and economic security.

Conflicts of interest

A. N. Egan is also an Associate Editor of the 'Advances in Legume Systematics 13' special issue. Despite this relationship, she did not at any stage have Associate Editor-level access to this manuscript while in peer review, as is the standard practice when handling manuscripts submitted by an editor to this journal. *Australian Systematic Botany* encourages its editors to publish in the journal and they are kept totally separate from the decision-making process for their manuscripts. The authors have no further conflicts of interest to declare.

Declaration of Funding

This research did not receive any specific funding. However, A. N. Egan was supported in part by a subaward from the Carlsberg Foundation to Henrik Balslev (grant #CF14-0245) and by the USA National Science Foundation (DEB-1352217).

Acknowledgements

A. N. Egan thanks Dr Henrik Balslev and the Department of Biosciences at Aarhus University for providing a sabbatical academic home from which to pursue this work and Dr Jeffrey J. Doyle and Dr Colin Hughes, as well as an anonymous reviewer, for helpful comments on the manuscript.

References

- Abdelkrim J, Robertson BC, Stanton JA, Gemmill NJ (2009) Fast, cost-effective development of species-specific microsatellite markers by genomic sequencing. *BioTechniques* **46**, 185–192. doi:10.2144/000113084
- Afkhami ME, Stinchcombe JR (2016) Multiple mutualist effects on genome-wide expression in the tripartite association between *Medicago truncatula*, nitrogen-fixing bacteria and mycorrhizal fungi. *Molecular Ecology* **25**, 4946–4962. doi:10.1111/mec.13809
- Ahrendsen DL, Aust SK, Kellar PR (2016) Biodiversity assessment using next-generation sequencing: comparison of phylogenetic and functional diversity between Nebraska grasslands. *Plant Systematics and Evolution* **302**, 89–108. doi:10.1007/s00606-015-1246-6
- Alverson AJ, Zhuo S, Rice DW, Sloan DB, Palmer JD (2011) The mitochondrial genome of the legume *Vigna radiata* and the analysis of recombination across short mitochondrial repeats. *PLoS One* **6**, e16404. doi:10.1371/journal.pone.0016404
- Andrews KR, Good JM, Miller MR, Luikart G, Hohenlohe PA (2016) Harnessing the power of RADseq for ecological and evolutionary genomics. *Nature Reviews. Genetics* **17**, 81–92. doi:10.1038/nrg.2015.28
- Andújar C, Arribas P, Ruzicka F, Crampton-Platt A, Timmermans MJTN, Vogler AP (2015) Phylogenetic community ecology of soil biodiversity using mitochondrial metagenomics. *Molecular Ecology* **24**, 3603–3617. doi:10.1111/mec.13195
- Annicchiarico P, Barrett B, Brummer EC, Julier B, Marshall AH (2015) Achievements and challenges in improving temperate perennial forage

- legumes. *Critical Reviews in Plant Sciences* **34**, 327–380. doi:10.1080/07352689.2014.898462
- Asaf S, Khan AL, Al-Harrasi A, Kim TH, Lee IJ (2018) The first complete mitochondrial genome of wild soybean (*Glycine soja*). *Mitochondrial DNA. Part B, Resources* **3**, 527–528. doi:10.1080/23802359.2018.1467228
- Azani N, Bruneau A, Wojciechowski MF, Zarre S (2017) Molecular phylogenetics of annual *Astragalus* (Fabaceae) and its systematic implications. *Botanical Journal of the Linnean Society* **184**, 347–365. doi:10.1093/botlinnean/box032
- Bagheri A, Maassoumi AA, Rahiminejad MR, Brassac J, Blattner FR (2017) Molecular phylogeny and divergence times of *Astragalus* section Hymenostegis: an analysis of a rapidly diversifying species group in Fabaceae. *Scientific Reports* **7**, 14033. doi:10.1038/s41598-017-14614-3
- Bakker FT, Lei D, Yu J, Mohammadin S, Wei Z, van de Kerke S, Gravendeel B, Nieuwenhuis M, Staats M, Alquezar-Planas DE, Holmer R (2016) Herbarium genomics: plastome sequence assembly from a range of herbarium specimens using an iterative organelle genome assembly pipeline. *Biological Journal of the Linnean Society. Linnean Society of London* **117**, 33–43. doi:10.1111/bj.12642
- Battenberg K, Potter D, Tabuloc CA, Chiu JC, Berry AM (2018) Comparative transcriptomic analysis of two actinorhizal plants and the legume *Medicago truncatula* supports the homology of root nodule symbioses and is congruent with a two-step process of evolution in the nitrogen-fixing clade of angiosperms. *Frontiers in Plant Science* **9**, 1256. doi:10.3389/fpls.2018.01256
- Bertioli DJ, Cannon SB, Froenicke L, Huang G, Farmer AD, Cannon EK, Liu X, Gao D, Clevenger J, Dash S, Ren L, Moretzsohn MC, et al. (2016) The genome sequences of *Arachis duranensis* and *Arachis ipaensis*, the diploid ancestors of cultivated peanut. *Nature Genetics* **48**, 438–446. doi:10.1038/ng.3517
- Bertioli DJ, Jenkins J, Clevenger J, Dudchenko O, Gao D, Seijo G, Leal-Bertioli SC, Ren L, Farmer AD, Pandey MK, Samoluk SS, et al. (2019) The genome sequence of segmental allotetraploid peanut *Arachis hypogaea*. *Nature Genetics* **51**, 877–884. doi:10.1038/s41588-019-0405-z
- Bertrand A, Bipifubusa M, Castonguay Y, Rocher S, Szopinska-Morawska A, Papadopoulos Y, Renault J (2016) A proteome analysis of freezing tolerance in red clover (*Trifolium pratense* L.). *BMC Plant Biology* **16**, 65. doi:10.1186/s12870-016-0751-2
- Bi C, Wang X, Xu Y, Wei S, Shi Y, Dai X, Yin T, Ye N (2016) The complete mitochondrial genome of *Medicago truncatula*. *Mitochondrial DNA – B. Resources* **1**, 122–123. doi:10.1080/23802359.2016.1144087
- Birnbaum C, Bissett A, Teste FP, Laliberté E (2018) Symbiotic N₂-fixer community composition, but not diversity, shifts in nodules of a single host legume across a 2-million-year dune chronosequence. *Microbial Ecology* **76**, 1009–1020. doi:10.1007/s00248-018-1185-1
- Blanco-Pastor JL, Bertrand YJ, Liberal IM, Wei Y, Brummer EC, Pfeil BE (2018) Robustness of RADseq for evolutionary network reconstruction from gene trees. *bioRxiv* **2018**, 414243.
- Bombarely A, Coate JE, Doyle JJ (2014) Mining transcriptomic data to study the origins and evolution of a plant allopolyploid complex. *PeerJ* **2**, e391. doi:10.7717/peerj.391
- Bonthala VS, Mayes K, Moreton J, Blythe M, Wright V, May ST, Massawe F, Mayes S, Twycross J (2016) Identification of gene modules associated with low temperatures response in bambara groundnut by network-based analysis. *PLoS One* **11**, e0148771. doi:10.1371/journal.pone.0148771
- Borges DB, Mariano-Neto E, Gaiotto FA (2015) Development of microsatellite primers for *Melanoxylon brauna* (Fabaceae): an endangered and endemic tree from the Brazilian Atlantic Forest. *Conservation Genetics Resources* **7**, 65–68. doi:10.1007/s12686-014-0288-3
- Bruneau A, Mercure M, Lewis GP, Herendeen PS (2008) Phylogenetic patterns and diversification in the caesalpinoid legumes. *Botany* **86**, 697–718. doi:10.1139/B08-058
- Bruneau A, Borges LM, Allkin R, Egan AN, de la Estrella M, Javadi F, Klitgård B, Miller JT, Murphy DJ, Sinou C, Vatanparast M, Zhang R (2019) Towards a new online species-information system for legumes. *Australian Systematic Botany* **32**(5–6), 495–518. doi:10.1071/SB19025
- Bybee SM, Bracken-Grissom H, Haynes BD, Hermansen RA, Byers RL, Clement MJ, Udall JA, Wilcox ER, Crandall KA (2011) Targeted amplicon sequencing (TAS): a scalable next-gen approach to multilocus, multitaxa phylogenetics. *Genome Biology and Evolution* **3**, 1312–1323. doi:10.1093/gbe/evr106
- Cai Z, Guisinger M, Kim HG, Ruck E, Blazier JC, McMurtry V, Kuehl JV, Boore J, Jansen RK (2008) Extensive reorganization of the plastid genome of *Trifolium subterraneum* (Fabaceae) is associated with numerous repeated sequences and novel DNA insertions. *Journal of Molecular Evolution* **67**, 696–704. doi:10.1007/s00239-008-9180-7
- Cannon SB, McKain MR, Harkess A, Nelson MN, Dash S, Deyholos MK, Peng Y, Joyce B, Stewart CN Jr, Rolf M, Kutchan T, et al. (2015) Multiple polyploidy events in the early radiation of nodulating and nonnodulating legumes. *Molecular Biology and Evolution* **32**, 193–210. doi:10.1093/molbev/msu296
- Cardoso D, de Queiroz LP, Pennington RT, de Lima HC, Fonty E, Wojciechowski MF, Lavin M (2012) Revisiting the phylogeny of papilionoid legumes: new insights from comprehensively sampled early-branching lineages. *American Journal of Botany* **99**, 1991–2013. doi:10.3732/ajb.1200380
- Chaintreuil C, Rivallan R, Bertioli DJ, Klopp C, Gouzy J, Courtois B, Leleux P, Martin G, Rami JF, Gully D, Parrinello H, Séverac D, et al. (2016) A gene-based map of the Nod factor-independent *Aeschynomene evenia* sheds new light on the evolution of nodulation and legume genomes. *DNA Research* **23**, 365–376. doi:10.1093/dnares/dsw020
- Chang S, Wang Y, Lu J, Gai J, Li J, Chu P, Guan R, Zhao T (2013) The mitochondrial genome of soybean reveals complex genome structures and gene evolution at intercellular and phylogenetic levels. *PLoS One* **8**, e56502. doi:10.1371/journal.pone.0056502
- Chang Y, Liu H, Liu M, Liao X, Sahu SK, Fu Y, Song B, Cheng S, Kariba R, Muthemba S, Hendre PS, Mayes S, et al. (2019) The draft genomes of five agriculturally important African orphan crops. *GigaScience* **8**, giy152. doi:10.1093/gigascience/giy152
- Chao Y, Xie L, Yuan J, Guo T, Li Y, Liu F, Han L (2018a) Transcriptome analysis of leaf senescence in red clover (*Trifolium pratense* L.). *Physiology and Molecular Biology of Plants* **24**, 753–765. doi:10.1007/s12298-018-0562-z
- Chao Y, Yuan J, Li S, Jia S, Han L, Xu L (2018b) Analysis of transcripts and splice isoforms in red clover (*Trifolium pratense* L.) by single-molecule long-read sequencing. *BMC Plant Biology* **18**, 300. doi:10.1186/s12870-018-1534-8
- Chapman MA (2015) Transcriptome sequencing and marker development for four underutilized legumes. *Applications in Plant Sciences* **3**, 1400111. doi:10.3732/apps.1400111
- Chau JH, Rahfeldt WA, Olmstead RG (2018) Comparison of taxon-specific versus general locus sets for targeted sequence capture in plant phylogenomics. *Applications in Plant Sciences* **6**, e1032. doi:10.1002/aps3.1032
- Chen X, Li H, Pandey MK, Yang Q, Wang X, Garg V, Li H, Chi X, Doddamani D, Hong Y, Upadhyaya H, Guo H, et al. (2016) Draft genome of the peanut A-genome progenitor (*Arachis duranensis*) provides insights into geocarpy, oil biosynthesis, and allergens. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 6785–6790. doi:10.1073/pnas.1600899113
- Chen X, Lu Q, Liu H, Zhang J, Hong Y, Lan H, Li H, Wang J, Liu H, Li S, Pandey MK, Zhang Z, et al. (2019) Sequencing of cultivated peanut, *Arachis hypogaea*, yields insights into genome evolution and oil improvement. *Molecular Plant* **12**(7), 920–934. doi:10.1016/j.molp.2019.03.005
- Cheng S, Melkonian M, Smith SA, Brockington S, Archibald JM, Delaux PM, Li FW, Melkonian B, Mavrodiev EV, Sun W, Fu Y, Yang H, et al. (2018) 10KP: a phylodiverse genome sequencing plan. *GigaScience* **7**, giy013. doi:10.1093/gigascience/giy013
- Chidebe IN, Jaiswal SK, Dakora FD (2018) Distribution and phylogeny of microsymbionts associated with cowpea (*Vigna unguiculata*) nodulation

- in three agroecological regions of Mozambique. *Applied and Environmental Microbiology* **84**, e01712–e01717.
- Choi HK, Luckow MA, Doyle J, Cook DR (2006) Development of nuclear gene-derived molecular markers linked to legume genetic maps. *Molecular Genetics and Genomics* **276**, 56–70. doi:10.1007/s00438-006-0118-8
- Chown SL, Hodgins KA, Griffin PC, Oakeshott JG, Byrne M, Hoffmann AA (2015) Biological invasions, climate change and genomics. *Evolutionary Applications* **8**, 23–46. doi:10.1111/eva.12234
- Contreras-Ortiz N, Atchison GW, Hughes CE, Madriñán S (2018) Convergent evolution of high elevation plant growth forms and geographically structured variation in Andean *Lupinus* (Fabaceae). *Botanical Journal of the Linnean Society* **187**, 118–136. doi:10.1093/botlinnean/box095
- Cortés AJ, Blair MW (2018) Genotyping by sequencing and genome–environment associations in wild common bean predict widespread divergent adaptation to drought. *Frontiers in Plant Science* **9**, 128. doi:10.3389/fpls.2018.00128
- Dash S, Campbell JD, Cannon EK, Cleary AM, Huang W, Kalberer SR, Karingula V, Rice AG, Singh J, Umale PE, Weeks NT, Wilkey AP, Farmer AD, Cannon SB (2016) Legume information system (LegumeInfo.org): a key component of a set of federated data resources for the legume family. *Nucleic Acids Research* **44**, D1181–D1188. doi:10.1093/nar/gkv1159
- de la Harpe M, Paris M, Karger DN, Rolland J, Kessler M, Salamin N, Lexer C (2017) Molecular ecology studies of species radiations: current research gaps, opportunities and challenges. *Molecular Ecology* **26**, 2608–2622. doi:10.1111/mec.14110
- de Queiroz LP, Pastore JF, Cardoso D, Snak C, Lima AL, Gagnon E, Vatanparast M, Holland AE, Egan AN (2015) A multilocus phylogenetic analysis reveals the monophyly of a circumscribed papilionoid legume tribe Diocleae with well-supported generic relationships. *Molecular Phylogenetics and Evolution* **90**, 1–9. doi:10.1016/j.ympev.2015.04.016
- de Sousa F, Bertrand YJ, Nylinder S, Oxelman B, Eriksson JS, Pfeil BE (2014) Phylogenetic properties of 50 nuclear loci in *Medicago* (Leguminosae) generated using multiplexed sequence capture and next-generation sequencing. *PLoS One* **9**, e109704. doi:10.1371/journal.pone.0109704
- De Vega JJ, Ayling S, Hegarty M, Kudrna D, Goicoechea JL, Ergon A, Rognli OA, Jones C, Swain M, Geurts R, Lang C, Mayer KFX, et al. (2015) Red clover (*Trifolium pratense* L.) draft genome provides a platform for trait improvement. *Scientific Reports* **5**, 17394. doi:10.1038/srep17394
- Deng LQ, Yu HQ, Liu YP, Jiao PP, Zhou SF, Zhang SZ, Li WC, Fu FL (2014) Heterologous expression of antifreeze protein gene *AnAFP* from *Ammopiptanthus nanus* enhances cold tolerance in *Escherichia coli* and tobacco. *Gene* **539**, 132–140. doi:10.1016/j.gene.2014.01.013
- Dohle S, Berny Mier y Teran JC, Egan AN, Kisha T, Khoury CK (2019) Wild beans (*Phaseolus* L.) of North America. In 'North American Crop Wild Relatives, Vol. 2: Important Species'. (Eds SL Greene, KA Williams, CK Khoury, MB Kantar, LF Marek) pp. 99–130. (Springer: Cham, Switzerland) doi:10.1007/978-3-319-97120-9
- Doyle JJ (2012) Polyploidy in legumes. In 'Polyploidy and Genome Evolution'. (Eds. P Soltis, D Soltis) pp. 147–180. (Springer: New York, NY, USA)
- Doyle JJ (2013) The promise of genomics for a next generation of advances in higher-level legume molecular systematics. *South African Journal of Botany* **89**, 10–18. doi:10.1016/j.sajb.2013.06.012
- Doyle JJ (2016) Chasing unicorns: nodulation origins and the paradox of novelty. *American Journal of Botany* **103**, 1865–1868. doi:10.3732/ajb.1600260
- Doyle JJ, Beachy RN (1985) Ribosomal gene variation in soybean (*Glycine*) and its relatives. *Theoretical and Applied Genetics* **70**, 369–376. doi:10.1007/BF00273741
- Dudchenko O, Pham M, Lui C, Batra SS, Hoeger M, Nyquist SK, Durand NC, Shamim MS, Machol I, Erskine W, Aiden EL, Kaur P (2018) Hi-C yields chromosome-length scaffolds for a legume genome, *Trifolium subterraneum*. *bioRxiv* **2018**, 473553.
- Egan AN, Doyle J (2010) A comparison of global, gene-specific, and relaxed clock methods in a comparative genomics framework: dating the polyploid history of soybean (*Glycine max*). *Systematic Biology* **59**, 534–547. doi:10.1093/sysbio/syq041
- Egan AN, Schlueter J, Spooner DM (2012) Applications of next-generation sequencing in plant biology. *American Journal of Botany* **99**, 175–185. doi:10.3732/ajb.1200020
- Egan AN, Vatanparast M, Cagle W (2016) Parsing polyphyletic *Pueraria*: delimiting distinct evolutionary lineages through phylogeny. *Molecular Phylogenetics and Evolution* **104**, 44–59. doi:10.1016/j.ympev.2016.08.001
- Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE (2011) A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS One* **6**, e19379. doi:10.1371/journal.pone.0019379
- Eriksson JS, Blanco-Pastor JL, Sousa F, Bertrand YJ, Pfeil BE (2017) A cryptic species produced by autopolyploidy and subsequent introgression involving *Medicago prostrata* (Fabaceae). *Molecular Phylogenetics and Evolution* **107**, 367–381. doi:10.1016/j.ympev.2016.11.020
- Eriksson JS, de Sousa F, Bertrand YJ, Antonelli A, Oxelman B, Pfeil BE (2018) Allele phasing is critical to revealing a shared allopolyploid origin of *Medicago arborea* and *M. strasseri* (Fabaceae). *BMC Evolutionary Biology* **18**, 9. doi:10.1186/s12862-018-1127-z
- Fan W, Xu JM, Wu P, Yang ZX, Lou HQ, Chen WW, Jin JF, Zheng SJ, Yang JL (2019) Alleviation by abscisic acid of AI toxicity in rice bean is not associated with citrate efflux but depends on ABI5-mediated signal transduction pathways. *Journal of Integrative Plant Biology* **61**, 140–154. doi:10.1111/jipb.12695
- Foyer CH, Lam HM, Nguyen HT, Siddique KH, Varshney RK, Colmer TD, Cowling W, Bramley H, Mori TA, Hodgson JM, Cooper JW, Miller AJ, et al. (2016) Neglecting legumes has compromised human health and sustainable food production. *Nature Plants* **2**, 16112. doi:10.1038/nplants.2016.112
- Gailing O, Staton ME, Lane T, Schlarbaum SE, Nipper R, Owusu SA, Carlson JE (2017) Construction of a framework genetic linkage map in *Gleditsia triacanthos* L. *Plant Molecular Biology Reporter* **35**, 177–187. doi:10.1007/s11105-016-1012-0
- Gao F, Wang X, Li X, Xu M, Li H, Abila M, Sun H, Wei S, Feng J, Zhou Y (2018) Long-read sequencing and de novo genome assembly of *Ammopiptanthus nanus*, a desert shrub. *GigaScience* **7**, 1–5. doi:10.1093/gigascience/giy074
- Gnirke A, Melnikov A, Maguire J, Rogov P, LeProust EM, Brockman W, Fennell T, Giannoukos G, Fisher S, Russ C, Gabriel S (2009) Solution hybrid selection with ultra-long oligonucleotides for massively parallel targeted sequencing. *Nature Biotechnology* **27**, 182–189. doi:10.1038/nbt.1523
- Graham PH, Vance CP (2003) Legumes: importance and constraints to greater use. *Plant Physiology* **131**, 872–877. doi:10.1104/pp.017004
- Grando C (2015) Mating system of *Piptadenia gonoacantha* (Mart) Macbr (Fabaceae) and genetic diversity at forest restoration sites. PhD dissertation, UNICAMP, Campinas, Sao Paulo, Brazil.
- Griesmann M, Chang Y, Liu X, Song Y, Haberer G, Crook MB, Billault-Penneteau B, Laressergues D, Keller J, Imanishi L, Roswanjaya YP, Kohlen W, et al. (2018) Phylogenomics reveals multiple losses of nitrogen-fixing root nodule symbiosis. *Science* **361**, eaat1743. doi:10.1126/science.aat1743
- Grillo MA, De Mita S, Burke PV, Solórzano-Lowell KL, Heath KD (2016) Intrapopulation genomics in a model mutualist: population structure and candidate symbiosis genes under selection in *Medicago truncatula*. *Evolution* **70**, 2704–2717. doi:10.1111/evo.13095

- Grover CE, Arick MA, Thrash A, Conover JL, Sanders WS, Peterson DG, Frelichowski JE, Scheffler JA, Scheffler BE, Wendel JF (2019) Insights into the evolution of the New World diploid cottons (*Gossypium*, subgenus *Houzingenia*) based on genome sequencing. *Genome Biology and Evolution* **11**, 53–71. doi:10.1093/gbe/evy256
- Gugger PF, Liang CT, Sork VL, Hodgskiss P, Wright JW (2018) Applying landscape genomic tools to forest management and restoration of Hawaiian koa (*Acacia koa*) in a changing environment. *Evolutionary Applications* **11**, 231–242. doi:10.1111/eva.12534
- Hane JK, Ming Y, Kamphuis LG, Nelson MN, Garg G, Atkins CA, Bayer PE, Bravo A, Bringans S, Cannon S, Edwards D, Foley R, *et al.* (2017) A comprehensive draft genome sequence for lupin (*Lupinus angustifolius*), an emerging health food: insights into plant–microbe interactions and legume evolution. *Plant Biotechnology Journal* **15**, 318–330. doi:10.1111/pbi.12615
- Harouna DV, Venkataramana PB, Ndakidemi PA, Matemu AO (2018) Under-exploited wild *Vigna* species potentials in human and animal nutrition: a review. *Global Food Security* **18**, 1–11. doi:10.1016/j.gfs.2018.06.002
- Harrison TL, Wood CW, Borges IL, Stinchcombe JR (2017) No evidence for adaptation to local rhizobial mutualists in the legume *Medicago lupulina*. *Ecology and Evolution* **7**, 4367–4376. doi:10.1002/ece3.3012
- Harrison JG, Forister ML, Mcknight SR, Nordin E, Parchman TL (2019) Rarity does not limit genetic variation or preclude subpopulation structure in the geographically restricted desert forb *Astragalus lentiginosus* var. *piscinensis*. *American Journal of Botany* **106**(2), 260–269. doi:10.1002/ajb2.1235
- Haynsen MS, Vatanparast M, Mahadwar G, Zhu D, Moger-Reischer RZ, Doyle JJ, Crandall KA, Egan AN (2018) *De novo* transcriptome assembly of *Pueraria montana* var. *lobata* and *Neustanthus phaseoloides* kudzu transcriptomes for the development of eSSR and SNP markers: narrowing the US origin(s) of the invasive kudzu. *BMC Genomics* **19**, 439. doi:10.1186/s12864-018-4798-3
- Helliwell EE, Faber-Hammond J, Lopez ZC, Garoutte A, von Wettberg E, Friesen ML, Porter SS (2018) Rapid establishment of a flowering cline in *Medicago polymorpha* after invasion of North America. *Molecular Ecology* **27**, 4758–4774. doi:10.1111/mec.14898
- Hiiesalu I, Öpik MA, Metsis M, Lilje L, Davison J, Vasar M, Moora M, Zobel M, Wilson SD, Pärtel M (2012) Plant species richness belowground: higher richness and new patterns revealed by next-generation sequencing. *Molecular Ecology* **21**, 2004–2016. doi:10.1111/j.1365-294X.2011.05390.x
- Hirakawa H, Kaur P, Shirasawa K, Nichols P, Nagano S, Appels R, Erskine W, Isobe SN (2016) Draft genome sequence of subterranean clover, a reference for genus *Trifolium*. *Scientific Reports* **6**, 30358. doi:10.1038/srep30358
- Hodel RG, Gitzendanner MA, Germain-Aubrey CC, Liu X, Crowl AA, Sun M, Landis JB, Segovia-Salcedo MC, Douglas NA, Chen S, Soltis DE (2016) A new resource for the development of SSR markers: millions of loci from a thousand plant transcriptomes. *Applications in Plant Sciences* **4**, 1600024. doi:10.3732/apps.1600024
- Hollingsworth PM, Li DZ, van der Bank M, Twyford AD (2016) Telling plant species apart with DNA: from barcodes to genomes. *Philosophical Transactions of the Royal Society of London – B. Biological Sciences* **371**, 20150338. doi:10.1098/rstb.2015.0338
- Honorio Coronado EN, Blanc-Jolivet C, Mader M, García-Dávila CR, Sebbenn AM, Meyer-Sand BR, Paredes-Villanueva K, Tysklind N, Troispoux V, Massot M, Degen B (2019) Development of nuclear and plastid SNP markers for genetic studies of *Dipteryx* tree species in Amazonia. *Conservation Genetics Resources* **11**(3), 333. doi:10.1007/s12686-019-01081-3
- Hughes C, Eastwood R (2006) Island radiation on a continental scale: exceptional rates of plant diversification after uplift of the Andes. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 10334–10339. doi:10.1073/pnas.0601928103
- Hughes CE, Nyffeler R, Linder HP (2015) Evolutionary plant radiations: where, when, why and how? *New Phytologist* **207**, 249–253. doi:10.1111/nph.13523
- International Human Genome Sequencing Consortium (2001) Initial sequencing and analysis of the human genome. *Nature* **409**, 860–921. doi:10.1038/35057062
- Ištvánec J, Jaroš M, Křenek A, Repková J (2014) Genome assembly and annotation for red clover (*Trifolium pratense*; Fabaceae). *American Journal of Botany* **101**, 327–337. doi:10.3732/ajb.1300340
- Jain M, Misra G, Patel RK, Priya P, Jhanwar S, Khan AW, Shah N, Singh VK, Garg R, Jeena G, Yadav M, Kant C, *et al.* (2013) A draft genome sequence of the pulse crop chickpea (*Cicer arietinum* L.). *The Plant Journal* **74**, 715–729. doi:10.1111/tbj.12173
- Jiang Z, Wang H, Michal JJ, Zhou X, Liu B, Woods LC, Fuchs RA (2016) Genome wide sampling sequencing for SNP genotyping: methods, challenges and future development. *International Journal of Biological Sciences* **12**, 100–108. doi:10.7150/ijbs.13498
- Jiao WB, Schneeberger K (2017) The impact of third generation genomic technologies on plant genome assembly. *Current Opinion in Plant Biology* **36**, 64–70. doi:10.1016/j.pbi.2017.02.002
- Jimenez-Madriral JP (2018) Next-generation sequencing technologies in tree improvement and conservation genetics of *Dipteryx oleifera* Benth. PhD dissertation, North Carolina State University, Raleigh, NC, USA.
- Johnson MG, Pokorny L, Dodsworth S, Botigué LR, Cowan RS, Devault A, Eiserhardt WL, Epitawalage N, Forest F, Kim JT, Leebens-Mack JH, Leitch IJ, *et al.* (2019) A universal probe set for targeted sequencing of 353 nuclear genes from any flowering plant designed using k-medoids clustering. *Systematic Biology* **68**, 594–606. doi:10.1093/sysbio/syy086
- Kadlec M, Bellstedt DU, Le Maitre NC, Pirie MD (2017) Targeted NGS for species level phylogenomics: ‘made to measure’ or ‘one size fits all’? *PeerJ* **5**, e3569. doi:10.7717/peerj.3569
- Kamutando CN, Vikram S, Kamgan-Nkuekam G, Makhalanyane TP, Greve M, Le Roux JJ, Richardson DM, Cowan D, Valverde A (2017) Soil nutritional status and biogeography influence rhizosphere microbial communities associated with the invasive tree *Acacia dealbata*. *Scientific Reports* **7**, 6472. doi:10.1038/s41598-017-07018-w
- Kang YJ, Kim SK, Kim MY, Lestari P, Kim KH, Ha BK, Jun TH, Hwang WJ, Lee T, Lee J, Shim S, Yoon MY, *et al.* (2014) Genome sequence of mungbean and insights into evolution within *Vigna* species. *Nature Communications* **5**, 5443. doi:10.1038/ncomms6443
- Kang YJ, Satyawati D, Shim S, Lee T, Lee J, Hwang WJ, Kim SK, Lestari P, Laosati K, Kim KH, Ha TJ, Chitikineni A, *et al.* (2015) Draft genome sequence of adzuki bean, *Vigna angularis*. *Scientific Reports* **5**, 8069. doi:10.1038/srep08069
- Kang SH, Lee HO, Ahn BO, Kim CK (2019a) The complete mitochondrial genome sequence of *Senna occidentalis* (Fabales: Fabaceae). *Mitochondrial DNA – B. Resources* **4**, 85–86. doi:10.1080/23802359.2018.1536472
- Kang SH, Won SY, Kim CK (2019b) The complete mitochondrial genome sequences of *Senna tora* (Fabales: Fabaceae). *Mitochondrial DNA – B. Resources* **4**, 1283–1284. doi:10.1080/23802359.2019.1591227
- Kaur P, Bayer PE, Milec Z, Vrána J, Yuan Y, Appels R, Edwards D, Batley J, Nichols P, Erskine W, Doležel J (2017) An advanced reference genome of *Trifolium subterraneum* L. reveals genes related to agronomic performance. *Plant Biotechnology Journal* **15**, 1034–1046. doi:10.1111/pbi.12697
- Kazakoff SH, Imelfort M, Edwards D, Koehorst J, Biswas B, Batley J, Scott PT, Gresshoff PM (2012) Capturing the biofuel wellhead and powerhouse: the chloroplast and mitochondrial genomes of the leguminous feedstock tree *Pongamia pinnata*. *PLoS One* **7**, e51687. doi:10.1371/journal.pone.0051687

- Kazempour Osaloo S, Maassoumi AA, Murakami N (2003) Molecular systematics of the genus *Astragalus* L. (Fabaceae): phylogenetic analyses of nuclear ribosomal DNA internal transcribed spacers and chloroplast gene *ndhF* sequences. *Plant Systematics and Evolution* **242**, 1–32. doi:10.1007/s00606-003-0014-1
- Kellar PR, Ahrensden DL, Aust SK, Jones AR, Pires JC (2015) Biodiversity comparison among phylogenetic diversity metrics and between three North American prairies. *Applications in Plant Sciences* **3**, 1400108. doi:10.3732/apps.1400108
- Keller J, Imperial J, Ruiz-Argüeso T, Privet K, Lima O, Michon-Coudouel S, Biget M, Salmon A, Ainouche A, Cabello-Hurtado F (2018) RNA sequencing and analysis of three *Lupinus* nodulomes provide new insights into specific host-symbiont relationships with compatible and incompatible *Bradyrhizobium* strains. *Plant Science* **266**, 102–116. doi:10.1016/j.plantsci.2017.10.015
- Kim MY, Lee S, Van K, Kim TH, Jeong SC, Choi IY, Kim DS, Lee YS, Park D, Ma J, Kim WY, Kim BC, *et al.* (2010) Whole-genome sequencing and intensive analysis of the undomesticated soybean (*Glycine soja* Sieb. and Zucc.) genome. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 22032–22037. doi:10.1073/pnas.1009526107
- Koboldt DC, Steinberg KM, Larson DE, Wilson RK, Mardis ER (2013) The next-generation sequencing revolution and its impact on genomics. *Cell* **155**, 27–38. doi:10.1016/j.cell.2013.09.006
- Koehn EJ, de Vos JM, Atchison GW, Simon MF, Schrire BD, de Souza ER, de Queiroz LP, Hughes CE (2013) Exploring the tempo of species diversification in legumes. *South African Journal of Botany* **89**, 19–30. doi:10.1016/j.sajb.2013.07.005
- Koehn EJ, Ojeda DI, Steeves R, Migliore J, Bakker FT, Wieringa JJ, Kidner C, Hardy O, Pennington RT, Herendeen PS, Bruneau A (2019) The origin and early evolution of the legumes are a complex paleopolyploid phylogenomic tangle closely associated with the Cretaceous–Paleogene (K–Pg) boundary. *bioRxiv* **2019**, 577957.
- Kovar L, Nageswara-Rao M, Ortega-Rodríguez S, Dugas DV, Straub S, Cronn R, Strickler SR, Hughes CE, Hanley KA, Rodríguez DN, Langhorst BW, Dimalanta ET, Bailey CD (2018) PacBio.-based mitochondrial genome assembly of *Leucaena trichandra* (Leguminosae) and an intragenomic assessment of mitochondrial RNA editing. *Genome Biology and Evolution* **10**, 2501–2517. doi:10.1093/gbe/evy179
- Kovi MR, Amdahl H, Alsheikh M, Rognli OA (2017) *De novo* and reference transcriptome assembly of transcripts expressed during flowering provide insight into seed setting in tetraploid red clover. *Scientific Reports* **7**, 44383. doi:10.1038/srep44383
- Kundu A, Paul S, Dey A, Pal A (2017) High throughput sequencing reveals modulation of microRNAs in *Vigna mungo* upon mungbean yellow mosaic India virus inoculation highlighting stress regulation. *Plant Science* **257**, 96–105. doi:10.1016/j.plantsci.2017.01.016
- Le Quéré A, Tak N, Gehlot HS, Lavire C, Meyer T, Chapulliot D, Rath S, Sakrouhi I, Rocha G, Rohmer M, Severac D, Filali-Maltouf A, Munive JA (2017) Genomic characterization of *Ensifer aridi*, a proposed new species of nitrogen-fixing rhizobium recovered from Asian, African and American deserts. *BMC Genomics* **18**, 85. doi:10.1186/s12864-016-3447-y
- Le Roux JJ, Keet JH, Mutiti B, Ellis AG (2017) Cultivation may not dramatically alter rhizobial community diversity or structure associated with rooibos tea (*Aspalathus linearis* Burm. f.) in South Africa. *South African Journal of Botany* **110**, 87–96. doi:10.1016/j.sajb.2017.01.014
- Legume Phylogeny Working Group (2013a) Legume phylogeny and classification in the 21st century: progress, prospects and lessons for other species-rich clades. *Taxon* **62**, 217–248. doi:10.12705/622.8
- Legume Phylogeny Working Group (2013b) Towards a new classification system for legumes: progress report from the 6th international legume conference. *South African Journal of Botany* **89**, 3–9. doi:10.1016/j.sajb.2013.07.022
- Legume Phylogeny Working Group (2017) A new subfamily classification of the Leguminosae based on a taxonomically comprehensive phylogeny. *Taxon* **66**, 44–77. doi:10.12705/661.3
- Lestari P, Kang YJ, Han KS, Gwag JG, Moon JK, Kim YH, Lee YH, Lee SH (2014) Genome-wide single nucleotide polymorphism discovery and validation in adzuki bean. *Molecular Breeding* **33**, 497–501. doi:10.1007/s11032-013-9962-5
- Lewis G, Schrire B, Mackinder B, Lock M (2005) 'Legumes of the World.' (Royal Botanic Gardens, Kew: London, UK)
- Li XW, Yang Y, Henry RJ, Rossetto M, Wang YT, Chen SL (2015) Plant DNA barcoding: from gene to genome. *Biological Reviews of the Cambridge Philosophical Society* **90**, 157–166. doi:10.1111/brv.12104
- Liu Q, Chang S, Hartman GL, Domier LL (2018) Assembly and annotation of a draft genome sequence for *Glycine latifolia*, a perennial wild relative of soybean. *The Plant Journal* **95**, 71–85. doi:10.1111/tbj.13931
- Liu H, Wei J, Yang T, Mu W, Song B, Yang T, Fu Y, Wang X, Hu G, Li W, Zhou H, Chang Y, Chen X, Liang X (2019) Molecular digitization of a botanical garden: high-depth whole genome sequencing of 689 vascular plant species from the Ruili Botanical Garden. *GigaScience* **8**, giz007. doi:10.1093/gigascience/giz007
- Lonardi S, Muñoz-Amatriain M, Liang Q, Shu S, Wanamaker SI, Lo S, Tanskanen J, Schulman AH, Zhu T, Luo MC, Alhakami H (2019) The genome of cowpea (*Vigna unguiculata* [L.] Walp.). *The Plant Journal* **98**, 767–782. doi:10.1111/tbj.14349
- Lu Q, Li H, Hong Y, Zhang G, Wen S, Li X, Zhou G, Li S, Liu H, Liu H, Liu Z, Varshney RK, Chen X, Liang X (2018) Genome sequencing and analysis of the peanut B-genome progenitor (*Arachis ipaensis*). *Frontiers in Plant Science* **9**, 604. doi:10.3389/fpls.2018.00604
- Mahato AK, Sharma AK, Sharma TR, Singh NK (2018) An improved draft of the pigeonpea (*Cajanus cajan* (L.) Millsp.) genome. *Data in Brief* **16**, 376–380. doi:10.1016/j.dib.2017.11.066
- Malyshev LI (2008) Phenetics of subgenera and sections in the genus *Oxytropis* DC. (Fabaceae) bearing on ecology and phylogeny. *Contemporary Problems of Ecology* **1**, 440–444. doi:10.1134/S1995425508040073
- Manzanilla V, Bruneau A (2012) Phylogeny reconstruction in the Caesalpinieae grade (Leguminosae) based on duplicated copies of the sucrose synthase gene and plastid markers. *Molecular Phylogenetics and Evolution* **65**, 149–162. doi:10.1016/j.ympev.2012.05.035
- Marques A, Moraes L, Aparecida dos Santos M, Costa I, Costa L, Nunes T, Melo N, Simon MF, Leitch AR, Almeida C, Souza G (2018) Origin and parental genome characterization of the allotetraploid *Stylosanthes scabra* Vogel (Papilionoideae, Leguminosae), an important legume pasture crop. *Annals of Botany* **122**, 1143–1159. doi:10.1093/aob/mcy113
- Marris E (2008) Soya genome sequenced. *NATNews* **2008**, 1294. [Published online 10 December 2008] doi:10.1038/news.2008.1294
- Masuta Y, Kawabe A, Nozawa K, Naito K, Kato A, Ito H (2018) Characterization of a heat-activated retrotransposon in *Vigna angularis*. *Breeding Science* **68**, 168–176. doi:10.1270/jsbbs.17085
- Matasci N, Hung LH, Yan Z, Carpenter EJ, Wickett NJ, Mirarab S, Nguyen N, Warnow T, Ayyampalayam S, Barker M, Burleigh JG (2014) Data access for the 1000 Plants (1KP) project. *GigaScience* **3**, 17. doi:10.1186/2047-217X-3-17
- McClellan PE, Burrige J, Beebe S, Rao IM, Porch TG (2011) Crop improvement in the era of climate change: an integrated, multi-disciplinary approach for common bean (*Phaseolus vulgaris*). *Functional Plant Biology* **38**, 927–933. doi:10.1071/FP11102
- McKenna P, Cannon N, Conway J, Dooley J, Davies WP (2018) Red clover (*Trifolium pratense*) in conservation agriculture: a compelling case for increased adoption. *International Journal of Agricultural Sustainability* **16**, 342–366. doi:10.1080/14735903.2018.1498442

- Meek MH, Larson WA (2019) The future is now: amplicon sequencing and sequence capture usher in the conservation genomics era. *Molecular Ecology Resources* **19**, 795–803. doi:10.1111/1755-0998.12998
- Mensous M, Van de Paer C, Manzi S, Bouchez O, Baâli-Cherif D, Besnard G (2017) Diversity and evolution of plastomes in Saharan mimosoids: potential use for phylogenetic and population genetic studies. *Tree Genetics & Genomes* **13**, 48. doi:10.1007/s11295-017-1131-2
- Miller MR, Dunham JP, Amores A, Cresko WA, Johnson EA (2007) Rapid and cost-effective polymorphism identification and genotyping using restriction site associated DNA (RAD) markers. *Genome Research* **17**, 240–248. doi:10.1101/gr.5681207
- Mochida K, Sakurai T, Seki H, Yoshida T, Takahagi K, Sawai S, Uchiyama H, Muranaka T, Saito K (2017) Draft genome assembly and annotation of *Glycyrrhiza uralensis*, a medicinal legume. *The Plant Journal* **89**, 181–194. doi:10.1111/tpj.13385
- Moore AJ, Vos JMD, Hancock LP, Goolsby E, Edwards EJ (2018) Targeted enrichment of large gene families for phylogenetic inference: phylogeny and molecular evolution of photosynthesis genes in the portulugo clade (Caryophyllales). *Systematic Biology* **67**, 367–383. doi:10.1093/sysbio/syx078
- Morris AB, Scalf C, Burleyson A, Johnson LT, Trostel K (2016) Development and characterization of microsatellite primers in the federally endangered *Astragalus bibullatus* (Fabaceae). *Applications in Plant Sciences* **4**, 1500126. doi:10.3732/apps.1500126
- Moura TM, Vatanparast M, Tozzi AMGA, Forest F, Wilmot-Dear MC, Simon MF, Mansano VF, Kajita T, Lewis GP (2016) A molecular phylogeny and new infrageneric classification of *Mucuna* Adans. (Leguminosae–Papilionoideae) including insights from morphology and hypotheses about biogeography. *International Journal of Plant Sciences* **177**, 76–89. doi:10.1086/684131
- Mousavi-Derazmahalleh M, Bayer PE, Hane JK, Valliyodan B, Nguyen HT, Nelson MN, Erskine W, Varshney RK, Papa R, Edwards D (2019) Adapting legume crops to climate change using genomic approaches. *Plant, Cell & Environment* **42**, 6–19. doi:10.1111/pce.13203
- Naito K, Kaga A, Tomooka N, Kawase M (2013) *De novo* assembly of the complete organelle genome sequences of azuki bean (*Vigna angularis*) using next-generation sequencers. *Breeding Science* **63**, 176–182. doi:10.1270/jsbbs.63.176
- Neale DB, Kremer A (2011) Forest tree genomics: growing resources and applications. *Nature Reviews. Genetics* **12**, 111. doi:10.1038/nrg2931
- Negrak V (2013) Mitochondrial genome sequence of the legume *Vicia faba*. *Frontiers in Plant Science* **4**, 128. doi:10.3389/fpls.2013.00128
- Nevado B, Atchison GW, Hughes CE, Filatov DA (2016) Widespread adaptive evolution during repeated evolutionary radiations in New World lupins. *Nature Communications* **7**, 12384. doi:10.1038/ncomms12384
- Nevado B, Contreras-Ortiz N, Hughes C, Filatov DA (2018) Pleistocene glacial cycles drive isolation, gene flow and speciation in the high-elevation Andes. *New Phytologist* **219**, 779–793. doi:10.1111/nph.15243
- Nicholls JA, Pennington RT, Koenen EJ, Hughes CE, Hearn J, Bunnefeld L, Dexter KG, Stone GN, Kidner CA (2015) Using targeted enrichment of nuclear genes to increase phylogenetic resolution in the neotropical rain forest genus *Inga* (Leguminosae: Mimosoideae). *Frontiers in Plant Science* **6**, 710. doi:10.3389/fpls.2015.00710
- Ojeda DI, Koenen E, Cervantes S, de la Estrella M, Banguera-Hinestroza E, Janssens SB, Migliore J, Demenou B, Bruneau A, Forest F, Hardy OJ (2019) Phylogenomic analyses reveal an exceptionally high number of evolutionary shifts in a florally diverse clade of African legumes. *Molecular Phylogenetics and Evolution* **137**, 156–167. doi:10.1016/j.ympev.2019.05.002
- Oliveira-Filho AT, Cardoso D, Schrire BD, Lewis GP, Pennington RT, Brummer TJ, Rotella J, Lavin M (2013) Stability structures tropical woody plant diversity more than seasonality: insights into the ecology of high legume–succulent-plant biodiversity. *South African Journal of Botany* **89**, 42–57. doi:10.1016/j.sajb.2013.06.010
- Palmer JD, Herbon LA (1988) Plant mitochondrial DNA evolved rapidly in structure but slowly in sequence. *Journal of Molecular Evolution* **28**, 87–97. doi:10.1007/BF02143500
- Palmer JD, Jorgensen RA, Thompson WF (1985) Chloroplast DNA variation and evolution in *Pisum*: patterns of change and phylogenetic analysis. *Genetics* **109**, 195–213.
- Pan L, Wang N, Wu Z, Guo R, Yu X, Zheng Y, Xia Q, Gui S, Chen C (2017) A high density genetic map derived from RAD sequencing and its application in QTL analysis of yield-related traits in *Vigna unguiculata*. *Frontiers in Plant Science* **8**, 1544. doi:10.3389/fpls.2017.01544
- Pang T, Ye CY, Xia X, Yin W (2013) *De novo* sequencing and transcriptome analysis of the desert shrub, *Ammopiptanthus mongolicus*, during cold acclimation using Illumina/Solexa. *BMC Genomics* **14**, 488. doi:10.1186/1471-2164-14-488
- Papadopoulou A, Taberlet P, Zinger L (2015) Metagenome skimming for phylogenetic community ecology: a new era in biodiversity research. *Molecular Ecology* **24**, 3515–3517. doi:10.1111/mec.13263
- Podlech D, Zarre S, Ekici M, Maassoumi AA, Sytin A (2014) A taxonomic revision of the genus *Astragalus* L. (Leguminosae) in the Old World. *Annalen des Naturhistorischen Museums in Wien – B. Botanik und Zoologie* **1**, 106
- Porch T, Beaver J, Debouck D, Jackson S, Kelly J, Dempewolf H (2013) Use of wild relatives and closely related species to adapt common bean to climate change. *Agronomy (Basel)* **3**, 433–461. doi:10.3390/agronomy3020433
- Ramirez-Villegas J, Khoury C, Jarvis A, Debouck DG, Guarino L (2010) A gap analysis methodology for collecting crop gene pools: a case study with *Phaseolus* beans. *PLoS One* **5**, e13497. doi:10.1371/journal.pone.0013497
- Reuter JA, Spacek DV, Snyder MP (2015) High-throughput sequencing technologies. *Molecular Cell* **58**, 586–597. doi:10.1016/j.molcel.2015.05.004
- Rice DW, Alverson AJ, Richardson AO, Young GJ, Sanchez-Puerta MV, Munzinger J, Barry K, Boore JL, Zhang Y, Knox EB, Palmer JD (2013) Horizontal transfer of entire genomes via mitochondrial fusion in the angiosperm *Amborella*. *Science* **342**, 1468–1473. doi:10.1126/science.1246275
- Richardson JE, Pennington TR, Pennington TD, Hollingsworth PM (2001) Rapid diversification of a species-rich genus of neotropical rain forest trees. *Science* **293**, 2242–2245. doi:10.1126/science.1061421
- Ruhfel BR, Gitzendanner MA, Soltis PS, Soltis DE, Burleigh JG (2014) From algae to angiosperms: inferring the phylogeny of green plants (Viridiplantae) from 360 plastid genomes. *BMC Evolutionary Biology* **14**, 23. doi:10.1186/1471-2148-14-23
- Sanchez-Puerta M, García LE, Wohlfeiler J, Ceriotti LF (2017) Unparalleled replacement of native mitochondrial genes by foreign homologs in a holoparasitic plant. *New Phytologist* **214**, 376–387. doi:10.1111/nph.14361
- Sanchez-Puerta MV, Edera A, Gandini CL, Williams AV, Howell KA, Nevill PG, Small I (2019) Genome-scale transfer of mitochondrial DNA from legume hosts to the holoparasite *Lophophytum mirabile* (Balanophoraceae). *Molecular Phylogenetics and Evolution* **132**, 243–250. doi:10.1016/j.ympev.2018.12.006
- Sanderson MJ, Wojciechowski MF (1996) Diversification rates in a temperate legume clade: are there ‘so many species’ of *Astragalus* (Fabaceae)? *American Journal of Botany* **83**, 1488–1502. doi:10.1002/j.1537-2197.1996.tb13942.x
- Sanger F, Nicklen S, Coulson AR (1977) DNA sequencing with chain-terminating inhibitors. *Proceedings of the National Academy of Sciences of the United States of America* **74**, 5463–5467. doi:10.1073/pnas.74.12.5463

- Sathyanarayana N, Pittala RK, Tripathi PK, Chopra R, Singh HR, Belamkar V, Bhardwaj PK, Doyle JJ, Egan AN (2017) Transcriptomic resources for the medicinal legume *Mucuna pruriens*: *de novo* transcriptome assembly, annotation, identification and validation of EST–SSR markers. *BMC Genomics* **18**, 409. doi:10.1186/s12864-017-3780-9
- Sato S, Nakamura Y, Kaneko T, Asamizu E, Kato T, Nakao M, Sasamoto S, Watanabe A, Ono A, Kawashima K, Fujishiro T, Katoh M, *et al.* (2008) Genome structure of the legume *Lotus japonicus*. *DNA Research* **15**, 227–239. doi:10.1093/dnares/dsn008
- Schafleitner R, Huang SM, Chu SH, Yen JY, Lin CY, Yan MR, Krishnan B, Liu MS, Lo HF, Chen CY, Chen L-FO, Wu DC, Thi Bui TG, Ramasamy S, Tung CW, Nair R (2016) Identification of single nucleotide polymorphism markers associated with resistance to bruchids (*Callosobruchus* spp.) in wild mungbean (*Vigna radiata* var. *sublobata*) and cultivated *V. radiata* through genotyping by sequencing and quantitative trait locus analysis. *BMC Plant Biology* **16**, 159. doi:10.1186/s12870-016-0847-8
- Scherson RA, Choi HK, Cook DR, Sanderson MJ (2005) Phylogenetics of New World *Astragalus*: screening of novel nuclear loci for the reconstruction of phylogenies at low taxonomic levels. *Brittonia* **57**, 354–366. doi:10.1663/0007-196X(2005)057[0354:PONWAS]2.0.CO;2
- Scherson RA, Vidal R, Sanderson MJ (2008) Phylogeny, biogeography and rates of diversification of New World *Astragalus* (Leguminosae) with an emphasis on South American radiations. *American Journal of Botany* **95**, 1030–1039. doi:10.3732/ajb.0800017
- Schmutz J, Cannon SB, Schlueter J, Ma J, Mitros T, Nelson W, Hyten DL, Song Q, Thelen JJ, Cheng J, Xu D, Hellsten U, *et al.* (2010) Genome sequence of the palaeopolyploid soybean. *Nature* **463**, 178–183. doi:10.1038/nature08670
- Schmutz J, McClean PE, Mamidi S, Wu GA, Cannon SB, Grimwood J, Jenkins J, Shu S, Song Q, Chavarro C, Torres-Torres M, Geffroy V, *et al.* (2014) A reference genome for common bean and genome-wide analysis of dual domestications. *Nature Genetics* **46**, 707–713. doi:10.1038/ng.3008
- Schwarz EN, Ruhlman TA, Sabir JSM, Hajrah NH, Alharbi NS, Al-Malki AL, Bailey CD, Jansen RK (2015) Plastid genome sequences of legumes reveal parallel inversions and multiple losses of *rps16* in papilionoids. *Journal of Systematics and Evolution* **53**, 458–468. doi:10.1111/jse.12179
- Shahi Shavvon R, Kazempour Osaloo S, Maassoumii AA, Moharrek F, Karaman Erkul S, Lemmon AR, Lemmon EM, Michalak I, Muellner-Riehl AN, Favre A (2017) Increasing phylogenetic support for explosively radiating taxa: the promise of high-throughput sequencing for *Oxytropis* (Fabaceae). *Journal of Systematics and Evolution* **55**, 385–404. doi:10.1111/jse.12269
- Shen Y, Liu J, Geng H, Zhang J, Liu Y, Zhang H, Xing S, Du J, Ma S, Tian Z (2018) *De novo* assembly of a Chinese soybean genome. *Science China. Life Sciences* **61**, 871–884. doi:10.1007/s11427-018-9360-0
- Sherman-Broyles S, Bombarely A, Doyle J (2017) Characterizing the allopolyploid species among the wild relatives of soybean: utility of reduced representation genotyping methodologies. *Journal of Systematics and Evolution* **55**, 365–376. doi:10.1111/jse.12268
- Shi Y, Liu Y, Zhang S, Zou R, Tang J, Mu W, Peng Y, Dong S (2018) Assembly and comparative analysis of the complete mitochondrial genome sequence of *Sophora japonica* ‘JinhuiJ2’. *PLoS One* **13**, e0202485. doi:10.1371/journal.pone.0202485
- Simon MF, Grether R, de Queiroz LP, Skema C, Pennington RT, Hughes CE (2009) Recent assembly of the Cerrado, a neotropical plant diversity hotspot, by *in situ* evolution of adaptations to fire. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 20359–20364. doi:10.1073/pnas.0903410106
- Singh NK, Gupta DK, Jayaswal PK, Mahato AK, Dutta S, Singh S, Bhutani S, Dogra V, Singh BP, Kumawat G, Pal JK, Pandit A, *et al.* (2012) The first draft of the pigeonpea genome sequence. *Journal of Plant Biochemistry and Biotechnology* **21**, 98–112. doi:10.1007/s13562-011-0088-8
- Snak C, Vatanparast M, Silva C, Lewis GP, Lavin M, Kajita T, de Queiroz LP (2016) A dated phylogeny of the papilionoid legume genus *Canavalia* reveals recent diversification by a pantropical liana lineage. *Molecular Phylogenetics and Evolution* **98**, 133–146. doi:10.1016/j.ympev.2016.02.001
- Soares TN, Melo DB, Resende LV, Vianello RP, Chaves LJ, Collevatti RG, Telles MP (2012) Development of microsatellite markers for the neotropical tree species *Dipteryx alata* (Fabaceae). *American Journal of Botany* **99**, e72–e73. doi:10.3732/ajb.1100377
- Soltis DE, Gitzendanner MA, Stull G, Chester M, Chandlerbali A, Chamala S, Jordon-Thaden I, Soltis PS, Schnable PS, Barbazuk BW (2013) The potential of genomics in plant systematics. *Taxon* **62**, 886–898. doi:10.12705/625.13
- Soltis DE, Moore MJ, Sessa EB, Smith SA, Soltis PS (2018) Using and navigating the plant tree of life. *American Journal of Botany* **105**, 287–290. doi:10.1002/ajb2.1071
- Spehn EM, Scherer-Lorenzen M, Schmid B, Hector A, Caldeira MC, Dimitrakopoulos PG, Finn JA, Jumpponen A, O’Donovan G, Pereira JS, Schulze ED, Troumbis AY, Körner C (2002) The role of legumes as a component of biodiversity in a cross-European study of grassland biomass nitrogen. *Oikos* **98**, 205–218. doi:10.1034/j.1600-0706.2002.980203.x
- Stai JS, Yadav A, Sinou C, Bruneau A, Doyle JJ, Fernández-Baca D, Cannon SB (2019) *Cercis*: a non-polyploid genomic relic within the generally polyploid legume family. *Frontiers in Plant Science* **10**, 345. doi:10.3389/fpls.2019.00345
- Stefanovic S, Pfeil BE, Palmer JD, Doyle JJ (2009) Relationships among phaseoloid legumes based on sequences from eight chloroplast regions. *Systematic Botany* **34**, 115–128. doi:10.1600/036364409787602221
- Stein JC, Yu Y, Copetti D, Zwickl DJ, Zhang L, Zhang C, Chougule K, Gao D, Iwata A, Goicoechea JL, Wei S (2018) Genomes of 13 domesticated and wild rice relatives highlight genetic conservation, turnover and innovation across the genus *Oryza*. *Nature Genetics* **50**, 285–296. doi:10.1038/s41588-018-0040-0
- Steinrücken TV (2017) Investigating the cause of dieback in the invasive plant, *Parkinsonia aculeata*. PhD thesis, Western Sydney University, Sydney, NSW, Australia.
- Straub SC, Parks M, Weitemier K, Fishbein M, Cronn RC, Liston A (2012) Navigating the tip of the genomic iceberg: next-generation sequencing for plant systematics. *American Journal of Botany* **99**, 349–364. doi:10.3732/ajb.1100335
- Sveinsson S, Cronk Q (2014) Evolutionary origin of highly repetitive plastid genomes within the clover genus (*Trifolium*). *BMC Evolutionary Biology* **14**, 228. doi:10.1186/s12862-014-0228-6
- Tang H, Krishnakumar V, Bidwell S, Rosen B, Chan A, Zhou S, Gentzbitell L, Childs KL, Yandell M, Gundlach H, Mayer KF, Schwartz DC, Town CD (2014) An improved genome release (version Mt4. 0) for the model legume *Medicago truncatula*. *BMC Genomics* **15**, 312. doi:10.1186/1471-2164-15-312
- Tian R, Parker M, Seshadri R, Reddy TB, Markowitz V, Ivanova N, Pati A, Woyke T, Baeshen MN, Baeshen NA, Kyrpidis N, Reeve W (2015) High-quality permanent draft genome sequence of *Bradyrhizobium* sp. Th. b2, a microsymbiont of *Amphicarpaea bracteata* collected in Johnson City, New York. *Standards in Genomic Sciences* **10**, 24. doi:10.1186/s40793-015-0008-y
- Tosso F, Hardy OJ, Doucet JL, Daïnou K, Kaymak E, Migliore J (2018) Evolution in the amphiatlantic tropical genus *Guibourtia* (Fabaceae, Detarioideae), combining NGS phylogeny and morphology. *Molecular Phylogenetics and Evolution* **120**, 83–93. doi:10.1016/j.ympev.2017.11.026

- Van de Peer Y, Mizrahi E, Marchal K (2017) The evolutionary significance of polyploidy. *Nature Reviews. Genetics* **18**, 411–424. doi:10.1038/nrg.2017.26
- van Velzen R, Holmer R, Bu F, Rutten L, van Zeijl A, Liu W, Santuari L, Cao Q, Sharma T, Shen D, Roswanjaya Y, Wardhani TAK, *et al.* (2018) Comparative genomics of the nonlegume *Parasponia* reveals insights into evolution of nitrogen-fixing rhizobium symbioses. *Proceedings of the National Academy of Sciences of the United States of America* **115**, E4700–E4709. doi:10.1073/pnas.1721395115
- van Velzen R, Doyle JJ, Geurts R (2019) A resurrected scenario: single gain and massive loss of nitrogen-fixing nodulation. *Trends in Plant Science* **24**, 49–57. doi:10.1016/j.tplants.2018.10.005
- Varshney RK (2016) Exciting journey of 10 years from genomes to fields and markets: some success stories of genomics-assisted breeding in chickpea, pigeonpea and groundnut. *Plant Science* **242**, 98–107. doi:10.1016/j.plantsci.2015.09.009
- Varshney RK, Chen W, Li Y, Bharti AK, Saxena RK, Schlueter JA, Donoghue MT, Azam S, Fan G, Whaley AM, Farmer AD, Sheridan J, *et al.* (2012) Draft genome sequence of pigeonpea (*Cajanus cajan*) an orphan legume crop of resource-poor farmers. *Nature Biotechnology* **30**, 83–89. doi:10.1038/nbt.2022
- Varshney RK, Song C, Saxena RK, Azam S, Yu S, Sharpe AG, Cannon S, Baek J, Rosen BD, Tar'an B, Millan T, Zhang X, *et al.* (2013) Draft genome sequence of chickpea (*Cicer arietinum*) provides a resource for trait improvement. *Nature Biotechnology* **31**, 240–246. doi:10.1038/nbt.2491
- Vatanparast M, Klitgård BB, Adema FA, Pennington RT, Yahara T, Kajita T (2013) First molecular phylogeny of the pantropical genus *Dalbergia*: implications for infrageneric circumscription and biogeography. *South African Journal of Botany* **89**, 143–149. doi:10.1016/j.sajb.2013.07.001
- Vatanparast M, Shetty P, Chopra R, Doyle JJ, Sathyanarayana N, Egan AN (2016) Transcriptome sequencing and marker development in winged bean (*Psophocarpus tetragonolobus*; Leguminosae). *Scientific Reports* **6**, 29070. doi:10.1038/srep29070
- Vatanparast M, Powell A, Doyle JJ, Egan AN (2018) Targeting legume loci: a comparison of three methods for targeted enrichment baits design in Leguminosae phylogenomics. *Applications in Plant Sciences* **6**, e1036. doi:10.1002/aps3.1036
- Verdu CF, Guichoux E, Quevauvillers S, De Thier O, Laizet YH, Delcamp A, Gévaudant F, Monty A, Porté AJ, Lejeune P, Lassois L, Mariette S (2016) Dealing with paralogy in RAD seq data: *in silico* detection and single nucleotide polymorphism validation in *Robinia pseudoacacia* L. *Ecology and Evolution* **6**, 7323–7333. doi:10.1002/ece3.2466
- Wang Z, Gerstein M, Snyder M (2009) RNA-Seq: a revolutionary tool for transcriptomics. *Nature Reviews. Genetics* **10**, 57–63. doi:10.1038/nrg2484
- Wang YH, Qu XJ, Chen SY, Li DZ, Yi TS (2017) Plastomes of Mimosoideae: structural and size variation, sequence divergence, and phylogenetic implication. *Tree Genetics & Genomes* **13**, 41. doi:10.1007/s11295-017-1124-1
- Wang YH, Wicke S, Wang H, Jin JJ, Chen SY, Zhang SD, Li DZ, Yi TS (2018) Plastid genome evolution in the early-diverging legume subfamily Cercidoideae (Fabaceae). *Frontiers in Plant Science* **9**, 138. doi:10.3389/fpls.2018.00138
- Wen J, Egan AN, Dikow R, Zimmer EA (2015) Utility of transcriptome sequencing for phylogenetic inference and character evolution. In 'Next-Generation Sequencing in Plant Systematics'. (Eds. E Hörandl, MS Appelhans) pp. 51–91. (Koeltz Scientific Books: Königstein, Germany)
- Werner GD, Cornwell WK, Sprent JI, Kattge J, Kiers ET (2014) A single evolutionary innovation drives the deep evolution of symbiotic N₂-fixation in angiosperms. *Nature Communications* **5**, 4087. doi:10.1038/ncomms5087
- Wickett NJ, Mirarab S, Nguyen N, Warnow T, Carpenter E, Matasci N, Ayyampalayam S, Barker MS, Burleigh JG, Gitzendanner MA, Ruhfel BR, Wafula E, *et al.* (2014) Phylotranscriptomic analysis of the origin and early diversification of land plants. *Proceedings of the National Academy of Sciences of the United States of America* **111**, E4859–E4868. doi:10.1073/pnas.1323926111
- Wigley K, Moot D, Wakelin SA, Laugraud A, Blond C, Seth K, Ridgway H (2017) Diverse bacterial taxa inhabit root nodules of lucerne (*Medicago sativa* L.) in New Zealand pastoral soils. *Plant and Soil* **420**, 253–262. doi:10.1007/s11104-017-3395-6
- Williams AV, Miller JT, Small I, Nevill PG, Boykin LM (2016) Integration of complete chloroplast genome sequences with small amplicon datasets improves phylogenetic resolution in *Acacia*. *Molecular Phylogenetics and Evolution* **96**, 1–8. doi:10.1016/j.ympev.2015.11.021
- Wojciechowski MF, Sanderson MJ, Steele KP, Liston A (2000) Molecular phylogeny of the 'temperate herbaceous tribes' of papilionoid legumes: a supertree approach. In 'Advances in Legume Systematics, Part 9'. (Eds PS Herendeen, A Bruneau) pp. 277–298. (Royal Botanic Gardens, Kew: London, UK)
- Wojciechowski M, Lavin M, Sanderson M (2004) A phylogeny of legumes (Leguminosae) based on analysis of the plastid *matK* gene resolves many well-supported subclades within the family. *American Journal of Botany* **91**, 1846–1862. doi:10.3732/ajb.91.11.1846
- Wong MM, Gujaria-Verma N, Ramsay L, Yuan HY, Caron C, Diapari M, Vandenberg A, Bett KE (2015) Classification and characterization of species within the genus *Lens* using genotyping-by-sequencing (GBS). *PLoS One* **10**, e0122025. doi:10.1371/journal.pone.0122025
- Xie M, Chung CY, Li MW, Wong FL, Wang X, Liu A, Wang Z, Leung AK, Wong TH, Tong SW, Xiao Z, Fan K, *et al.* (2019) A reference-grade wild soybean genome. *Nature Communications* **10**, 1216. doi:10.1038/s41467-019-09142-9
- Xu P, Wu X, Muñoz-Amatrián M, Wang B, Wu X, Hu Y, Huynh BL, Close TJ, Roberts PA, Zhou W, Lu Z, Li G (2017) Genomic regions, cellular components and gene regulatory basis underlying pod length variations in cowpea (*V. unguiculata* L. Walp). *Plant Biotechnology Journal* **15**, 547–557. doi:10.1111/pbi.12639
- Yahara T, Javadi F, Onoda Y, de Queiroz LP, Faith DP, Prado DE, Akasaka M, Kadoya T, Ishihama F, Davies S, Slik JW, Yi T, *et al.* (2013) Global legume diversity assessment: concepts key indicators and strategies. *Taxon* **62**, 249–266. doi:10.12705/622.12
- Yan J, Yan H, Liu LX, Chen WF, Zhang XX, Verástegui-Valdés MM, Wang ET, Han XZ (2017) *Rhizobium hidalgonense* sp. nov., a nodule endophytic bacterium of *Phaseolus vulgaris* in acid soil. *Archives of Microbiology* **199**, 97–104. doi:10.1007/s00203-016-1281-x
- Yang H, Tao Y, Zheng Z, Zhang Q, Zhou G, Sweetingham MW, Howieson JG, Li C (2013) Draft genome sequence, and a sequence-defined genetic linkage map of the legume crop species *Lupinus angustifolius* L. *PLoS One* **8**, e64799. doi:10.1371/journal.pone.0064799
- Yang JY, Ojeda DI, Santos-Guerra A, Molina RJ, Caujapé-Castells J, Cronk Q (2018) Population differentiation in relation to conservation: nuclear microsatellite variation in the Canary Island endemic *Lotus sessilifolius* (Fabaceae). *Conservation Genetics Resources* **10**, 219–227. doi:10.1007/s12686-017-0778-1
- Yates SA, Swain MT, Hegarty MJ, Chernukin I, Lowe M, Allison GG, Ruttink T, Abberton MT, Jenkins G, Skot L (2014) *De novo* assembly of red clover transcriptome based on RNA-seq data provides insight into drought response, gene discovery and marker identification. *BMC Genomics* **15**, 453. doi:10.1186/1471-2164-15-453
- Young ND, Debellé F, Oldroyd GE, Geurts R, Cannon SB, Udvardi MK, Benedito VA, Mayer KF, Gouzy J, Schoof H, Van de Peer Y, Proost S, *et al.* (2011) The *Medicago* genome provides insight into the evolution of rhizobial symbioses. *Nature* **480**, 520–524. doi:10.1038/nature10625
- Yu HQ, Wang YG, Yong TM, She YH, Fu FL, Li WC (2014) Heterologous expression of betaine aldehyde dehydrogenase gene from *Ammopiptanthus nanus* confers high salt and heat tolerance to *Escherichia coli*. *Gene* **549**, 77–84. doi:10.1016/j.gene.2014.07.049

- Yu T, Sun L, Cui H, Liu S, Men J, Chen S, Chen Y, Lu C (2018) The complete mitochondrial genome of a tertiary relict evergreen woody plant *Ammopiptanthus mongolicus*. *Mitochondrial DNA – B. Resources* **3**, 9–11. doi:[10.1080/23802359.2017.1413301](https://doi.org/10.1080/23802359.2017.1413301)
- Zalapa JE, Cuevas H, Zhu H, Steffan S, Senalik D, Zeldin E, McCown B, Harbut R, Simon P (2012) Using next-generation sequencing approaches to isolate simple sequence repeat (SSR) loci in the plant sciences. *American Journal of Botany* **99**, 193–208. doi:[10.3732/ajb.1100394](https://doi.org/10.3732/ajb.1100394)
- Zhan L, Paterson IG, Fraser BA, Fraser BA, Watson B, Bradbury IR, Nadukkalam Ravindran P, Reznick D, Beiko RG, Bentzen P (2017) Megasat: automated inference of microsatellite genotypes from sequence data. *Molecular Ecology Resources* **17**, 247–256. doi:[10.1111/1755-0998.12561](https://doi.org/10.1111/1755-0998.12561)
- Zhang J, Chiodini R, Badr A, Zhang G (2011) The impact of next-generation sequencing on genomics. *Journal of Genetics and Genomics* **38**, 95–109. doi:[10.1016/j.jgg.2011.02.003](https://doi.org/10.1016/j.jgg.2011.02.003)
- Zhang ML, Huang JF, Sanderson SC, Yan P, Wu YH, Pan BR (2015) Molecular biogeography of tribe Thermopsidae (Leguminosae): a Madrean–Tethyan disjunction pattern with an African origin of core genistoides. *BioMed Research International* **2015**, 864804. doi:[10.1155/2015/864804](https://doi.org/10.1155/2015/864804)
- Zhang B, Jin J, Moore MJ, Yi T-S (2019) Assembly and comparative analyses of the mitochondrial genome of *Castanospermum australe* (Papilionoideae, Leguminosae). *Australian Systematic Botany* **32**(5–6), 484–494. doi:[10.1071/SB19014](https://doi.org/10.1071/SB19014)
- Zhou Y, Gao F, Liu R, Feng J, Li H (2012) *De novo* sequencing and analysis of root transcriptome using 454 pyrosequencing to discover putative genes associated with drought tolerance in *Ammopiptanthus mongolicus*. *BMC Genomics* **13**, 266. doi:[10.1186/1471-2164-13-266](https://doi.org/10.1186/1471-2164-13-266)
- Zimmer EA, Wen J (2013) Using nuclear gene data for plant phylogenetics: progress and prospects. *Molecular Phylogenetics and Evolution* **66**, 539–550. doi:[10.1016/j.ympev.2013.01.005](https://doi.org/10.1016/j.ympev.2013.01.005)
- Zimmer EA, Wen J (2015) Using nuclear gene data for plant phylogenetics: progress and prospects II: next-gen approaches. *Journal of Systematics and Evolution* **53**, 371–379. doi:[10.1111/jse.12174](https://doi.org/10.1111/jse.12174)

Handling editor: Colin Hughes