



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

22nd International Grassland Congress

PROGARDES™: A Legume for Tropical/Subtropical Semi-Arid Clay Soils

Chris Gardiner
James Cook University, Australia

Nick Kempe
Agrimix Pty Ltd., Australia

Iain Hannah
Agrimix Pty Ltd., Australia

Jim McDonald
Agrimix Pty Ltd., Australia

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1-2/10>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

PROGARDES™: a legume for tropical/subtropical semi-arid clay soils

Chris Gardiner^A, Nick Kempe^B, Iain Hannah^B and Jim McDonald^B

^A School of Veterinary Science, James Cook University, Townsville, Qld 4811 Australia

^B Agrimix Pty Ltd, 19 Chapman Place, Eagle Farm, Qld 4009 Australia

Contact email: christopher.gardiner@jcu.edu.au

Keywords: Progardes™, plant establishment, grazing and feeding trials.

Introduction

The range of available sown pasture legumes for the vast heavy clay soil regions of northern Australia has long been regarded as being deficient (Burt 1993; Jones and Clem 1997; Pengelly and Conway 2000). Indeed immense areas of northern Australia's semi-arid clay soil regions have no sown pasture legume with proven adaptation and persistence through the long annual dry seasons (Gardiner and Swan 2008). The genus *Desmanthus* is a Mimosaceae legume containing some 24 species which are native to the Americas and range from being herbaceous to suffruticose in habit (Luckow 1993). *Desmanthus* is one of the very few legumes consistently observed to persist under heavy grazing on clay soils in their native environments (Pengelly and Conway 2000). Numerous accessions of *Desmanthus* were collected and introduced into Australia by various institutions, notably CSIRO and QDPI, over the past 50 years (Reid 1983; Pengelly and Liu 2001), as potential legumes for clay soils. After years of multi-site field evaluation of *Desmanthus* and other species, in 1991 QDPI released 3 *Desmanthus* cultivars, cvv. Marc (*D. virgatus*), Bayamo (*D. leptophyllus*) and Uman (*D. pubescens*), which were marketed as a blend named "Jaribu" (Cook *et al.* 1993). Currently, only cv. Marc is available commercially with a focus on southern subtropical Queensland markets. However, Pengelly and Conway (2000) state that, owing to Marc's low dry matter production, its contribution to animal diets and soil N is limited.

Development of Progardes™

In the 1990s Chris Gardiner at James Cook University, Townsville started to survey legumes at a number of old abandoned trial sites across remote northern and central western Queensland's semi-arid clay soil regions (<500 mm AAR). He found that various *Desmanthus* accessions were the only surviving and thriving legumes of all species sown in those environments. These particular accessions had persisted for a decade or more (now more than 2 decades) and had survived grazing, drought, flooding, fire and frosts and had therefore withstood the full gambit of environmental stresses (Gardiner and Swan 2008). The selection and breeding of plants from these survivors and others from other similar old sites and their subsequent evaluation in new trials and seed multiplication has led to the development of new lines of *Desmanthus* for northern Australia and similar environments.

The best of these varieties have been released by Agrimix Pty Ltd, JCU's commercialisation partner, as a blend

named Progardes™ (www.progardes.com.au). The blend comprises new selections of the species *D. virgatus*, *D. leptophyllus* and *D. bicornutus*, giving it a broad range of early to late maturity types, plus a range of habits (herbaceous to suffruticose) and edaphic and climatic tolerances. In recent times Progardes™ has been sown in trials and demonstration areas totalling some 2000 ha in a wide range of environments including: open Downs, cleared Gidgee/Boree and Brigalow land types across Queensland. A number of sowing techniques have been used including aerial, blade ploughing, stick raking, broadcasting onto cultivated seedbeds in a buffel grass renovation and broadcasting onto unprepared native grass pasture, all with considerable success. Agrimix expects to have some 10,000 ha commercially sown in the 2012/13 season.

As with most crops and pastures, the key to successful establishment is one of timing. With Progardes™ in northern Australia planting at the end of the dry season/start of the wet season gives the establishing plants a good opportunity to come away with the sown or native grasses. Annual rainfall on our rangelands is highly variable (McKeon 2006), so the common practice is to plant a mixture of hard and soft (scarified) seed, which gives some insurance against false seasonal starts. Scarification typically enhances germination from about 10% to 70-80%. Experience is showing that 2 or 3 wet seasons may be required for the Progardes™ to become well established in the harsher drier environments such as on the Mitchell Grass Downs, but only one season in the more favourable Brigalow region of Queensland.

Grazing and feeding trials with both sheep and cattle have been undertaken. Steers grazing a buffel grass Progardes™ mixed pasture in central Queensland gained an additional 40 kg live weight over a 90-day study period compared with steers on an adjacent buffel grass-only pasture during a cool dry winter (Gardiner and Parker 2012). In a feeding study with sheep, adding Progardes™ to a Flinders grass diet resulted in a positive N balance and significantly improved intake and weight gain over that of sheep fed only Flinders grass (Ngo 2012). Concentrations of crude protein in leaf, stem and pods with seeds of Progardes™ have been recorded as being 20.2, 11.9 and 17%, respectively, while energy (ME) levels of the whole plant were 10.3 MJ/kg DM.

Commercial seed crops are being grown successfully on the Atherton Tablelands and the Burdekin Irrigation areas; some hay has also been effectively made from Progardes™. It is expected that even small amounts of Progardes™ will

improve animal production in the targeted dry environments owing to what Bell (2009) terms the marginal value of feed that such nutritious plants offer. Kretschmer and Pitman (2001) stated that as little as 10% of a legume in poor quality grass pasture would make significant impacts on animal intake and therefore production. In a 250 ha trial in central Queensland's Brigalow region, sowing Progardes™ into renovated buffel grass resulted in a legume density of 7 plants/m², with a frequency of 75% at the end of the 1st growing season. This indicates potential compatibility with buffel grass, where few legumes, as stated by Robertson *et al.* (1997), have been successful. Reid (1983) noted, when collecting *Desmanthus* in Mexico, that in certain regions on clay soils it often occurred with the introduced buffel grass. Its compatibility with buffel grass suggests too that it has potential to alleviate buffel grass run-down, a widespread problem which reduces animal production.

Conclusions

Progardes™ appears to be an appropriate legume for improving livestock production in terms of increased liveweight gains, reproductive rates, and turn-off, particularly in the semi-arid clay soil regions of northern Australia, where few, if any, other herbaceous sown pasture legumes are currently adapted. Widespread plantings by graziers will confirm whether this potential can be realised and what contribution the legume can make to sustainability of pastures in these regions.

References

- Bell L (2009) Building better feed systems. *Tropical Grasslands* **43**, 199-206.
- Burt RL (1993) *Desmanthus*: A tropical and subtropical forage legume. Part 1. General review. *Herbage Abstracts* **63**, 401-413.
- Cook BG, Graham TWG, Clem RL, Hall TJ, Quirk MF (1993) Evaluation and development of *Desmanthus virgatus* on medium- to heavy-textured soils in Queensland, Australia. Proceedings of the XVII International Grassland Congress, Rockhampton. pp. 2148-2149.
- Gardiner CP, Bielig LM, Schlink AC, Coventry RJ, Waycott M (2004) *Desmanthus* a new pasture legume for the dry tropics. Proceedings of the 4th International Crop Science Congress, Brisbane, Australia, 26 Sep - 1 Oct 2004.
- Gardiner C, Parker A (2012) Steer liveweight gains on Progardes™/buffel pastures in Qld. Proceedings of the 29th Biennial Conference of the Australian Society of Animal Production, Christchurch, NZ, July 2012.
- Gardiner C, Swan S (2008) Abandoned pasture legumes offer potential economic and environmental benefits in semiarid clay soil rangelands. Proceedings of the Australian Rangeland and Conference, Charters Towers, 28th Sept. - 2nd October 2008.
- Jones RM, Clem RL (1997) The role of genetic resources in developing improved pastures in semiarid and subhumid northern Australia. *Tropical Grasslands* **31**, 315-319.
- Kretschmer AE Jr, Pitman WD (2001) Germplasm resources of tropical forage legumes. In: 'Tropical Forage Plants'. (Eds A Sotomayor-Rios, WD Pitman) pp. 41-57. (CRC Press: Boca Raton)
- Luckow M (1993) Monograph of *Desmanthus* (Leguminosae-Mimosoideae). *Systematic Botany Monographs* Vol. **38**. The American Society of Plant Taxonomists.
- McKeon G (2006) Living in a variable climate. <http://www.environment.gov.au/soe/2006/publications/integrative/climate/index.html>
- Ngo T (2012) The effects of diet preference on feed intake, digestibility and nitrogen balance of sheep given Flinders grass (*Iseilema* spp.) hay and/or *Desmanthus leptophyllus ad libitum*. MSc Thesis (submitted 2012) JCU, Townsville.
- Pengelly BC, Conway MJ (2000) Pastures on cropping soils: which tropical pasture legume to use? *Tropical Grasslands* **34**, 162-168.
- Pengelly BC, Liu CJ (2001) Genetic relationships and variation in the tropical mimosoid legume *Desmanthus* assessed by random amplified polymorphic DNA. *Genetic Resources and Crop Evolution* **48**, 91-99.
- Reid R (1983) Pasture plant collecting in Mexico with emphasis on legumes for dry regions. *Australian Plant Introduction Review* **15**, 1-11.
- Robertson FA, Myers RJK, Saffigna PG (1997) Nitrogen cycling in brigalow clay soils under pasture and cropping. *Australian Journal of Soil Research* **35**, 1323-1339.