

OPEN

Expression of microRNA in follicular fluid in women with and without PCOS

Alexandra E. Butler^{1*}, Vimal Ramachandran², Shahina Hayat², Soha R. Dargham², Thomas Keith Cunningham³, Manasi Benurwar², Thozhukat Sathyapalan³, S. Hani Najafi-Shoushtari² & Stephen L. Atkin^{2,4}

Several studies have shown the expression of small non-coding microRNA (miRNA) changes in PCOS and their expression in follicular fluid has been described, though the number of studies remains small. In this prospective cohort study, miRNA were measured using quantitative polymerase chain reaction (qPCR) in 29 weight and aged matched anovulatory women with PCOS and 30 women without from follicular fluid taken at the time of oocyte retrieval who were undergoing *in vitro* fertilization (IVF); miRNA levels were determined from a miRNA data set. 176 miRNA were detected, of which 29 differed significantly between normal women and PCOS women. Of these, the top 7 ($p < 0.015$) were miR-381-3p, miR-199b-5p, miR-93-3p, miR-361-3p, miR-127-3p, miR-382-5p, miR-425-3p. In PCOS, miR-382-5p correlated with age and free androgen index (FAI), miR-199b-5p correlated with anti-mullerian hormone (AMH) and miR-93-3p correlated with C-reactive protein (CRP). In normal controls, miR-127-3p, miR-382-5p and miR-425-3p correlated with the fertilisation rate; miR-127-3p correlated with insulin resistance and miR-381-3p correlated with FAI. Ingenuity pathway assessment revealed that 12 of the significantly altered miRNA related to reproductive pathways, 12 miRNA related to the inflammatory disease pathway and 6 were implicated in benign pelvic disease. MiRNAs differed in the follicular fluid between PCOS and normal control women, correlating with age, FAI, inflammation and AMH in PCOS, and with BMI, fertilization rate (3 miRNA), insulin resistance, FAI and inflammation in control women, according to Ingenuity Pathway Analysis.

Polycystic ovarian syndrome (PCOS) affects 9-21% of the female population and results in anovulatory infertility¹, and is associated with clinical and biochemical hyperandrogenism, and insulin resistance (IR), hypercholesterolemia and type 2 diabetes². The underlying pathophysiology of PCOS is unclear and complex metabolic pathways underlying PCOS, involving deregulated carbohydrate, fat and protein metabolism, have been described³. A single unifying mechanism to account for these disparate physiological and clinical observations has not been described, though noncoding RNA expression regulating gene expression may be a candidate⁴.

MicroRNAs (miRNAs) are small noncoding single-stranded RNA molecules 18-24 nucleotides in length that directly regulate gene expression post-transcriptionally. Canonical miRNA binding occurs at the 3' untranslated regions of the target messenger RNAs (mRNAs), that leads to the repression of mRNA expression resulting in protein translation inhibition. Several critical regulatory functions common to many biological processes are regulated by miRNAs that include cell growth and turnover, and the stress response⁵. The miRBase miRNA registry now includes over 2600 mature human miRNAs, of which only approximately 600 are confidentially annotated and can be considered as validated microRNAs⁶. A single miRNA may modulate the expression and function of hundreds of downstream target genes⁷. Amplification or inhibition of the miRNA signal through feedback on gene regulation may cause dramatic deregulation of physiologic cellular functions by just a few miRNAs, such as that seen following obesity surgery⁸. There is increasing evidence showing the involvement of miRNA in the pathogenesis of both type 1 and type 2 diabetes, and their identification as novel disease biomarkers⁹. Circulating miRNA have been described in PCOS¹⁰⁻¹⁵ and the expression of 3 miRNAs have been suggested to have utility as

¹Diabetes Research Center (DRC), Qatar Biomedical Research Institute (QBRI), Hamad Bin Khalifa University (HBKU), Qatar Foundation (QF), PO Box 34110, Doha, Qatar. ²Weill Cornell Medical College Qatar, Education City, PO Box 24144, Doha, Qatar. ³Academic Diabetes, Endocrinology and Metabolism, Hull York Medical School, Hull, UK. ⁴Royal College of Surgeons in Ireland Bahrain, Bahrain, Bahrain. *email: abutler@hbku.edu.qa

	Control (n = 30)	PCOS (n = 29)	p-value
	Mean data (±SD)	Mean data (±SD)	
Age	32.6 ± 4.7	30.9 ± 4.8	0.14
BMI	25.5 ± 3.6	26.0 ± 3.8	0.56
Menarche	13.0 ± 2.0	13.0 ± 1.1	0.99
Total Antral Follicle Count	17.2 ± 6.8	38.4 ± 17.8	0.0001*
Testosterone (nmol/l)	0.8 ± 0.4	1.4 ± 0.8	0.0004*
Free androgen index	1.35 ± 0.6	4.21 ± 2.9	0.0001*
Fasting Insulin (mIU/ml)	7.68 ± 4.0	8.13 ± 4.7	0.69
Fasting Glucose (mmol/L)	4.81 ± 0.4	4.62 ± 0.4	0.06
HOMA-IR	1.71 ± 1.0	1.72 ± 1.0	0.97
SHBG (mmol/l)	110.9 ± 82.4	63.9 ± 49.8	0.01*
AMH (pg/ml)	21.9 ± 10.4	55.8 ± 14.5	0.001*

Table 1. Mean Demographic and biochemical data. BMI = body mass index; HOMA-IR = insulin resistance; SHBG = sex hormone binding globulin; AMH = antimullerian hormone. * = statistically significant result.

PCOS biomarkers¹⁰. However, whilst miRNA-21, miRNA-27b, miRNA-103, and miRNA-155 have been reported to be differentially expressed in obesity and in PCOS¹³, their regulation is likely to be complex.

The intrafollicular environment is critical for oocyte development, maturation and quality; constituents of the follicular fluid reflect local secretory activity of the oocyte, granulosa and theca cells¹⁶. Studies on miRNAs in follicular fluid are limited and conflicting with, for example, one study finding miR-132 and miR-320 expression reduced in women with PCOS¹⁷, another finding miR-320 expression increased in PCOS¹⁸ and a third reporting no change in miR-320a in PCOS¹⁹. Another study found five miRNAs (miR-9, miR-18b, miR-32, miR-34c and miR-135a) to have increased expression in women with PCOS¹². When PCOS was phenotypically stratified, decreased levels of miR-24-3p, miR-29a, miR-151-3p and miR-574-3p were found in PCOS women versus controls, and differentially high expression of miR-518f-3p was found in hyperandrogenic versus normo-androgenic PCOS women²⁰.

Given the differing results of miRNA expression reported that could reflect PCOS alone, obesity, insulin resistance, androgen levels and/or a metabolic phenotype among other factors, this study was undertaken to assess miRNA in the follicular fluid in age and BMI matched women with and without PCOS who underwent the same IVF protocol for subfertility.

Materials and Methods

“This prospective cohort study was performed from January 2014 to January 2016 within the Hull IVF Unit, UK following approval by the Yorkshire and the Humber NRES ethical committee, UK and all patients gave their written informed consent. Subject demographics are shown in Table 1. The PCOS subjects were recruited using the revised 2003 criteria²¹, namely any 2 out of 3 criteria were met; menstrual disturbance (oligo or amenorrhoea), clinical and/or biochemical signs of androgenism and polycystic ovaries on ultrasound, with the exclusion of other conditions. All women were on folic acid 400 mcg daily but not any other medication. Exclusion criteria were patients with diabetes, renal or liver insufficiency, acute or chronic infections, systemic inflammatory diseases, age <20 or >45 years, or having known immunological disease. Liver ultrasound was performed at the same time to exclude non alcoholic fatty liver disease. Study participants had no concurrent illness and were not on any medication for the preceding nine months except study medications. None of the patients had had a successful pregnancy or miscarriage in at least five years prior to study entry. Diabetes was excluded by a 75 g oral glucose tolerance test. Non-classical 21-hydroxylase deficiency, hyperprolactinaemia, Cushing’s disease and androgen-secreting tumours were excluded by appropriate tests”²².

Sample Collection. “All patients underwent a standard IVF antagonist protocol. The patients commenced their rFSH stimulation on day 2 of their menstrual cycle using either Merional (Pharmasure) or Gonal-F (Merck Serono). A GnRH antagonist (Cetrotide: Merck Serono) was used to prevent a premature LH surge²²”.

“The patients underwent ultrasound scans from day 7 to observe the ovarian response to stimulation and these were repeated every 48 hours. The scans were used to measure the diameters of the follicles, thus observing response and follicle numbers. Final maturation was triggered when two or more leading follicles were ≥18 mm using human chorionic gonadotrophin (hCG, Pregnyl (Merck Sharp and Dohme). Follicular fluid was collected at the same time as oocyte harvesting, was centrifuged at 3500 g for 15 minutes and placed into aliquots and frozen at −80 °C until analysis²²”.

“Transcervical embryo transfer was performed and embryos were classified using standard criteria²³ at the cleavage stage (day 2 to 3 after egg collection) and for blastocyst stage (day 5 to 6 after egg collection). Top Quality embryos were graded on Day 3 as per Alpha Consensus²⁴. Embryo transfers were performed on either day 3 or ideally at day 5 (blastocyst) to give the best chance for implantation, as this timing is similar when compared to natural cycle embryos moving into the uterus”²².

“A fasting blood sample was taken on day 21 of the luteal phase of the cycle before commencing IVF treatment, and centrifuged at 3500 g for 15 minutes and placed into aliquots and frozen at −80 °C until analysis.

	Control (N = 30)	PCOS (N = 29)	p-value
	Mean (± S.D.)	Mean (± S.D.)	
Endometrium at oocyte retrieval (mm)	10.31 ± 1.78	10.72 ± 2.06	0.42
Follicles aspirated	11.47 ± 5.11	15.96 ± 5.30	0.002**
Eggs Retrieved	8.47 ± 5.08	11.29 ± 5.02	0.04 [†]
Fertilisation rate	4.82 ± 2.65	8.43 ± 3.87	0.0003***
Cleavage	4.68 ± 2.72	7.26 ± 4.40	0.01 [†]
G3D3	3.00 ± 2.29	4.17 ± 3.47	0.16
Blastocyst	1.46 ± 1.77	2.91 ± 3.01	0.05 [†]
Clinical Pregnancy	10	7	0.24

Table 2. Mean outcome data for stimulated ovarian cycle for Control and PCOS groups. G3D3: Top Quality embryos on Day 3 as per Alpha Consensus²⁴. * = statistically significant result.

All PCOS women were oligomenorrheic, none having had a period within the preceding 6 weeks. Serum and plasma was stored frozen at -80°C pending analysis. Serum testosterone was measured by isotope dilution liquid chromatography- tandem mass spectrometry (Waters Corporation, Manchester, UK). Sex hormone binding globulin (SHBG) was determined by an immunometric assay with fluorescence detection on the DPC Immulite 2000 analyzer using the manufacturer's recommended protocol. The free androgen index was obtained as the total testosterone $\times 100/\text{SHBG}$. Serum insulin was assayed using a competitive chemiluminescent immunoassay performed on the manufacturer's DPC Immulite 2000 analyzer (Euro/DPC, Llanberis, UK). The analytical sensitivity of the insulin assay was $2\ \mu\text{U}/\text{ml}$, the coefficient of variation was 6%, and there was no stated cross-reactivity with proinsulin. Plasma glucose was measured using a Synchron LX 20 analyzer (Beckman-Coulter), using the manufacturer's recommended protocol. The coefficient of variation for the assay was 1.2% at a mean glucose value of $5.3\ \text{mmol}/\text{liter}$ during the study period. The insulin resistance was calculated using the HOMA method [$\text{HOMA-IR} = (\text{insulin} \times \text{glucose})/22.5$], and pancreatic beta cell sensitivity measured by $\text{HOMA-}\beta$ [$\text{HOMA-}\beta = (20 \times \text{insulin})/\text{glucose} - 3.5$]²².

MicroRNA analysis

“Currently, due to accuracy, simplicity, and greater reproducibility, quantitative polymerase chain reaction (qPCR) is the favored method for determining microRNA expression when compared to other hybridization or sequencing-based technologies²⁵. MicroRNA was isolated from $200\ \mu\text{l}$ EDTA plasma using the miRCURY[™] RNA Isolation Kit (Exiqon A/S, Denmark) following the manufacturer's instructions and applying an RNA Spike-in kit (Exiqon A/S, Denmark) for assessment of hemolysis and contamination with blood cells and RNA isolation efficiency. The yield of microRNA preparation from plasma samples was monitored by including carrier RNA from the bacteriophage MS2. The quality and the integrity of the RNA were assessed by a 2100 Bioanalyzer instrument (Agilent Technologies), equipped with the small RNA specific chip. The expression of microRNA levels was retrieved from a microRNA data set generated by RNA sequencing of more than 390 different human tissue and primary cell samples^{8,26}”

Statistics

There was no information regarding changes in miR on which to base a sample size calculation. For such pilot studies, Birkett and Day²⁷ suggest a minimum of 20 degrees of freedom to estimate variance from which a larger trial could be powered, hence 24 subjects in each group were recruited. Statistical analysis was performed using SPSS (v22, Chicago, Illinois). Descriptive data is presented as mean \pm SD for continuous data and n (%) for categorical data. T-tests or Mann Whitney tests were used to compare means/medians where appropriate. Linear associations were assessed using the Spearman's correlation test. Mean normalization was performed before analysis using the global mean of each miRNA. Using GenEx software that accompanied the Bioanalyzer, normalization was performed to achieve a global mean of all miRNAs with Ct below 35. An unpaired t-test was used to test the paired changes between PCOS and controls in miRNA levels. False discovery rates (FDR) of $q < 0.05$ ²⁸ were considered significant. Pathway analysis was performed using Ingenuity Pathway Analysis (Qiagen), and those pathways over-represented by the FDR-significant miRNA changes at $q < 0.05$ ²⁸ were identified.

Results

Baseline characteristics of the 59 patients, 29 PCOS and 30 normal controls, are shown in Table 1 where it can be seen that patients were non-obese, age and weight matched and did not differ in their insulin resistance. There were significant differences in ovarian reserve parameters antral follicle count (AFC) and anti-Mullerian Hormone, (AMH), and androgen status between the groups; however, there was no significant difference in fasting insulin or HOMA-IR (Table 1).

IVF parameters are shown in Table 2, where it can be seen that more follicles were aspirated and eggs retrieved ($p < 0.05$) and the fertilization rate was higher for PCOS ($p < 0.0003$).

PCOS vs Control	Fold change	P-Value
hsa-miR-381-3p	1.95136	0.0018
hsa-miR-199b-5p	2.2861	0.0040
has-miR-650	-2.45385	0.0053
hsa-miR-93-3p	3.47102	0.0060
has-miR-663b	-2.47917	0.0095
hsa-miR-361-3p	2.10148	0.0104
hsa-miR-127-3p	2.15942	0.0132
has-miR-1238-3p	1.78509	0.0149
hsa-miR-382-5p	2.4991	0.0150
hsa-miR-425-3p	2.41375	0.0161
hsa-miR-212-3p	2.91193	0.0183
hsa-miR-423-3p	1.59576	0.0189
hsa-miR-891b	-2.95618	0.0198
hsa-miR-187-3p	1.41842	0.0229
hsa-miR-513c-5p	2.21283	0.0262
hsa-miR-510-5p	1.1463	0.0300
hsa-miR-507	2.07081	0.0305
hsa-miR-509-3-5p	1.62364	0.0366
hsa-miR-638	1.18991	0.0375
hsa-miR-532-5p	1.93401	0.0375
hsa-miR-1271-5p	1.55672	0.0375
hsa-miR-193a-5p	1.58387	0.0377
hsa-miR-32-5p	-2.14067	0.0396
hsa-miR-200c-3p	2.08148	0.0459
hsa-miR-33b-3p	1.23812	0.0460
hsa-miR-675-5p	1.94931	0.0462
hsa-miR-125b-5p	1.31108	0.0481
hsa-miR-206	-1.23035	0.0486
hsa-miR-9-5p	-1.53421	0.0500

Table 3. Significant differences (fold change) in the miRNA in follicular fluid between PCOS (n = 29) versus control women (n = 30).

176 miRNAs were detected, of which 29 differed significantly (Table 3, Fig. 1) between normal women and PCOS; all of the miRNA detected are shown in Supplementary Table 1.

The top 7 of these ($p < 0.015$) were miR-381-3p, miR-199b-5p, miR-93-3p, miR-361-3p, miR-127-3p, miR-382-5p, miR-425-3p. In PCOS, miR-382-5p correlated with age and FAI, miR-199b-5p correlated with AMH and miR-93-3p correlated with C-reactive protein (CRP). In normal controls, miR-127-3p, miR382-5p and miR-425-3p correlated with the fertilisation rate; miR-127-3p correlated with insulin resistance, and miR-381-3p correlated with FAI (Table 4).

Ingenuity pathway assessment. When Ingenuity pathway analysis was undertaken to look at the functional relationships, it was seen that 12 of the significantly altered miRNA related to reproductive pathways, as is shown in Fig. 2. Twelve miRNAs were related to the inflammatory disease pathway, as is shown in Fig. 3. Six miRNAs were related to the pathways implicated in benign pelvic disease: miR-212-3p, miR-32-5p, miR-200c-3p, miR-125b-5p, miR-206 and miR-9-5p.

Discussion

Ovulatory dysfunction, increased IR, an elevated FAI and inflammation are hallmarks of PCOS and, in line with the objective, this study has shown that miRNA found in the follicular fluid were related to these parameters. In PCOS, miR-199b-5p correlated with AMH that is recognized to be elevated in PCOS and, indeed, AMH has been suggested by some studies, but not others, to be a biomarker of PCOS^{29,30} where it has a four fold association with a diagnosis of PCOS although with low sensitivity. miR-382-5p correlated with FAI that is classically elevated, and miR-93-3p correlated with C-reactive protein (CRP), a marker of inflammation, that has consistently been shown to be greater in PCOS than controls^{31,32}. In addition, in normal subjects, miR-127-3p, miR-382-5p and miR-425-3p correlated with the fertilisation rate, whilst miR-127-3p correlated with IR and miR-381-3p correlated with FAI. Whether these miRNAs are involved in the associated processes would seem to be supported by the Ingenuity pathways showing the involvement of these miRNA specifically in the reproductive pathways (12 miRNAs) and inflammation (12 miRNAs). However, given that miRNA action is essentially inhibitory by inducing degradation or inhibition of protein translation³, their roles, either direct or




Figure 1. Volcano plot of microRNA detected in the follicular fluid between those women with PCOS (n = 29) versus normal women without PCOS (n = 30). Altered expression of microRNAs in a comparison between PCOS and controls are shown as fold change on the x-axis and the significance (p-value) of that fold change on the y-axis. Both are shown in log scale. Red dots: miRs with >2-fold change (+ or -) but not changing significantly. Yellow dots: miRs with >2-fold change and also changing significantly (p < 0.05).


	PCOS													
	miR-199b-5p		miR-127-3p		miR-361-3p		miR-382-5p		miR-425-3p		miR-381-3p		miR-93-3p	
	R	P-value	R	P-value	R	P-value	R	P-value	R	P-value	R	P-value	R	P-value
Age	0.348	0.095	0.215	0.29	-0.039	0.875	0.436	0.023	0.198	0.333	-0.088	0.744	-0.363	0.105
BMI	0.075	0.728	-0.091	0.658	-0.172	0.481	-0.149	0.459	-0.124	0.547	-0.088	0.745	0.04	0.862
AMH	-0.591	0.002	0.023	0.911	0.069	0.779	-0.152	0.449	-0.143	0.485	-0.073	0.789	-0.106	0.647
Fertilisation rate	0.191	0.408	0.152	0.51	-0.133	0.638	-0.095	0.674	0.023	0.92	-0.159	0.586	0.156	0.536
HOMA-IR	-0.163	0.445	-0.08	0.699	-0.244	0.314	-0.148	0.462	-0.286	0.156	-0.2	0.458	0.143	0.537
CRP	0.101	0.655	-0.056	0.796	0.221	0.393	0.247	0.233	-0.147	0.492	0.089	0.751	0.637	0.003
FAI	0.012	0.955	-0.077	0.708	-0.366	0.124	-0.419	0.03	-0.306	0.129	-0.395	0.13	-0.321	0.156
	CONTROLS													
	miR-199b-5p		miR-127-3p		miR-361-3p		miR-382-5p		miR-425-3p		miR-381-3p		miR-93-3p	
	R	P-value	R	P-value	R	P-value	R	P-value	R	P-value	R	P-value	R	P-value
Age	0.034	0.887	0.06	0.77	0.191	0.533	0.382	0.072	0.133	0.535	0.629	0.028	-0.265	0.405
BMI	0.342	0.14	-0.087	0.674	-0.41	0.164	0.142	0.517	0.167	0.435	-0.109	0.737	-0.196	0.54
AMH	-0.186	0.431	-0.168	0.413	0.022	0.943	-0.019	0.93	-0.16	0.455	0.224	0.484	-0.182	0.572
Fertilisation rate	0.395	0.104	0.449	0.024	0.119	0.712	0.537	0.012	0.424	0.049	0.436	0.18	0.073	0.83
HOMA-IR	-0.108	0.65	-0.387	0.050	0.121	0.694	0.013	0.954	-0.065	0.762	0.196	0.542	0.126	0.697
CRP	0.198	0.402	0.123	0.56	0.33	0.271	0.099	0.66	0.166	0.448	-0.209	0.537	0.333	0.29
FAI	-0.002	0.993	-0.213	0.296	-0.06	0.847	-0.308	0.153	-0.005	0.983	-0.655	0.021	0.27	0.395

Table 4. Spearman’s correlation of the top 7 microRNA (p < 0.015) in follicular fluid taken at the time of oocyte retrieval during IVF in 59 age and weight matched women, 29 with PCOS and 30 normal control women. BMI = body mass index; AMH = Antimullarian Hormone; HOMA-IR = insulin resistance; FAI = free androgen index; CRP = C reactive protein.

indirect, require further clarification. It is recognized that miRNA may be affected by BMI¹³ and, to account for this, PCOS and normal controls were weight matched in this study; however, despite this, in normal controls, miR-1238-3p correlated with BMI.

Recent reports have indicated that dysregulated miRNAs may be key players in the ovulatory dysfunction of women with PCOS. MiRNAs have been reported to be differentially expressed in follicular fluid of women with PCOS^{12,17}. Upregulated miRNAs include miRNA-9, miRNA-18b, miRNA-32, miRNA-34c and miRNA-135a, and, by analysis of target genes, these have been shown to be involved in carbohydrate metabolism, beta-cell

Inflammation



© 2000-2019 QIAGEN. All rights reserved.

Figure 3. Ingenuity pathway analysis of the significantly altered miRNA found in the follicular fluid showing that 12 of these miRNA were related to the genes involved in the Inflammatory disease pathway. Strongly upregulated and downregulated miRNA are represented with dark red and green color and vice versa.

In conclusion, miR differed in the follicular fluid between PCOS and normal control women, correlating with age, FAI, inflammation and AMH in PCOS, and with fertilization rate, insulin resistance and FAI and inflammation in control women, in accord with the Ingenuity pathway assessment.

Data availability

The data that support the findings of this study are available from the corresponding author upon request.

Received: 18 August 2019; Accepted: 22 October 2019;

Published online: 08 November 2019

References

1. Azziz, R. *et al.* The prevalence and features of the polycystic ovary syndrome in an unselected population. *The Journal of clinical endocrinology and metabolism*. **89**(6), 2745–9 (2004).
2. Franks, S. Polycystic ovary syndrome. *The New England journal of medicine*. **333**(13), 853–61 (1995).
3. Halama, A. *et al.* Metabolomics of Dynamic Changes in Insulin Resistance Before and After Exercise in PCOS. *Front Endocrinol (Lausanne)*. **10**, 116 (2019).

4. Butler, A. E. *et al.* Alterations in long non-coding RNAs in women with and without Polycystic Ovarian Syndrome. *Clinical endocrinology* (2019).
5. Flynt, A. S. & Lai, E. C. Biological principles of microRNA-mediated regulation: shared themes amid diversity. *Nature reviews Genetics*. **9**(11), 831–42 (2008).
6. The microRNA database [Internet]. www.mirbase.org (2013).
7. Lim, L. P. *et al.* Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature*. **433**(7027), 769–73 (2005).
8. Atkin, S. L. *et al.* Changes in Blood microRNA Expression and Early Metabolic Responsiveness 21 Days Following Bariatric Surgery. *Front Endocrinol (Lausanne)*. **9**, 773 (2018).
9. Guay, C. & Regazzi, R. Circulating microRNAs as novel biomarkers for diabetes mellitus. *Nature reviews. Endocrinology*. **9**(9), 513–21 (2013).
10. Long, W. *et al.* Characterization of serum microRNAs profile of PCOS and identification of novel non-invasive biomarkers. *Cellular physiology and biochemistry: international journal of experimental cellular physiology, biochemistry, and pharmacology*. **33**(5), 1304–15 (2014).
11. Chen, Y. H. *et al.* miRNA-93 inhibits GLUT4 and is overexpressed in adipose tissue of polycystic ovary syndrome patients and women with insulin resistance. *Diabetes*. **62**(7), 2278–86 (2013).
12. Roth, L. W. *et al.* Altered microRNA and gene expression in the follicular fluid of women with polycystic ovary syndrome. *J Assist Reprod Genet*. **31**(3), 355–62 (2014).
13. Murri, M., Insenser, M., Fernandez-Duran, E., San-Millan, J. L. & Escobar-Morreale, H. F. Effects of polycystic ovary syndrome (PCOS), sex hormones, and obesity on circulating miRNA-21, miRNA-27b, miRNA-103, and miRNA-155 expression. *The Journal of clinical endocrinology and metabolism*. **98**(11), E1835–44 (2013).
14. Xu, B., Zhang, Y. W., Tong, X. H. & Liu, Y. S. Characterization of microRNA profile in human cumulus granulosa cells: Identification of microRNAs that regulate Notch signaling and are associated with PCOS. *Molecular and cellular endocrinology*. **404**, 26–36 (2015).
15. Lin, L., Du, T., Huang, J., Huang, L. L. & Yang, D. Z. Identification of differentially expressed microRNAs in the ovary of polycystic ovary syndrome with hyperandrogenism and insulin resistance. *Chin Med J (Engl)*. **128**(2), 169–74 (2015).
16. Revelli, A. *et al.* Follicular fluid content and oocyte quality: from single biochemical markers to metabolomics. *Reprod Biol Endocrinol*. **7**, 40 (2009).
17. Sang, Q. *et al.* Identification of microRNAs in human follicular fluid: characterization of microRNAs that govern steroidogenesis *in vitro* and are associated with polycystic ovary syndrome *in vivo*. *J Clin Endocrinol Metab*. **98**(7), 3068–79 (2013).
18. Yin, M. *et al.* Transactivation of microRNA-320 by microRNA-383 regulates granulosa cell functions by targeting E2F1 and SF-1 proteins. *J Biol Chem*. **289**(26), 18239–57 (2014).
19. Scalici, E. *et al.* Circulating microRNAs in follicular fluid, powerful tools to explore *in vitro* fertilization process. *Sci Rep*. **6**, 24976 (2016).
20. Sorensen, A. E., Wissing, M. L., Englund, A. L. & Dalgaard, L. T. MicroRNA Species in Follicular Fluid Associating With Polycystic Ovary Syndrome and Related Intermediary Phenotypes. *J Clin Endocrinol Metab*. **101**(4), 1579–89 (2016).
21. Rotterdam ESHRE/ASRM-Sponsored PCOS consensus workshop group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome (PCOS). *Hum Reprod*. **19**(1), 41–7 (2004).
22. Cunningham, T. C. The effect of environmental, physical, and nutritional factors on *in vitro* fertilisation: Hull; (2016).
23. Cutting, R., Morroll, D., Roberts, S. A., Pickering, S. & Rutherford, A. Elective single embryo transfer: guidelines for practice British Fertility Society and Association of Clinical Embryologists. *Human fertility (Cambridge, England)*. **11**(3), 131–46 (2008).
24. Istanbul consensus workshop on embryo assessment: proceedings of an expert meeting. *Reprod Biomed Online*. **22**(6), 632–46 (2011).
25. Git, A. *et al.* Systematic comparison of microarray profiling, real-time PCR, and next-generation sequencing technologies for measuring differential microRNA expression. *RNA*. **16**(5), 991–1006 (2010).
26. de Rie, D. *et al.* An integrated expression atlas of miRNAs and their promoters in human and mouse. *Nat Biotechnol*. **35**(9), 872–8 (2017).
27. Birkett, M. A. & Day, S. J. Internal pilot studies for estimating sample size. *Stat Med*. **13**(23–24), 2455–63 (1994).
28. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B*. **57**(1), 289–300 (1995).
29. Iliodromiti, S., Kelsey, T. W., Anderson, R. A. & Nelson, S. M. Can anti-Mullerian hormone predict the diagnosis of polycystic ovary syndrome? A systematic review and meta-analysis of extracted data. *The Journal of clinical endocrinology and metabolism*. **98**(8), 3332–40 (2013).
30. Quinn, M. M. *et al.* Age-stratified thresholds of anti-Mullerian hormone improve prediction of polycystic ovary syndrome over a population-based threshold. *Clinical endocrinology* (2017).
31. Boulman, N. *et al.* Increased C-reactive protein levels in the polycystic ovary syndrome: a marker of cardiovascular disease. *J Clin Endocrinol Metab*. **89**(5), 2160–5 (2004).
32. Verit, F. F. High sensitive serum C-reactive protein and its relationship with other cardiovascular risk factors in normoinsulinemic polycystic ovary patients without metabolic syndrome. *Arch Gynecol Obstet*. **281**(6), 1009–14 (2010).
33. Aye, M. M. *et al.* Acute hypertriglyceridemia induces platelet hyperactivity that is not attenuated by insulin in polycystic ovary syndrome. *Journal of the American Heart Association*. **3**(1), e000706 (2014).
34. Yang, Y. M., Seo, S. Y., Kim, T. H. & Kim, S. G. Decrease of microRNA-122 causes hepatic insulin resistance by inducing protein tyrosine phosphatase 1B, which is reversed by licorice flavonoid. *Hepatology*. **56**(6), 2209–20 (2012).
35. Giardina, S. *et al.* Changes in circulating miRNAs in healthy overweight and obese subjects: Effect of diet composition and weight loss. *Clin Nutr*. **38**(1), 438–43 (2019).
36. Liu, S. X. *et al.* Exercise Reduces Insulin Resistance in Type 2 Diabetes Mellitus via Mediating the lncRNA MALAT1/MicroRNA-382-3p/Resistin Axis. *Molecular therapy. Nucleic acids*. **18**, 34–44 (2019).
37. Diez-Fraile, A. *et al.* Age-associated differential microRNA levels in human follicular fluid reveal pathways potentially determining fertility and success of *in vitro* fertilization. *Human fertility (Cambridge, England)*. **17**(2), 90–8 (2014).
38. Sorensen, A. E., Udesen, P. B., Wissing, M. L., Englund, A. L. M. & Dalgaard, L. T. MicroRNAs related to androgen metabolism and polycystic ovary syndrome. *Chem Biol Interact*. **259**(Pt A), 8–16 (2016).

Acknowledgements

This work was part of an MD thesis and its publishing is in line with the University of Hull policy, and can be accessed online <https://hydra.hull.ac.uk/resources/hull:14051>²². The publication of this article was funded by the Qatar National Library.

Author contributions

A.E.B. was involved in data analysis and wrote the first draft of the paper. V.R. and M.B. were involved in miR measurement. T.K.C. and T.S. were involved in the study design, acquisition of data and paper drafting. S.H. performed the I.P.A. and S.R.D. performed the statistical analysis. S.H.N.S. and S.L.A. designed the study and manuscript preparation and interpretation of the data. All authors read and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-019-52856-5>.

Correspondence and requests for materials should be addressed to A.E.B.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019