

Supplementary data for the article:

Marković, J. M.; Trišović, N. P.; Tóth-Katona, T.; Milčić, M. K.; Marinković, A. D.; Zhang, C.; Jákli, A. J.; Fodor-Csorba, K. A Structure-Property Relationship Study of Bent-Core Mesogens with Pyridine as the Central Unit. *New Journal of Chemistry* **2014**, 38 (4), 1751–1760. <https://doi.org/10.1039/c3nj01430d>

Structure–property relationship study of bent-core mesogens with pyridine as central unit

J.M. Marković,^a N.P. Trišović,^{*a} T. Tóth-Katona,^b M.K. Milčić,^c A.D. Marinković,^a C. Zhang,^d
A.J. Jáklí^{b,d} and K. Fodor-Csorba^b

^a*Faculty of Technology and Metallurgy, University of Belgrade, Belgrade, Serbia*

^b*Wigner Research Centre for Physics, Institute for Solid State Physics and Optics of the Hungarian Academy of
Sciences H-1525 Budapest, P.O.Box 49, Hungary*

^c*Faculty of Chemistry, University of Belgrade, Belgrade, Serbia*

^d*Liquid Crystal Institute, Kent State University, Kent, Ohio 44242, USA*

New Journal of Chemistry

Electronic Supplementary Information




Figure S1 DSC curves of compounds IIIa–IIIc




Figure S2 Optical photomicrographs (crossed polarizers) of compound **IIc**: a) $T=238^{\circ}\text{C}$: mesophase formation on heating; b) $T=241^{\circ}\text{C}$: mesophase transition on heating;