

ORIGINAL

Electrons released from both flavins of NADPH-P450 reductase contribute to the reductive mobilization of iron from ferritin

Kazumi Takaishi^a and Hiroshi Kitahata^b

^aDepartment of Dental Anesthesiology, Tokushima University Hospital, Tokushima, Japan

^bDepartment of Dental Anesthesiology, Tokushima University Graduate School of Biomedical Sciences, Tokushima, Japan

Abstract: Ferritin, an iron storage protein, plays an important role in iron homeostasis. The mechanism of reductive mobilization of iron from ferritin has not been clarified yet despite many studies. The aim of this study was to assess the mechanisms of the mobilization of iron from ferritin by NADPH P-450 reductase. Nucleotide-dependent flavoenzymes generated significant mobilization of iron from ferritin. The possibility of reductive mobilization of iron from ferritin by electrons released from flavin sites or heme site of two flavoenzymes was investigated to elucidate the mediator-independent mechanisms of such reductive mobilization. The mobilization by NADPH-P450 reductase in the presence of ferricyanide increased threefold, while in the presence of cytochrome C increased thirteen-fold. These results indicate that electrons released from both flavins of NADPH-P450 reductase contribute to the reductive mobilization of iron from ferritin. The mechanism of the mobilization of iron from ferritin is discussed. *J. Med. Invest.* 66:230-232, August, 2019

Keywords: Ferritin, Iron, Electron, Flavoenzyme, Reductive mobilization

INTRODUCTION

Ferritin is an iron storage protein in many mammalian cells (1). Cellular iron homeostasis is maintained by regulation of the expression of ferritin and the transferrin receptor.

The ferritin molecule consists of a mineral core of hydrated ferric oxide and 24 protein shells, and the iron content in the core consists of a maximum of 4500 atoms. Two types of channels, six four-fold hydrophobic channels and eight three-fold hydrophilic channels, are generated by assembly of the protein shells. The hydrophobic channel is 0.3-0.4 nm wide and 1.2 nm long, and the hydrophilic channel is 0.3-0.4 nm wide and 0.5 nm long, but is blocked by metal ions (2, 3). The route of iron deposition and mobilization via the hydrophobic channel has been the subject of investigations. The iron in the core can be mobilized from ferritin by reduction of ferric iron, and iron mobilization has been observed on the addition of biological reductants such as dihydroflavin (4-7), dihydrolipoate (7, 8), superoxide anion radical (9, 10), and nitric oxide (11). In recent studies, it was shown that the iron release from ferritin occurred through lysosomal proteolysis or Nuclear Receptor Coactivator-4 – mediated ferritinophagy (12, 13). However, this process is relatively slow from the perspective of rapid reactivity of iron ion and is related to degradation of ferritin cage (12, 14). It has been suggested that the reductant must penetrate into the interior of the channel in order to reduce the iron.

Whether ferritin iron reduction involves direct access by the reductant into the central cavity of the molecule or electron tunneling via the hydrophobic channel has remained unclear. The purpose of our study was to investigate the molecular mechanisms of the mobilization of iron from ferritin by the nucleotide-dependent flavoenzymes.

Received for publication February 8, 2018; accepted February 2, 2019.

Address correspondence and reprint requests to Kazumi Takaishi, Department of Dental Anesthesiology, Tokushima University Hospital, 3-18-15 Kuramoto, Tokushima 770-8504, Japan and Fax: +81-88-633-7368.

MATERIALS AND METHODS

Ferritin (horse spleen) and NADPH-P 450 reductase (rabbit liver) were obtained from Sigma-Aldrich Co. (St. Louis, MO).

The standard reaction mixture for iron mobilization from ferritin contains 1.9 μ M ferritin, 1U NADPH-P 450 reductase, 20 μ M bathophenanthroline disulfonic acid 2Na salt in 20 mM phosphate buffer, pH 7.4. Bathophenanthroline is used as a quantity reagent of iron by binding to ferrous iron and forming orange-red chelate. We measured the quantity of released ferrous iron in real time using the spectrophotometer (U-3900, Hitachi High-Tech Science Corporation, Japan) after adding 100 μ M NADPH to the standard reaction mixture. The quantity of released ferrous iron was calculated from absorbance at 530 nm by using a molar absorbance coefficient of $22.1 \times 10^3 \text{ M}^{-1} \text{ cm}^{-1}$ for bathophenanthroline.

To investigate the effect of superoxide dismutase on the reductive mobilization of iron, 10 μ M superoxide dismutase added to the standard reaction mixture. In another experiment to elucidate the mechanism of ferritin iron reduction, 2 mM ferricyanide or 6.7 μ M cytochrome C was added each to the standard reaction mixture.

No less than 4 experiments were performed separately. The results are expressed as the representative data for all the experiments.

RESULTS

Mobilization of iron from ferritin by NADPH-P 450 reductase

NADPH-P 450 reductase mobilized iron from ferritin as shown in Fig.1. This reductive mobilization was reduced by 6% in the presence of superoxide dismutase (Table 1). It was not significantly reduced in the presence of superoxide dismutase, indicating that the superoxide anion radical did not mediate the reduction of ferritin iron.




Fig 1. Time course of iron mobilization from ferritin by means of NADPH P-450 reductase.

The reaction mixture contains 1.9 μ M ferritin, 1U NADPH-P 450 reductase, 100 μ M NADPH, 20 μ M bathophenanthroline disulfonic acid 2Na salt in 20mM phosphate buffer, pH 7.4. The quantity of released ferrous iron was calculated from absorbance at 530 nm by using a molar absorbance coefficient of $22.1 \times 10^{-3} \text{ M}^{-1} \text{ cm}^{-1}$ for bathophenanthroline.

Table 1 Effects of Superoxide dismutase on iron mobilization from ferritin by NADPH P-450 reductase

Rate of iron mobilization (Fe ²⁺ μ M / 10min)	
Control	2.71
Superoxide dismutase (10 μ M)	2.55

The effects of ferricyanide and cytochrome C on the mobilization of iron from ferritin by NADPH-P 450 reductase

We performed another study to elucidate the mechanism of ferritin iron reduction. It is recognized that the external electron acceptors of FAD and FMN of diflavin reductase family are ferricyanide and cytochrome C, respectively. Here, the reductive mobilization of iron from ferritin is investigated in the presence of external electron acceptors. The mobilization in the presence of ferricyanide increased threefold, while in the presence of cytochrome C increased thirteen-fold as shown in Table 2.

Table 2 Effects of ferricyanide and Cytochrome C on iron mobilization from ferritin by NADPH-P450 reductase

Rate of iron mobilization (Fe ²⁺ μ M / 10min)	
Control	2.71
Ferricyanide (2 mM)	8.58
Cytochrome C (6.7 μ M)	34.29

DISCUSSION

In this study, NADPH-P450 reductase was found to be able to catalyze the mobilization of iron from ferritin. It was reported that the reductive mobilization of iron from ferritin is dependent on the relative rate of NADH oxidations, dissolved O₂ consumption and mineral core reduction in the study using recombinant frog M ferritin (15). It was also reported that the iron mobilization from ferritin is dependent on the concentration of dissolved oxygen in solution (16). However, Johnson LE *et al.* reported that

the reductive mobilization of iron by reduced flavin mononucleotide (FMNH₂) was limited by the concentration of FMNH₂ under their experimental conditions excluded the possibility of oxygen diffusion into the reaction mixture, and that iron reductive mobilization was independent on the presence of chaotropes (17). In our experiment, the influence of oxygen diffusion on iron release was extremely limited because the iron mobilization was reduced by only 6% in the presence of superoxide dismutase. Moreover, Johnson LE *et al.* showed that the diffusion of FMNH₂ through the ferritin pores was unlikely mechanism by the experiments about encapsulation of FMN inside the ferritin cavity (17).

A further investigation is needed to clarify the mechanism of ferritin iron reduction by the nucleotide-dependent flavoenzymes. One possibility is thought to be that the electron from either enzyme could gain direct access into the core through the ferritin channel and thus activate the reduction of ferritin iron.

NADPH-P 450 reductase exerts their ferricyanide reductive activity by means of electrons from the FAD site and the cytochrome C reductive activity by means of electrons from the FMN site (18, 19). In our experimental conditions, the reductive mobilization of iron from ferritin is about three folds by the addition of ferricyanide. On the other hand, the reductive mobilization of iron from ferritin is about thirteen folds by the addition of cytochrome C, indicating that electrons released from both flavins of NADPH-P450 reductase contribute to the reductive mobilization of iron from ferritin.

The hop distance of electron transfer from site to site of the metal complex model is 1-2 nm. The rate constant is 8×10 sec at a distance of 1.13 nm and drops during metal ion separation by 10 seconds for each 0.3 nm increment (20). While it can be assumed that a hop distance of 2 nm is close to the limit of usefulness, it can occur in proteins over a distance of more than 1 nm (21).

Access to the interior of the shell of the ferritin molecule is possible only via a channel. There are two types of channels through which small molecules can gain access to the central core of the molecule. The hydrophobic channel is 0.3-0.4 nm in diameter and 1.2 nm long, while the hydrophilic channel is 0.3-0.4 nm in diameter and 0.5 nm long (2, 3). In theory, therefore, the electrons could gain direct access via one of these channels to the central core of the ferritin molecule.

Ferritin iron cores can be reduced by one electron per iron accompanied by an uptake of two protons per electron from the surrounding medium (22). The midpoint potential of ferritin iron is $E = -190\text{mV}$ (23). The midpoint potential of NADPH-P 450 reductase is $E = -274 \text{ mV}$ (ox/sq), $E = -371 \text{ mV}$ (sq/hq) for FAD and $E = -109 \text{ mV}$ (ox/sq), $E = -279 \text{ mV}$ (sq/hq) for FMN (24). Since the midpoint potential for FAD of NADPH-P 450 reductase is much lower than the midpoint potential of ferritin iron, flavin semiquinone of the enzyme is provided thermodynamically for the mechanism of electron transfer to ferritin iron. For these reasons, the reductive mobilization of iron from ferritin may be induced by direct electron transfer from the flavin site of NADPH-P 450 reductase to the ferritin core.

Iron homeostasis is related to a variety of important physiological functions (25). A large quantity of free iron released from ferritin generates highly reactive hydroxyl radical from superoxide and hydrogen peroxide. It can lead the control of the pathophysiology related to free radicals to explore the mechanism of iron release from ferritin. It is mandatory to investigate more about the mobilization of iron from ferritin.

CONCLUSION

The iron mobilization by NADPH-P450 reductase from ferritin in the presence of ferricyanide or cytochrome C increased markedly. It is suggested that electrons released from both flavins of NADPH-P450 reductase contribute to the reductive mobilization of iron from ferritin.

CONFLICT OF INTEREST

None of the authors have any conflicts of interest to declare.

ACKNOWLEDGEMENT

We are greatly indebted and wish to express our profound gratitude to Prof. Emeritus Aki K for his advice and direct guidance throughout the entire course of this study.

This study was supported in part by a Gran-in-Aid for Scientific Research from Japan Society for the Promotion of Science (#23792350).

REFERENCES

1. Arosio P, Ingrassia R, Cavadini P: Ferritins : a family of molecules for iron storage, antioxidation and more. *Biochim Biophys Acta* 1790 : 589-599, 2009
2. Crichton RR, Charlotiaux-Wauters M : Iron transport and storage. *Eur J Biochem* 164 : 485-506, 1987
3. Crichton RR, Ward RJ : Iron metabolism--new perspectives in view. *Biochemistry* 31 : 11255-11264, 1992
4. Sirivech S, Frieden E, Osaki S : The release of iron from horse spleen ferritin by reduced flavins. *Biochem J* 143 : 311-315, 1974
5. Jones T, Spencer R, Walsh C : Mechanism and kinetics of iron release from ferritin by dihydroflavins and dihydroflavin analogues. *Biochemistry* 17 : 4011-4017, 1978
6. Funk F, Lenders JP, Crichton RR, Schneider W : Reductive mobilisation of ferritin iron. *Eur J Biochem* 152 : 167-172, 1985
7. Bonomi F, Pagani S : Removal of ferritin-bound iron by DL-dihydrolipoate and DL-dihydrolipoamide. *Eur J Biochem* 155 : 295-300, 1986
8. Bonomi F, Cerioli A, Pagani S : Molecular aspects of the removal of ferritin-bound iron by DL-dihydrolipoate. *Biochim Biophys Acta* 994 : 180-186, 1989
9. Biemond P, Swaak AJ, Beindorff CM, Koster JF : Superoxide-dependent and -independent mechanisms of iron mobilization from ferritin by xanthine oxidase. Implications for oxygen-free-radical-induced tissue destruction during ischaemia and inflammation. *Biochem J* 239 : 169-173, 1986
10. Bolann BJ, Ulvik RJ : Release of iron from ferritin by xanthine oxidase. Role of the superoxide radical. *Biochem J* 243 : 55-59, 1987
11. Reif DW, Simmons RD : Nitric oxide mediates iron release from ferritin. *Arch Biochem Biophys* 283 : 537-541, 1990
12. Kidane TZ, Sauble E, Linder MC : Release of iron from ferritin requires lysosomal activity. *Am J Physiol Cell Physiol* 291 : C445-C455, 2006
13. Gryzik M, Srivastava A, Longhi G, Bertuzzi M, Gianoncelli A, Carmona F, Poli M, Arosio P : Expression and characterization of the ferritin binding domain of Nuclear Receptor Co-activator-4 (NCOA4). *Biochim Biophys Acta Gen Subj* 1861 : 2710-2716, 2017.
14. De Domenico I, Ward DM, Kaplan J : Specific iron chelators determine the route of ferritin degradation. *Blood* 114 : 4546-4551, 2009.
15. Koochana PK, Mohantly A, Das S, Subhadarshane B, Satpati S, Dixit A, Sabat SC, Behera RK: Releasing iron from ferritin protein nanocage by reductive method : The role of electron transfer mediator. *Biochim Biophys Acta Gen Subj* 1862 : 1190-1198, 2018.
16. Melman G, Bou-Abdallah F, Vane E, Maura P, Arosio P, Melman A : Iron release from ferritin by flavin nucleotides. *Biochim Biophys Acta* 1830 : 4669-4674, 2013.
17. Johnson LE, Wilkinson T, Arosio P, Melman A, Bou-Abdallah F : Effect of chaotropes on the kinetics of iron release from ferritin by flavin nucleotides. *Biochim Biophys Acta Gen Subj* 1861 : 3257-3262, 2017.
18. Vermilion JL, Coon MJ : Identification of the high and low potential flavins of liver microsomal NADPH-cytochrome P-450 reductase. *J Biol Chem* 253 : 8812-8819, 1978
19. Vermilion JL, Ballou DP, Massey V, Coon MJ : Separate roles for FMN and FAD in catalysis by liver microsomal NADPH-cytochrome P-450 reductase. *J Biol Chem* 256 : 266-277, 1981
20. Marcus RA, Sutin N : Electron transfers in chemistry and biology. *Biochim Biophys Acta* 811 : 265-322, 1985
21. Williams RJ : Electron transfer in biology. *Biochem Int* 18 : 475-499, 1989
22. Jacobs DL, Watt GD, Frankel RB, Papaefthymiou GC : Redox reactions associated with iron release from mammalian ferritin. *Biochemistry* 28 : 1650-1655, 1989
23. Watt GD, Frankel RB, Papaefthymiou GC : Reduction of mammalian ferritin. *Proc Natl Acad Sci U S A* 82 : 3640-3643, 1985
24. Iyanagi T, Makino N, Mason HS : Redox properties of the reduced nicotinamide adenine dinucleotide phosphate-cytochrome P-450 and reduced nicotinamide adenine dinucleotide-cytochrome b5 reductases. *Biochemistry* 13 : 1701-1710, 1974
25. Bogdan AR, Miyazawa M, Hashimoto K, Tsuji Y : Regulators of iron homeostasis : New players in metabolism, cell death, and disease. *Trends Biochem Sci* 41 : 274-286, 2016