

# Neutrosophic Sets and Systems

---

Volume 27

Article 3

---

1-1-2019

## Neutrosophic General Finite Automata

J. Kavikumar

D. Nagarajan

Said Broumi

Florentin Smarandache

Follow this and additional works at: [https://digitalrepository.unm.edu/nss\\_journal](https://digitalrepository.unm.edu/nss_journal)

---

### Recommended Citation

Kavikumar, J.; D. Nagarajan; Said Broumi; and Florentin Smarandache. "Neutrosophic General Finite Automata." *Neutrosophic Sets and Systems* 27, 1 (2019). [https://digitalrepository.unm.edu/nss\\_journal/vol27/iss1/3](https://digitalrepository.unm.edu/nss_journal/vol27/iss1/3)

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact [amywinter@unm.edu](mailto:amywinter@unm.edu).

# Neutrosophic General Finite Automata

J. Kavikumar<sup>1</sup>, D. Nagarajan<sup>2</sup>, Said Broumi<sup>3,\*</sup>, F. Smarandache<sup>4</sup>,  
M. Lathamaheswari<sup>2</sup>, Nur Ain Ebas<sup>1</sup>

<sup>1</sup> Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia, 86400 Malaysia.

E-mail: kavi@uthm.edu.my; nurainebas@gmail.com

<sup>2</sup>Department of Mathematics, Hindustan Institute of Technology & Science, Chennai 603 103, India.

E-mail: dnrmsu2002@yahoo.com; lathamax@gmail.com

<sup>3</sup>Laboratory of Information Processing, Faculty of Science Ben M'Sik, University Hassan II, Casablanca, Morocco.

E-mail: s.broumi@flbenmsik.ma

<sup>4</sup> Department of Mathematics, University of New Mexico, 705 Gurley Avenue, Gallup, NM 87301, USA.

E-mail: smarand@unm.edu.

\*Correspondence: J. Kavikumar (kavi@uthm.edu.my)

**Abstract:** The constructions of finite switchboard state automata is known to be an extension of finite automata in the view of commutative and switching automata. In this research, the idea of a neutrosophic is incorporated in the general fuzzy finite automata and general fuzzy finite switchboard automata to introduce neutrosophic general finite automata and neutrosophic general finite switchboard automata. Moreover, we define the notion of the neutrosophic subsystem and strong neutrosophic subsystem for both structures. We also establish the relationship between the neutrosophic subsystem and neutrosophic strong subsystem.

**Keywords:** Neutrosophic set, General fuzzy automata; switchboard; subsystems.

## 1 Introduction

It is well-known that the simplest and most important type of automata is finite automata. After the introduction of fuzzy set theory by [47] Zadeh in 1965, the first mathematical formulation of fuzzy automata was proposed by [46] Wee in 1967, considered as a generalization of fuzzy automata theory. Consequently, numerous works have been contributed towards the generalization of finite automata by many authors such as Cao and Ezawac [9], Jin et al [18], Jun [20], Li and Qiu [27], Qiu [34], Sato and Kuroki [36], Srivastava and Tiwari [41], Santos [35], Jun and Kavikumar [21], Kavikumar et al, [22, 23, 24] especially the simplest one by Mordeson and Malik [29]. In 2005, the theory of general fuzzy automata was firstly proposed by Doostfateme and Kermer [11] which is used to resolve the problem of assigning membership values to active states of the fuzzy automaton and its multi-membership. Subsequently, as a generalization, the concept of intuitionistic general fuzzy automata has been introduced and studied by Shamsizadeh and Zahedi [37], while Abolpour and Zahedi [6] proposed general fuzzy automata theory based on the complete residuated lattice-valued. As a further

extension, Kavikumar et al [25] studied the notions of general fuzzy switchboard automata. For more details see the recent literature as [5, 12, 13, 14, 15, 16, 17].

The notions of neutrosophic sets was proposed by Smarandache [38, 39], generalizing the existing ordinary fuzzy sets, intuitionistic fuzzy sets and interval-valued fuzzy set in which each element of the universe has the degrees of truth, indeterminacy and falsity and the membership values are lies in  $]0^-, 1^+[$ , the nonstandard unit interval [40] it is an extension from standard interval  $[0,1]$ . It has been shown that fuzzy sets provides limited platform for computational complexity but neutrosophic sets is suitable for it. The neutrosophic sets is an appropriate mechanism for interpreting real-life philosophical problems but not for scientific problems since it is difficult to consolidate. In neutrosophic sets, the degree of indeterminacy can be defined independently since it is quantified explicitly which led to different from intuitionistic fuzzy sets. Single-valued neutrosophic set and interval neutrosophic set are the subclasses of the neutrosophic sets which was introduced by Wang et al. [44, 45] in order to examine kind of real-life and scientific problems. The applications of fuzzy sets have been found very useful in the domain of mathematics and elsewhere. A number of authors have been applied the concept of the neutrosophic set to many other structures especially in algebra [19, 28], decision-making [1, 2, 10, 30], medical [3, 4, 8], water quality management [33] and traffic control management [31, 32].

## 1.1 Motivation

In view of exploiting neutrosophic sets, Tahir et al. [43] introduced and studied the concept of single valued Neutrosophic finite state machine and switchboard state machine. Moreover, the fuzzy finite switchboard state machine is introduced into the context of the interval neutrosophic set in [42]. However, the realm of general structure of fuzzy automata in the neutrosophic environment has not been studied yet in the literature so far. Hence, it is still open to many possibilities for innovative research work especially in the context of neutrosophic general automata and its switchboard automata. The fundamental advantage of incorporating neutrosophic sets into general fuzzy automata is the ability to bring indeterminacy membership and nonmembership in each transitions and active states which help us to overcome the uncertain situation at the time of predicting next active state. Motivated by the work of [11], [36] and [38] the concept of neutrosophic general automata and neutrosophic general switchboard automata are introduced in this paper.

## 1.2 Main Contribution

The purpose of this paper is to introduce the primary algebraic structure of neutrosophic general finite automata and neutrosophic switchboard finite automata. The subsystem and strong subsystem of neutrosophic general finite automata and neutrosophic general finite switchboard f automata are exhibited. The relationship between these subsystems have been discussed and the characterizations of switching and commutative are discussed in the neutrosophic backdrop. We prove that the implication of a strong subsystem is a subsystem of neutrosophic general finite automata. The remainder of this paper is organised as follows. Section 2 provides the results and definitions concerning the general fuzzy automata. Section 3 describes the algebraic properties of the neutrosophic general finite automata. Finally, in section 4, the notion of the neutrosophic general finite switchboard automata is introduced. The paper concludes with Section 5.

## 2 Preliminaries

”For a nonempty set  $X$ ,  $\tilde{P}(X)$  denotes the set of all fuzzy sets on  $X$ .

**Definition 2.1.** [11] A general fuzzy automaton (GFA) is an eight-tuple machine  $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \tilde{\delta}, \omega, F_1, F_2)$  where

- (a)  $Q$  is a finite set of states,  $Q = \{q_1, q_2, \dots, q_n\}$ ,
- (b)  $\Sigma$  is a finite set of input symbols,  $\Sigma = \{a_1, a_2, \dots, a_m\}$ ,
- (c)  $\tilde{R}$  is the set of fuzzy start states,  $\tilde{R} \subseteq \tilde{P}(Q)$ ,
- (d)  $Z$  is a finite set of output symbols,  $Z = \{b_1, b_2, \dots, b_k\}$ ,
- (e)  $\omega : Q \rightarrow Z$  is the non-fuzzy output function,
- (f)  $F_1 : [0, 1] \times [0, 1] \rightarrow [0, 1]$  is the membership assignment function,
- (g)  $\tilde{\delta} : (Q \times [0, 1]) \times \Sigma \times Q \xrightarrow{F_1(\mu, \delta)} [0, 1]$  is the augmented transition function,
- (h)  $F_2 : [0, 1]^* \rightarrow [0, 1]$  is a multi-membership resolution function.

Noted that the function  $F_1(\mu, \delta)$  has two parameters  $\mu$  and  $\delta$ , where  $\mu$  is the membership value of a predecessor and  $\delta$  is the weight of a transition. In this definition, the process that takes place upon the transition from state  $q_i$  to  $q_j$  on input  $a_k$  is represented as:

$$\mu^{t+1}(q_j) = \tilde{\delta}((q_i, \mu^t(q_i)), a_k, q_j) = F_1(\mu^t(q_i), \delta(q_i, a_k, q_j)).$$

This means that the membership value of the state  $q_j$  at time  $t + 1$  is computed by function  $F_1$  using both the membership value of  $q_i$  at time  $t$  and the weight of the transition. The usual options for the function  $F(\mu, \delta)$  are  $\max\{\mu, \delta\}$ ,  $\min\{\mu, \delta\}$  and  $(\mu + \delta)/2$ . The multi-membership resolution function resolves the multi-membership active states and assigns a single membership value to them.

Let  $Q_{act}(t_i)$  be the set of all active states at time  $t_i, \forall i \geq 0$ . We have  $Q_{act}(t_0) = \tilde{R}$ ,

$$Q_{act}(t_i) = \{(q, \mu^{t_i}(q)) : \exists q' \in Q_{act}(t_{i-1}), \exists a \in \Sigma, \delta(q', a, q) \in \Delta\}, \forall i \geq 1.$$

Since  $Q_{act}(t_i)$  is a fuzzy set, in order to show that a state  $q$  belongs to  $Q_{act}(t_i)$  and  $T$  is a subset of  $Q_{act}(t_i)$ , we should write:  $q \in \text{Domain}(Q_{act}(t_i))$  and  $T \subset \text{Domain}(Q_{act}(t_i))$ . Hereafter, we simply denote them as:  $q \in Q_{act}(t_i)$  and  $T \subset Q_{act}(t_i)$ . The combination of the operations of functions  $F_1$  and  $F_2$  on a multi-membership state  $q_j$  leads to the multi-membership resolution algorithm.

**Algorithm 2.2.** [11] (Multi-membership resolution) If there are several simultaneous transitions to the active state  $q_j$  at time  $t + 1$ , the following algorithm will assign a unified membership value to it:

1. Each transition weight  $\tilde{\delta}(q_i, a_k, q_j)$  together with  $\mu^t(q_i)$ , will be processed by the membership assignment function  $F_1$ , and will produce a membership value. Call this  $v_i$ ,

$$v_i = \tilde{\delta}((q_i, \mu^t(q_i)), a_k, q_j) = F_1(\mu^t(q_i), \delta(q_i, a_k, q_j)).$$

2. These membership values are not necessarily equal. Hence, they need to be processed by the multi-membership resolution function  $F_2$ .

3. The result produced by  $F_2$  will be assigned as the instantaneous membership value of the active state  $q_j$ ,

$$\mu^{t+1}(q_j) = F_{2_{i=1}}^n[v_i] = F_{2_{i=1}}^n[F_1(\mu^t(q_i), \delta(q_i, a_k, q_j))],$$

where

- $n$  is the number of simultaneous transitions to the active state  $q_j$  at time  $t + 1$ .
- $\delta(q_i, a_k, q_j)$  is the weight of a transition from  $q_i$  to  $q_j$  upon input  $a_k$ .
- $\mu^t(q_i)$  is the membership value of  $q_i$  at time  $t$ .
- $\mu^{t+1}(q_j)$  is the final membership value of  $q_j$  at time  $t + 1$ .

**Definition 2.3.** Let  $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \tilde{\delta}, \omega, F_1, F_2)$  be a general fuzzy automaton, which is defined in Definition 2.1. The max-min general fuzzy automata is defined of the form:

$$\tilde{F}^* = (Q, \Sigma, \tilde{R}, Z, \tilde{\delta}^*, \omega, F_1, F_2),$$

where  $Q_{act} = \{Q_{act}(t_0), Q_{act}(t_1), \dots\}$  and for every  $i, i \geq 0$ :

$$\tilde{\delta}^*((q, \mu^{t_i}(q)), \Lambda, p) = \begin{cases} 1, & q = p \\ 0, & \text{otherwise} \end{cases}$$

and for every  $i, i \geq 1$ :  $\tilde{\delta}^*((q, \mu^{t_{i-1}}(q)), u_i, p) = \tilde{\delta}((q, \mu^{t_{i-1}}(q)), u_i, p)$ ,

$$\tilde{\delta}^*((q, \mu^{t_{i-1}}(q)), u_i u_{i+1}, p) = \bigvee_{q' \in Q_{act}(t_i)} (\tilde{\delta}((q, \mu^{t_{i-1}}(q)), u_i, q') \wedge \tilde{\delta}((q', \mu^{t_i}(q')), u_{i+1}, p))$$

and recursively

$$\tilde{\delta}^*((q, \mu^{t_0}(q)), u_1 u_2 \dots u_n, p) = \bigvee \{ \tilde{\delta}((q, \mu^{t_0}(q)), u_1, p_1) \wedge \tilde{\delta}((p_1, \mu^{t_1}(p_1)), u_2, p_2) \wedge \dots \wedge \tilde{\delta}((p_{n-1}, \mu^{t_{n-1}}(p_{n-1})), u_n, p) \mid p_1 \in Q_{act}(t_1), p_2 \in Q_{act}(t_2), \dots, p_{n-1} \in Q_{act}(t_{n-1}) \},$$

in which  $u_i \in \Sigma, \forall 1 \leq i \leq n$  and assuming that the entered input at time  $t_i$  be  $u_i, \forall 1 \leq i \leq n - 1$ .

**Definition 2.4.** [13] Let  $\tilde{F}^*$  be a max-min GFA,  $p \in Q, q \in Q_{act}(t_i), i \geq 0$  and  $0 \leq \alpha < 1$ . Then  $p$  is called a successor of  $q$  with threshold  $\alpha$  if there exists  $x \in \Sigma^*$  such that  $\tilde{\delta}^*((q, \mu^{t_j}(q)), x, p) > \alpha$ .

**Definition 2.5.** [13] Let  $\tilde{F}^*$  be a max-min GFA,  $q \in Q_{act}(t_i), i \geq 0$  and  $0 \leq \alpha < 1$ . Also let  $S_\alpha(q)$  denote the set of all successors of  $q$  with threshold  $\alpha$ . If  $T \subseteq Q$ , then  $S_\alpha(T)$  the set of all successors of  $T$  with threshold  $\alpha$  is defined by  $S_\alpha(T) = \bigcup \{S_\alpha(q) : q \in T\}$ .

**Definition 2.6.** [38] Let  $X$  be an universe of discourse. The neutrosophic set is an object having the form  $A = \{\langle x, \mu_1(x), \mu_2(x), \mu_3(x) \rangle \mid \forall x \in X\}$  where the functions can be defined by  $\mu_1, \mu_2, \mu_3 : X \rightarrow ]0, 1[$  and  $\mu_1$  is the degree of membership or truth,  $\mu_2$  is the degree of indeterminacy and  $\mu_3$  is the degree of non-membership or false of the element  $x \in X$  to the set  $A$  with the condition  $0 \leq \mu_1(x) + \mu_2(x) + \mu_3(x) \leq 3$ ."

### 3 Neutrosophic General Finite Automata

**Definition 3.1.** An eight-tuple machine  $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \tilde{\delta}, \omega, F_1, F_2)$  is called neutrosophic general finite automata (NGFA for short), where

1.  $Q$  is a finite set of states,  $Q = \{q_1, q_2, \dots, q_n\}$ ,
2.  $\Sigma$  is a finite set of input symbols,  $\Sigma = \{u_1, u_2, \dots, u_m\}$ ,
3.  $\tilde{R} = \{(q, \mu_1^{t_0}(q), \mu_2^{t_0}(q), \mu_3^{t_0}(q)) | q \in R\}$  is the set of fuzzy start states,  $R \subseteq \tilde{P}(Q)$ ,
4.  $Z$  is a finite set of output symbols,  $Z = \{b_1, b_2, \dots, b_k\}$ ,
5.  $\tilde{\delta} : (Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma \times Q \xrightarrow{F_1(\mu, \delta)} [0, 1] \times [0, 1] \times [0, 1]$  is the neutrosophic augmented transition function,
6.  $\omega : (Q \times [0, 1] \times [0, 1] \times [0, 1]) \rightarrow Z$  is the non-fuzzy output function,
7.  $F_1 = (F_1^\wedge, F_1^{\wedge\vee}, F_1^\vee)$ , where  $F_1^\wedge : [0, 1] \times [0, 1] \rightarrow [0, 1]$ ,  $F_2^{\wedge\vee} : [0, 1] \times [0, 1] \rightarrow [0, 1]$  and  $F_3^\vee : [0, 1] \times [0, 1] \rightarrow [0, 1]$  are the truth, indeterminacy and false membership assignment functions, respectively.  $F_1^\wedge(\mu_1, \tilde{\delta}_1)$ ,  $F_2^{\wedge\vee}(\mu_2, \tilde{\delta}_2)$  and  $F_3^\vee(\mu_3, \tilde{\delta}_3)$  are motivated by two parameters  $\mu_1, \mu_2, \mu_3$  and  $\tilde{\delta}_1, \tilde{\delta}_2, \tilde{\delta}_3$  where  $\mu_1, \mu_2$  and  $\mu_3$  are the truth, indeterminacy and false membership value of a predecessor and  $\tilde{\delta}_1, \tilde{\delta}_2$  and  $\tilde{\delta}_3$  are the truth, indeterminacy and false membership value of a transition,
8.  $F_2 = (F_2^\wedge, F_2^{\wedge\vee}, F_2^\vee)$ , where  $F_2^\wedge : [0, 1]^* \rightarrow [0, 1]$ ,  $F_2^{\wedge\vee} : [0, 1]^* \rightarrow [0, 1]$  and  $F_2^\vee : [0, 1]^* \rightarrow [0, 1]$  are the truth, indeterminacy and false multi-membership resolution function.

**Remark 3.2.** In Definition 3.1, the process that takes place upon the transition from the state  $q_i$  to  $q_j$  on an input  $u_k$  is represented by

$$\begin{aligned} \mu_1^{t_{k+1}}(q_j) &= \tilde{\delta}_1((q_i, \mu_1^{t_k}(q_i)), u_k, q_j) = F_1^\wedge(\mu_1^{t_k}(q_i), \delta_1(q_i, u_k, q_j)) = \bigwedge(\mu_1^{t_k}(q_i), \delta_1(q_i, u_k, q_j)), \\ \mu_2^{t_{k+1}}(q_j) &= \tilde{\delta}_2((q_i, \mu_2^{t_k}(q_i)), u_k, q_j) = F_1^{\wedge\vee}(\mu_2^{t_k}(q_i), \delta_2(q_i, u_k, q_j)) = \begin{cases} \bigvee(\mu_2^{t_k}(q_i), \delta_2(q_i, u_k, q_j)) & \text{if } t_k < t_{k+1} \\ \bigwedge(\mu_2^{t_k}(q_i), \delta_2(q_i, u_k, q_j)) & \text{if } t_k \geq t_{k+1} \end{cases}, \\ \mu_3^{t_{k+1}}(q_j) &= \tilde{\delta}_3((q_i, \mu_3^{t_k}(q_i)), u_k, q_j) = F_1^\vee(\mu_3^{t_k}(q_i), \delta_3(q_i, u_k, q_j)) = \bigvee(\mu_3^{t_k}(q_i), \delta_3(q_i, u_k, q_j)), \end{aligned}$$

where

$$\begin{aligned} \tilde{\delta}((q_i, \mu^t(q_i)), u_k, q_j) &= (\tilde{\delta}_1((q_i, \mu_1^t(q_i)), u_k, q_j), \tilde{\delta}_2((q_i, \mu_2^t(q_i)), u_k, q_j), \tilde{\delta}_3((q_i, \mu_3^t(q_i)), u_k, q_j)) \text{ and} \\ \delta(q_i, u_k, q_j) &= (\delta_1(q_i, u_k, q_j), \delta_2(q_i, u_k, q_j), \delta_3(q_i, u_k, q_j)). \end{aligned}$$

**Remark 3.3.** The algorithm for truth, indeterminacy and false multi-membership resolution for transition function is same as Algorithm 2.2 but the computation depends (see Remark 3.2) on the truth, indeterminacy and false membership assignment function.

**Definition 3.4.** Let  $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \tilde{\delta}, \omega, F_1, F_2)$  be a NGFA. We define the max-min neutrosophic general fuzzy automaton  $\tilde{F}^* = (Q, \Sigma, \tilde{R}, Z, \tilde{\delta}^*, \omega, F_1, F_2)$ , where  $\tilde{\delta}^* : (Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma^* \times Q \rightarrow [0, 1] \times [0, 1] \times [0, 1]$  and define a neutrosophic set  $\tilde{\delta}^* = \langle \tilde{\delta}_1^*, \tilde{\delta}_2^*, \tilde{\delta}_3^* \rangle$  in  $(Q \times [0, 1] \times [0, 1] \times [01]) \times \Sigma^* \times Q$  and for every  $i, i \geq 0$  :

$$\begin{aligned} \tilde{\delta}_1^*((q, \mu^{t_i}(q)), \Lambda, p) &= \begin{cases} 1, & q = p \\ 0, & q \neq p \end{cases} , \\ \tilde{\delta}_2^*((q, \mu^{t_i}(q)), \Lambda, p) &= \begin{cases} 0, & q = p \\ 1, & q \neq p \end{cases} , \\ \tilde{\delta}_3^*((q, \mu^{t_i}(q)), \Lambda, p) &= \begin{cases} 0, & q = p \\ 1, & q \neq p \end{cases} , \end{aligned}$$

and for every  $i, i \geq 1$ :

$$\begin{aligned} \tilde{\delta}_1^*((q, \mu^{t_{i-1}}(q)), u_i, p) &= \tilde{\delta}_1((q, \mu^{t_{i-1}}(q)), u_i, p), \tilde{\delta}_2^*((q, \mu^{t_{i-1}}(q)), u_i, p) = \tilde{\delta}_2((q, \mu^{t_{i-1}}(q)), u_i, p) \\ \tilde{\delta}_3^*((q, \mu^{t_{i-1}}(q)), u_i, p) &= \tilde{\delta}_3((q, \mu^{t_{i-1}}(q)), u_i, p) \end{aligned}$$

and recursively,

$$\begin{aligned} \tilde{\delta}_1^*((q, \mu^{t_0}(q)), u_1 u_2 \cdots u_n, p) &= \bigvee \{ \tilde{\delta}_1((q, \mu^{t_0}(q)), u_1, p_1) \wedge \tilde{\delta}_1((p_1, \mu^{t_1}(p_1)), u_2, p_2) \wedge \cdots \wedge \\ &\quad \tilde{\delta}_1((p_{n-1}, \mu^{t_{n-1}}(p_{n-1})), u_n, p) | p_1 \in Q_{act}(t_1), p_2 \in Q_{act}(t_2), \cdots, p_{n-1} \in Q_{act}(t_{n-1}) \}, \\ \tilde{\delta}_2^*((q, \mu^{t_0}(q)), u_1 u_2 \cdots u_n, p) &= \bigwedge \{ \tilde{\delta}_2((q, \mu^{t_0}(q)), u_1, p_1) \vee \tilde{\delta}_2((p_1, \mu^{t_1}(p_1)), u_2, p_2) \vee \cdots \vee \\ &\quad \tilde{\delta}_2((p_{n-1}, \mu^{t_{n-1}}(p_{n-1})), u_n, p) | p_1 \in Q_{act}(t_1), p_2 \in Q_{act}(t_2), \cdots, p_{n-1} \in Q_{act}(t_{n-1}) \}, \\ \tilde{\delta}_3^*((q, \mu^{t_0}(q)), u_1 u_2 \cdots u_n, p) &= \bigwedge \{ \tilde{\delta}_3((q, \mu^{t_0}(q)), u_1, p_1) \vee \tilde{\delta}_3((p_1, \mu^{t_1}(p_1)), u_2, p_2) \vee \cdots \vee \\ &\quad \tilde{\delta}_3((p_{n-1}, \mu^{t_{n-1}}(p_{n-1})), u_n, p) | p_1 \in Q_{act}(t_1), p_2 \in Q_{act}(t_2), \cdots, p_{n-1} \in Q_{act}(t_{n-1}) \}, \end{aligned}$$

in which  $u_i \in \Sigma, \forall 1 \leq i \leq n$  and assuming that the entered input at time  $t_i$  be  $u_i, \forall 1 \leq i \leq n - 1$ .

**Example 3.5.** Consider the NGFA in Figure 1 with several transition overlaps. Let  $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \tilde{\delta}, \omega, F_1, F_2)$ , where

- $Q = \{q_0, q_1, q_2, q_3, q_4, q_5, q_6, q_7, q_8, q_9\}$  be a set of states,
- $\Sigma = \{a, b\}$  be a set of input symbols,
- $\tilde{R} = \{(q_0, 0.7, 0.5, 0.2), (q_4, 0.6, 0.2, 0.45)\}$ , set of initial states,
- the operation of  $F_1^\wedge, F_1^{\wedge \vee}$  and  $F_1^\vee$  are according to Remark 3.2,
- $Z = \emptyset$  and  $\omega$  are not applicable (output mapping is not of our interest in this paper),
- $\tilde{\delta} : (Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma \times Q \xrightarrow{F_1(\mu, \delta)} [0, 1] \times [0, 1] \times [0, 1]$ , the neutrosophic augmented transition function.

Assuming that  $\tilde{F}$  starts operating at time  $t_0$  and the next three inputs are  $a, b, b$  respectively (one at a time), active states and their membership values at each time step are as follows:




Figure 1: The NGFA of Example 3.5

- At time  $t_0$ :  $Q_{act}(t_0) = \tilde{R} = \{(q_0, 0.7, 0.5, 0.2), (q_4, 0.6, 0.2, 0.45)\}$
- At time  $t_1$ , input is  $a$ . Thus  $q_1, q_5$  and  $q_8$  get activated. Then:

$$\begin{aligned} \mu^{t_1}(q_1) &= \tilde{\delta}((q_0, \mu_1^{t_0}(q_0), \mu_2^{t_0}(q_0), \mu_3^{t_0}(q_0)), a, q_1) \\ &= [F_1^\wedge(\mu_1^{t_0}(q_0), \delta_1(q_0, a, q_1)), F_1^{\wedge\vee}(\mu_2^{t_0}(q_0), \delta_2(q_0, a, q_1)), F_1^\vee(\mu_3^{t_0}(q_0), \delta_3(q_0, a, q_1))] \\ &= [F_1^\wedge(0.7, 0.4), F_1^{\wedge\vee}(0.5, 0.2), F_1^\vee(0.2, 0.3)] = (0.4, 0.2, 0.3), \end{aligned}$$

$$\begin{aligned} \mu^{t_1}(q_8) &= \tilde{\delta}((q_0, \mu_1^{t_0}(q_0), \mu_2^{t_0}(q_0), \mu_3^{t_0}(q_0)), a, q_8) \\ &= [F_1^\wedge(\mu_1^{t_0}(q_0), \delta_1(q_0, a, q_8)), F_1^{\wedge\vee}(\mu_2^{t_0}(q_0), \delta_2(q_0, a, q_8)), F_1^\vee(\mu_3^{t_0}(q_0), \delta_3(q_0, a, q_8))] \\ &= [F_1^\wedge(0.7, 0.7), F_1^{\wedge\vee}(0.5, 0.1), F_1^\vee(0.2, 0.2)] = (0.7, 0.1, 0.2), \end{aligned}$$

but  $q_5$  is multi-membership at  $t_1$ . Then

$$\begin{aligned} \mu^{t_1}(q_5) &= F_2 \underset{i=0\&4}{[F_1[\mu^{t_0}(q_i), \delta(q_i, a, q_5)]]} \\ &= F_2 [F_1[\mu^{t_0}(q_0), \delta(q_0, a, q_5)], F_1[\mu^{t_0}(q_4), \delta(q_4, a, q_5)]] \\ &= F_2 [F_1[(0.7, 0.5, 0.2), (0.3, 0.4, 0.1)], F_1[(0.6, 0.2, 0.45), (0.4, 0.6, 0.5)]] \\ &= (F_2^\wedge [F_1^\wedge(0.7, 0.3), F_1^\wedge(0.6, 0.4)], F_2^{\wedge\vee} [F_1^{\wedge\vee}(0.5, 0.4), F_1^{\wedge\vee}(0.2, 0.6)], \\ &\hspace{15em} F_2^\vee [F_1^\vee(0.2, 0.1), F_1^\vee(0.45, 0.5)]) \\ &= (F_2^\wedge(0.3, 0.4), F_2^{\wedge\vee}(0.4, 0.2), F_2^\vee(0.2, 0.5)) = (0.3, 0.2, 0.5). \end{aligned}$$



Then we have:

$$\begin{aligned} Q_{act}(t_1) &= \{(q_1, \mu^{t_1}(q_1)), (q_5, \mu^{t_1}(q_5)), (q_8, \mu^{t_1}(q_8))\} \\ &= \{(q_1, 0.4, 0.2, 0.3), (q_5, 0.3, 0.2, 0.5), (q_8, 0.7, 0.1, 0.2)\}. \end{aligned}$$

- At  $t_2$  input is  $b$ .  $q_2, q_5, q_6$  and  $q_9$  get activated. Then

$$\begin{aligned} \mu^{t_2}(q_5) &= \tilde{\delta}((q_1, \mu^{t_1}(q_1), \mu^{t_1}(q_1), \mu^{t_1}(q_1)), b, q_5) \\ &= [F_1^\wedge(\mu^{t_1}(q_1), \delta_1(q_1, b, q_5)), F_1^{\wedge\vee}(\mu^{t_1}(q_1), \delta_2(q_1, b, q_5)), F_1^\vee(\mu^{t_1}(q_1), \delta_3(q_1, b, q_5))] \\ &= [F_1^\wedge(0.4, 0.1), F_1^{\wedge\vee}(0.2, 0.4), F_1^\vee(0.3, 0.6)] = (0.1, 0.2, 0.6), \end{aligned}$$

$$\begin{aligned} \mu^{t_2}(q_6) &= \tilde{\delta}((q_5, \mu^{t_1}(q_5), \mu^{t_1}(q_5), \mu^{t_1}(q_5)), b, q_6) \\ &= [F_1^\wedge(\mu^{t_1}(q_5), \delta_1(q_5, b, q_6)), F_1^{\wedge\vee}(\mu^{t_1}(q_5), \delta_2(q_5, b, q_6)), F_1^\vee(\mu^{t_1}(q_5), \delta_3(q_5, b, q_6))] \\ &= [F_1^\wedge(0.3, 0.5), F_1^{\wedge\vee}(0.2, 0.6), F_1^\vee(0.5, 0.2)] = (0.3, 0.2, 0.5), \end{aligned}$$

$$\begin{aligned} \mu^{t_2}(q_9) &= \tilde{\delta}((q_8, \mu^{t_1}(q_8), \mu^{t_1}(q_8), \mu^{t_1}(q_8)), b, q_9) \\ &= [F_1^\wedge(\mu^{t_1}(q_8), \delta_1(q_8, b, q_9)), F_1^{\wedge\vee}(\mu^{t_1}(q_8), \delta_2(q_8, b, q_9)), F_1^\vee(\mu^{t_1}(q_8), \delta_3(q_8, b, q_9))] \\ &= [F_1^\wedge(0.7, 0.5), F_1^{\wedge\vee}(0.1, 0.3), F_1^\vee(0.2, 0.7)] = (0.5, 0.1, 0.7), \end{aligned}$$

but  $q_2$  is multi-membership at  $t_2$ . Then:

$$\begin{aligned} \mu^{t_2}(q_2) &= F_2 \underset{i=1\&5}{[F_1[\mu^{t_1}(q_i), \delta(q_i, b, q_2)]]} \\ &= F_2 [F_1[\mu^{t_1}(q_1), \delta(q_1, b, q_2)], F_1[\mu^{t_1}(q_5), \delta(q_5, b, q_2)]] \\ &= F_2 [F_1[(0.4, 0.2, 0.3), (0.5, 0.3, 0.45)], F_1[(0.3, 0.2, 0.5), (0.1, 0.4, 0.6)]] \\ &= (F_2^\wedge[F_1^\wedge(0.4, 0.5), F_1^\wedge(0.3, 0.1)], F_2^{\wedge\vee}[F_1^{\wedge\vee}(0.2, 0.3), F_1^{\wedge\vee}(0.2, 0.4)], \\ &\quad F_2^\vee[F_1^\vee(0.3, 0.45), F_1^\vee(0.5, 0.6)]) \\ &= (F_2^\wedge(0.4, 0.1), F_2^{\wedge\vee}(0.2, 0.2), F_2^\vee(0.3, 0.5)) = (0.1, 0.2, 0.5). \end{aligned}$$

Then we have:

$$\begin{aligned} Q_{act}(t_2) &= \{(q_2, \mu^{t_2}(q_2)), (q_5, \mu^{t_2}(q_5)), (q_6, \mu^{t_2}(q_6)), (q_9, \mu^{t_2}(q_9))\} \\ &= \{(q_2, 0.1, 0.2, 0.5), (q_5, 0.1, 0.2, 0.6), (q_6, 0.3, 0.2, 0.5), (q_9, 0.5, 0.1, 0.7)\}. \end{aligned}$$

- At  $t_3$  input is  $b$ .  $q_2, q_6, q_7$  and  $q_9$  get activated and none of them is multi-membership. It is easy to verify that:

$$\begin{aligned} Q_{act}(t_3) &= \{(q_2, \mu^{t_3}(q_2)), (q_6, \mu^{t_3}(q_6)), (q_7, \mu^{t_3}(q_7)), (q_9, \mu^{t_3}(q_9))\} \\ &= \{(q_2, 0.1, 0.1, 0.6), (q_6, 0.1, 0.2, 0.6), (q_7, 0.3, 0.1, 0.5), (q_9, 0.3, 0.1, 0.5)\}. \end{aligned}$$

**Proposition 3.6.** *Let  $\tilde{F}$  be a NGFA, if  $\tilde{F}^*$  is a max-min NGFA, then for every  $i \geq 1$ ,*

$$\begin{aligned}\tilde{\delta}_1^*((q, \mu^{t_{i-1}}(q)), xy, p) &= \bigvee_{r \in Q_{act}(t_i)} \left[ \tilde{\delta}_1^*((p, \mu^{t_{i-1}}(p)), x, r) \wedge \tilde{\delta}_1^*((r, \mu^{t_{i-1}}(r)), y, q) \right], \\ \tilde{\delta}_2^*((q, \mu^{t_{i-1}}(q)), xy, p) &= \bigwedge_{r \in Q_{act}(t_i)} \left[ \tilde{\delta}_2^*((p, \mu^{t_{i-1}}(p)), x, r) \vee \tilde{\delta}_2^*((r, \mu^{t_{i-1}}(r)), y, q) \right], \\ \tilde{\delta}_3^*((q, \mu^{t_{i-1}}(q)), xy, p) &= \bigwedge_{r \in Q_{act}(t_i)} \left[ \tilde{\delta}_3^*((p, \mu^{t_{i-1}}(p)), x, r) \vee \tilde{\delta}_3^*((r, \mu^{t_{i-1}}(r)), y, q) \right],\end{aligned}$$

for all  $p, q \in Q$  and  $x, y \in \Sigma^*$ .

*Proof.* Since  $p, q \in Q$  and  $x, y \in \Sigma^*$ , we prove the result by induction on  $|y| = n$ . First, we assume that  $n = 0$ , then  $y = \Lambda$  and so  $xy = x\Lambda = x$ . Thus, for all  $r \in Q_{act}(t_i)$

$$\begin{aligned}\bigvee \left[ \tilde{\delta}_1^*((p, \mu^{t_{i-1}}(p)), x, r) \wedge \tilde{\delta}_1^*((r, \mu^{t_{i-1}}(r)), y, q) \right] &= \bigvee \left[ \tilde{\delta}_1^*((p, \mu^{t_{i-1}}(p)), x, r) \wedge \tilde{\delta}_1^*((r, \mu^{t_{i-1}}(r)), \Lambda, q) \right] \\ &= \tilde{\delta}_1^*((p, \mu^{t_{i-1}}(p)), x, r) = \tilde{\delta}_1^*((q, \mu^{t_{i-1}}(q)), xy, p), \\ \bigwedge \left[ \tilde{\delta}_2^*((p, \mu^{t_{i-1}}(p)), x, r) \vee \tilde{\delta}_2^*((r, \mu^{t_{i-1}}(r)), y, q) \right] &= \bigwedge \left[ \tilde{\delta}_2^*((p, \mu^{t_{i-1}}(p)), x, r) \vee \tilde{\delta}_2^*((r, \mu^{t_{i-1}}(r)), \Lambda, q) \right] \\ &= \tilde{\delta}_2^*((p, \mu^{t_{i-1}}(p)), x, r) = \tilde{\delta}_2^*((q, \mu^{t_{i-1}}(q)), xy, p), \\ \bigwedge \left[ \tilde{\delta}_3^*((p, \mu^{t_{i-1}}(p)), x, r) \vee \tilde{\delta}_3^*((r, \mu^{t_{i-1}}(r)), y, q) \right] &= \bigwedge \left[ \tilde{\delta}_3^*((p, \mu^{t_{i-1}}(p)), x, r) \vee \tilde{\delta}_3^*((r, \mu^{t_{i-1}}(r)), \Lambda, q) \right] \\ &= \tilde{\delta}_3^*((p, \mu^{t_{i-1}}(p)), x, r) = \tilde{\delta}_3^*((q, \mu^{t_{i-1}}(q)), xy, p).\end{aligned}$$

The result holds for  $n = 0$ . Now, continue the result is true for all  $u \in \Sigma^*$  with  $|u| = n - 1$ , where  $n > 0$ . Let  $y = ua$ , where  $a \in \Sigma$  and  $u \in \Sigma^*$ . Then

$$\begin{aligned}\tilde{\delta}_1^*((q, \mu^{t_{i-1}}(q)), xy, p) &= \tilde{\delta}_1^*((q, \mu^{t_{i-1}}(q)), xua, p) = \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1^*((q, \mu^{t_{i-1}}(q)), xu, r) \wedge \tilde{\delta}_1^*((r, \mu^{t_i}(r)), a, p) \right) \\ &= \bigvee_{r \in Q_{act}(t_i)} \left( \bigvee_{s \in Q_{act}(t_i)} \left( \tilde{\delta}_1^*((q, \mu^{t_{i-1}}(q)), x, s) \wedge \tilde{\delta}_1^*((s, \mu^{t_{i-1}}(s)), u, r) \right) \wedge \tilde{\delta}_1^*((r, \mu^{t_i}(r)), a, p) \right) \\ &= \bigvee_{r, s \in Q_{act}(t_i)} \left( \tilde{\delta}_1^*((q, \mu^{t_{i-1}}(q)), x, s) \wedge \tilde{\delta}_1^*((s, \mu^{t_{i-1}}(s)), u, r) \wedge \tilde{\delta}_1^*((r, \mu^{t_i}(r)), a, p) \right) \\ &= \bigvee_{s \in Q_{act}(t_i)} \left( \tilde{\delta}_1^*((q, \mu^{t_{i-1}}(q)), x, s) \wedge \left( \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1^*((s, \mu^{t_{i-1}}(s)), u, r) \wedge \tilde{\delta}_1^*((r, \mu^{t_i}(r)), a, p) \right) \right) \right) \\ &= \bigvee_{s \in Q_{act}(t_i)} \left( \tilde{\delta}_1^*((q, \mu^{t_{i-1}}(q)), x, s) \wedge \tilde{\delta}_1^*((s, \mu^{t_i}(r)), ua, p) \right) = \bigvee_{s \in Q_{act}(t_i)} \left( \tilde{\delta}_1^*((q, \mu^{t_{i-1}}(q)), x, s) \wedge \tilde{\delta}_1^*((s, \mu^{t_i}(r)), y, p) \right), \\ \tilde{\delta}_2^*((q, \mu^{t_{i-1}}(q)), xy, p) &= \tilde{\delta}_2^*((q, \mu^{t_{i-1}}(q)), xua, p) = \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2^*((q, \mu^{t_{i-1}}(q)), xu, r) \vee \tilde{\delta}_2^*((r, \mu^{t_i}(r)), a, p) \right) \\ &= \bigwedge_{r \in Q_{act}(t_i)} \left( \bigwedge_{s \in Q_{act}(t_i)} \left( \tilde{\delta}_2^*((q, \mu^{t_{i-1}}(q)), x, s) \vee \tilde{\delta}_2^*((s, \mu^{t_{i-1}}(s)), u, r) \right) \vee \tilde{\delta}_2^*((r, \mu^{t_i}(r)), a, p) \right) \\ &= \bigwedge_{r, s \in Q_{act}(t_i)} \left( \tilde{\delta}_2^*((q, \mu^{t_{i-1}}(q)), x, s) \vee \tilde{\delta}_2^*((s, \mu^{t_{i-1}}(s)), u, r) \vee \tilde{\delta}_2^*((r, \mu^{t_i}(r)), a, p) \right) \\ &= \bigwedge_{s \in Q_{act}(t_i)} \left( \tilde{\delta}_2^*((q, \mu^{t_{i-1}}(q)), x, s) \vee \left( \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2^*((s, \mu^{t_{i-1}}(s)), u, r) \vee \tilde{\delta}_2^*((r, \mu^{t_i}(r)), a, p) \right) \right) \right) \\ &= \bigwedge_{s \in Q_{act}(t_i)} \left( \tilde{\delta}_2^*((q, \mu^{t_{i-1}}(q)), x, s) \vee \tilde{\delta}_2^*((s, \mu^{t_i}(r)), ua, p) \right) = \bigwedge_{s \in Q_{act}(t_i)} \left( \tilde{\delta}_2^*((q, \mu^{t_{i-1}}(q)), x, s) \vee \tilde{\delta}_2^*((s, \mu^{t_i}(r)), y, p) \right),\end{aligned}$$

$$\begin{aligned}
 \tilde{\delta}_3^*((q, \mu^{t_{i-1}}(q)), xy, p) &= \tilde{\delta}_3^*((q, \mu^{t_{i-1}}(q)), xua, p) = \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_3((q, \mu^{t_{i-1}}(q)), xu, r) \vee \tilde{\delta}_3((r, \mu^{t_i}(r)), a, p) \right) \\
 &= \bigwedge_{r \in Q_{act}(t_i)} \left( \bigwedge_{s \in Q_{act}(t_i)} (\tilde{\delta}_3((q, \mu^{t_{i-1}}(q)), x, s) \vee \tilde{\delta}_3((s, \mu^{t_{i-1}}(s)), u, r)) \vee \tilde{\delta}_3((r, \mu^{t_i}(r)), a, p) \right) \\
 &= \bigwedge_{r, s \in Q_{act}(t_i)} (\tilde{\delta}_3((q, \mu^{t_{i-1}}(q)), x, s) \vee \tilde{\delta}_3((s, \mu^{t_{i-1}}(s)), u, r) \vee \tilde{\delta}_3((r, \mu^{t_i}(r)), a, p)) \\
 &= \bigwedge_{s \in Q_{act}(t_i)} (\tilde{\delta}_3((q, \mu^{t_{i-1}}(q)), x, s) \vee (\bigwedge_{r \in Q_{act}(t_i)} (\tilde{\delta}_3((s, \mu^{t_{i-1}}(s)), u, r) \vee \tilde{\delta}_3((r, \mu^{t_i}(r)), a, p)))) \\
 &= \bigwedge_{s \in Q_{act}(t_i)} (\tilde{\delta}_3((q, \mu^{t_{i-1}}(q)), x, s) \vee \tilde{\delta}_3((s, \mu^{t_i}(r)), ua, p)) = \bigwedge_{s \in Q_{act}(t_i)} (\tilde{\delta}_3((q, \mu^{t_{i-1}}(q)), x, s) \vee \tilde{\delta}_3((s, \mu^{t_i}(r)), y, p)).
 \end{aligned}$$

Hence the result is valid for  $|y| = n$ . This completes the proof. □

**Definition 3.7.** Let  $\tilde{F}^*$  be a max-min NGFA,  $p \in Q, q \in Q_{act}(t_i), i \geq 0$  and  $0 \leq \alpha < 1$ . Then  $p$  is called a successor of  $q$  with threshold  $\alpha$  if there exists  $x \in \Sigma^*$  such that  $\tilde{\delta}_1^*((q, \mu_1^{t_j}(q)), x, p) > \alpha, \tilde{\delta}_2^*((q, \mu_2^{t_j}(q)), x, p) < \alpha$  and  $\tilde{\delta}_3^*((q, \mu_3^{t_j}(q)), x, p) < \alpha$ .

**Definition 3.8.** Let  $\tilde{F}^*$  be a max-min NGFA,  $q \in Q_{act}(t_i), i \geq 0$  and  $0 \leq \alpha < 1$ . Also let  $S_\alpha(q)$  denote the set of all successors of  $q$  with threshold  $\alpha$ . If  $T \subseteq Q$ , then  $S_\alpha(T)$  the set of all successors of  $T$  with threshold  $\alpha$  is defined by  $S_\alpha(T) = \bigcup \{S_\alpha(q) : q \in T\}$ .

**Definition 3.9.** Let  $\tilde{F}^*$  be a max-min NGFA. Let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  and  $\tilde{\delta}^* = \langle \tilde{\delta}_1^*, \tilde{\delta}_2^*, \tilde{\delta}_3^* \rangle$  in  $(Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma^* \times Q$  be a neutrosophic set in  $Q$ . Then  $\mu$  is a neutrosophic subsystem of  $\tilde{F}^*$ , say  $\mu \subseteq \tilde{F}^*$  if for every  $j, 1 \leq j \leq k$  such that  $\mu_1^{t_j}(p) \geq \tilde{\delta}_1^*((q, \mu_1^{t_j}(q)), x, p), \mu_2^{t_j}(p) \leq \tilde{\delta}_2^*((q, \mu_2^{t_j}(q)), x, p), \mu_3^{t_j}(p) \leq \tilde{\delta}_3^*((q, \mu_3^{t_j}(q)), x, p). \forall q, p \in Q$  and  $x \in \Sigma^*$ .

**Example 3.10.** Let  $Q = \{p, q\}, \Sigma = \{a\}$ . Let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  and  $\tilde{\delta}^* = \langle \tilde{\delta}_1^*, \tilde{\delta}_2^*, \tilde{\delta}_3^* \rangle$  in  $(Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma^* \times Q$  be a neutrosophic set in  $Q$  such that  $\mu_1^{t_j}(p) = 0.8, \mu_2^{t_j}(p) = 0.7, \mu_3^{t_j}(p) = 0.5, \mu_1^{t_j}(q) = 0.5, \mu_2^{t_j}(q) = 0.6, \mu_3^{t_j}(q) = 0.8, \delta_1(q, x, p) = 0.7, \delta_2(q, x, p) = 0.9$  and  $\delta_3(q, x, p) = 0.7$ . Then

$$\begin{aligned}
 \tilde{\delta}_1^*((q, \mu_1^{t_j}(q)), x, p) &= F_1^\wedge(\mu_1^{t_j}(q), \delta_1(q, x, p)) = \min\{0.5, 0.7\} = 0.5 \leq \mu_1^{t_j}(p), \\
 \tilde{\delta}_2^*((q, \mu_2^{t_j}(q)), x, p) &= F_2^{\wedge \vee}(\mu_2^{t_j}(q), \delta_2(q, x, p)) = \max\{0.6, 0.9\} = 0.9 \geq \mu_2^{t_j}(p), \text{ (since } t < t_j) \\
 \tilde{\delta}_3^*((q, \mu_3^{t_j}(q)), x, p) &= F_3^\vee(\mu_3^{t_j}(q), \delta_3(q, x, p)) = \max\{0.8, 0.7\} = 0.8 \geq \mu_3^{t_j}(p).
 \end{aligned}$$

Hence  $\mu$  is a neutrosophic subsystem of  $\tilde{F}^*$ .

**Theorem 3.11.** Let  $\tilde{F}^*$  be a NGFA and let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  and  $\tilde{\delta}^* = \langle \tilde{\delta}_1^*, \tilde{\delta}_2^*, \tilde{\delta}_3^* \rangle$  in  $(Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma^* \times Q$  be a neutrosophic set in  $Q$ . Then  $\mu$  is a neutrosophic subsystem of  $\tilde{F}^*$  if and only if  $\mu_1^{t_j}(p) \geq \tilde{\delta}_1^*((q, \mu_1^{t_j}(q)), x, p), \mu_2^{t_j}(p) \leq \tilde{\delta}_2^*((q, \mu_2^{t_j}(q)), x, p), \mu_3^{t_j}(p) \leq \tilde{\delta}_3^*((q, \mu_3^{t_j}(q)), x, p),$  for all  $q \in Q_{act}(t_j), p \in Q$  and  $x \in \Sigma^*$ .

*Proof.* Suppose that  $\mu$  is a neutrosophic subsystem of  $\tilde{F}^*$ . Let  $q \in Q_{act}(t_j), p \in Q$  and  $x \in \Sigma^*$ . The proof is by induction on  $|x| = n$ . If  $n = 0$ , then  $x = \Lambda$ . Now if  $q = p$ , then  $\tilde{\delta}_1^*((p, \mu_1^{t_i}(p)), \Lambda, p) = F_1^\wedge(\mu_1^{t_i}(p), \tilde{\delta}_1(p, \Lambda, p)) = \mu_1^{t_i}(p), \tilde{\delta}_2^*((p, \mu_2^{t_i}(p)), \Lambda, p) = F_1^{\wedge \vee}(\mu_2^{t_i}(p), \tilde{\delta}_2(p, \Lambda, p)) = \mu_2^{t_i}(p), \tilde{\delta}_3^*((p, \mu_3^{t_i}(p)), \Lambda, p) = F_1^\vee(\mu_3^{t_i}(p), \tilde{\delta}_3(p, \Lambda, p)) = \mu_3^{t_i}(p).$

If  $q \neq p$ , then  $\tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), \Lambda, p) = F_1^\wedge(\mu_1^{t_i}(q), \tilde{\delta}_1(q, \Lambda, p)) = 0 \leq \mu_1^{t_j}(p), \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), \Lambda, p) = F_1^{\wedge \vee}(\mu_2^{t_i}(q), \tilde{\delta}_2(q, \Lambda, p)) = 1 \geq \mu_2^{t_j}(p), \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), \Lambda, p) = F_1^\vee(\mu_3^{t_i}(q), \tilde{\delta}_3(q, \Lambda, p)) = 1 \geq \mu_3^{t_j}(p).$

Hence the result is true for  $n = 0$ . For now, we assume that the result is valid for all  $y \in \Sigma^*$  with  $|y| = n - 1$ ,  $n > 0$ . For the  $y$  above, let  $x = u_1 \cdots u_n$  where  $u_i \in \Sigma, i = 1, 2, \dots, n$ . Then

$$\begin{aligned} \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), x, p) &= \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), u_1 \cdots u_n, p) = \bigvee \left( \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), u_1, r_1) \wedge \cdots \wedge \tilde{\delta}_1^*((r_{n-1}, \mu_1^{t_{i+n}}(r_{n-1})), u_n, p) \right) \\ &\leq \bigvee \left( \tilde{\delta}_1^*((r_{n-1}, \mu_1^{t_{i+n}}(r_{n-1})), u_n, p) \mid r_{n-1} \in Q_{(act)}(t_{i+n}) \right) \leq \bigvee \mu_1^{t_j}(p) = \mu_1^{t_j}(p), \end{aligned}$$

$$\begin{aligned} \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), x, p) &= \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), u_1 \cdots u_n, p) = \bigwedge \left( \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), u_1, r_1) \vee \cdots \vee \tilde{\delta}_2^*((r_{n-1}, \mu_2^{t_{i+n}}(r_{n-1})), u_n, p) \right) \\ &\leq \bigwedge \left( \tilde{\delta}_2^*((r_{n-1}, \mu_2^{t_{i+n}}(r_{n-1})), u_n, p) \mid r_{n-1} \in Q_{(act)}(t_{i+n}) \right) \leq \bigwedge \mu_2^{t_j}(p) = \mu_2^{t_j}(p), \end{aligned}$$

$$\begin{aligned} \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), x, p) &= \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), u_1 \cdots u_n, p) = \bigwedge \left( \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), u_1, r_1) \vee \cdots \vee \tilde{\delta}_3^*((r_{n-1}, \mu_3^{t_{i+n}}(r_{n-1})), u_n, p) \right) \\ &\leq \bigwedge \left( \tilde{\delta}_3^*((r_{n-1}, \mu_3^{t_{i+n}}(r_{n-1})), u_n, p) \mid r_{n-1} \in Q_{(act)}(t_{i+n}) \right) \leq \bigwedge \mu_3^{t_j}(p) = \mu_3^{t_j}(p), \end{aligned}$$

where  $r_1 \in Q_{(act)}(t_{i+1}) \cdots r_{n-1} \in Q_{(act)}(t_{i+n})$ . Hence  $\mu_1^{t_j}(p) \geq \tilde{\delta}_1^*((q, \mu_1^{t_j}(q)), x, p)$ ,  $\mu_2^{t_j}(p) \leq \tilde{\delta}_2^*((q, \mu_2^{t_j}(q)), x, p)$ ,  $\mu_3^{t_j}(p) \leq \tilde{\delta}_3^*((q, \mu_3^{t_j}(q)), x, p)$ . The converse is trivial. This proof is completed.  $\square$

**Definition 3.12.** Let  $\tilde{F}^*$  be a NGFA. Let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  and  $\tilde{\delta}^* = \langle \tilde{\delta}_1^*, \tilde{\delta}_2^*, \tilde{\delta}_3^* \rangle$  in  $(Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma^* \times Q$  be a neutrosophic set in  $Q$ . Then  $\mu$  is a neutrosophic strong subsystem of  $\tilde{F}^*$ , say  $\mu \subseteq \tilde{F}^*$ , if for every  $i, 1 \leq i \leq k$  such that  $p \in S_\alpha(q)$ , then for  $q, p \in Q$  and  $x \in \Sigma$ ,  $\mu_1^{t_j}(p) \geq \mu_1^{t_j}(q)$ ,  $\mu_2^{t_j}(p) \leq \mu_2^{t_j}(q)$ ,  $\mu_3^{t_j}(p) \leq \mu_3^{t_j}(q)$ , for every  $1 \leq j \leq k$ .

**Theorem 3.13.** Let  $\tilde{F}^*$  be a NGFA and let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  and  $\tilde{\delta}^* = \langle \tilde{\delta}_1^*, \tilde{\delta}_2^*, \tilde{\delta}_3^* \rangle$  in  $(Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma^* \times Q$  be a neutrosophic set in  $Q$ . Then  $\mu$  is a strong neutrosophic subsystem of  $\tilde{F}^*$  if and only if there exists  $x \in \Sigma^*$  such that  $p \in S_\alpha(q)$ , then  $\mu_1^{t_j}(p) \geq \mu_1^{t_j}(q)$ ,  $\mu_2^{t_j}(p) \leq \mu_2^{t_j}(q)$ ,  $\mu_3^{t_j}(p) \leq \mu_3^{t_j}(q)$ , for all  $q \in Q_{(act)}(t_j), p \in Q$ .

*Proof.* Suppose that  $\mu$  is a strong neutrosophic subsystem of  $\tilde{F}^*$ . Let  $q \in Q_{(act)}(t_j), p \in Q$  and  $x \in \Sigma^*$ . The proof is by induction on  $|x| = n$ . If  $n = 0$ , then  $x = \Lambda$ . Now if  $q = p$ , then  $\delta_1^*((p, \mu_1^{t_i}(p)), \Lambda, p) = 1$ ,  $\delta_2^*((p, \mu_2^{t_i}(p)), \Lambda, p) = 0$ ,  $\delta_3^*((p, \mu_3^{t_i}(p)), \Lambda, p) = 0$  and  $\mu_1^{t_j}(p) = \mu_1^{t_j}(p)$ ,  $\mu_2^{t_j}(p) = \mu_2^{t_j}(p)$ ,  $\mu_3^{t_j}(p) = \mu_3^{t_j}(p)$ . If  $q \neq p$ , then  $\tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), \Lambda, p) = F_1^\wedge(\mu_1^{t_i}(q), \tilde{\delta}_1(q, \Lambda, p)) = c \leq \mu_1^{t_j}(p)$ ,  $\tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), \Lambda, p) = F_1^\vee(\mu_2^{t_i}(q), \tilde{\delta}_2(q, \Lambda, p)) = d \geq \mu_2^{t_j}(p)$ ,  $\tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), \Lambda, p) = F_1^\vee(\mu_3^{t_i}(q), \tilde{\delta}_3(q, \Lambda, p)) = e \geq \mu_3^{t_j}(p)$ . Hence the result is true for  $n = 0$ . For now, we assume that the result is valid for all  $u \in \Sigma^*$  with  $|u| = n - 1$ ,  $n > 0$ . For the  $u$  above, let  $x = u_1 \cdots u_n$  where  $u_i \in \Sigma^*, i = 1, 2, \dots, n$ . Suppose that  $\tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), x, p) > c$ ,  $\tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), x, p) < d$ ,  $\tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), x, p) < e$ . Then

$$\begin{aligned} \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), u_1 \cdots u_n, p) &= \bigvee \left\{ \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), u_1, p_1) \wedge \cdots \wedge \tilde{\delta}_1^*((p_{n-1}, \mu_1^{t_{i+n}}(p_{n-1})), u_n, p) \right\} > c, \\ \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), u_1 \cdots u_n, p) &= \bigwedge \left\{ \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), u_1, p_1) \vee \cdots \vee \tilde{\delta}_2^*((p_{n-1}, \mu_2^{t_{i+n}}(p_{n-1})), u_n, p) \right\} < d, \\ \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), u_1 \cdots u_n, p) &= \bigwedge \left\{ \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), u_1, p_1) \vee \cdots \vee \tilde{\delta}_3^*((p_{n-1}, \mu_3^{t_{i+n}}(p_{n-1})), u_n, p) \right\} < e, \end{aligned}$$

where  $p_1 \in Q_{(act)}(t_i), \dots, p_{n-1} \in Q_{(act)}(t_{i+n})$ .

This implies that  $\tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), u_1, p_1) > c, \dots, \tilde{\delta}_1^*((p_{n-1}, \mu_1^{t_{i+n}}(p_{n-1})), u_n, p) > c, \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), u_1, p_1) < d, \dots, \tilde{\delta}_2^*((p_{n-1}, \mu_2^{t_{i+n}}(p_{n-1})), u_n, p) < d, \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), u_1, p_1) < e, \dots, \tilde{\delta}_3^*((p_{n-1}, \mu_3^{t_{i+n}}(p_{n-1})), u_n, p) < e$ . Hence  $\mu_1^{t_j}(p) \geq \mu_1^{t_{i+n}}(p_{n-1}), \mu_1^{t_{i+n}}(p) \geq \mu_1^{t_{i+n-1}}(p_{n-2}), \dots, \mu_1^{t_i}(p_1) \geq \mu_1^{t_j}(q), \mu_2^{t_j}(p) \leq \mu_2^{t_{i+n}}(p_{n-1}), \mu_2^{t_{i+n}}(p) \leq \mu_2^{t_{i+n-1}}(p_{n-2}), \dots, \mu_2^{t_i}(p_1) \leq \mu_2^{t_j}(q), \mu_3^{t_j}(p) \leq \mu_3^{t_{i+n}}(p_{n-1}), \mu_3^{t_{i+n}}(p) \leq \mu_3^{t_{i+n-1}}(p_{n-2}), \dots, \mu_3^{t_i}(p_1) \leq \mu_3^{t_j}(q)$ . Thus  $\mu_1^{t_j}(p) \geq \mu_1^{t_j}(q), \mu_2^{t_j}(p) \leq \mu_2^{t_j}(q), \mu_3^{t_j}(p) \leq \mu_3^{t_j}(q)$ . The converse is trivial. The proof is completed.  $\square$

## 4 Neutrosophic General Finite Switchboard Automata

**Definition 4.1.** Let  $\tilde{F}^*$  be a max-min NGFA. Let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  and  $\tilde{\delta}^* = \langle \tilde{\delta}_1^*, \tilde{\delta}_2^*, \tilde{\delta}_3^* \rangle$  be a neutrosophic set in  $(Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma \times Q$  in  $Q$ . Then

1.  $\tilde{F}^*$  is switching, if it satisfies  $\forall p, q \in Q, a \in \Sigma$  and for every  $i, i \geq 0$ ,  

$$\tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), a, p) = \tilde{\delta}_1^*((p, \mu_1^{t_i}(p)), a, q), \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), a, p) = \tilde{\delta}_2^*((p, \mu_2^{t_i}(p)), a, q),$$

$$\tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), a, p) = \tilde{\delta}_3^*((p, \mu_3^{t_i}(p)), a, q).$$
2.  $\tilde{F}^*$  is commutative, if it satisfies  $\forall p, q \in Q, a, b \in \Sigma$  and for every  $i, i \geq 1$ ,  $\tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), ab, p) = \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), ba, p), \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), ab, p) = \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), ba, p),$   

$$\tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), ab, p) = \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), ba, p).$$
3.  $\tilde{F}^*$  is Neutrosophic General Finite Switchboard Automata (NGFSA, for short), if  $\tilde{F}^*$  satisfies both switching and commutative.

**Proposition 4.2.** Let  $\tilde{F}$  be a NGFA, if  $\tilde{F}^*$  is a commutative NGFSA, then for every  $i \geq 1$ ,

$$\begin{aligned} \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), xa, p) &= \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), ax, p), \\ \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), xa, p) &= \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), ax, p), \\ \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), xa, p) &= \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), ax, p), \end{aligned}$$

for all  $q \in Q_{act}(t_{i-1}), p \in S_c(q), a \in \Sigma$  and  $x \in \Sigma^*$ .

*Proof.* Since  $p \in S_c(q)$  then  $q \in Q_{act}(t_{i-1})$  and  $|x| = n$ . If  $n = 0$ , then  $x = \Lambda$ . Thus

$$\begin{aligned} \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), xa, p) &= \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), \Lambda a, p) = \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), a, p) = \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), a\Lambda, p) \\ &= \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), ax, p), \\ \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), xa, p) &= \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), \Lambda a, p) = \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), a, p) = \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), a\Lambda, p) \\ &= \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), ax, p), \\ \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), xa, p) &= \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), \Lambda a, p) = \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), a, p) = \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), a\Lambda, p) \\ &= \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), ax, p). \end{aligned}$$

Suppose the result is true for all  $u \in \Sigma^*$  with  $|u| = n - 1$ , where  $n > 0$ . Let  $x = ub$ , where  $b \in \Sigma$ . Then

$$\begin{aligned} \tilde{\delta}_1^*((q, \mu_1^{t_i-1}(q)), xa, p) &= \tilde{\delta}_1^*((q, \mu_1^{t_i-1}(q)), uba, p) = \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1((q, \mu_1^{t_i-1}(q)), u, r) \wedge \tilde{\delta}_1((r, \mu_1^{t_i}(r)), ba, p) \right) \\ &= \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1((q, \mu_1^{t_i-1}(q)), u, r) \wedge \tilde{\delta}_1((r, \mu_1^{t_i}(r)), ab, p) \right) = \tilde{\delta}_1^*((q, \mu_1^{t_i-1}(q)), uab, p) \\ &= \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1((q, \mu_1^{t_i-1}(q)), ua, r) \wedge \tilde{\delta}_1((r, \mu_1^{t_i}(r)), b, p) \right) \\ &= \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1((q, \mu_1^{t_i-1}(q)), au, r) \wedge \tilde{\delta}_1((r, \mu_1^{t_i}(r)), b, p) \right) = \tilde{\delta}_1^*((q, \mu_1^{t_i-1}(q)), aub, p) = \tilde{\delta}_1^*((q, \mu_1^{t_i-1}(q)), ax, p), \end{aligned}$$

$$\begin{aligned} \tilde{\delta}_2^*((q, \mu_2^{t_i-1}(q)), xa, p) &= \tilde{\delta}_2^*((q, \mu_2^{t_i-1}(q)), uba, p) = \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2((q, \mu_2^{t_i-1}(q)), u, r) \vee \tilde{\delta}_2((r, \mu_2^{t_i}(r)), ba, p) \right) \\ &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2((q, \mu_2^{t_i-1}(q)), u, r) \vee \tilde{\delta}_2((r, \mu_2^{t_i}(r)), ab, p) \right) = \tilde{\delta}_2^*((q, \mu_2^{t_i-1}(q)), uab, p) \\ &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2((q, \mu_2^{t_i-1}(q)), ua, r) \vee \tilde{\delta}_2((r, \mu_2^{t_i}(r)), b, p) \right) \\ &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2((q, \mu_2^{t_i-1}(q)), au, r) \vee \tilde{\delta}_2((r, \mu_2^{t_i}(r)), b, p) \right) = \tilde{\delta}_2^*((q, \mu_2^{t_i-1}(q)), aub, p) = \tilde{\delta}_2^*((q, \mu_2^{t_i-1}(q)), ax, p), \end{aligned}$$

$$\begin{aligned} \tilde{\delta}_3^*((q, \mu_3^{t_i-1}(q)), xa, p) &= \tilde{\delta}_3^*((q, \mu_3^{t_i-1}(q)), uba, p) = \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_3((q, \mu_3^{t_i-1}(q)), u, r) \vee \tilde{\delta}_3((r, \mu_3^{t_i}(r)), ba, p) \right) \\ &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_3((q, \mu_3^{t_i-1}(q)), u, r) \vee \tilde{\delta}_3((r, \mu_3^{t_i}(r)), ab, p) \right) = \tilde{\delta}_3^*((q, \mu_3^{t_i-1}(q)), uab, p) \\ &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_3((q, \mu_3^{t_i-1}(q)), ua, r) \vee \tilde{\delta}_3((r, \mu_3^{t_i}(r)), b, p) \right) \\ &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_3((q, \mu_3^{t_i-1}(q)), au, r) \vee \tilde{\delta}_3((r, \mu_3^{t_i}(r)), b, p) \right) = \tilde{\delta}_3^*((q, \mu_3^{t_i-1}(q)), aub, p) = \tilde{\delta}_3^*((q, \mu_3^{t_i-1}(q)), ax, p). \end{aligned}$$

This completes the proof. □

**Proposition 4.3.** *Let  $\tilde{F}$  be a NGFA, if  $\tilde{F}^*$  is a switching NGFSA, then for every  $i \geq 0$ ,  $\tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), x, p) = \tilde{\delta}_1^*((p, \mu_1^{t_i}(p)), x, q)$ ,  $\tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), x, p) = \tilde{\delta}_2^*((p, \mu_2^{t_i}(p)), x, q)$ ,  $\tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), x, p) = \tilde{\delta}_3^*((p, \mu_3^{t_i}(p)), x, q)$ , for all  $p, q \in Q_{act}(t_i)$  and  $x \in \Sigma^*$ .*

*Proof.* Since  $p, q \in Q_{act}(t_i)$  and  $x \in \Sigma^*$ , we prove the result by induction on  $|x| = n$ . First, we assume that  $x = \Lambda$ , whenever  $n = 0$ . Then we have  $\tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), x, p) = \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), \Lambda, p) = \tilde{\delta}_1^*((p, \mu_1^{t_i}(p)), \Lambda, q) = \tilde{\delta}_1^*((p, \mu_1^{t_i}(p)), x, q)$ ,  $\tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), x, p) = \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), \Lambda, p) = \tilde{\delta}_2^*((p, \mu_2^{t_i}(p)), \Lambda, q) = \tilde{\delta}_2^*((p, \mu_2^{t_i}(p)), x, q)$ ,  $\tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), x, p) = \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), \Lambda, p) = \tilde{\delta}_3^*((p, \mu_3^{t_i}(p)), \Lambda, q) = \tilde{\delta}_3^*((p, \mu_3^{t_i}(p)), x, q)$ . Thus, the theorem holds for  $x = \Lambda$ . Now, we assume that the results holds for all  $u \in \Sigma^*$  such that  $|u| = n - 1$  and  $n > 0$ . Let

$a \in \Sigma$  and  $x \in \Sigma^*$  be such that  $x = ua$ . Then

$$\begin{aligned} \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), x, p) &= \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), ua, p) = \bigvee_{r \in Q_{act}(t_{i+1})} \left( \tilde{\delta}_1((q, \mu_1^{t_i}(q)), u, r) \wedge \tilde{\delta}_1((r, \mu_1^{t_{i+1}}(r)), a, p) \right) \\ &= \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1((r, \mu_1^{t_i}(r)), u, q) \wedge \tilde{\delta}_1((p, \mu_1^{t_{i+1}}(p)), a, r) \right) = \bigvee_{r \in Q_{act}(t_{i+1})} \left( \tilde{\delta}_1((p, \mu_1^{t_i}(p)), a, r) \wedge \tilde{\delta}_1((r, \mu_1^{t_{i+1}}(r)), u, q) \right) \\ &= \tilde{\delta}_1^*((p, \mu_1^{t_i}(p)), au, q) = \tilde{\delta}_1^*((p, \mu_1^{t_i}(p)), ua, q) = \tilde{\delta}_1^*((p, \mu_1^{t_i}(p)), x, q), \end{aligned}$$

$$\begin{aligned} \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), x, p) &= \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), ua, p) = \bigwedge_{r \in Q_{act}(t_{i+1})} \left( \tilde{\delta}_2((q, \mu_2^{t_i}(q)), u, r) \vee \tilde{\delta}_2((r, \mu_2^{t_{i+1}}(r)), a, p) \right) \\ &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2((r, \mu_2^{t_i}(r)), u, q) \vee \tilde{\delta}_2((p, \mu_2^{t_{i+1}}(p)), a, r) \right) = \bigwedge_{r \in Q_{act}(t_{i+1})} \left( \tilde{\delta}_2((p, \mu_2^{t_i}(p)), a, r) \vee \tilde{\delta}_2((r, \mu_2^{t_{i+1}}(r)), u, q) \right) \\ &= \tilde{\delta}_2^*((p, \mu_2^{t_i}(p)), au, q) = \tilde{\delta}_2^*((p, \mu_2^{t_i}(p)), ua, q) = \tilde{\delta}_2^*((p, \mu_2^{t_i}(p)), x, q), \end{aligned}$$

$$\begin{aligned} \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), x, p) &= \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), ua, p) = \bigwedge_{r \in Q_{act}(t_{i+1})} \left( \tilde{\delta}_3((q, \mu_3^{t_i}(q)), u, r) \vee \tilde{\delta}_3((r, \mu_3^{t_{i+1}}(r)), a, p) \right) \\ &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_3((r, \mu_3^{t_i}(r)), u, q) \vee \tilde{\delta}_3((p, \mu_3^{t_{i+1}}(p)), a, r) \right) = \bigwedge_{r \in Q_{act}(t_{i+1})} \left( \tilde{\delta}_3((p, \mu_3^{t_i}(p)), a, r) \vee \tilde{\delta}_3((r, \mu_3^{t_{i+1}}(r)), u, q) \right) \\ &= \tilde{\delta}_3^*((p, \mu_3^{t_i}(p)), au, q) = \tilde{\delta}_3^*((p, \mu_3^{t_i}(p)), ua, q) = \tilde{\delta}_3^*((p, \mu_3^{t_i}(p)), x, q). \end{aligned}$$

Hence, the result is true for  $|u| = n$ . This completes the proof.  $\square$

**Proposition 4.4.** *Let  $\tilde{F}$  be a NGFA, if  $\tilde{F}^*$  is a NGFSA, then for every  $i \geq 1$ ,  $\tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), xy, p) = \tilde{\delta}_1^*((p, \mu_1^{t_{i-1}}(p)), yx, q)$ ,  $\tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), xy, p) = \tilde{\delta}_2^*((p, \mu_2^{t_{i-1}}(p)), yx, q)$ ,  $\tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), xy, p) = \tilde{\delta}_3^*((p, \mu_3^{t_{i-1}}(p)), yx, q)$  for all  $p, q \in Q$  and  $x, y \in \Sigma^*$ .*

*Proof.* Since  $p, q \in Q$  and  $x, y \in \Sigma^*$ , we prove the result by induction on  $|x| = n$ . First, we assume that  $n = 0$ , then  $x = \Lambda$ . Thus

$$\begin{aligned} \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), xy, p) &= \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), x\Lambda, p) = \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), \Lambda x, p) = \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), yx, p), \\ \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), xy, p) &= \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), x\Lambda, p) = \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), \Lambda x, p) = \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), yx, p), \\ \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), xy, p) &= \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), x\Lambda, p) = \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), \Lambda x, p) = \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), yx, p). \end{aligned}$$

Suppose that

$$\begin{aligned} \tilde{\delta}_1^*((q, \mu_1^{t_{i-1}}(q)), xu, p) &= \tilde{\delta}_1^*((p, \mu_1^{t_{i-1}}(p)), ux, q), \tilde{\delta}_2^*((q, \mu_2^{t_{i-1}}(q)), xu, p) = \tilde{\delta}_2^*((p, \mu_2^{t_{i-1}}(p)), ux, q), \\ \tilde{\delta}_3^*((q, \mu_3^{t_{i-1}}(q)), xu, p) &= \tilde{\delta}_3^*((p, \mu_3^{t_{i-1}}(p)), ux, q), \text{ for every } u \in \Sigma^*. \end{aligned}$$

Now, continue the result is true for all  $u \in \Sigma^*$  with  $|u| = n - 1$ , where  $n > 0$ . Let  $y = ua$ , where  $a \in \Sigma$

and  $u \in \Sigma^*$ . Then

$$\begin{aligned}
 \tilde{\delta}_1^*((q, \mu_1^{t_i-1}(q)), xy, p) &= \tilde{\delta}_1^*((q, \mu_1^{t_i-1}(q)), xua, p) = \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1((q, \mu_1^{t_i-1}(q)), xu, r) \wedge \tilde{\delta}_1((r, \mu_1^{t_i}(r)), a, p) \right) \\
 &= \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1((q, \mu_1^{t_i-1}(q)), ux, r) \wedge \tilde{\delta}_1((r, \mu_1^{t_i}(r)), a, p) \right) \\
 &= \bigvee_{r \in Q_{act}(t_{i-1})} \left( \tilde{\delta}_1((r, \mu_1^{t_i-1}(r)), ux, q) \wedge \tilde{\delta}_1((p, \mu_1^{t_i}(p)), a, r) \right) \\
 &= \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1((p, \mu_1^{t_i-1}(p)), a, r) \wedge \tilde{\delta}_1((r, \mu_1^{t_i}(r)), ux, q) \right) \\
 &= \tilde{\delta}_1^*((p, \mu_1^{t_i-1}(p)), aux, q) = \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1((p, \mu_1^{t_i-1}(p)), au, r) \wedge \tilde{\delta}_1((r, \mu_1^{t_i}(r)), x, q) \right) \\
 &= \bigvee_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_1((p, \mu_1^{t_i-1}(p)), ua, r) \wedge \tilde{\delta}_1((r, \mu_1^{t_i}(r)), x, q) \right) \\
 &= \tilde{\delta}_1^*((p, \mu_1^{t_i-1}(p)), uax, q) = \tilde{\delta}_1^*((q, \mu_1^{t_i-1}(q)), uax, p) = \tilde{\delta}_1^*((q, \mu_1^{t_i-1}(q)), yx, p),
 \end{aligned}$$

$$\begin{aligned}
 \tilde{\delta}_2^*((q, \mu_2^{t_i-1}(q)), xy, p) &= \tilde{\delta}_2^*((q, \mu_2^{t_i-1}(q)), xua, p) = \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2((q, \mu_2^{t_i-1}(q)), xu, r) \vee \tilde{\delta}_2((r, \mu_2^{t_i}(r)), a, p) \right) \\
 &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2((q, \mu_2^{t_i-1}(q)), ux, r) \vee \tilde{\delta}_2((r, \mu_2^{t_i}(r)), a, p) \right) \\
 &= \bigwedge_{r \in Q_{act}(t_{i-1})} \left( \tilde{\delta}_2((r, \mu_2^{t_i-1}(r)), ux, q) \vee \tilde{\delta}_2((p, \mu_2^{t_i}(p)), a, r) \right) \\
 &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2((p, \mu_2^{t_i-1}(p)), a, r) \vee \tilde{\delta}_2((r, \mu_2^{t_i}(r)), ux, q) \right) \\
 &= \tilde{\delta}_2^*((p, \mu_2^{t_i-1}(p)), aux, q) = \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2((p, \mu_2^{t_i-1}(p)), au, r) \wedge \tilde{\delta}_2((r, \mu_2^{t_i}(r)), x, q) \right) \\
 &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_2((p, \mu_2^{t_i-1}(p)), ua, r) \vee \tilde{\delta}_2((r, \mu_2^{t_i}(r)), x, q) \right) \\
 &= \tilde{\delta}_2^*((p, \mu_2^{t_i-1}(p)), uax, q) = \tilde{\delta}_2^*((q, \mu_2^{t_i-1}(q)), uax, p) = \tilde{\delta}_2^*((q, \mu_2^{t_i-1}(q)), yx, p),
 \end{aligned}$$

$$\begin{aligned}
 \tilde{\delta}_3^*((q, \mu_3^{t_i-1}(q)), xy, p) &= \tilde{\delta}_3^*((q, \mu_3^{t_i-1}(q)), xua, p) = \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_3((q, \mu_3^{t_i-1}(q)), xu, r) \vee \tilde{\delta}_3((r, \mu_3^{t_i}(r)), a, p) \right) \\
 &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_3((q, \mu_3^{t_i-1}(q)), ux, r) \vee \tilde{\delta}_3((r, \mu_3^{t_i}(r)), a, p) \right) \\
 &= \bigwedge_{r \in Q_{act}(t_{i-1})} \left( \tilde{\delta}_3((r, \mu_3^{t_i-1}(r)), ux, q) \vee \tilde{\delta}_3((p, \mu_3^{t_i}(p)), a, r) \right) \\
 &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_3((p, \mu_3^{t_i-1}(p)), a, r) \vee \tilde{\delta}_3((r, \mu_3^{t_i}(r)), ux, q) \right) \\
 &= \tilde{\delta}_3^*((p, \mu_3^{t_i-1}(p)), aux, q) = \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_3((p, \mu_3^{t_i-1}(p)), au, r) \wedge \tilde{\delta}_3((r, \mu_3^{t_i}(r)), x, q) \right) \\
 &= \bigwedge_{r \in Q_{act}(t_i)} \left( \tilde{\delta}_3((p, \mu_3^{t_i-1}(p)), ua, r) \vee \tilde{\delta}_3((r, \mu_3^{t_i}(r)), x, q) \right) \\
 &= \tilde{\delta}_3^*((p, \mu_3^{t_i-1}(p)), uax, q) = \tilde{\delta}_3^*((q, \mu_3^{t_i-1}(q)), uax, p) = \tilde{\delta}_3^*((q, \mu_3^{t_i-1}(q)), yx, p).
 \end{aligned}$$

This completes the proof. □

**Definition 4.5.** Let  $\tilde{F}^*$  be a GNFSFA. Let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  and  $\tilde{\delta}^* = \langle \tilde{\delta}_1^*, \tilde{\delta}_2^*, \tilde{\delta}_3^* \rangle$  in  $(Q \times [0, 1]) \times \Sigma^* \times Q$  be



a neutrosophic set in  $Q$ . Then  $\mu$  is a neutrosophic switchboard subsystem of  $\tilde{F}^*$ , say  $\mu \subseteq \tilde{F}^*$ , if for every  $j$ ,  $1 \leq j \leq k$  such that  $\mu_1^{t_j}(p) \geq \tilde{\delta}_1^*((q, \mu_1^{t_j}(q)), x, p)$ ,  $\mu_2^{t_j}(p) \leq \tilde{\delta}_2^*((q, \mu_2^{t_j}(q)), x, p)$ ,  $\mu_3^{t_j}(p) \leq \tilde{\delta}_3^*((q, \mu_3^{t_j}(q)), x, p)$ .  $\forall q, p \in Q$  and  $x \in \Sigma$ .

**Theorem 4.6.** Let  $\tilde{F}^*$  be a NGFSA and let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  and  $\tilde{\delta}^* = \langle \tilde{\delta}_1^*, \tilde{\delta}_2^*, \tilde{\delta}_3^* \rangle$  in  $(Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma^* \times Q$  be a neutrosophic set in  $Q$ . Then  $\mu$  is a neutrosophic switchboard subsystem of  $\tilde{F}^*$  if and only if  $\mu_1^{t_j}(p) \geq \tilde{\delta}_1^*((q, \mu_1^{t_j}(q)), x, p)$ ,  $\mu_2^{t_j}(p) \leq \tilde{\delta}_2^*((q, \mu_2^{t_j}(q)), x, p)$ ,  $\mu_3^{t_j}(p) \leq \tilde{\delta}_3^*((q, \mu_3^{t_j}(q)), x, p)$ , for all  $q \in Q_{(act)}(t_j)$ ,  $p \in Q$  and  $x \in \Sigma^*$ .

*Proof.* The proof of the theorem is similar to Theorem 3.11 and it is clear that  $\mu$  satisfies switching and commutative, since  $\tilde{F}^*$  is NGFSA. This proof is completed.  $\square$

**Definition 4.7.** Let  $\tilde{F}^*$  be a NGFSA. Let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  and  $\tilde{\delta}^* = \langle \tilde{\delta}_1^*, \tilde{\delta}_2^*, \tilde{\delta}_3^* \rangle$  in  $(Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma^* \times Q$  be a neutrosophic set in  $Q$ . Then  $\mu$  is a neutrosophic strong switchboard subsystem of  $\tilde{F}^*$ , say  $\mu \subseteq \tilde{F}^*$ , if for every  $i$ ,  $1 \leq i \leq k$  such that  $p \in S_\alpha(q)$ , then for  $q, p \in Q$  and  $x \in \Sigma$ ,  $\mu_1^{t_j}(p) \geq \mu_1^{t_j}(q)$ ,  $\mu_2^{t_j}(p) \leq \mu_2^{t_j}(q)$ ,  $\mu_3^{t_j}(p) \leq \mu_3^{t_j}(q)$ , for every  $1 \leq j \leq k$ .

**Theorem 4.8.** Let  $\tilde{F}^*$  be a NGFA and let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  and  $\tilde{\delta}^* = \langle \tilde{\delta}_1^*, \tilde{\delta}_2^*, \tilde{\delta}_3^* \rangle$  in  $(Q \times [0, 1] \times [0, 1] \times [0, 1]) \times \Sigma^* \times Q$  be a neutrosophic set in  $Q$ . Then  $\mu$  is a strong neutrosophic switchboard subsystem of  $\tilde{F}^*$  if and only if there exists  $x \in \Sigma^*$  such that  $p \in S_\alpha(q)$ , then  $\mu_1^{t_j}(p) \geq \mu_1^{t_j}(q)$ ,  $\mu_2^{t_j}(p) \leq \mu_2^{t_j}(q)$ ,  $\mu_3^{t_j}(p) \leq \mu_3^{t_j}(q)$ , for all  $q \in Q_{(act)}(t_j)$ ,  $p \in Q$ .

*Proof.* The proof of the theorem is similar to Theorem 3.13 and it is clear that  $\mu$  satisfies switching and commutative, since  $\tilde{F}^*$  is NGFSA. The proof is completed.  $\square$

**Theorem 4.9.** Let  $\tilde{F}^*$  be a NGFSA and let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  be a neutrosophic subset of  $Q$ . If  $\mu$  is a neutrosophic switchboard subsystem of  $\tilde{F}^*$ , then  $\mu$  is a strong neutrosophic switchboard subsystem of  $\tilde{F}^*$ .

*Proof.* Assume that  $\tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), x, p) > 0$ ,  $\tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), x, p) < 1$  and  $\tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), x, p) < 1$ , for all  $x \in \Sigma$ . Since  $\mu$  is a neutrosophic switchboard subsystem of  $\tilde{F}^*$ , we have

$$\mu_1^{t_j}(p) \geq \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), x, p), \quad \mu_2^{t_j}(p) \leq \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), x, p), \quad \mu_3^{t_j}(p) \leq \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), x, p).$$

for all  $q \in Q_{(act)}(t_j)$ ,  $p \in Q$  and  $x \in \Sigma$ . As  $\mu$  is switching, then we have

$$\begin{aligned} \mu_1^{t_j}(p) &\geq \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), x, p) = \tilde{\delta}_1^*((p, \mu_1^{t_i}(p)), x, q) = \mu_1^{t_j}(q), \\ \mu_2^{t_j}(p) &\leq \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), x, p) = \tilde{\delta}_2^*((p, \mu_2^{t_i}(p)), x, q) = \mu_2^{t_j}(q), \\ \mu_3^{t_j}(p) &\leq \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), x, p) = \tilde{\delta}_3^*((p, \mu_3^{t_i}(p)), x, q) = \mu_3^{t_j}(q). \end{aligned}$$

As  $\mu$  is commutative, then  $x = uv$ , we have

$$\begin{aligned}
 \mu_1^{t_j}(p) &\geq \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), x, p) = \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), uv, p) \\
 &= \bigvee \left\{ \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), u, r) \wedge \tilde{\delta}_1^*((r, \mu_1^{t_{i+1}}(r)), v, p) \mid r \in Q_{1(act)}(t_{i+1}) \right\} \\
 &= \bigvee \left\{ \tilde{\delta}_1^*((r, \mu_1^{t_i}(r)), u, q) \wedge \tilde{\delta}_1^*((p, \mu_1^{t_{i+1}}(p)), v, r) \mid r \in Q_{1(act)}(t_{i+1}) \right\} \\
 &= \bigvee \left\{ \tilde{\delta}_1^*((p, \mu_1^{t_{i+1}}(p)), v, r) \wedge \tilde{\delta}_1^*((r, \mu_1^{t_i}(r)), u, q) \mid r \in Q_{1(act)}(t_{i+1}) \right\} \\
 &= \tilde{\delta}_1^*((p, \mu_1^{t_{i+1}}(p)), vu, q) = \tilde{\delta}_1^*((p, \mu_1^{t_{i+1}}(p)), uv, q) = \tilde{\delta}_1^*((p, \mu_1^{t_{i+1}}(p)), x, q) \geq \mu_1^{t_j}(q), \\
 \mu_2^{t_j}(p) &\leq \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), x, p) = \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), uv, p) \\
 &= \bigwedge \left\{ \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), u, r) \vee \tilde{\delta}_2^*((r, \mu_2^{t_{i+1}}(r)), v, p) \mid r \in Q_{1(act)}(t_{i+1}) \right\} \\
 &= \bigwedge \left\{ \tilde{\delta}_2^*((r, \mu_2^{t_i}(r)), u, q) \vee \tilde{\delta}_2^*((p, \mu_2^{t_{i+1}}(p)), v, r) \mid r \in Q_{1(act)}(t_{i+1}) \right\} \\
 &= \bigwedge \left\{ \tilde{\delta}_2^*((p, \mu_2^{t_{i+1}}(p)), v, r) \vee \tilde{\delta}_2^*((r, \mu_2^{t_i}(r)), u, q) \mid r \in Q_{1(act)}(t_{i+1}) \right\} \\
 &= \tilde{\delta}_2^*((p, \mu_2^{t_{i+1}}(p)), vu, q) = \tilde{\delta}_2^*((p, \mu_2^{t_{i+1}}(p)), uv, q) = \tilde{\delta}_2^*((p, \mu_2^{t_{i+1}}(p)), x, q) \leq \mu_2^{t_j}(q), \\
 \mu_3^{t_j}(p) &\leq \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), x, p) = \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), uv, p) \\
 &= \bigwedge \left\{ \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), u, r) \vee \tilde{\delta}_3^*((r, \mu_3^{t_{i+1}}(r)), v, p) \mid r \in Q_{1(act)}(t_{i+1}) \right\} \\
 &= \bigwedge \left\{ \tilde{\delta}_3^*((r, \mu_3^{t_i}(r)), u, q) \vee \tilde{\delta}_3^*((p, \mu_3^{t_{i+1}}(p)), v, r) \mid r \in Q_{1(act)}(t_{i+1}) \right\} \\
 &= \bigwedge \left\{ \tilde{\delta}_3^*((p, \mu_3^{t_{i+1}}(p)), v, r) \vee \tilde{\delta}_3^*((r, \mu_3^{t_i}(r)), u, q) \mid r \in Q_{1(act)}(t_{i+1}) \right\} \\
 &= \tilde{\delta}_3^*((p, \mu_3^{t_{i+1}}(p)), vu, q) = \tilde{\delta}_3^*((p, \mu_3^{t_{i+1}}(p)), uv, q) = \tilde{\delta}_3^*((p, \mu_3^{t_{i+1}}(p)), x, q) \leq \mu_3^{t_j}(q).
 \end{aligned}$$

Hence  $\mu$  is a strong neutrosophic switchboard subsystem of  $\tilde{F}^*$ . □

**Theorem 4.10.** *Let  $\tilde{F}^*$  be a NGFSA and let  $\mu = \langle \mu_1, \mu_2, \mu_3 \rangle$  be a neutrosophic subset of  $Q$ . If  $\mu$  is a strong neutrosophic switchboard subsystem of  $\tilde{F}^*$ , then  $\mu$  is a neutrosophic switchboard subsystem of  $\tilde{F}^*$ .*

*Proof.* Let  $q, p \in Q$ . Since  $\mu$  is a strong neutrosophic switchboard subsystem of  $\tilde{F}^*$  and  $\mu$  is switching, we have for all  $x \in \Sigma$ , since  $\tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), x, p) > 0$ ,  $\tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), x, p) < 1$  and  $\tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), x, p) < 1$ ,  $\forall x \in \Sigma$ ,

$$\begin{aligned}
 \mu_1^{t_j}(p) &\geq \mu_1^{t_j}(q) \geq \tilde{\delta}_1^*((p, \mu_1^{t_i}(p)), x, q) \geq \tilde{\delta}_1^*((q, \mu_1^{t_i}(q)), x, p), \\
 \mu_2^{t_j}(p) &\leq \mu_2^{t_j}(q) \leq \tilde{\delta}_2^*((p, \mu_2^{t_i}(p)), x, q) \leq \tilde{\delta}_2^*((q, \mu_2^{t_i}(q)), x, p), \\
 \mu_3^{t_j}(p) &\leq \mu_3^{t_j}(q) \leq \tilde{\delta}_3^*((p, \mu_3^{t_i}(p)), x, q) \leq \tilde{\delta}_3^*((q, \mu_3^{t_i}(q)), x, p).
 \end{aligned}$$

It is clear that  $\mu$  is commutative. Thus  $\mu$  is a neutrosophic switchboard subsystem of  $\tilde{F}^*$ . □

## 5 Conclusions

This paper attempt to develop and present a new general definition for neutrosophic finite automata. The general definition for (strong) subsystem also examined and discussed their properties. A comprehensive analysis and an appropriate methodology to manage the essential issues of output mapping in general fuzzy

automata were studied by Doostfatemen and Kremer [11]. Their approach is consistent with the output which is either associated with the states (Moore model) or with the transitions (Mealy model). Interval-valued fuzzy subsets have many applications in several areas. The concept of interval-valued fuzzy sets have been studied in various algebraic structures, see [7, 26]. On the basis [11] and [7], the future work will focus on general interval-valued neutrosophic finite automata with output respond to input strings.

## 6 Acknowledgments

This research work is supported by the Fundamental Research Grant Schemes (Vote No: 1562), Ministry of Higher Education, Malaysia.

## References

- [1] M. Abdel-Basset, M.Saleh, A. Gamal, F. Smarandache. An approach of TOPSIS technique for developing supplier selection with group decision making under type-2 neutrosophic number. *Applied Soft Computing*, 77 (2019), 438-452. doi:10.1016/j.asoc.2019.01.035
- [2] M. Abdel-Basset, V. Chang, A. Gamal, F. Smarandache. An integrated neutrosophic ANP and VIKOR method for achieving sustainable supplier selection: A case study in importing field. *Computers in Industry*, 106 (2019), 94-110. doi:10.1016/j.compind.2018.12.017
- [3] M. Abdel-Basset, G. Manogaran, A. Gamal, F. Smarandache. A group decision making framework based on neutrosophic TOPSIS approach for smart medical device selection. *Journal of Medical Systems*, 43:38 (2019), 1-13. doi:10.1007/s10916-019-1156-1
- [4] M. Abdel-Basset, M. Mohamed. A novel and powerful framework based on neutrosophic sets to aid patients with cancer. *Future Generation Computer Systems*, 98 (2019), 144-153. doi:10.1016/j.future.2018.12.019
- [5] K. Abolpour and M. M. Zahedi, L-general fuzzy automata. 2015 4th Iranian Joint Congress on Fuzzy and Intelligent Systems (CFIS), Zahedan, 2015, p 1-5. doi: 10.1109/CFIS.2015.7391673
- [6] K. Abolpour and M.M. Zahedi. General fuzzy automata theory based on the complete residuated lattice-valued. *Iranian Journal of Fuzzy Systems*, 14(5)(2017), 103-121. doi: 10.22111 / IJFS.2017.3435
- [7] M. Akram, N. Yaqoob, J. Kavikumar. Interval-valued  $(\tilde{\theta}, \tilde{\delta})$ -fuzzy KU ideals of KU-algebras. *International Journal of Pure and Applied Mathematics*, 92(3) (2014), 335-349. doi:10.12732/ijpam.v92i3.2
- [8] M. Ali, L. H. Son, M. Khan, N. T. Tung. Segmentation of dental X-ray images in medical imaging using neutrosophic orthogonal matrices. *Expert Systems with Applications*, 91 (2018), 434-441. doi:10.1016/j.eswa.2017.09.027
- [9] Y. Cao and Y. Ezawac. Nondeterministic fuzzy automata. *Information Sciences*, 191(2012), 86-97. doi: 10.1016/j.ins.2011.12.024
- [10] I. Deli. Some operators with IVGSVTrN-numbers and their applications to multiple criteria group decision making, *Neutrosophic Sets and Systems*, 25 (2019), 33-53. doi: 10.5281/zenodo.2631496
- [11] M. Doostfateme and S. C. Kremer. New directions in fuzzy automata. *International Journal of Approximate Reasoning*, 38(2005), 175-214. doi: 10.1016/j.ijar.2004.08.001
- [12] M. Doostfateme and S. C. Kremer. General Fuzzy Automata, New Efficient Acceptors for Fuzzy Languages. 2006 IEEE International Conference on Fuzzy Systems, Vancouver, BC, 2006, p. 2097-2103. doi: 10.1109/FUZZY.2006.1681991

- [13] M. Horry and M. M. Zahedi. Some (fuzzy) topologies on general fuzzy automata. *Iranian Journal of Fuzzy Systems*, 10(2013), 73-89. doi: 10.22111/IJFS.2013.1317.
- [14] M. Horry. Irreducibility on general fuzzy automata. *Iranian Journal of Fuzzy Systems*, 13(2016), 131-144. doi:10.22111/IJFS.2016.2363.
- [15] M. Horry. Bipolar general fuzzy automata. *Journal of Linear and Topological Algebra*, 5(2)(2016), 83-91.
- [16] M. Horry. Applications of a group in general fuzzy automata. *Journal of Algebraic Structures and their Applications*, 4(2)(2017), 57-69. doi: 10.29252/asta.4.2.57
- [17] A. T. Jahromi, P. Karimaghaye, S. G. Haghghi, M. Doostfateme and M. J. Er. Intelligent selective packet discarding using general fuzzy automata. 2008 IEEE International Conference on Systems, Man and Cybernetics, Singapore, 2008, p. 3714-3720. doi: 10.1109/ICSMC.2008.4811877
- [18] J. Jin, Q. Li and Y. Li. Algebraic properties of L-fuzzy finite automata. *Information Sciences*, 234(2013), 182-202. doi: 10.1016/j.ins.2013.01.018
- [19] Y. B. Jun, Seon Jeong Kim, Florentin Smarandache. Interval neutrosophic sets with applications in BCK/BCI-algebra. *Axioms*, 7(2) (2018), 23.
- [20] Y. B. Jun. Intuitionistic fuzzy finite switchboard state machines. *Journal of Applied Mathematics and Computation*, 20(1-2)(2006), 315-325. doi: 10.1007/BF02831941
- [21] Y. B. Jun and J. Kavikumar. Bipolar fuzzy finite state machines. *Bulletin of the Malaysian Mathematical Sciences Society*, 34(1)(2011), 181-188.
- [22] J. Kavikumar, A. Khamis and R. Roslan. Bipolar-valued fuzzy finite switchboard state machines. *Lecture Notes in Engineering and Computer Science*, 2200(1)(2012), 571-576.
- [23] J. Kavikumar, A. Khamis and M. S. Rusiman. N-structures applied to finite state machines. *IAENG International Journal of Applied Mathematics*, 43(3)(2013), 233-237.
- [24] J. Kavikumar, S. P. Tiwari, Nor Shamsidah Amir Hamzah and S. Sharan. Restricted cascade and wreath products of fuzzy finite switchboard state machines. *Iranian Journal of Fuzzy Systems*, 16(1)(2019), 75-88. doi: 10.22111/IJFS.2019.4485
- [25] J. Kavikumar, S. P. Tiwari, Nur Ain Ebas and Nor Shamsidah Amir Hamzah. General fuzzy switchboard automata. *New Mathematics and Natural Computation (Article in Press)* doi: 10.1142S1793005719500157
- [26] M. Kaviyarasu, K. Indhira, V.M. Chandrasekaran, J. Kavikumar. Interval-valued fuzzy subalgebra and fuzzy INK-ideal in INK-algebra. In: Madhu V., Manimaran A., Easwaramoorthy D., Kalpanapriya D., Mubashir Unnissa M. (eds) *Advances in Algebra and Analysis*. Trends in Mathematics. Birkhäuser, Cham, 2018, 19-25. doi:10.1007/978-3-030-01120-8\_3
- [27] L. Li and D. W. Qiu. On the state minimization of fuzzy automata. *IEEE Transactions on Fuzzy System*, 23(2)(2015), 434-443. doi: 10.1109/TFUZZ.2014.2315620
- [28] K. Mohana, V.Christy, F. Smarandache. On multi-criteria decision making problem via bipolar single-valued neutrosophic settings, *Neutrosophic Sets and Systems*, 25 (2019), 125-135. doi: 10.5281/zenodo.2631512
- [29] J.N. Mordeson and D.S. Malik, *Fuzzy automata and languages: Theory and Applications*, Chapman and Hall/CRC, London/Boca Raton, 2002.
- [30] N.A. Nabeeh, N. A, F. Smarandache, M. Abdel-Basset, H.A. El-Ghareeb, A. Aboelfetouh. An integrated neutrosophic-TOPSIS approach and its application to personnel selection: A new trend in brain processing and analysis. *IEEE Access*, 7 (2019), 29734-29744.

- [31] D. Nagarajana, M. Lathamaheswari, Said Broumi, J. Kavikumar. A new perspective on traffic control management using triangular interval type-2 fuzzy sets and interval neutrosophic sets. *Operations Research Perspectives*, (2019). doi: 10.1016/j.orp.2019.100099
- [32] D. Nagarajan, M. Lathamaheswari, Said Broumi, J. Kavikumar. Dombi interval valued neutrosophic graph and its role in traffic control management. *Neutrosophic Sets and Systems*, 24 (2019), 114-133. doi: 10.5281/zenodo.2593948
- [33] R. G. Ortega, M. L. Vazquez, J. A. S. Figueiredo, A. Guijarro-Rodriguez. Sinos river basin social-environmental prospective assessment of water quality management using fuzzy cognitive maps and neutrosophic AHP-TOPSIS, *Neutrosophic Sets and Systems*, 23 (2018), 60-171. doi: 10.5281/zenodo.2158501
- [34] D. W. Qiu. Automata theory based on complete residuated lattice-valued logic. *Science in China (Series F: Information Sciences)*, 44(6)(2001), 419-429. doi: 10.1007/BF02713945
- [35] E.S. Santos. Fuzzy automata and languages. *Information Sciences*, 10(1981), 193-197. doi: 10.1016/0020-0255(76)90040-2
- [36] Y. Sato and N. Kuroki. Fuzzy finite switchboard state machines. *J. Fuzzy Mathematics*, 10(4) (2002), 863-874.
- [37] M. Shamsizadeh and M.M. Zahedi. Intuitionistic general fuzzy automata. *Soft Computing*, 20(9) (2016), 3505-3519. doi: 10.1007/s00500-015-1969-x
- [38] F. Smarandache. *A unifying Field in logics: Neutrosophy, neutrosophic probability, set and logic*, Rehoboth, American Research Press, 1999.
- [39] F. Smarandache. Neutrosophic set - a generalization of the intuitionistic fuzzy set. *Granular Computing 2006 IEEE International Conference*, 8-42, 2006. doi:10.1109/GRC.2006.1635754
- [40] U. Rivieccio. Neutrosophic logics: Prospects and problems. *Fuzzy sets and systems*, 159(14)(2018), 1860-1868. doi:10.1016/j.fss.2007.11.011
- [41] A. K. Srivastava and S. P. Tiwari. On relationships among fuzzy approximation operators, fuzzy topology, and fuzzy automata. *Fuzzy Sets and Systems*, 138(1)(2003), 197-204. doi: 10.1016/S0165-0114(02)00442-6
- [42] M. Tahir and Q. Khan. Interval neutrosophic finite switchboard state machine. *Afrika Matematika*, 27(2016), 1361-1376. doi: 10.1007/s13370-016-0416-1
- [43] M. Tahir, Q. Khan, K. Ullah and N. Jan. Single valued neutrosophic finite state machine and switchboard state machine. *New Trends in Neutrosophic Theory and Applications, II*, 2018, p. 384-402.
- [44] H. Wang, F. Smarandache, Y.Q. Zhang, R. Sunderraman. Single-valued neutrosophic sets. *Multispace and Multistructure*, 4 (2010), 410-413.
- [45] H. Wang, F. Smarandache, R. Sunderraman, Y. Q. Zhang. *Interval neutrosophic sets and logic: Theory and applications in computing*, Hexis, Phoenix, AZ, 2005.
- [46] W. G. Wee. *On generalizations of adaptive algorithm and application of the fuzzy sets concept to pattern classification*; Ph.D. Thesis, Purdue University, 1967.
- [47] L. A. Zadeh. Fuzzy sets. *Information and Control*, 8(1965), 338-353. doi: 10.1016/S0019-9958(65)90241-X

Received: March 12, 2019.

Accepted: April 30, 2019.