



UWS Academic Portal

Current evidence and practical applications of flywheel eccentric overload exercises as post-activation potentiation protocols

Beato, Marco; McErlain-Naylor, Stuart A. ; Halperin, Israel; Dello Iacono, Antonio

Published in:
International Journal of Sports Physiology and Performance

DOI:
[10.1123/ijsp.2019-0476](https://doi.org/10.1123/ijsp.2019-0476)

E-pub ahead of print: 18/11/2019

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Beato, M., McErlain-Naylor, S. A., Halperin, I., & Dello Iacono, A. (2019). Current evidence and practical applications of flywheel eccentric overload exercises as post-activation potentiation protocols: a brief review. *International Journal of Sports Physiology and Performance*. <https://doi.org/10.1123/ijsp.2019-0476>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted author manuscript version reprinted, by permission, from International Journal of Sports Physiology and Performance, 2019 (ahead of print). © Human Kinetics, Inc

1 **Current evidence and practical applications of flywheel eccentric overload exercises as**
2 **post-activation potentiation protocols: A brief review**

3
4
5 **Abstract**

6 **Purpose:** This review summarizes the evidence on post-activation potentiation (PAP)
7 protocols using flywheel eccentric overload (EOL) exercises.

8 **Methods:** Studies were searched using the electronic databases PubMed, Scopus and ISI Web
9 of Knowledge.

10 **Results:** Seven eligible studies were identified, identifying the following results: First,
11 practitioners can use different inertia intensities (*e.g.* 0.03 to 0.88 kg·m²), based on the exercise
12 selected, to enhance sport specific performance. Second, the PAP time window following EOL
13 exercise seems to be consistent with traditional PAP literature, where acute fatigue is dominant
14 in the early part of the recovery period (*e.g.* 30 seconds) and PAP is dominant in the second
15 part (*e.g.* 3 and 6 minutes). Third, since EOL exercises require large force and power outputs,
16 a volume of 3 sets with the conditioning activity (*e.g.* half squat or lunge) seems to be a sensible
17 approach. This could reduce the transitory muscular fatigue and thereby allow for a stronger
18 potentiation effect compared to larger exercise volumes. Fourth, athletes should gain
19 experience performing EOL exercises prior to utilizing the tool as part of a PAP protocol (3-4
20 sessions of familiarization). Finally, the dimensions of common flywheel devices offer useful
21 and practical solutions to induce PAP effects outside of normal training environments and prior
22 to competitions.

23 **Conclusions:** EOL exercise can be utilized to stimulate PAP responses in order to obtain
24 performance advantages in various sports. However, future research is needed to determine
25 what EOL exercises, intensity, volume, and rest intervals optimally induce the PAP
26 phenomenon and facilitate transfer effects on athletic performances.

27
28
29
30
31
32
33
34

35

36 **Introduction to the topic**

37 This review summarizes the current evidence regarding post-activation potentiation (PAP)
38 strategies using flywheel eccentric overload (EOL) exercises. The first section covers the PAP
39 phenomenon, its underpinning neurophysiological mechanisms, and commonly used PAP
40 protocols. The second section describes the characteristics of flywheel ergometers and the
41 rationale for using EOL to induce PAP effects. The third section summarizes the growing
42 literature, which has evaluated the onset, time course, and magnitude of PAP effects on athletic
43 performance using EOL exercises. Lastly, this review reports some practical recommendations
44 on how PAP effects can be elicited using EOL exercises in applied settings and proposes future
45 research directions.

46

47 **1.1 Post-activation potentiation (PAP)**

48 PAP is defined as “the phenomena by which muscular performance characteristics are acutely
49 enhanced as a result of their contractile history”.¹⁻³ This term is generally used when the
50 enhanced muscular response following a potentiation activity can be verified with a twitch
51 interpolation technique.^{2,4,5} However, among sport scientists and coaches, PAP is commonly
52 interpreted as an enhancement of athletic performance measured in voluntary exercise
53 requiring rapid or maximal force production.^{3,6} Two underpinning pathways are thought to
54 account for the PAP effects: peripheral and central. Myosin regulatory light chain (RLC)
55 phosphorylation is suggested to be the main peripheral mechanism associated with PAP. The
56 augmented phosphorylation of RLC is mediated via the enzyme myosin light chain kinase,
57 which leads to a greater rate of cross-bridge attachment.^{1,7,8} This is due to an increased
58 sensitivity of the contractile proteins to calcium (Ca^{2+}), which is released from the sarcoplasmic
59 reticulum.^{3,9,10} This mechanism facilitates the force and rate of force development of low and
60 high frequency contractions.^{11,12}

61 PAP may also result from spinal and supraspinal pathways. It is speculated that through
62 increases in synaptic efficiency induced by residual elevation of presynaptic Ca^{2+} , and
63 decreases in transmitter failure occurring at higher order motoneurons responsible for fast-
64 twitch motor units.^{13,14} These central effects may contribute to a sustained recruitment of higher
65 threshold motor units and increases in fast-twitch fiber contribution to muscular contraction.¹⁵
66 However, a recent review does not support this central explanation underpinning PAP.² Hence,
67 it could be concluded that RLC phosphorylation is considered the primary mechanism for PAP,
68 while other influences at the central level remain to be clarified.

69

70 **1.2 Methodological approaches for PAP protocol design**

71 There are a number of variables that need to be accounted for when designing PAP protocols:
72 type of muscular contraction, time interval between the PAP conditioning activity and
73 subsequent performance test, biomechanical similarities, and intensity of load. PAP methods
74 are commonly classified as either static or dynamic, according to the muscular contraction
75 mode of the conditioning activity.¹ Examples of static potentiating protocols include isometric
76 continuous or intermittent maximal voluntary contractions (MVC), while dynamic protocols
77 include loaded jumping, sprinting, throwing movements, and resistance exercises.³ Although
78 both methods can potentiate subsequent athletic performances, they induce dissimilar fatigue
79 and potentiation responses. The different nature of the underpinning PAP mechanisms induced
80 by static and dynamic methods has specific implications for the methodological design of PAP
81 protocols. Static PAP protocols implement volumes (1-5 sets x 3-10 s) of isometric contractions
82 executed at high intensity (> 90% MVC).¹⁶⁻¹⁹ PAP protocols using dynamic contractions
83 require greater volumes and are commonly designed as multiple-set configurations (2-3 sets x
84 3-8 repetitions) and executed at submaximal intensities (60-90% 1RM).^{3,9,20-22}

85 Another key variable affected by the specific potentiation method is the necessary time interval
86 between the PAP conditioning activity and the subsequent performance test. Whereas the
87 majority of the PAP studies suggest a recovery interval of 3 to 11 min to elicit the greatest PAP
88 effect,³ the exact PAP onset time and duration vary and depend on the type of the conditioning
89 activity. Isometric contractions evoke PAP earlier (≤ 3 min) when compared with dynamic
90 conditions,^{16,23} which require longer rest intervals (≥ 3 min).⁶ However, PAP effects induced
91 by dynamic protocols persist for longer durations compared to static protocols, and can be
92 maintained up to 12 min after protocol completion.^{1,24} Thus, it is likely that each potentiation
93 complex achieves the PAP via different pathways, affecting the onset, magnitude and duration
94 of the potentiation effects.^{7,13,25} Finally, the contemporary literature recommends practitioners
95 to select conditioning exercises with biomechanical similarity to the subsequent athletic
96 performance intended to improve (*e.g.* squat exercises for jump tasks or hip thrusts for sprint
97 tasks).^{15,26-28} Indeed, high kinematic and kinetic specificity seem to play a favorable role in
98 optimizing the potentiation effects.^{6,27}

99

100 **2.1 Flywheel devices and eccentric overload (EOL) training**

101 Flywheel ergometers have been present in the scientific literature since the early twentieth
102 century.²⁹ They were developed as resistance training devices for space travellers exposed to

103 non-gravity environments and became popular in the early 1990s as a tool for high intensity
104 resistance training without the requirement for gravitational resistance.^{30,31} During the
105 concentric phase, the rotational acceleration of the flywheel develops inertial torque, initially
106 accumulated and then returned back during the eccentric phase, allowing for repetitive
107 concentric-eccentric cycles.³² Skeletal muscle is able to develop greater forces during eccentric
108 than concentric activities,³³ and such flywheel exercises can determine a more demanding
109 eccentric phase due to the augmented mechanical load necessary to absorb the kinetic energy
110 stored in the flywheel and to decelerate it. This is not achievable by performing traditional
111 isotonic weightlifting exercises.³⁴⁻³⁶ As a consequence, flywheel resistance devices allow for
112 maximal force development throughout the full range of motion, with short periods of greater
113 eccentric than concentric force demands. This observation has led to subsequent increased
114 utilisation of these devices to obtain acute responses and chronic adaptations (*e.g.* for strength,
115 hypertrophy, power, injury prevention, and rehabilitation) in both amateur and professional
116 sporting settings.^{9,33,37-40} Moreover, due to the portability of these devices, practitioners can
117 use them outdoors or bring them out from weight rooms, further increasing their practical
118 sporting applications.

119

120 **2.2 Evidence and hypothesis supporting EOL training as a PAP strategy**

121 EOL training has been consistently used to induce chronic adaptations, however a few studies
122 have investigated the acute potentiation benefits offered by this exercise modality.^{34,41} The
123 rationale for utilizing flywheel EOL protocols to facilitate PAP responses is based on the to
124 central and peripheral mechanisms underpinning PAP.¹³ EOL actions, as well as eccentric
125 contractions in general, are believed to selectively recruit higher order motor units to a greater
126 extent than concentric contractions.⁴²⁻⁴⁶ This results from higher motor unit discharge rate and
127 synchrony.^{1,47} This relatively greater contribution of motor unit activation may be augmented
128 even more during compound multi-joint movements, commonly executed during EOL
129 exercises (*e.g.* squat).⁴⁸⁻⁵⁰ A further advantage of EOL exercises as potentiating activities are
130 the consistent greater eccentric force, power and derivative outputs produced.^{51,52} These greater
131 eccentric kinetic outputs can contribute to improving stretch-shortening cycle performance,
132 which may induce stronger transfer effects on the fast mixed eccentric/concentric actions of
133 athletic tasks such as jumps, sprinting and changing direction.^{51,53} These tasks may benefit from
134 the prior execution of EOL exercises which functionally overload the musculo-tendinous
135 system in a specific manner (*e.g.* eccentric contraction) and with a high degree of similarity in
136 terms of muscle actions and joint kinematics used.^{15,26-28}

137

138 **3.1 Current knowledge related to EOL exercise and PAP**

139 Knowledge on the PAP effects of EOL exercises is relatively new to the scientific community.
140 The first investigation on this topic was published in 2014 and seven studies have examined
141 the PAP effects of EOL exercises on athletic tasks performance to date (Table 1).⁹ These
142 studies were identified through searches using Pubmed, Scopus and ISI Web of Knowledge
143 databases using the following terms “eccentric overload”, “eccentric overload exercise”,
144 “flywheel”, “iso-inertial”, “flywheel resistance”, and “post-activation potentiation”.
145 Additionally, the references of all the identified articles were searched for other relevant
146 articles.

147 In the selected studies, changes in performance following PAP protocols were calculated as
148 percentage differences (%) using the following formula: $\frac{(\text{post-PAP}i - \text{baseline})}{\text{baseline}} \times 100$, with i
149 representing any post-PAP assessment time point. Hedges' g effect sizes (ES) were calculated
150 from the original to examine the extent of the PAP effects. Specifically, ES were determined
151 for each PAP protocol as for within-group analyses and calculated relatively to baseline or
152 control conditions absent of any PAP intervention.

153

154 The equation $d = \frac{M_{\text{diff}}}{S_{\text{av}}}$ (M_{diff} : mean difference; S_{av} : average standard deviation) with the
155 adjustment factor: $g = \left(1 - \frac{3}{4df-1}\right) \times d$ were used for this purpose.

156

157 This approach enabled estimation of unbiased effects as well as standardized comparisons
158 between protocols. ES were then interpreted as *trivial* (< 0.2), *small* (0.2 - 0.5), *medium* (0.5 -
159 0.8), or *large* (> 0.8).^{54,55}

160

161 Despite the low number of studies, the summary of their results provides preliminary evidence
162 about methodological guidelines for practical applications. PAP protocols designed with
163 flywheel EOL exercises using either single or multiple sets, performed at varying intensities
164 (0.03 kg·m² to 0.88 kg·m²), with brief rest period durations (3-9 min) seem effective to induce
165 PAP effects (Table 1).^{6,9,34,56-59} Moreover, the potentiation was found to be of a greater extent
166 on athletic tasks having higher biomechanically similarity with the potentiating EOL exercise.

167

168

Table 1 near here

169

170 With regard to the volume of EOL exercise implemented as PAP protocols, both single and
171 multiple sets can induce potentiation resulting in augmented kinetic outputs (*e.g.* force,
172 impulse, power) and enhanced athletic performances (*e.g.* vertical and horizontal jumps,
173 sprints, changes of direction, and swimming kick start).^{34,56,57} Although no study has
174 specifically compared the PAP effects of different EOL exercise volumes, this review suggests,
175 based on previous PAP literature, possible advantages in protocols using multiple sets
176 compared with a single set.³ This assumption is supported by the relative greater range of ES
177 on athletic performances reported in studies implementing multiple sets (*small to large*)
178 compared with those using single set protocols (*small*) (Table 1). Based on the contemporary
179 scientific literature, multiple set protocols seem relatively preferable, though this interpretation
180 must be taken with caution. It is known that even the same PAP conditioning activity and
181 stimulus may induce varying responses between individuals and on different athletic tasks.^{3,34}
182 In contrast to traditional PAP methods, where onset, magnitude and duration of the potentiation
183 are modulated by the different intensities of the conditioning activity, it seems that consistent
184 PAP effects can be induced by EOL exercises using a broader range of intensities.^{3,20,60,61} On
185 one hand, the present review confirms the relationship between fatigue and PAP and the
186 evidence that both are present at PAP protocol completion. In fact, EOL exercises using
187 different inertial loads (*e.g.* 0.03 kg·m² or 0.06 kg·m²) initially induce a transient state of fatigue
188 where athletic performance is impaired. However, it is interesting to note that following EOL
189 exercise PAP outweighs fatigue after relatively short rest intervals (<6 min) regardless of the
190 exercise intensity. In a recent study, Beato et al. compared the PAP effects of “moderate” (0.03
191 kg·m²) and “high” (0.06 kg·m²) inertial flywheel half squat intensities on countermovement
192 jump, long jump, and change of direction performance.³⁴ The authors did not find any
193 difference between the protocols on the onset and magnitude of the resulting PAP effects, thus
194 concluding that both exercise intensities may be used equivalently.

195 The present review reconfirms exercise specificity and similarity between the potentiation
196 protocol and the subsequent athletic tasks for exploiting optimal PAP effects following EOL
197 exercises. This assumption is supported by two main observations. First, greater potentiation
198 ES were consistently found on athletic tasks with kinematic characteristics and ground reaction
199 force orientation profiles similar to those of the EOL exercise. Most of the EOL exercises used
200 in the reviewed studies were performed as half squat movements, which are characterized by
201 a predominant vertical orientation of the associated kinetic (*e.g.* ground reaction force)
202 responses. Therefore, it is not surprising that EOL half squats potentiated vertical-oriented
203 tasks like squat jumps and countermovement jump to a greater extent (*small to medium*) than

204 horizontal-oriented ones like sprinting (*trivial*) and change of direction (*small*).^{6,34} Second,
205 similarly greater effects were found on athletic tasks executed as coupled eccentric-concentric
206 movements compared with concentric-only movements or isokinetic actions.⁵⁹ Specifically,
207 *small to large* effects were reported on countermovement jump performance following EOL
208 half squats,^{6,9} whereas the same potentiation stimulus and rest intervals only induced *trivial to*
209 *small* effects on either swimming kick start performance⁵⁸ or isokinetic concentric knee
210 extension and concentric and eccentric flexion peak torque outputs.⁶ These findings support
211 the rationale of prescribing potentiating exercises in which muscle actions and joint kinematic
212 and kinetic profiles are similar to those in the subsequent activity to optimize the PAP effects.
213 Nevertheless, this interpretation must be taken with caution and needs to be further verified
214 since limited literature currently exists on the topic. Future research comparing PAP effects of
215 horizontal and vertical based EOL exercises are needed.

216

217 **4.1 Practical applications**

218 Implementing EOL exercises is a novel PAP inducing strategy that can be used by applied
219 practitioners. Until further research is conducted to provide precise evidence-based guidelines,
220 the following preliminary practical recommendations can be suggested. First, EOL using
221 different loads can stimulate similar magnitudes of PAP response, therefore practitioners may
222 use a broader range of inertial intensities (*e.g.* 0.03 to 0.88 kg·m²) to enhance the subsequent
223 athletic performances (*e.g.* countermovement jump, long jump, change of direction). However,
224 greater intensity may be accompanied with greater levels of acute fatigue, which should be
225 taken into account when planning the rest period between the conditioning stimulus and
226 subsequent activity. Second, the rest period needed following EOL exercises seems to be
227 consistent with the gravitational loading-based PAP literature: muscular fatigue is dominant
228 immediately following the PAP stimulus (up to 3 minutes), whereas PAP is dominant in the
229 minutes thereafter (after 3 minutes). Third, since EOL exercises require large force and power
230 outputs, low volumes (*e.g.* 2-3 sets) of the conditioning activity seems to be a sensible
231 approach. In fact, higher volumes could induce greater acute fatigue and potentially delay or
232 even restrict the onset of the PAP effects on performance. Due to the heavy eccentric muscular
233 strain and the specificity of the EOL exercises, it is suggested that athletes gain experience
234 performing 3-4 EOL conditioning sessions prior to utilizing this training method as part of a
235 PAP protocol. Furthermore, the dimensions of common flywheel devices offer useful and
236 practical solutions to induce PAP effects outside normal training environments and in
237 competitions. While mobilizing barbells and weight plates can be challenging, such challenges

238 are minimized with flywheel devices, making them a logistically excellent PAP inducing tool
239 for such situations.

240

241 **4.2 Limitations and future directions**

242 A few limitations emerged from the existing literature which should be acknowledged and
243 discussed in view of future research directions. In particular, none of the studies reported in
244 this review have enrolled professional senior team-sport or female athletes which causes
245 uncertainty about the beneficial application of EOL-based PAP protocols to enhance athletic
246 performances in these populations. The potentiation responses induced by traditional PAP
247 protocols are clearly mediated by the participants' training background, strength and power
248 capabilities. Conversely, there is no evidence about the concurrent role of individual subjects'
249 physical characteristics or any of the EOL-related performances (*e.g.* maximal and average
250 force and power outputs) on the potentiating effects on subsequent athletic performance. These
251 aspects should be addressed and investigated through dedicated research designs. Additionally,
252 EOL requires large force and power output during execution, thus a relatively lower volumes
253 (*e.g.* 3 sets) of the PAP conditioning activity seem to be a viable approach. This could also
254 reduce the transitory muscular fatigue, and thereby allowing potentiation effects to be realized
255 earlier (*e.g.* < 3 min *vs.* > 6 min) and to a greater extent (*e.g.* *moderate vs. small* effects)
256 compared with higher conditioning volumes (> 3 sets) but future research is needed to clarify
257 this statement. The relatively greater mechanical demands and the specificity of the EOL
258 exercises also highlight the importance of longer familiarization periods compared to
259 traditional resistance exercises before their implementation as PAP protocols.⁴² Indeed, it may
260 be the case that the PAP effects will increase with experience gained in performing EOL
261 exercises. EOL exercise is commonly performed through a variety of brands and flywheel
262 models having different designs, inertial mechanisms, manufacturing materials and friction
263 coefficients. This is the main reason behind the lack of gold standard valid and reliable
264 procedures that objectively determine the magnitude of inertial loads and associated intensities.

265

266 Future studies are warranted to determine what EOL exercise modalities among intensity
267 (inertias), volume (sets and repetitions), rest interval, and exercise type optimally induce the
268 PAP phenomenon and enhance athletic performances. For example, using metrics such as mean
269 velocity, could provide objective feedback on both concentric and eccentric outputs during the
270 flywheel exercise for more precise intensity prescription and monitoring. This could also
271 enable relative intensities to be quantified between athletes or within-athlete at a given inertial

272 load. Another research direction worth perusing is the usefulness of self-regulating the output
273 produced with flywheel devices to better manage accumulating fatigue, and thus, to optimize
274 the PAP response. Furthermore, in all studies the same PAP inducing exercise (half squats and
275 lunges) was utilized. It would thus be of value to study other exercises (*e.g.* horizontal
276 dominant) as well in future studies. Finally, only two studies compared EOL to traditional
277 gravitational resistance protocols as the PAP inducing modality. Given the extensive
278 knowledge of gravitational resistance exercise on PAP, a comparison of EOL to such exercise
279 would shed further light on the overall usefulness of EOL as a tool to induce PAP.

280

281 **Conclusions**

282 EOL exercises performed through inertial flywheel devices can be used as an alternative PAP
283 method to acutely potentiate athletic performance. This review describes the theoretical
284 rationale of using EOL exercises to induce potentiation effects and the underpinning
285 mechanisms favoring enhanced performance. The contemporary literature provides
286 preliminary methodological guidelines for coaches and practitioners intending to design PAP
287 protocols by using EOL exercises. Future research is required to clarify the acute effects
288 induced by EOL exercises so to optimize their use as a PAP methodology in sport.

289

290

291 **References**

292

- 293 1. Tillin NA, Bishop D. Factors modulating post-activation potentiation and its effect on
294 performance of subsequent explosive activities. *Sports Med.* 2009;39(2):147-166.
295 doi:10.2165/00007256-200939020-00004
- 296 2. Boulosa D, Del Rosso S, Behm DG, Foster C. Post-activation potentiation (PAP) in
297 endurance sports: A review. *Eur J Sport Sci.* 2018;18(5):595-610.
298 doi:10.1080/17461391.2018.1438519
- 299 3. Seitz LB, Haff GG. Factors modulating post-activation potentiation of jump, sprint,
300 throw, and upperbody ballistic performances: a systematic review with meta-analysis.
301 *Sports Med.* 2016;46(2):231-240. doi:10.1007/s40279-015-0415-7
- 302 4. MacIntosh BR, Grange RW, Cory CR, Houston ME. Myosin light chain
303 phosphorylation during staircase in fatigued skeletal muscle. *Pflugers Arch.*
304 1993;425(1-2):9-15. <http://www.ncbi.nlm.nih.gov/pubmed/8272388>.
- 305 5. MacIntosh BR. Cellular and whole muscle studies of activity dependent potentiation.

- 306 *Adv Exp Med Biol.* 2010;682:315-342. doi:10.1007/978-1-4419-6366-6_18
- 307 6. Beato M, Stiff A, Coratella G. Effects of postactivation potentiation after an eccentric
308 overload bout on countermovement jump and lower-limb muscle strength. *J Strength*
309 *Cond Res.* 2019;in print:1. doi:10.1519/JSC.0000000000003005
- 310 7. Baudry S, Duchateau J. Postactivation potentiation in a human muscle: effect on the
311 rate of torque development of tetanic and voluntary isometric contractions. *J Appl*
312 *Physiol.* 2007;102(4):1394-1401. doi:10.1152/jappphysiol.01254.2006
- 313 8. Sweeney HL, Bowman BF, Stull JT. Myosin light chain phosphorylation in vertebrate
314 striated muscle: regulation and function. *Am J Physiol.* 1993;264(5 Pt 1):C1085-95.
315 doi:10.1152/ajpcell.1993.264.5.C1085
- 316 9. de Hoyo M, de la Torre A, Pradas F, et al. Effects of eccentric overload bout on
317 change of direction and performance in soccer players. *Int J Sports Med.*
318 2014;36(04):308-314. doi:10.1055/s-0034-1395521
- 319 10. Abbate F, Sargeant AJ, Verdijk PWL, de Haan A. Effects of high-frequency initial
320 pulses and posttetanic potentiation on power output of skeletal muscle. *J Appl Physiol.*
321 2000;88(1):35-40. doi:10.1152/jappl.2000.88.1.35
- 322 11. Vandenoorn R, Grange RW, Houston ME. Threshold for force potentiation associated
323 with skeletal myosin phosphorylation. *Am J Physiol Physiol.* 1993;265(6):C1456-
324 C1462. doi:10.1152/ajpcell.1993.265.6.C1456
- 325 12. Bauer P, Sansone P, Mitter B, Makivic B, Seitz LB, Tschan H. Acute effects of back
326 squats on countermovement jump performance across multiple sets of a contrast
327 training protocol in resistance-trained men. *J Strength Cond Res.* 2019;33(4):995-
328 1000. doi:10.1519/JSC.0000000000002422
- 329 13. Wallace BJ, Shapiro R, Wallace KL, Abel MG, Symons TB. Muscular and neural
330 contributions to postactivation potentiation. *J strength Cond Res.* 2019;33(3):615-625.
331 doi:10.1519/JSC.0000000000003011
- 332 14. Zucker RS, Regehr WG. Short-term synaptic plasticity. *Annu Rev Physiol.*
333 2002;64(1):355-405. [http://www.scholarpedia.org/article/Short-](http://www.scholarpedia.org/article/Short-term_synaptic_plasticity)
334 [term_synaptic_plasticity.](http://www.scholarpedia.org/article/Short-term_synaptic_plasticity)
- 335 15. Suchomel TJ, Nimphius S, Stone MH. The importance of muscular strength in athletic
336 performance. *Sports Med.* 2016;46(10):1419-1449. doi:10.1007/s40279-016-0486-0
- 337 16. Skurvydas A, Jurgelaitiene G, Kamandulis S, et al. What are the best isometric
338 exercises of muscle potentiation? *Eur J Appl Physiol.* 2019;119(4):1029-1039.
339 doi:10.1007/s00421-019-04092-y

- 340 17. Guellich A, Güllich A, Schmidtbleicher D. Potentiation of Explosive Force. *Iaaf*.
341 1996;4(October):67-80.
- 342 18. Rixon KP, Lamont HS, Bembem MG. Influence of type of muscle contraction, gender,
343 and lifting experience on postactivation potentiation performance. *J strength Cond*
344 *Res*. 2007;21(2):500-505. doi:10.1519/R-18855.1
- 345 19. Behm DG, Button DC, Barbour G, Butt JC, Young WB. Conflicting effects of fatigue
346 and potentiation on voluntary force. *J Strength Cond Res*. 2004;18(2):365-372.
347 doi:10.1519/R-12982.1
- 348 20. Wilson JM, Duncan NM, Marin PJ, et al. Meta-analysis of postactivation potentiation
349 and power: effects of conditioning activity, volume, gender, rest periods, and training
350 status. *J strength Cond Res*. 2013;27(3):854-859. doi:10.1519/JSC.0b013e31825c2bdb
- 351 21. Maloney SJ, Turner AN, Fletcher IM. Ballistic exercise as a pre-activation stimulus: a
352 review of the literature and practical applications. *Sport Med*. 2014;44(10):1347-1359.
353 doi:10.1007/s40279-014-0214-6
- 354 22. Dello Iacono A, Beato M, Halperin I. The effects of cluster-set and traditional-set post-
355 activation potentiation protocols on vertical jump performance. *Int J Sports Physiol*
356 *Perform*. 2019:Epub ahead of print.
- 357 23. Hamada T, Sale DG, MacDougall JD, Tarnopolsky MA. Postactivation potentiation,
358 fiber type, and twitch contraction time in human knee extensor muscles. *J Appl*
359 *Physiol*. 2000;88(6):2131-2137. doi:10.1152/jappl.2000.88.6.2131
- 360 24. Dobbs WC, Toluoso D V, Fedewa M V, Esco MR. Effect of postactivation
361 potentiation on explosive vertical jump: a Systematic Review and Meta-Analysis. *J*
362 *strength Cond Res*. August 2018. doi:10.1519/JSC.0000000000002750
- 363 25. Docherty D, Hodgson MJ. The application of postactivation potentiation to elite sport.
364 *Int J Sports Physiol Perform*. 2007;2(4):439-444.
365 <http://www.ncbi.nlm.nih.gov/pubmed/19171961>.
- 366 26. Dello Iacono A, Martone D, Padulo J. Acute effects of drop-jump protocols on
367 explosive performances of elite handball players. *J strength Cond Res*.
368 2016;30(11):3122-3133. doi:10.1519/JSC.0000000000001393
- 369 27. Dello Iacono A, Seitz LB. Hip thrust-based PAP effects on sprint performance of
370 soccer players: heavy-loaded versus optimum-power development protocols. *J Sports*
371 *Sci*. 2018;36(20):2375-2382. doi:10.1080/02640414.2018.1458400
- 372 28. Dello Iacono A, Padulo J, Seitz LD. Loaded hip thrust-based PAP protocol effects on
373 acceleration and sprint performance of handball players. *J Sports Sci*.

- 374 2018;36(11):1269-1276. doi:10.1080/02640414.2017.1374657
- 375 29. Hansen TE, Lindhard J. On the maximum work of human muscles especially the
376 flexors of the elbow. *J Physiol.* 1923;57(5):287-300.
377 doi:10.1113/jphysiol.1923.sp002066
- 378 30. Berg HE, Tesch A. A gravity-independent ergometer to be used for resistance training
379 in space. *Aviat Space Environ Med.* 1994;65(8):752-756.
380 <http://www.ncbi.nlm.nih.gov/pubmed/7980338>.
- 381 31. Dudley GA, Tesch PA, Miller BJ, Buchanan P. Importance of eccentric actions in
382 performance adaptations to resistance training. *Aviat Space Environ Med.*
383 1991;62(6):543-550. <http://www.ncbi.nlm.nih.gov/pubmed/1859341>.
- 384 32. Vicens-Bordas J, Esteve E, Fort-Vanmeerhaeghe A, Bandholm T, Thorborg K.
385 Skeletal muscle functional and structural adaptations after eccentric overload flywheel
386 resistance training: a systematic review and meta-analysis. *J Sci Med Sport.*
387 2018;21(1):2-3. doi:10.1016/j.jsams.2017.09.001
- 388 33. Colliander EB, Tesch PA. Effects of eccentric and concentric muscle actions in
389 resistance training. *Acta Physiol Scand.* 1990;140(1):31-39. doi:10.1111/j.1748-
390 1716.1990.tb08973.x
- 391 34. Beato M, De Keijzer KL, Leskauskas Z, Allen WJ, Dello Iacono A, McErlain-Naylor
392 SA. Effect of postactivation potentiation after medium vs. high inertia eccentric
393 overload exercise on standing long jump, countermovement jump, and change of
394 direction performance. *J strength Cond Res.* June 2019:1.
395 doi:10.1519/JSC.0000000000003214
- 396 35. Núñez FJ, Suarez-Arrones LJ, Cater P, Mendez-Villanueva A. The high-pull exercise:
397 a comparison between a versapulley flywheel device and the free weight. *Int J Sports*
398 *Physiol Perform.* 2017;12(4):527-532. doi:10.1123/ijsp.2016-0059
- 399 36. Piqueras-Sanchiz F, Martín-Rodríguez S, Martínez-Aranda LM, et al. Effects of
400 moderate vs. high iso-inertial loads on power, velocity, work and hamstring contractile
401 function after flywheel resistance exercise. Sacchetti M, ed. *PLoS One.*
402 2019;14(2):e0211700. doi:10.1371/journal.pone.0211700
- 403 37. Gonzalo-Skok O, Tous-Fajardo J, Valero-Campo C, et al. Eccentric-overload training
404 in team-sport functional performance: constant bilateral vertical versus variable
405 unilateral multidirectional movements. *Int J Sports Physiol Perform.* 2017;12(7):951-
406 958. doi:10.1123/ijsp.2016-0251
- 407 38. Tous-Fajardo J, Maldonado RA, Quintana JM, Pozzo M, Tesch PA. The flywheel leg-

- 408 curl machine: offering eccentric overload for hamstring development. *Int J Sports*
409 *Physiol Perform.* 2006;1(3):293-298. doi:10.1123/ijsp.2013-0547
- 410 39. Tous-Fajardo J, Gonzalo-Skok O, Arjol-Serrano JL, Tesch P. Enhancing change-of-
411 direction speed in soccer players by functional inertial eccentric overload and vibration
412 training. *Int J Sports Physiol Perform.* 2016;11(1):66-73. doi:10.1123/ijsp.2015-0010
- 413 40. Norrbrand L, Fluckey JD, Pozzo M, Tesch PA. Resistance training using eccentric
414 overload induces early adaptations in skeletal muscle size. *Eur J Appl Physiol.*
415 2008;102(3):271-281. doi:10.1007/s00421-007-0583-8
- 416 41. Vicens-Bordas J, Esteve E, Fort-Vanmeerhaeghe A, Bandholm T, Thorborg K. Is
417 inertial flywheel resistance training superior to gravity-dependent resistance training in
418 improving muscle strength? A systematic review with meta-analyses. *J Sci Med Sport.*
419 2017;21(1):75-83. doi:10.1016/j.jsams.2017.10.006
- 420 42. Sabido R, Hernández-Davó JL, Pereyra-Gerber GT. Influence of different inertial
421 loads on basic training variables during the flywheel squat exercise. *Int J Sports*
422 *Physiol Perform.* 2018;13(4):482-489. doi:10.1123/ijsp.2017-0282
- 423 43. Vázquez-Guerrero J, Moras G, Baeza J, Rodríguez-Jiménez S. Force outputs during
424 squats performed using a rotational inertia device under stable versus unstable
425 conditions with different loads. *PLoS One.* 2016;11(4).
426 doi:10.1371/journal.pone.0154346
- 427 44. Douglas J, Pearson S, Ross A, McGuigan M. Effects of accentuated eccentric loading
428 on muscle properties, strength, power, and speed in resistance-trained rugby players. *J*
429 *strength Cond Res.* 2018;32(10):2750-2761. doi:10.1519/JSC.0000000000002772
- 430 45. Nardone A, Romanò C, Schieppati M. Selective recruitment of high-threshold human
431 motor units during voluntary isotonic lengthening of active muscles. *J Physiol.*
432 1989;409:451-471. <http://www.ncbi.nlm.nih.gov/pubmed/2585297>.
- 433 46. Coratella G, Bellin G, Beato M, Schena F. Fatigue affects peak joint torque angle in
434 hamstrings but not in quadriceps. *J Sports Sci.* 2015;33(12):1276-1282.
435 doi:10.1080/02640414.2014.986185
- 436 47. Higbie EJ, Cureton KJ, Warren GL, Prior BM. Effects of concentric and eccentric
437 training on muscle strength, cross-sectional area, and neural activation. *J Appl Physiol.*
438 1996;81(5):2173-2181. doi:10.1152/jappl.1996.81.5.2173
- 439 48. Fahs CA, Rossow LM, Zourdos MC. Analysis of factors related to back squat
440 concentric velocity. *J strength Cond Res.* 2018;32(9):2435-2441.
441 doi:10.1519/JSC.0000000000002295

- 442 49. Fang Y, Siemionow V, Sahgal V, Xiong F, Yue GH. Greater movement-related
443 cortical potential during human eccentric versus concentric muscle contractions. *J*
444 *Neurophysiol.* 2001;86(4):1764-1772. doi:10.1152/jn.2001.86.4.1764
- 445 50. Norrbrand L, Pozzo M, Tesch PA. Flywheel resistance training calls for greater
446 eccentric muscle activation than weight training. *Eur J Appl Physiol.* 2010;110(5):997-
447 1005. doi:10.1007/s00421-010-1575-7
- 448 51. Cormie P, McGuigan MR, Newton RU. Developing Maximal Neuromuscular Power.
449 *Sport Med.* 2011;41(2):125-146. doi:10.2165/11538500-000000000-00000
- 450 52. Maroto-Izquierdo S, García-López D, Fernandez-Gonzalo R, Moreira OC, González-
451 Gallego J, de Paz JA. Skeletal muscle functional and structural adaptations after
452 eccentric overload flywheel resistance training: a systematic review and meta-analysis.
453 *J Sci Med Sport.* 2017;20(10):943-951. doi:10.1016/j.jsams.2017.03.004
- 454 53. Markovic G, Mikulic P. Neuro-musculoskeletal and performance adaptations to lower-
455 extremity plyometric training. *Sports Med.* 2010;40(10):859-895.
456 doi:10.2165/11318370-000000000-00000
- 457 54. Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a
458 practical primer for t-tests and ANOVAs. *Front Psychol.* 2013;4(November):1-12.
459 doi:10.3389/fpsyg.2013.00863
- 460 55. Cohen J. *Statistical Power Analysis for the Behavioral Sciences.* New York, NY:
461 Routledge Academic; 1988.
- 462 56. Cuenca-Fernández F, López-Contreras G, Arellano R. Effect on swimming start
463 performance of two types of activation protocols: lunge and YoYo squat. *J strength*
464 *Cond Res.* 2015;29(3):647-655. doi:10.1519/JSC.0000000000000696
- 465 57. Cuenca-Fernández F, Ruiz-Teba A, López-Contreras G, Arellano R. Effects of 2 types
466 of activation protocols based on postactivation potentiation on 50-m freestyle
467 performance. *J strength Cond Res.* 2018;00(00):1.
468 doi:10.1519/JSC.0000000000002698
- 469 58. Cuenca-Fernández F, López-Contreras G, Mourão L, et al. Eccentric flywheel post-
470 activation potentiation influences swimming start performance kinetics. *J Sports Sci.*
471 2019;37(4):443-451. doi:10.1080/02640414.2018.1505183
- 472 59. Timon R, Allemano S, Martinez-guardado I, Olcina G. Post-activation potentiation on
473 squat jump following two different protocols : Traditional vs Inertial flywheel. *J Hum*
474 *Kinet.* 2019. doi:10.2478/hukin-2019-0017
- 475 60. Häggglund M, Waldén M, Ekstrand J. Risk factors for lower extremity muscle injury in

476 professional soccer. *Am J Sports Med.* 2013;41(2):327-335.
477 doi:10.1177/0363546512470634
478 61. Gouvêa AL, Fernandes IA, César EP, Silva WAB, Gomes PSC. The effects of rest
479 intervals on jumping performance: A meta-analysis on post-activation potentiation
480 studies. *J Sports Sci.* 2013;31(5):459-467. doi:10.1080/02640414.2012.738924
481