

Int. J. Aquat. Biol. (2015) 3(2): 119-128  
E-ISSN: 2322-5270; P-ISSN: 2383-0956  
Journal homepage: [www.ij-aquaticbiology.com](http://www.ij-aquaticbiology.com)  
© 2015 Iranian Society of Ichthyology

## Original Article

# Population dynamic parameters of the highly endemic fish, *Alburnoides qanati* Coad and Bogutskaya 2009, (Teleostei: Cyprinidae) in the Kor River Basin, Iran

Mohadeseh Sadat Tahami<sup>1</sup>, Hamid Reza Esmaeili\*<sup>1</sup>, Mohsen Safaie<sup>2</sup>

<sup>1</sup>Ichthyology Lab., Department of Biology, College of Science, Shiraz University, Shiraz, Iran, 71454.

<sup>2</sup>Fisheries Department, Hormozgan University, Bandar Abbas, Iran.

**Abstract:** The present study provides the data on the population parameters of an endemic qanat spiralin, *Alburnoides qanati* from the endorheic Kor River basin of Iran to assess the stock status of this cyprinid fish species, which is highly important from management and conservation points of views. Fish specimens (387) were collected from Moshkan Stream, Kor River basin, Iran in 2011-2012. Asymptotic length ( $L_{\infty}$ ) and growth coefficient ( $K$ ) were estimated at 123.9 mm and 0.31/year for females and 93 mm and 0.49/year for males, respectively. Growth performance index ( $\phi'$ ) was calculated as 8.47 and 8.35 for female and male specimens, respectively. Total mortality ( $Z$ ) of females (1.56/year) was higher than males (1.14/year) whereas natural mortality ( $M$ ) of female specimens (0.44/year) was lower than male specimens (0.65/year). Data on growth and mortality parameters and also length-weight relationship of *A. qanati* revealed significant differences with corresponding data from other *Alburnoides* species from Iran and other countries, which could be attributed to habitat's differences and specific species characteristics.

### Article history:

Received 25 January 2015

Accepted 17 March 2015

Available online 25 April 2015

### Keywords:

Asymptotic length

Growth coefficient

Growth performance index

Management

Conservation

## Introduction

*Alburnoides bipunctatus* (Bloch, 1782) was the name applied to most populations of spiralins (riffle minnows or tailor fish) across Europe and the Middle East from France north of the Alps eastwards to the Black, Caspian and Aral Sea basins (Coad and Bogutskaya, 2012) and reportedly to be one of the abundant and widespread groups in Iranian endorheic and exoreic basins (Esmaeili et al., 2010). However, ongoing research was revealed a greater diversity and presence of at least seven species in different basins of Iran, including the Caspian Sea, Lake Orumiyeh (Urmia), Tedzhen River, Kavir, Namak Lake, Tigris River, Persian Gulf drainage and Kor River basins (Bogutskaya and Coad 2009, Coad and Bogutskaya, 2009, 2012; Esmaeili et al., 2010, 2014a) of which the Kor River populations have been assigned to *Alburnoides qanati*, qanat

tailor fish by Coad and Bogutskaya (2009).

Members of the genus *Alburnoides* are lithophilic and rheophilic fishes, which inhabit in barbell and grayling zones and spawn on gravel and rubble (Breitenstein and Kirchhofer, 2000). The spiralins are very sensitive to human activities and levels of dissolved oxygen. Having low tolerance to water polluted by industrial, agriculture or urban wastes makes these cyprinid fishes a good biological indicator of the environment quality (Čihār, 1999). In European waters, spiralins are extremely threatened and nearly close to extinction because of this sensitivity (Kirchhofer, 1997; Lusk et al., 1998). The freshwater fishes of Iran are also faced to recent severe droughts, climate change, pollutions, introduction of exotic fishes and anthropogenic impacts, and as a consequence, many fish populations have been intensively affected

\* Corresponding author: Hamid Reza Esmaeili  
E-mail address: [esmaeili@susc.ac.ir](mailto:esmaeili@susc.ac.ir)

especially sensitive fishes, like the spirilins (Esmaeili et al., 2014a, b). Hence, there is a constant need for an increase in knowledge on different aspects of biology of these fishes, including estimating growth parameters as an indicator of fish health and habitat quality to implant effective management and conservation measures.

However, many of these aspects on *A. qanati* have not been documented in Iran, partly because of narrow endemic range and no commercial importance due to its slow growth rate and small size.

Therefore, due to the information scarcity on the population dynamics of *A. qanati*, coupled with the need to provide much-needed scientific data for the management and rational exploitation of this valuable resource, the present study was undertaken to evaluate and to estimate the growth characteristics (growth parameters) of this species to assess the stock status of this endemic fish species from a spring-stream system in Kor River basin of Iran for the first time.

### Material and Methods

The samples were collected from Moshkan spring-stream system in Kor River basin located at N 30°36'16.9" and E 52°56'40.1". Sampling was performed monthly from March 2011 to April 2012 by electrofishing device. Samples were fixed in 10% formalin solution and transported to the laboratory. Some morphological characters, including: total length (TL), fork length (FL), standard length (SL) and maximum and minimum body width were measured to the nearest 0.01 mm and fish weight (gr) to the nearest 0.1 gr. Sexes were determined by investigation of the gonads. The relationship between the length and weight were determined by fitting the data to a potential relationship in the form of:  $W = \alpha L^b$ , where  $W$  is the fish weight;  $L$ , fish length; and  $\alpha$  and  $b$  are the parameters to be estimated, with  $b$  being the coefficient of allometry based on the test given by Pauly (1980). Prior to regression analyses, log-log plots of length and weight values were performed for visual inspection

of outliers (Froese, 2006). One-way analysis of variance (ANOVA) was used to test significant relationship between length and weight.

To calculate growth performance and measure the back-calculation of growth in length, data were put in to the FISATII software. The von Bertalanffy growth function was used to study the overall growth performance using the values of growth in length and Phi-Prime ( $\Phi'$ ) (Bertalanffy, 1934; Sparre and Venema, 1992), as follows:

$$L_t = L_\infty(1 - e^{-k(t-t_0)})$$

$$\Phi' = \ln K + \ln L_\infty$$

where  $L_t$  is total length at age  $t$ ,  $L_\infty$  is the ultimate total length that an average fish should achieve if it continues to live and growth,  $K$  is the growth coefficient that determines how fast the fish approaches to  $L_\infty$ ,  $t_0$  is hypothetical age  $L_t=0$ , and  $\Phi'$  is overall growth performance.

Estimations of the mean total mortality rate ( $Z$ ) was obtained from using length converted catch curve analysis. Natural mortality ( $M$ ) was calculated using the equation of Pauly (1980):

$$\text{Log}_{10}M = 0.0066 - 0.279\text{Log}_{10}L_\infty + 0.6543\text{Log}_{10}K + 0.4634\text{Log}_{10}T$$

Where,  $M$  is the natural mortality;  $L_\infty$  is the asymptotic length;  $K$  is the growth co-efficient of the von Bertalanffy growth function (VBGF) and  $T$  is the mean annual water temperature (°C) of sampling site. Mortality rate was calculated by Length Converted Catch Curve Method (Pauly, 1980):

$$F = Z - M, E = F/Z = F/F + M$$

Where  $Z$  is total mortality,  $F$  is fishing mortality,  $E$  is exploitation coefficient and  $M$  is natural mortality.

### Results

The descriptive statistics on *A. qanati* specimens, collected from Moshkan spring-stream, Kor River basin are presented in Table 1. Minimum and maximum total length (TL) for female and male specimens were 24.76 and 118.66 and 24.88 and 87.07, respectively (Table 1). One-way analysis of variance (ANOVA,  $P < 0.05$ ) showed a high significant relationship between length and weight of *A. qanati* in both sexes and the estimates of the

Table 1. Descriptive statistics and estimated parameters of the length-weight relationships (LWRs) for females and males of *Alburnoides qanati* collected from Kor River Basin, during 2011-2012.

Length parameter (mm)	Sex	Min	Max	$b$	$a$	$r^2$	N
TL	Female	24.76	118.66	3.45	0.0033	0.98	209
	Male	24.88	87.07	3.44	0.0033	0.98	155
FL	Female	23.00	110.1	3.35	0.0044	0.97	196
	Male	23.42	81.86	3.32	0.0047	0.97	141
SL	Female	21.02	102.96	3.30	0.0055	0.98	219
	Male	20.76	75.06	3.35	0.0047	0.98	168

TL, total length; FL, fork length; SL, standard length;  $b$ , regression slope;  $a$ , intercept;  $r^2$ , coefficient of determination; N, number of specimens.




Figure 1. Length- Weight relationship of *Alburnoides qanati* from Kor River Basin, Iran.


parameter  $b$ , varied between 3.30 and 3.45 for females and 3.32 and 3.44 for males with high  $r^2$  values of 0.97 to 0.98 for both sexes (Table 1, Figs. 1a, b). The parameters of growth including growth coefficient ( $K$ ), asymptotic length ( $L_\infty$ ), overall growth performance ( $\phi'$ ), and total, fishing, natural

and exploitation mortality ( $Z$ ,  $F$ ,  $M$ ,  $E$ ) for females and males of *A. qanati* are given in Table 2 and von Bertalanffy growth curves are shown in Figure 2. It can be realized that infinity length ( $L_\infty$ ) of females is greater than males (123.9 vs. 93 mm, respectively) and value of  $K$  parameter for females is lower than

Table 2. Growth parameters for females and males of *Alburnoides qanati* from Kor River basin, Iran.

Growth Parameters	$L_{\infty}$ (mm)	$K$	$\emptyset'$	$Z$	$M$	$F$	$E$
Female	123.9	0.31	8.47	1.56	0.44	1.12	0.72
Male	93	0.49	8.35	1.14	0.65	0.49	0.43

von Bertalanffy growth parameters ( $L_{\infty}$ ,  $K$ ), overall growth performance ( $\emptyset'$ ), total mortality ( $Z$ ), fishing mortality ( $F$ ), natural mortality ( $M$ ) and exploitation mortality ( $E$ ).

Figure 2. Von Bertalanffy growth curves of *Alburnoides qanati* (a., female; b. male).

males (0.31 vs. 0.49, respectively) (Table 2).

### Discussion

The ranges of *A. qanati* length obtained in this study is in the ranges reported for other *Alburnoides* species from different regions. It indicates that females are almost larger than males (Table 3). This attribute could be interpreted as an inter-population pattern related to nature of water body (river, stream, spring, qanat), different habitat quality, growth rate and some intrinsic factors. Length-weight relationships

(LWRs) showed highly significant positive relationships for *A. qanati* as it can be seen in other reports of this genus (Table 4) in the range of 2.5-3.6 (Treer et al., 2000; Patimar et al., 2012; Tabatabaei et al., 2014). The estimation of  $b$  parameter is remained within the expected range of 2.5-4 reported by Tesch (1971). The upper limit showing deep body form and that fish getting more weight as its length increases. The reasons for the variation of  $b$  in the different regions are reported to be due to seasonal fluctuations in environmental parameters,

Table 3. Maximum observed length (TL, cm) and age (years) for different *Alburnoides* species in the European region and Iran. A.B., *A. bipunctatus*; A.E., *A. eichwaldii*; A.N., *A. namaki*; A.Q., *A. qanati*; F, female; M, male; U, unsexed; \* mean TL.

Species	Sex	TL	Age	Country	Reference
A.B.	F	10.3	6+	Velika Morava River, former Yugoslavia	Soric and Ilic (1985)
A.B.	F	12.53	6+	Oltu stream, Coruh Basin, Turkey	Yıldırım et al. (1999)
A.B.	U	11.72	---	Rudava River, Slovakia	Siryova (2004)
A.B.	F	11.0	4+	Sava River, Croatia	Treer et al. (2006)
A.B.	U	13.0 (SL)	---	European freshwater systems	Kottelat and Freyhof (2007)
A.B.	U	16.0	3+	Azerbaijan	Abdurahmanov (1962)
A.B.	U	12.5	---	Madarsu Stream, National Park of Golestan, northern Iran	Akbaripasand (1999)
A.E.	U	15.0	---	Sardabroud River, northern Iran	Abdoli (2000)
A.E.	M	9.5	4+	Zarrin-Gol River, northern Iran	Patimar and Dowlati (2007)
A.E.	F	11.0	4+	Zarrin-Gol River, northern Iran	Patimar and Dowlati (2007)
A.E.	U	14.0	6+	Rivers of northern Iran	Abdoli and Naderi (2009)
A.E.	M	11.0	4+	Qanat of Uzineh, northern Iran	Patimar et al. (2012)
A.E.	F	11.1	4+	Qanat of Uzineh, northern Iran	Patimar et al. (2012)
A.N.	M	6.06*		Gharachai, Namk basin, Iran	Tabatabaei et al. (2014)
A.N.	F	7.45*		Gharachai, Namk basin, Iran	Tabatabaei et al. (2014)
A.N.	M	6.49*		Jajrud, Namak basin, Iran	Tabatabaei et al. (2014)
A.N.	F	6.91*		Jajrud, Namak basin, Iran	Tabatabaei et al. (2014)
A.N.	M	8.41*		Cheshme Ali, Kavir basin, Iran	Tabatabaei et al. (2014)
A.N.	F	8.62*		Cheshme Ali, Kavir basin, Iran	Tabatabaei et al. (2014)
A.E.	M	8.01*		Tajan River, Caspian basin, Iran	Tabatabaei et al. (2014)
A.E.	F	8.70*		Tajan River, Caspian basin, Iran	Tabatabaei et al. (2014)
A.Q.	M	8.71		Kor River basin, Iran	Present study
A.Q.	F	11.87		Kor River basin, Iran	Present study

physiological conditions of the fish at the time of collection, sex, gonad development, food availability and quality in habitat, diet, stomach fullness, health and the preservation techniques of the samples (Tesch, 1971; Esmaeili, 2001; Esmaeili and Ebrahimi, 2006; Esmaeili et al., 2014c).

Fitting the von Bertalanffy growth formula (VBGF) to back-calculated lengths resulted in the estimation of higher values of L-infinity than the maximum observed total lengths for both sexes. This length is seen as a capacity for growth (Bagenal and Tesch, 1978) and inter-sex differences in this parameter of the VBGF correspond to different growth rates of the sexes. On the other hand, a different VBGF can be a result of resource allocation between growth and reproduction. Therefore, differences between sexes in the VBGF parameters could also reflect differences in the reproductive effort of the fish (see

Bagenal and Tesch, 1978; Patimar et al., 2012). It is theoretically admitted that growth parameters  $L_{\infty}$  and  $K$  are negatively correlated (Gaertner et al., 2008). The parameters of von Bertalanffy's equation obtained for linear growth of *A. qanati* confirm the theoretical assumption explained in Raikova-Petrova et al. (2011) that the smaller ( $L_{\infty}$ ), shows a faster growth rate ( $K$ ) which is remarkably exhibited in male and female populations of *A. qanati*. The  $K$  parameter of male specimens is higher than females (0.49 vs. 0.31, respectively) revealing that the male grows rapidly initially and approaches its asymptotic length ( $L_{\infty}=93$  mm) earlier in life. The same as other species of the family Cyprinidae, the females of *A. qanati* were found to have reasonably larger  $L_{\infty}$  (123.9 vs. 93 mm, respectively) and smaller  $K$ -values than males (0.31 vs.0.49, respectively). A comparison between estimates of von Bertalanffy

Table 4. Parameters of the length-weight relationship of spirilins, *Alburnoides* species. A.B., *A. bipunctatus*; A.E., *A. eichwaldii*; A.N., *A. namaki*; A.Q., *A. qanati*; F, female; M, male; C, combined sex; P, population.

Species	Area	Sex	Length	$a$	$b$	Reference
A.B.	Dobra River, Croatia	C	TL	0.0059	3.2245	Treer et al. (2000)
A.B.	Bednja River, Croatia	C	TL	0.0150	2.7970	Treer et al. (2000)
A.B.	Middle Korana River, Croatia	C	TL	0.0088	3.1043	Treer et al. (2000)
A.B.	Lower Korana River, Croatia	C	TL	0.0030	3.5567	Treer et al. (2000)
A.B.	Sava River, Croatia	C	TL	0.0044	3.4032	Treer et al. (2000)
A.B.	Çoruh River, Turkey	C	FL	0.0249	2.79	Torcu-Koç et al. (2006)
A.B.	Çoruh River, Turkey	F	FL	0.0375	2.62	Torcu-Koç et al. (2006)
A.B.	Çoruh River, Turkey	M	FL	0.0166	2.95	Torcu-Koç et al. (2006)
A.B.	Sava River, Croatia	C	TL	0.0083	3.025	Treer et al. (2006)
A.B.	Seyhan Dam Lake, Turkey	C	TL	0.0028	2.72	Ergüden and Goksu (2009)
A.B.	Emajõgi River Basin, Estonia	C	SL	0.0103	3.251	www.fishbase.org
A.E.	Zarrin-Gol River, northern Iran	M	TL	0.0054	2.59	Patimar and Dowlati (2007)
A.E.	Zarrin-Gol River, northern Iran	F	TL	0.0088	2.52	Patimar and Dowlati (2007)
A.E.	Qanat of Uzineh, northern Iran	M	TL	0.0068	3.2559	Patimar et al. (2012)
A.E.	Qanat of Uzineh, northern Iran	F	TL	0.0079	3.2067	Patimar et al. (2012)
A.E.	Qanat of Uzineh, northern Iran	P	TL	0.0072	3.2387	Patimar et al. (2012)
A.Q.	Gharachai, Namk basin, Iran	F	TL	0.01	2.87	Tabatabaei et al. (2014)
A.Q.	Gharachai, Namk basin, Iran	M	TL	0.01	3.05	Tabatabaei et al. (2014)
A.Q.	Jajrud, Namak basin, Iran	F	TL	0.01	3.26	Tabatabaei et al. (2014)
A.Q.	Jajrud, Namak basin, Iran	M	TL	0.01	3.20	Tabatabaei et al. (2014)
A.Q.	Cheshme Ali, Kavir basin, Iran	F	TL	0.02	2.77	Tabatabaei et al. (2014)
A.Q.	Cheshme Ali, Kavir basin, Iran	M	TL	0.01	3.05	Tabatabaei et al. (2014)
A.E.	Tajan River, Caspian basin, Iran	F	TL	0.01	3.26	Tabatabaei et al. (2014)
A.E.	Tajan River, Caspian basin, Iran	M	TL	0.01	3.30	Tabatabaei et al. (2014)
A.Q.	Kor River basin, Iran	F	TL	0.0033	3.45	Present study
A.Q.	Kor River basin, Iran	M	TL	0.0033	3.44	Present study

length-at-age growth parameters for different species and populations of spirilins (Skorva, 1972; Breitenstein and Kirchhofer, 2000; Treer et al., 2000; Treer et al., 2006; Raikova-Petrova et al., 2011; Seifali et al., 2012; Patimar et al., 2012; Tabatabaei et al., 2014 and present study) reveals that these parameters vary for different populations, species at different water bodies (e.g. qanat, stream, river) and different distribution ranges (Table 5). The  $L_{\infty}$  and  $K$ -values ranges from 93 to 123.9 and 0.31 to 0.49, respectively. Many studies have revealed that under different environmental conditions, biological features such as life span (Abdoli et al., 2007; Mann et al., 1984; Ricker, 1975), age at maturity (Lobón-Cerviá et al., 1996), age structure and growth rate (Abdoli et al., 2007; Kamal et al., 2009; Patimar et al., 2012b; Basilone et al., 2004; Naddafi et al., 2005;

Tabatabaei et al., 2014), the maximum length ( $L_{\infty}$ ) (Basilone et al., 2004; Kamal et al., 2009; Naddafi et al., 2005; Tabatabaei et al., 2014) can change. For example, according to Saifali et al. (2012), population parameters for south Caspian spirilin (*A. eichwaldii*) are as follows: Asymptotic length ( $L_{\infty}$ ) = 104.48 mm; Growth co-efficient ( $K/\text{year}$ ) = 1.19; Natural mortality ( $M/\text{year}$ ) = 0.97; Fishing mortality ( $F/\text{year}$ ) = 2.43; Total mortality ( $Z/\text{year}$ ) = 3.4 and Exploitation level ( $E$ ) = 0.71.

In the present work,  $\Phi'$  was estimated to be 3.86 and 3.75 for female and male (based on the  $L_{\infty}$  of 9.3 cm and 12.39 cm) specimens of *A. qanati*, respectively, which is less than other spirilins (Table 6). According to Pauly (1979), phi prime values are very similar within related taxa and have narrow normal distributions. However, similarity of this index in

Table 5. Estimates of von Bertalanffy length-at-age growth parameters for populations of different *Alburnoides* species in the distribution area (Iran and European regions). A.B., *A. bipunctatus*; A.E., *A. eichwaldii*; A.N., *A. namaki*; A.Q., *A. qanati*; F, female; M, male; U, unsexed.

Species	Location	Sex	$L_{\infty}$ (mm)	$K$ (year <sup>-1</sup> )	$T_0$ (year)	Reference
A.B.	Dunajec River, Czechoslovakia	U	20.1	0.15	---	Skora (1972)
A.B.	Turiec River, former Czechoslovakia	U	15.6	0.28	---	Bastl et al. (1975)
A.B.	Radimna River, Romania	U	14.4	0.30	---	Papadopol and Cristofor (1980)
A.B.	Dobra River, Croatia	U	20.5	0.16	-1.38	Treer et al. (2000)
A.B.	Bednja River, Croatia	U	15.5	0.33	-0.42	Treer et al. (2000)
A.B.	Middle Korana River, Croatia	U	15.1	0.28	-0.86	Treer et al. (2000)
A.B.	Lower Korana River, Croatia	U	17.7	0.19	-1.47	Treer et al. (2000)
A.B.	Sava River, Croatia	U	11.5	0.59	-0.47	Treer et al. (2000)
A.B.	Sava River, Croatia	U	12.0	0.59	-0.14	Treer et al. (2006)
A.E.	Zarrin-Gol River, northern Iran	M	99.64	0.51	-0.715	Patimar and Dowlati (2007)
A.E.	Zarrin-Gol River, northern Iran	F	107.23	0.55	-0.548	Patimar and Dowlati (2007)
A.E.	Qanat of Uzineh, northern Iran	M	140.7	0.27	-0.92	Patimar et al. (2012)
A.E.	Qanat of Uzineh, northern Iran	F	153.7	0.23	-1.08	Patimar et al. (2012)
A.E.	Qanat of Uzineh, northern Iran	U	148.3	0.24	-1.04	Patimar et al. (2012)
A.E.	Kesselian Stream, Caspian Sea, Iran	U	104.5	1.19	---	Seifali et al. (2012)
A.N.	Gharachai, Namk basin, Iran	U	122.43	0.27	-0.76	Tabatabaei et al. (2014)
A.N.	Jajrud, Namak basin, Iran	U				
A.N.	Cheshme Ali, Kavir basin, Iran	U	112.66 120	0.54 0.29	0.18 -1	Tabatabaei et al. (2014) Tabatabaei et al. (2014)
A.E.	Tajan River, Caspian basin, Iran	U	123.01	0.29	-1	Tabatabaei et al. (2014)
A.Q.	Kor River basin, Iran	F	123.9	0.31	---	Present study
A.Q.	Kor River basin, Iran	M	93	0.49	---	Present study

Table 6. Phi prime ( $\phi'$ ) values of different species of *Alburnoides* from Europe and Iran.

Locality	Species	$\phi'$	Reference
Turiec ,Slovenian	<i>A. bipunctatus</i>	4.22	Bastl et al. (1975)
Radimna, Slovenian	<i>A. bipunctatus</i>	4.13	Papadopol and Cristofor (1980)
Croatian- Slovenian	<i>A. bipunctatus</i>	4.17	Treer et al. (2000)
Caspian Sea, Iran	<i>Alburnoides</i> sp.	4.87	Saifali et al. (2012)
Kor river, Iran	<i>A. qanati</i> (F)	3.86	Present study
Kor river, Iran	<i>A. qanati</i> (M)	3.75	Present study

northern populations of *Alburnoides* (ex. south Caspian Sea spiralin and the European populations) and its difference with *A. qanati* may reflect the distinctiveness of northern and southern populations of *Alburnoides*.

Mortality parameters are higher for Caspian spiralin than *A. qanati* from Kor Basin; it can be due to more sympatric predator species and over-fishing activities in Caspian Basin. Being small spring, Moshkan has no predatory fish, and *A. qanati* and

*Oxynemacheilus persa* are the only fish species inhabit in this stream. It also has been reported that the spiralin population reflects the changes in habitats (Jurajda et al., 1996); therefore, water dam construction can critically endanger this species (Lusk, 1995) which can be considered for all species of genus *Alburnoides* including *A. qanati* species (Kirchhofer, 1997; Lusk et al., 1998; Treer et al., 2006).

Based on the available data, it can be concluded that many factors affect growth parameters of fish populations. The commonly proposed explanations include the effects of changes in environment conditions, (most notably temperature and food availability), density-dependent effects, eutrophication, habitat disruption and specific species characteristics. These factors can operate singly or in combination and have positive or negative effects on growth parameters. According to Ma et al. (2010), the slow-growing and long-lived fish tend to be, particularly, vulnerable to excessive mortality and rapid stock collapse and population turnover may be lower than expected recovery. Study of these parameters for the endemic fish, *A. qanati* is highly recommended.

### Acknowledgments

We wish to thank R. Khaefi, B. Parsi, G. Sayadzadeh and S. Mirghiasi for their valuable helps with the fish collections and Shiraz University for the financial support.

### References

Abdoli A., Pont D., Sagnes P. (2007). Intrabasin variations in age and growth of bullhead: the effects of temperature. *Journal of Fish Biology*, 70: 1224-1238.

Abdurahmanov Y.A. (1962). *Ryby presnykh vod Azerbidjana*. Baku. 224 pp. In Azerbaijani.

Akbaripasand A. (1999). Ecological investigation of fishes of Gorganroud River in the National Park of Golestan. MSc thesis, Tarbiat Modarres University, Tehran, pp: 114.

Basilone G., Guisande C., Patti B., Mazzola S., Cuttitta A., Bonanno A., Kallianiotis A. (2004). Linking habitat conditions and growth in the European

anchovy (*Engraulis encrasicolus*). *Fisheries Research*, 68: 9-19.

Bastl I., Holèik J., Kirka A. (1975). Ichthyologic investigation of the protected habitat of the Danubian salmon (*Hucho hucho* L.) on the River Turiec (Czechoslovakia) and suggestions for its management. Academia Research. National Natural Museum of Slovak, Bratislava, 21: 191-224. [In Slovakian with English summary.]

Bagenal T., Tesch F. (1978). *Methods for assessment of fish production in fresh waters*, IBP Handbook 3 Blackwell, Oxford.

Bertalanffy L.V. (1934). Untersuchungen über die Gesetzlichkeit des Wachstums I. Allgemeine Grundlagen der Theorie. Mathematisch-physiologische Gesetzlichkeiten des Wachstums bei Wassertieren, Wilhelm Roux' Archiv für Entwicklungsmechanik der Organismen, 131: 613-652.

Breitenstein M.E., Kirchhofer A. (2000). Growth, age structure and species association of the cyprinid *Alburnoides bipunctatus* in River Aare, Switzerland. *Folia Zoologica*, 49(1): 59-68.

Bogutskaya N.G., Coad B.W. (2009). A review of vertebral and fin-ray counts in the genus *Alburnoides* (Teleostei: Cyprinidae) with a description of six new species. *Zoosystema Rossica*, 18(1): 126-173.

Čihár J. (1999). *A field guide in color to freshwater fish*. Blitz ed., Leicester.

Coad B.W., Bogutskaya N.G. (2009). *Alburnoides qanati*, a new species of cyprinid fish from southern Iran (Actinopterygii, Cyprinidae). *ZooKeys*, 13: 67-77.

Coad B.W., Bogutskaya N.G. (2012). New species of riffle minnow, *Alburnoides holciki*, from the Hari River basin in Afghanistan and Iran (Actinopterygii: Cyprinidae). *Zootaxa*, 3453: 43-55.

Coad B.W. (2012). *Freshwater Fishes of Iran*. Available at [www.briancoad.com](http://www.briancoad.com) (accessed October 2012).

Ergüden S.A., Goksu M.Z.L. (2009). Length-weight relationships for 12 fish species caught in Seyhan Dam Lake in southern Anatolia, Adana, Turkey. *Journal of Applied Ichthyology*, 25(4): 501-502.

Esmaili H.R. (2001). Biology of an exotic fish, silver carp, *Hypophthalmichthys molitrix*, from Gobindsagar reservoir, India. PhD thesis, Punjab University, India.

Esmaili H.R., Ebrahimi M. (2006). Length-weight relationships of some freshwater fishes of Iran.



- Journal of Applied Ichthyology, 22: 328-329.
- Esmaeili H.R., Coad B.W., Gholamifard A., Nazari N., Teimori A. (2010). Annotated checklist of the freshwater fishes of Iran. *Zoosystematica Rossica*, 19: 361-386.
- Esmaeili H.R., Coad B.W., Mehraban H.R., Masoudi M., Khaefi R., Abbasi K., Mostafavi H., Vatandoust S. (2014a). An updated checklist of fishes of the Caspian Sea basin of Iran with a note on their zoogeography. *Iranian Journal of Ichthyology*, 1(3): 152-184.
- Esmaeili H.R., Teimori A., Owfi F., Abbasi K., Coad B.W. (2014b). Alien and invasive freshwater fish species in Iran: Diversity, environmental impacts and management. *Iranian Journal of Ichthyology*, 1(2): 62-72.
- Esmaeili H., Gholamifard A., Vatandoust S., Sayyadzadeh G., Zare R., Babaei S. (2014c). Length-weight relationships for 37 freshwater fish species of Iran. *Journal of Applied Ichthyology*, 30(5): 1073-1076.
- Froese R. (2006). Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22: 241-253.
- Gaertner D., Delgado de Molina A., Ariz J., Pianet R., Hallier J.P. (2008). Variability of the growth parameters of the skipjack tuna (*Katsuwonus pelamis*) among areas in the eastern Atlantic: analysis from tagging data within a meta-analysis approach. *Aquatic Living resources*, 21: 349-356.
- Jurajda P., Hohausova E., Prasek V., Dvorak M. (1996). Fishes of Polana Mountain streams. *Biologia*, 51: 173-178.
- Kamal S., Bakhtiyari M., Abdoli A., Eagderi S., Karami M. (2009). Life-history variations of killifish (*Aphanius sophiae*) populations in two environmentally different habitats in central Iran. *Journal of Applied Ichthyology*, 25: 474-478.
- Kirchhofer A. (1997). The assessment of Fish vulnerability in Switzerland based on distribution data. *Biological Conservation*, 80: 1-8.
- Kottelat M., Freyhof J. (2007). Handbook of European Freshwater Fishes. Kottelat Cornol, Switzerland and Freyhof, Berlin. xiii + 646 pp.
- Lobón-Cerviá J., Dgebuadze Y., Utrilla C., Rincon P., Granado-Lorencio C. (1996). The reproductive tactics of dace in central Siberia: evidence for temperature regulation of the spatio-temporal variability of its life history. *Journal of Fish Biology*, 48: 1074-1087.
- Lusk S. (1995). Influence of valley dams on the changes in fish communities inhabiting streams in the Dyje drainage area. *Folia Zoologica*, 44: 45-56.
- Lusk S., Lusková V., Halačka K., Šlechta V., Šlechtova V. (1998). Trends and production of fish communities of the barbell zone in a stream of the Czech Republic. *Folia Zoologica*, 47:67-72.
- Ma B.S., Xie C.X., Huo B., Yang X.F., Huang H.P. (2010). Age and growth of a long-lived fish *Schizothorax o'connori* in the Yarlung Tsangpo River, Tibet. *Zoological Studies*, 49: 749-759.
- Mann R., Mills C., Crisp D. (1984). Geographical variation in the life-history tactics of some species of freshwater fish. In: G.W. Potts, R.J. Wootton (eds.). *Fish reproduction: strategies and tactics*. London: Academic Press. pp: 171-186.
- Naddafi R., Abdoli A., Hassanzadeh Kiabi B., Mojazi Amiri B., Karami M. (2005). Age, growth and reproduction of the Caspian roach (*Rutilus rutilus caspicus*) in the Anzali and Gomishan wetlands, North Iran. *Journal of Applied Ichthyology*, 21: 492-497.
- Papadopol M., Cristofor S. (1980). Recherches sur l'ecologie de deux populations de spirilin, *Alburnoides bipunctatus*, des eaux de la Roumanie. *Travaux de Museum d'Histoire Naturelle "Grigore Antipa"*, 22: 483-493.
- Patimar R., Dowlati F. (2007). Investigation on age, growth and reproduction of riffle minnow *Alburnoides bipunctatus* (Bloch, 1782) in Zarrin-Gol River, east Alborz Mountain. *Journal of Fisheries*, 1(1): 55-62.
- Patimar R., Zare M., Hesam M. (2012). On the life history of spirilin *Alburnoides bipunctatus* (Bloch, 1782) in the qanat of Uzineh, northern Iran. *Turkish Journal of Zoology*, 36: 383-393.
- Pauly D., Gaschütz G. (1979). A simple method for fitting oscillating length growth data, with a program for pocket calculators. Pp. 26. In: International Council for the Exploration of the Sea. Council meeting 1979/G: 24. Demersal Fish Committee.
- Pauly D. (1980). On the interrelationships between natural mortality, growth parameters and mean environmental temperature in 175 fish stocks. *Journal of Conservation and International Exploration of Maritime*, 39(3): 175-192.
- Raikova-Petrova G.N., Petrov I.K., Hamwi N.I., Marinova D.M. (2011). Growth Rate and Condition of

- Riffle Minnow (*Alburnoides bipunctatus* Bloch) from the Middle Stream of Iskar River (Bulgaria). *Acta Zoologica Bulgarica*, 63(3): 295-300.
- Ricker W.E. (1975). Computation and interpretation of biological statistics of fish populations. *Bullitan of the Fisheries Resources Board of Canada* 191, p. 382.
- Seifali M., Arshad A., Nurul Amin S.M., Hasanzadeh Kiabi B., Esmaili H.R., Yazdani Moghaddam F. (2012). Population dynamics of Caspian spiralin (*Actinopterygii: Cyprinidae*) in the Kesselian Stream, Iran. *African Journal of Biotechnology*, 11(38): 9214-9222.
- Siryova S. (2004). External morphology of spiralin *Alburnoides bipunctatus* (Bloch). *Acta Zoologica*, 46: 113-122
- Skora S. (1972). The Cyprinid *Alburnoides bipunctatus* Bloch from the basins of the rivers Upper San and Dunajec. *Acta Hydrobiologica*, 14: 173-204.
- Soric V.M., Ilic K.R. (1985). Systematical and ecological characteristics of *Alburnoides bipunctatus* (Bloch) in some waters of Yugoslavia. *Ichthyologica* 17(1): 47-58.
- Sparre P., Venema S.C. (1992). Introduction to tropical fish stock assessment Part 1, Rome, FAO, pp: 376.
- Tabatabaei S.N., Hashemzadeh-Segherloo I., Abdoli A., Milani M., Mirzaei R. (2014). Age and growth of spiralins, *Alburnoides eichwaldii* and *A. namaki*, from the Caspian, Kavir and Namak basins of Iran. *Iranian Journal of Ichthyology*, 1 (4).
- Tesch F.W. (1971). Age and growth. In: W.E. Ricker (ed.). *Methods for Assessment of Fish Production in Freshwaters*. Blackwell Scientific Publications: Oxford. Pp. 98-100.
- Treer T., Habekovi D., Anic I.I., Safner R., Piria M. (2000). Growth of five spiralin (*Alburnoides bipunctatus*) populations from the Croatian rivers. *Agriculturae Conspectus Scientificus*, 65(3): 175-180.
- Treer T., Piria M., Aničić I., Safner R., Tomljanović T. (2006). Diet and growth of spiralin, *Alburnoides bipunctatus*, in the barbell Zone of Sava River. *Folia Zoologica*, 55(1): 97-106.
- Torcu-Koç H., Erdoğan Z., Treer T. (2006). A review of length-weight relationships of fishes from freshwaters of Turkey. *Journal of Applied Ichthyology*, 22(4): 264-270.
- Yıldırım A., Erdoğan O., Turkmen M., Demir B.C. (1999). The investigation of some reproduction characteristics of the *Alburnoides bipunctatus faciatii* (Nordman, 1840) living in Oltu stream, Coruh basin. *Turkish Journal of Veterinary and Animal Sciences*, 4: 679-686.