

**Int. J. Aquat. Biol.** (2015) 3(5): 346-351  
 E-ISSN: 2322-5270; P-ISSN: 2383-0956  
 Journal homepage: [www.ij-aquaticbiology.com](http://www.ij-aquaticbiology.com)  
 © 2015 Iranian Society of Ichthyology

## Original Article

### Acute toxicity of *Euphorbia turcomanica* on *Aphanius dispar*

Homa Zare<sup>1</sup>, Ahmad Noori\*<sup>1</sup>, Morteza Yusefzadi<sup>2</sup>, Mahdi Banaee<sup>3</sup>

<sup>1</sup>Department of Fisheries Science, Faculty of Marine Science and Technology, Hormozgan University, Bandar Abbas, Iran.

<sup>2</sup>Department of Marine Biology, Faculty of Marine Science and Technology, Hormozgan University, Bandar Abbas, Iran.

<sup>3</sup>Department of Aquaculture, Faculty of Natural Resources and Environment, Behbahan Khatam Alanbia University of Technology, Behbahan, Iran.

**Abstract:** Piscicidal and molluscicidal activities of aqueous extracts of many members of the family Euphorbiaceae are well-known, but the toxicity potential of *Euphorbia turcomanica* was not yet studied on any aquatic animals. An acute toxicity test was performed by using a four-day static renewal test to determine the LC<sub>50</sub> value of dried powder of *E. turcomanica* for the euryhaline fish, *Aphanius dispar*. The LC<sub>50</sub> values at various exposure periods are 0.177±0.039 g/L for 24 hrs, 0.131±0.030 g/L for 48 hrs, 0.073±0.018 g/L for 72 hrs, and 0.052±0.013 g/L for 96 hrs. The toxicity of dried powder of *E. turcomanica* exhibits a positive correlation between fish mortality and exposure periods. As this is the first report about toxicity of *E. turcomanica* on *A. dispar*, the results could be only compared to that of other Euphorbiaceae as well as other fishes. It is concluded that the toxicity potential of *E. turcomanica* is comparable and close to that of well-documented Euphorbiaceae. It has been suggested that *E. turcomanica* products cannot be used directly in fish-inhabiting water reservoirs.

#### Article history:

Received 10 July 2015

Accepted 25 October 2015

Available online 25 November 2015

#### Keywords:

*Euphorbia turcomanica*

Dried powder

Piscicidal

Killifish

## Introduction

Increased awareness of the negative effects caused by overexposure to synthetic organo-piscicides (Reidinger and Russell, 1976) has led to efforts for finding products from plant origin to substitute. Being the products of biosynthesis, they are potentially biodegradable (Marston and Hostettmann, 1985). The Euphorbiaceae is a large families with about 300 genera and 7500 species (Vasas and Hohmann, 2014). Of the member of this family, the genus *Euphorbia* with about 2000 species (Frodin, 2004), is one of the five most species-rich genera of flowering plants (Govaerts et al., 2000). They have a poisonous milky white latex-like sap and unique kind of floral structures. The chemical constituents of these plants, include triterpenoids and related compounds (sterols, alcohols and hydrocarbons), phenolic compounds (flavonoids, lignans, coumarins, tannins, phenanthrenes,

quinones, phenolic acids, etc.), alkaloids, cyanogenic glucosides, and glucosinolates (Abdel-Fattah, 1987; Neuwinger, 2004; Kumar et al., 2010) that are poisonous to target and non-target aquatic organisms (Singh et al., 1996; Ebenso, 2004; Prasad et al., 2010). Many of these plants are cosmopolitan distributing in subtropics and temperate regions (Horn et al., 2012). In the flora of Iran, this genus comprises 70 species, of which 17 species are endemic (Mozaffarian, 1996). *Euphorbia turcomanica* Boiss, is an annual herb, which grows wild in plains of Iran (Mozaffarian, 1996; Pahlevani and Riina, 2011).

Different species of the genus *Euphorbia* are used as insecticide, piscicide, and molluscicide (Sastry and Siddiqui, 1983; Oliveira-Filho and Paumgarten, 2000; Tiwari and Singh, 2004; Tiwari et al., 2004; Singh et al., 2005; Oliveira-Filho et al., 2010; Hassan et al., 2011). Since, there is no information available

\* Corresponding author: Ahmad Noori  
 E-mail address: [nooryahmad@gmail.com](mailto:nooryahmad@gmail.com)




Figure 1. Percent mortality of the fish *Aphanis dispar* after 24, 48, 72 and 96 hrs exposures to different concentrations of *Euphorbia turcomanica* powder (g/L).

regarding the effect of *E. turcomanica* on fishes as piscicid. Therefore, due to wide use of various parts of the members of the family Euphorbiaceae (Bani et al., 1997; Abdel-Hamid, 2003; Srivastava et al., 2004; Tiwari et al., 2004), this study was conducted to assess the acute toxicity of the lethal concentrations of aqueous extracts of *E. turcomanica*'s aerial parts on *Aphanis dispar*, an euryhaline fish of Iranian inland water.

### Materials and Methods

A total of 240 individuals of *A. dispar* (both sexes: with mean body weight and total length of  $2.03 \pm 0.5$  g, and  $47.7 \pm 0.45$  mm, respectively) were collected from seasonal rivers in Bandar Abbas and transported to the Hormozgan University Fisheries laboratory. In the laboratory, healthy fish were introduced into a 100 L tank with continuous aeration, where they were acclimatize for 14 days to the laboratory conditions. The fish were considered fully acclimatized when no death was observed for seven successive days. The fish were fed 2-3 times a day during this period with commercial pellets,

containing protein >28%, lipid >3%, fiber <4% and moisture <10%.

The stem, branches and leaves of *E. turcomanica* were dried in room temperature away from direct sunlight. Then, they were powdered and mixed with water to obtain the required plant concentrations. For the determination of LC<sub>50</sub> of *E. turcomanica* on *A. dispar*, the four-day static renewal acute toxicity test was performed based on Clesceri et al. (1998). Fish were exposed to 0.001, 0.01, 0.1, 1, 2, 4 and 6 g/L of the dried powder of *E. turcomanica* with three replicates each containing 10 fish (kept in 4 L plastic aquaria). In addition, control group with three replicates in similar stocking density and aquaria were considered for this experiment.

The water was renewed every day and required dried plant was added after water renewal to keep experimental concentrations. The fish were not feed 24 hrs before and during the experiment. Dead fish were counted and removed from the treatments immediately. A toxic effect was determined by a statistically significant decrease in the survival of fish exposed to the plant relative to the survival of

Table 1. Effective dose, confidence limits, and slope function for aqueous extract of *Euphorbia turcomanica* at different intervals on *Aphanius dispar*.

Exposure periods	Effective dose (g/L)	SE	limits		Slope function	't' ratio
			LCL	UCL		
24 hrs	LC <sub>1</sub> =0.010	0.004	0.004	0.024	1.414±0.229	6.183
	LC <sub>5</sub> =0.024	0.008	0.012	0.047		
	LC <sub>10</sub> =0.037	0.011	0.020	0.067		
	LC <sub>20</sub> =0.063	0.017	0.038	0.106		
	<b>LC<sub>50</sub>=0.177</b>	<b>0.039</b>	<b>0.115</b>	<b>0.272</b>		
	LC <sub>80</sub> =0.496	0.117	0.313	0.789		
	LC <sub>90</sub> =0.851	0.228	0.503	1.439		
	LC <sub>95</sub> =1.327	0.401	0.734	2.400		
LC <sub>99</sub> =3.056	1.155	1.457	6.411			
48 hrs	LC <sub>1</sub> =0.005	0.002	0.002	0.013	1.485±0.226	6.556
	LC <sub>5</sub> =0.014	0.005	0.007	0.028		
	LC <sub>10</sub> =0.023	0.007	0.012	0.043		
	LC <sub>20</sub> =0.041	0.011	0.024	0.071		
	<b>LC<sub>50</sub>=0.131</b>	<b>0.030</b>	<b>0.084</b>	<b>0.204</b>		
	LC <sub>80</sub> =0.414	0.104	0.253	0.678		
	LC <sub>90</sub> =0.757	0.219	0.430	1.334		
	LC <sub>95</sub> =1.246	0.409	0.654	2.373		
LC <sub>99</sub> =3.172	1.317	1.406	7.156			
72 hrs	LC <sub>1</sub> =0.001	0.001	0.001	0.004	1.570±0.219	7.163
	LC <sub>5</sub> =0.005	0.002	0.002	0.011		
	LC <sub>10</sub> =0.009	0.003	0.004	0.018		
	LC <sub>20</sub> =0.018	0.006	0.010	0.033		
	<b>LC<sub>50</sub>=0.073</b>	<b>0.018</b>	<b>0.045</b>	<b>0.118</b>		
	LC <sub>80</sub> =0.296	0.083	0.171	0.512		
	LC <sub>90</sub> =0.618	0.202	0.325	1.173		
	LC <sub>95</sub> =1.133	0.427	0.541	2.373		
LC <sub>99</sub> =3.540	1.714	1.371	9.144			
96 hrs	LC <sub>1</sub> =0.001	0.000	0.000	0.003	1.675±0.228	7.332
	LC <sub>5</sub> =0.003	0.001	0.001	0.007		
	LC <sub>10</sub> =0.005	0.002	0.003	0.012		
	LC <sub>20</sub> =0.012	0.004	0.006	0.023		
	<b>LC<sub>50</sub>=0.052</b>	<b>0.013</b>	<b>0.032</b>	<b>0.087</b>		
	LC <sub>80</sub> =0.230	0.067	0.130	0.407		
	LC <sub>90</sub> =0.500	0.172	0.255	0.981		
	LC <sub>95</sub> =0.949	0.378	0.435	2.070		
LC <sub>99</sub> =3.155	1.621	1.153	8.635			

fish in a control. The physico-chemical parameters of the water during experiment, including temperature (23.5±1°C), dissolved oxygen (4.17±0.1 mg/L), electrical conductivity (840.41±2.46 µS/cm), and pH (8.12±0.03) were measured daily.

At different exposure periods (24, 48, 72 and 96 hrs),

the mortality of the fish was subjected to probit analysis using Minitab software (Minitab®16.2.4) to calculate the LC values, their slope functions, and confidence limits.

### Results

The percent mortality of the exposed *A. dispar* to

various concentrations of the plant extract of *E. turcomanica* for 24, 48, 72 and 96 hrs are depicted in Figure 1. The LC<sub>50</sub> values at various exposure periods were 0.177 g/L for 24 hrs, 0.131 g/L for 48 hrs, 0.073 g/L for 72 hrs, and 0.052 g/L for 96 hrs. The LC values, their upper and lower confidence limits, and slope functions are given in Table 1.

The toxicity of dried powder of *E. turcomanica* was found to be time and dose-dependent ( $P < 0.05$ ). The regression coefficient demonstrated a significant positive correlation ( $P < 0.05$ ) between mortality percent and concentration of *E. turcomanica*. Also a significant negative correlation ( $P < 0.05$ ) was found between the exposure time and different LC values.

### Discussion

The results revealed that the dried powder of *E. turcomanica* has a high piscicidal activity causing more mortality with increasing its concentration. The toxicity was both time and dose dependent. A significant negative correlation demonstrated between different LC values and exposure time. LC<sub>50</sub> value of *E. turcomanica* decrease with increasing exposure time in *A. dispar* from 0.177 g/L after 24 hrs to 0.052 g/L after 96 hrs.

To my best knowledge, there is no report on the toxicity effects of *E. turcomanica* on aquatic animals. Although, some reports are present on the toxicity effects of the aqueous and latex extracts of other members of the family Euphorbiaceae on some animals, including fish and molluscs (Singh and Singh, 2002; Singh and Singh, 2005; Tiwari and Singh, 2006; dos Santos et al., 2007; Oliveira-Filho et al., 2010).

The LC<sub>50</sub> value for 24 hrs of dried powder of *E. tirucalli* stem bark latex for *Colisa fasciatus* was 8.14 mg/L, whereas this value for *Channa punctatus* was 9.01 mg/L (Tiwari et al., 2003). In another experiment, the toxicity of four plants belonging to members of Euphorbiaceae and Apocynaceae on *C. punctatus* evaluated (Singh and Singh, 2005). In this study, 96 hrs LC<sub>50</sub> values of *E. royleana*, *Nerium indicum*, *Jatropha gossypifolia*, and *Thevetia peruviana* were 0.020, 0.041, 4.34, and 3.17 g/L,

respectively (Singh and Singh, 2005). This value for *E. turcomanica* was 0.052 g/L on *A. dispar*. In general, juicy and latex-bearing Euphorbiaceae are more potent in their toxic effects than rotenone-yielding plants (Neuwinger, 2004). In the present study, the dried powder of whole plant is used directly and nevertheless, the LC<sub>50</sub> values of *E. turcomanica* is comparable in potent toxicity to the other studies. Also, it should be noted that the experimental fish in this study is very hardy comparing to other studies. Therefore, it is suggested that *E. turcomanica* has more toxicity than other species such as *J. gossypifolia* and *T. peruviana*.

It is concluded that dried powder of *E. turcomanica* has a potent piscicidal activity on *A. dispar*. Therefore, enough precautions must be exercised when derivatives of *E. turcomanica* is being used near fish-inhabiting water reservoirs.

### References

- Abdel-Hamid H. (2003). Molluscicidal and in-vitro schistosomicidal activities of the latex and some extracts of some plants belonging to Euphorbiaceae. Journal of the Egyptian Society of Parasitology, 33: 947-954.
- Abdel-Fattah M.R. (1987). The chemical constituents and economic plants of the Euphorbiaceae. Botanical Journal of the Linnean Society, 94: 293-326.
- Bani S., Suri K., Suri O., Sharma O. (1997). Analgesic and antipyretic properties of *Euphorbia royleana* latex. Phytotherapy Research, 11: 597-599.
- Clesceri L.S., Greenberg A.E., Eaton A.D. (1998). Standard methods for the examination of water and wastewater 20th ed. American Public Health Association. Washington DC, 2671 p.
- dos Santos A.F., de Azevedo D.P.L., dos Santos Mata R.d.C., de Mendonça D.I.M.D., Sant'Ana A.E.G. (2007). The lethality of *Euphorbia conspicua* to adults of *Biomphalaria glabrata*, cercaria of *Schistosoma mansoni* and larvae of *Artemia salina*. Bioresource Technology, 98: 135-139.
- Ebenso I.E. (2004). Molluscicidal effects of neem (*Azadirachta indica*) extracts on edible tropical land snails. Pest Management Science, 60: 178-182.
- Frodin D.G. (2004). History and concepts of big plant genera. Taxon, 53: 753-776.

- Govaerts R., Frodin D.G., Radcliffe-Smith A., Carter S. (2000). World checklist and bibliography of Euphorbiaceae (with Pandaceae). Royal Botanic Gardens, Kew. 1661 p.
- Hassan A.A., Mahmoud A.E., Hassan R.A., Husein E. (2011). Evaluation of *Euphorbia aphylla*, *Ziziphus spina-christi* and *Enterolobium contortisiliquum* as Molluscicidal Agents. Journal of American Science, 7: 511-520.
- Horn J.W., van Ee B.W., Morawetz J.J., Riina R., Steinmann V.W., Berry P.E., Wurdack K.J. (2012). Phylogenetics and the evolution of major structural characters in the giant genus *Euphorbia* L. (Euphorbiaceae). Molecular Phylogenetics and Evolution, 63: 305-326.
- Kumar S., Malhotra R., Kumar D. (2010). *Euphorbia hirta*: Its chemistry, traditional and medicinal uses, and pharmacological activities. Pharmacognosy Reviews, 4: 58-61.
- Marston A., Hostettmann K. (1985). Review article number 6: Plant molluscicides. Phytochemistry, 24: 639-652.
- Mozaffarian V. (1996). A Dictionary of Iranian plant names. Farhang Moaser Publishers. Tehran, Iran. (In Farsi)
- Neuwinger H.D. (2004). Plants used for poison fishing in tropical Africa. Toxicon, 44: 417-430.
- Oliveira-Filho E.C., Geraldino B.R., Coelho D.R., De-Carvalho R.R., Paumgarten F.J. (2010). Comparative toxicity of *Euphorbia milii* latex and synthetic molluscicides to *Biomphalaria glabrata* embryos. Chemosphere, 81: 218-227.
- Oliveira-Filho E.C., Paumgarten F.J. (2000). Toxicity of *Euphorbia milii* latex and niclosamide to snails and nontarget aquatic species. Ecotoxicology and Environmental Safety, 46: 342-350.
- Prasad M., Kumar A., Mishra D., Srivastav S.K., Srivastav A.K. (2010). Acute toxicity of *Euphorbia royleana* boiss (euphorbiaceae) latex on fresh water catfish, *Heteropneustes fossilis* (siluriformes, heteropneustidae). Acta Toxicológica Argentina, 18: 5-9.
- Reidinger J., Russell F. (1976). Organochlorine residues in adults of six southwestern bat species. Journal of Wildlife Management, 40: 677-680.
- Sastry K., Siddiqui A. (1983). Metabolic changes in the snake head fish *Channa punctatus* chronically exposed to endosulfan. Water, Air, and Soil Pollution, 19: 133-141.
- Singh D., Singh A. (2002). Piscicidal effect of some common plants of India commonly used in freshwater bodies against target animals. Chemosphere, 49: 45-49.
- Singh D., Singh A. (2005). The toxicity of four native Indian plants: Effect on AChE and acid/alkaline phosphatase level in fish *Channa marulius*. Chemosphere, 60: 135-140.
- Singh K., Singh A., Singh D.K. (1996). Molluscicidal activity of neem (*Azadirachta indica* A.Juss). Journal of Ethnopharmacology, 52: 35-40.
- Singh S.K., Yadav R.P., Tiwari S., Singh A. (2005). Toxic effect of stem bark and leaf of *Euphorbia hirta* plant against freshwater vector snail *Lymnaea acuminata*. Chemosphere, 59: 263-270.
- Srivastava V., Singh S.K., Rai M., Singh A. (2004). Toxicity of *Nerium indicum* and *Euphorbia royleana* lattices against *Culex quinquefasciatus* mosquito larvae. Nigerian Journal of Natural Products and Medicine, 7: 61-64.
- Tiwari S., Singh A. (2004). Piscicidal and anti-acetylcholinesterase activity of *Euphorbia royleana* stem bark extracts against freshwater common predatory fish *Channa punctatus*. Environmental Toxicology and Pharmacology, 18: 47-53.
- Tiwari S., Singh A. (2006). Biochemical stress response in freshwater fish *Channa punctatus* induced by aqueous extracts of *Euphorbia tirucalli* plant. Chemosphere, 64: 36-42.
- Tiwari S., Singh P., Singh A. (2003). Toxicity of *Euphorbia tirucalli* plant against freshwater target and non-target organisms. Pakistan Journal of Biological Sciences, 6: 1423-1429.
- Tiwari S., Singh S., Singh A. (2004). Toxicological effect and biochemical alterations induced by different fractions of *Euphorbia royleana* latex in freshwater harmful vector snail *Lymnaea acuminata*. Indian Journal of Experimental Biology, 42: 1220-1225.
- Vasas A., Hohmann J. (2014). *Euphorbia Diterpenes*: isolation, structure, biological activity, and synthesis (2008–2012). Chemical Reviews, 114: 8579-8612.

## تعیین غلظت های کشنده گیاه فرفیون ترکمنی (*Euphorbia turcomanica*) بر روی ماهی آفانیوس گورخری (*Aphanius dispar*)

هما زارع<sup>۱</sup>، احمد نوری<sup>۱\*</sup>، مرتضی یوسف زادی<sup>۲</sup>، مهدی بنایی<sup>۳</sup>

<sup>۱</sup>گروه شیلات، دانشکده علوم و فنون دریایی، دانشگاه هرمزگان، بندرعباس، ایران.

<sup>۲</sup>گروه زیست دریا، دانشکده علوم و فنون دریایی، دانشگاه هرمزگان، بندرعباس، ایران.

<sup>۳</sup>گروه شیلات، دانشکده محیط زیست و منابع طبیعی، دانشگاه خاتم الانبیاء، بهبهان، ایران.

### چکیده:

تاکنون تاثیر عصاره های آبی حاصل از بسیاری از گونه های متعلق به خانواده Euphorbiaceae به عنوان سم کشنده ماهی و نرم تنان مورد مطالعه قرار گرفته است، با این وجود اطلاعاتی در مورد توان سمیت و کشندگی گیاه فرفیون ترکمنی *E. turcomanica* بر روی موجودات آبی در دسترس نمی باشد. آزمون تعیین غلظت های کشنده این گیاه با استفاده از روش آزمایش چهار روزه در محیط محصور به منظور تعیین غلظت های LC<sub>50</sub> پودر خشک شده گیاه *E. turcomanica* بر روی ماهی آفانیوس گورخری *A. dispar* یک گونه با دامنه تحمل بالای شوری، به اجرا درآمد. مقادیر LC<sub>50</sub> در زمان های مختلف در معرض قرارگیری به ترتیب  $0.177 \pm 0.039$  گرم بر لیتر برای ۲۴ ساعت،  $0.131 \pm 0.030$  گرم بر لیتر برای ۴۸ ساعت،  $0.073 \pm 0.018$  گرم بر لیتر برای ۷۲ ساعت و  $0.052 \pm 0.013$  گرم بر لیتر برای ۹۶ ساعت به دست آمد. سمیت پودر خشک شده گیاه *E. turcomanica* همبستگی مثبتی را بین تلفات ماهی و مدت زمان تاثیرگذاری نشان داد. با توجه به اینکه نتایج این تحقیق اولین گزارش در مورد سمیت گیاه *E. turcomanica* بر روی ماهی *A. dispar* می باشد، این نتایج تنها قابل مقایسه با نتایج سایر گونه های Euphorbiaceae بر روی ماهیان دیگر است. از نتایج حاصل می توان استنباط نمود که توان سمیت گیاه *E. turcomanica* قابل مقایسه و نزدیک به سایر گونه های Euphorbiaceae است که تاکنون مورد بررسی قرار گرفته اند. همچنین پیشنهاد می شود که ترکیبات تولیدی از گیاه *E. turcomanica* با توجه به سمیت بالای آن به طور مستقیم در محیط های آبی که محل زیست ماهیان مختلف است، استفاده نگردد.

کلمات کلیدی: فرفیون ترکمنی، پودر خشک، سم کشنده ماهی، ماهی گورخری.