



Chemical and morphological characterization of *Allium tuncelianum* (Amaryllidaceae) and its antioxidant and anticholinesterase potentials

Songul KARAKAYA^{1,*}, Gulnur EKSI², Mehmet KOCA³, Betul DEMIRCI⁴,
Haluk Caglar KAYMAK⁵, Mehmet Emin KAPLAN⁶ & Ozkan AKSAKAL⁷

^{1,6}Department of Pharmacognosy, Faculty of Pharmacy, Ataturk University, Erzurum, Turkey.

²Department of Pharmaceutical Botany, Faculty of Pharmacy, Ankara University, Ankara, Turkey.

³Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Ataturk University, Erzurum, Turkey.

⁴Department of Pharmacognosy, Faculty of Pharmacy, Anadolu University, Eskisehir, Turkey.

⁵Department of Horticulture, Faculty of Agriculture Ataturk University, Erzurum, Turkey.

⁷Department of Biology, Faculty of Science, Ataturk University, Erzurum, Turkey.

*Corresponding author: ecz-songul@hotmail.com, <https://orcid.org/0000-0002-3268-721X>

²gulnur_eksi@yahoo.com, <https://orcid.org/0000-0002-6869-8703>

³kocamehmet@atauni.edu.tr, <https://orcid.org/0000-0002-1517-5925>

⁴betuldemirci@gmail.com, <https://orcid.org/0000-0003-2343-746X>

⁵hckaymak@atauni.edu.tr, <https://orcid.org/0000-0002-0836-7654>

⁶ecz.eminkaplan@gmail.com, <https://orcid.org/0000-0001-6833-4451>

⁷ozkanaksakal@atauni.edu.tr, <https://orcid.org/0000-0003-0760-7502>

Abstract. Alzheimer's disease is the main reason for dementia, which increases with age. Cholinesterase inhibition and antioxidant potentials of extracts and essential oils from bulbs of *A. tuncelianum* (Kollmann) Özhatay & al., an endemic species to Tunceli (eastern Turkey), were evaluated. The fraction extracted of ethyl acetate had the highest phenolics level, 1,1-diphenyl-2-picrylhydrazyl, and thiobarbituric acid antioxidant capacity. Also, the ethyl acetate fraction presented the highest acetylcholinesterase (15.98 ± 2.76%), and butyrylcholinesterase inhibition (47.33 ± 3.27%). Diallyl disulfide (49.8%), diallyl trisulfide (27.9%) and allyl methyl trisulfide (6.9%) were found to be the major components of essential oil. This paper shows that the ethyl acetate fraction of *A. tuncelianum* could be a potent source of antioxidant and anticholinesterase components.

Keywords. *Allium tuncelianum*, anticholinesterase, endemic, essential oil, morphology.

Resumen. La enfermedad de Alzheimer es la causa principal de la demencia, cuya aparición aumenta según la edad. Se evaluaron la inhibición de la colinesterasa y el potencial antioxidante de los extractos y los aceites esenciales de los bulbos de *A. tuncelianum* (Kollmann) Özhatay & al., una especie endémica de Tunceli (este de Turquía). La fracción extraída de acetato de etilo presentó los niveles más altos de fenoles, 1,1-difenil-2-picrilhidrazilo y capacidad antioxidante, ácido tiobarbitúrico. Asimismo, la fracción de etil acetato presentó la mayor capacidad de inhibición de acetilcolinesterasa (15.98 ± 2.76%) y butirilcolinesterasa (47.33 ± 3.27%). El disulfuro de dialilo (49.8%), el trisulfuro de dialilo (27.9%) y el trisulfuro de metil alilo (6.9%) fueron los componentes principales del aceite esencial. Este artículo muestra que la fracción de etil acetato de *A. tuncelianum* podría ser una fuente potencial novedosa de componentes antioxidantes y anticolinesterasa.

Palabras clave. Aceite esencial, *Allium tuncelianum*, anticolinesterasa, endémico, morfología.

How to cite this article: Karakaya S., Eksi G., Koca M., Demirci B., Kaymak H.C., Kaplan M.E. & Aksakal O. 2019. Chemical and morphological characterization of *Allium tuncelianum* (Amaryllidaceae) and its antioxidant and anticholinesterase potentials. *Anales del Jardín Botánico de Madrid* 76 (2): e085. <https://doi.org/10.3989/ajbm.2523>

Title in Spanish: Caracterización química y morfológica de *Allium tuncelianum* (Amaryllidaceae) y potenciales antioxidantes y anticolinesterasa.

Received: 25–III–2019; accepted: 28–V–2019; published on-line: 28–X–2019; Associate Editors: M.C. Sánchez Mata & R. Morales.

INTRODUCTION

Alzheimer's disease (AD) is a neurodegenerative disease characterized by an accumulation of extracellular amyloid-beta peptide (A β) and intracellular neurofibrils resulting in a loss of memory. A β is the principal constituent of senile plaques that are thought to play a central role in the healing and progression of oligomer and fibril forms of AD. Besides, many studies have shown that oxidative stress

and mitochondrial dysfunction might have a substantial role in AD and that they are suppressed or reduced by using antioxidant agents, suggesting a therapeutic intervention for AD patients. It is reported that numerous antioxidant components protect the brain from A β neurotoxicity (Sgarbossa & al. 2015). Pharmacological treatments available for AD occur by relaxation of the symptoms rather than targeting the etiological mechanisms (Ng & al. 2015).

Recent investigations have indicated that numerous medicinal herbs are a significant source of antioxidants. Plants include a large number of free radical scavenging molecules, such as alkaloids, tannins, steroids, rotenoids, carotenoids, dietary glutathione, anthocyanins, saponins, terpenoids, and flavonoids. Consequently, medical herbs become popular as a cheap source to discover novel antioxidants (Chand & al. 2018).

Volatile oils are found to have several pharmacological activities such as hepatoprotective, carminative, antispasmodic, antiviral, and anti-tumor, etc. Nowadays, many volatile oils have been characterized as natural antioxidants and proposed as potential sources for food preservation. Additionally, biologically active natural compounds are of interest to the pharmaceuticals industry for the control of human ailments of microbial origin and for the containment of lipid peroxidative damage, which has been implicated in specific pathological diseases such as AD, ischemia-reperfusion injury, cancer, coronary, atherosclerosis, and aging (Mimica & al. 2004).

The representatives of family *Amaryllidaceae* J.St.-Hil. have been characterized by high phenolics content (Resetár & al. 2017) and some members of this family have been used in the treatment of AD. Galanthamine is an alkaloid of the *Amaryllidaceae* that is a competitive selective, long-acting, and reversible acetylcholinesterase inhibitor that maintains the beneficial effects even after treatment (López & al. 2002).

Garlic has been produced and eaten worldwide and has attracted attention due to its preservative potentials against various diseases. Previous investigations indicated that garlic has numerous biological and pharmacological effective compounds. It is used for medical aims since ancient times, and its usage for cancer treatment dates back to 3,500 years ago (Özkan & al. 2013).

The genus *Allium* L. (*Amaryllidaceae*; cf. APGIII 2009) consists of more than 900 species which naturally grow in the northern hemisphere (Duman & al. 2017; Ekşi & al. 2016). According to phylogenetic analyses, *A. tuncelianum* (Kollmann) Özhatay, B.Mathew & Şiraneci belongs to the *Allium* sect. *Allium*. Fritsch & Friesen (2002) have proposed that the wild progenitor of garlic species should grow in the district from South Central Asia to the Mediterranean, according to their taxonomic investigations. Mathew (1996) proposed that *A. tuncelianum* might be the wild progenitor of *A. sativum* L. and both share general characters such as the odor of leaves and bulbs.

Because of its similarity to the widespread garlic, it is locally named as 'Ovacik garlic' or 'Tunceli garlic' in the district. *Allium tuncelianum* generally consists of a single cloved white bulb, unlike garlic, which has bulbs with multiple cloves (Kıralan & al. 2013). Even though

A. tuncelianum has been considered as a close relative of *A. sativum*, the precise phylogenetic or genetic relationships are not well known yet.

In this regard, the aim of this paper was to report the cholinesterase inhibitory and antioxidant activity of the methanol, hexane, dichloromethane, ethyl acetate, butanol, and aqueous extracts and essential oils of bulbs of *A. tuncelianum*. The total phenolic content of extracts and essential oil, as well as the essential oil composition and morphology of *A. tuncelianum*, were assessed.

MATERIAL AND METHODS

Plant specimen

Allium tuncelianum was gathered from Ovacık, Tunceli province (Eastern Turkey). The voucher specimens have been preserved at the Herbarium of Ataturk University, Faculty of Science under the code ATA-9877.

Extraction

Dried bulbs of *A. tuncelianum* (50 g) were crushed and macerated with methanol (3 times, 8 h) in a water-bath not exceeding 35°C (3 × 100 ml) by 200 rpm with using of a mechanical mixer. Combined bulbs extracts were filtered and concentrated by rotary evaporator to dryness, then dissolved in methanol: water (1 : 9) and fractionated three times with 150 ml of n-hexane, dichloromethane, ethyl acetate, and n-butanol, respectively.

On the other hand, 50 g of bulbs of *A. tuncelianum* were crushed and macerated with 200 ml of distilled water for 8 h/3 days at 30 to 35°C. The aqueous extract was filtered, frozen and lyophilized to attain aqueous extracts of bulbs. Amounts of the powdered parts of *A. tuncelianum* and acquired extracts/fractions are shown in Table 1.

Isolation of the essential oil, GC-FID and GC/MS analyses

The essential oil isolation, GC-FID, and GC/MS assays were done in accordance with Karakaya & al. (2016). The crushed part, the percentage of essential oil, and the color of the essential oil are displayed in Table 2.

Determination of total phenolic content

The total phenolic content of the samples was done in accordance with Karakaya & al. (2018). The procedure was repeated three times for each sample.

Antioxidant activity

The quantitative 1,1-diphenyl-2-picrylhydrazyl (DPPH) of the samples was done in Karakaya & al. (2018). The IC₅₀ values of samples were established by linear regression analysis in triplicate.

Table 1. Amounts of the powdered parts of *A. tuncelianum* (Kollmann) Özhatay & al. and acquired extracts/fractions.

| Extracts/Fractions (g) | Aerial part |
|---------------------------------|-------------|
| MeOH | 14.21 |
| Hexane | 2.08 |
| CH ₂ Cl ₂ | 5.11 |
| EtOAc | 1.03 |
| BuOH | 3.23 |
| Methanolic residue | 2.96 |
| Lyophilised aqueous extract | 15.44 |

Table 2. The crushed part, essential oil % yield *A. tuncelianum* (Kollmann) Özhatay & al. and color of essential oil (w/v, %).

| Part | Crushed | Yield | Colour | Collection |
|-------|---------|--------|--------|------------|
| bulbs | 220 g | 0.0046 | White | 2018 |

Table 3. Total phenolic contents of the extracts, fractions and essential oil from *A. tuncelianum* (Kollmann) Özhatay & al. [The data present the mean \pm SD of three independent experiments ($p < 0.05$)].

| Tested samples | Total phenolic contents (mg/g) \pm SD |
|---------------------------------|---|
| MeOH | 377.25 \pm 4.78 |
| Hexane | 67.78 \pm 1.56 |
| CH ₂ Cl ₂ | 434.61 \pm 4.12 |
| EtOAc | 666.45 \pm 3.21 |
| BuOH | 124.05 \pm 2.44 |
| Methanolic residue | 338.14 \pm 5.34 |
| Lyophilised aqueous extract | 567.20 \pm 3.07 |
| Essential oil | 209.01 \pm 2.13 |

Anti-lipid peroxidation activity

The anti-lipid peroxidation activity of the samples was done according to Karakaya & al. (2018). The IC₅₀ values were established through linear regression assay.

Evaluation of AChE and BuChE inhibition activities

The evaluation of AChE and BuChE inhibition followed Karakaya & al. (2018). The procedure was repeated three times for each plate. All data were denoted as mean \pm SE of three independent tests.

Statistical analysis

Overall indications are denoted as mean \pm SE and statistically analyzed through ANOVA one-way analysis followed by way of complementary analysis of Bonferroni ($P < 0.05$), planned to determine statistical significance.

RESULTS

General morphology

Allium tuncelianum (Kollmann) Özhatay, B.Mathew & Şiraneci, Kew Bull. 50 (4): 723 (Özhatay & Mathew 1995). *A. macrochetum* subsp. *tuncelianum* Kollmann, Notes Roy. Bot. Gard. Edinburgh 41: 262 (Kollmann & al. 1983), basion. Tipo: [Turkey] Tunceli, Munzur Da., Aksu Dere above Ovacik, 1800 m a.s.l., 21 Jul. 1957, Davis 31498 leg. (holo-: E!; iso-: K!). Figs. 1 and 2.

Bulb 1.5–5.5 cm in diameter, ovoid; outer tunics thick, membranous, brownish to dirty yellowish-white; inner tunics thin, membranous, white; bulblets 1–2. Scape 50–150 cm. Leaves 4–8, 1–2.5 cm wide, flat, canaliculate, glabrous, shorter than scape. Spathe 10–20 cm, 1-valved, deciduous. Umbel 2–8 cm in diameter, spherical, dense (100–200 flowers), bracteolate. Perigone 2.5–3.5 mm, campanulate, pale pinkish to white, smooth. Stamens longer than perigone. Capsule 3–4 mm. Chromosome number (2n) 16.

Distribution and habitat.—East Anatolia; rocky areas, calcareous soils; 1000–2200 m a.s.l.

Phenology.—Flowering time from June to August.

Experimental section

The methanol extracts of bulbs of *A. tuncelianum* were fragmented using solvents with different polarities (C₆H₁₄, CH₂Cl₂, C₄H₈O₂, and C₄H₁₀O). Also, the lyophilised aqueous extract of bulbs was obtained. The extracts, fractions and essential oil were assessed for antioxidant and cholinesterase inhibitory activities. Also, the morphological study of *A. tuncelianum* was also evaluated.

The essential oils, extracts, and fractions of bulbs were estimated with regard to antioxidant capacity effect. The data of samples with regard to total phenolics content are displayed in Table 3. The fraction extracted of ethyl acetate (EtOAc) had the highest level of total phenolic (666.45 mg GAE g⁻¹ DW) however the hexane fraction got the lowest phenolic content levels (67.78 mg GAE g⁻¹ DW). DPPH analysis data were presented in Table 4 and the EtOAc fraction got the highest antioxidant activity (17.21 \pm 4.33 μ g/ml) and the hexane fraction had the lowest phenolic content (128.45 \pm 3.56 μ g/ml). The findings of the analysis of thiobarbituric acid (TBA) were exhibited in Table 5 as IC₅₀ (μ g/ml). The EtOAc fraction and lyophilized aqueous extract had the highest antioxidant potential (IC₅₀ = 54.67 and 65.15 μ g/ml, respectively) in TBA analysis.

Cholinesterase inhibitory activity of samples was revealed via colorimetric Ellman's method (Ellman & al. 1961), some changes were done following the mentioned method and donepezil was used as a standard (Yerdelen & Tosun 2015). In vitro cholinesterase inhibitory activity of samples at 100 μ g/ml is displayed in Table 6. The EtOAc



Fig. 1. *Allium tuncelianum* (Kollmann) Özhatay, B.Mathew & Şiraneci: **a, b**, plant during the anthesis; **c**, plant during the pre-anthesis [AEF 24116; illustrated by Gülnur Ekşi].

Table 4. DPPH radical scavenging activity of the the extracts, fractions and essential oil from *A. tuncelianum* (Kollmann) Özhatay & al. ($\mu\text{g/ml}$) [The data present the mean \pm SD of three independent experiments ($p < 0.05$)].

| Tested samples | IC ₅₀ values ($\mu\text{g/ml}$) \pm SD |
|---------------------------------|---|
| MeOH | 87.35 \pm 3.42 |
| Hexane | 105.41 \pm 3.34 |
| CH ₂ Cl ₂ | 31.43 \pm 2.56 |
| EtOAc | 17.21 \pm 4.33 |
| BuOH | 93.22 \pm 1.66 |
| Methanolic residue | 128.45 \pm 3.56 |
| Lyophilised aqueous extract | 52.57 \pm 2.68 |
| Essential oil | 55.09 \pm 3.22 |
| Chlorogenic acid | 2.41 \pm 0.58 |
| Propyl gallate | 0.005 \pm 0.21 |
| Rutin | 3.05 \pm 0.89 |

Table 5. Antioxidant activities of the the the extracts, fractions and essential oil from *A. tuncelianum* (Kollmann) Özhatay & al. in TBA test [The data present the mean \pm SD of four independent experiments ($p < 0.05$)].

| Tested samples | IC ₅₀ values ($\mu\text{g/ml}$) \pm SD |
|---------------------------------|---|
| MeOH | 451.61 \pm 2.98 |
| Hexane | >500 |
| CH ₂ Cl ₂ | 76.67 \pm 3.77 |
| EtOAc | 54.67 \pm 4.24 |
| BuOH | 166.29 \pm 2.77 |
| Methanolic residue | >500 |
| Lyophilised aqueous extract | 65.15 \pm 1.79 |
| Essential oil | 89.25 \pm 2.78 |
| Chlorogenic acid | 12.98 \pm 4.89 |
| Propyl gallate | 3.44 \pm 2.05 |
| Rutin | 9.65 \pm 3.09 |

and essential oil indicated remarkable inhibition against BuChE (47.33 \pm 3.27 and 28.65 \pm 2.58%, respectively) at 100 $\mu\text{g/ml}$. Also, EtOAc and CH₂Cl₂ fractions displayed inhibition against AChE (15.98 \pm 2.76 and 14.12 \pm 2.76%, respectively) at 100 $\mu\text{g/ml}$. On the other side, the methanolic residue fraction had no activity against both enzymes. Moreover, MeOH, lyophilized aqueous extracts and hexane, BuOH fractions had no activity against AChE. EtOAc fraction has been characterized by substantially higher total phenolic content than other samples.

The percentage yield of essential oil of *A. tuncelianum* and color of essential oil are presented in Table 2. The color of the essential oil was white. A total of five compounds making up 88.9% of the oil were defined in the bulbs of *A. tuncelianum*. Diallyl disulfide, diallyl trisulfide and allyl methyl trisulfide were the major components, amounting to 49.8%, 27.9 and 6.9%, respectively. Many of the defined

Table 6. In vitro AChE and BuChE inhibitory activities of samples from *A. tuncelianum* (Kollmann) Özhatay & al. at 100 $\mu\text{g/ml}$. [Superscript: a, standard error mean; b, no activity; c, not detected because of turbidity in the wells of microplates. The data present the mean \pm SD of three independent experiments ($p < 0.05$)

| Samples | Enzymes | Percentile of inhibition \pm S.E.M ^a against AChE and BuChE |
|---------------------------------|---------|--|
| MeOH | AChE | - ^b |
| | BuChE | 9.68 \pm 1.67 |
| Hexane | AChE | ND ^c |
| | BuChE | 2.56 \pm 2.55 |
| CH ₂ Cl ₂ | AChE | 14.12 \pm 2.76 |
| | BuChE | 23.56 \pm 2.65 |
| EtOAc | AChE | 15.98 \pm 2.76 |
| | BuChE | 47.33 \pm 3.27 |
| BuOH | AChE | - ^b |
| | BuChE | 3.98 \pm 2.56 |
| Methanolic residue | AChE | - ^b |
| | BuChE | - ^b |
| Lyophilised aqueous extract | AChE | ND ^c |
| | BuChE | 13.58 \pm 1.93 |
| Essential oils | AChE | 7.59 \pm 2.90 |
| | BuChE | 28.65 \pm 2.58 |
| Donepezil | AChE | 82.45 \pm 2.64 |
| | BuChE | 90.33 \pm 4.16 |

Table 7. The essential oil composition of *A. tuncelianum* (Kollmann) Özhatay & al. RRI Relative retention indices calculated against n-alkanes. % calculated from FID data.

| RRI | Compound | % |
|--------------|--------------------------|-------------|
| 1292 | Allyl methyl disulfide | 1.9 |
| 1438 | Allyl propenyl disulfide | 2.4 |
| 1492 | Diallyl disulfide | 49.8 |
| 1607 | Allyl methyl trisulfide | 6.9 |
| 1811 | Diallyl trisulfide | 27.9 |
| Total | | 88.9 |

compounds were sulphur compounds. The compositions of essential oil are presented in Table 7.

DISCUSSION

Previously, it was observed that MeOH extract of the bulb of *A. tuncelianum* showed high antioxidant activities with the DPPH method (51.1 \pm 5.5%) and greater content of total phenols (Yumrutaş & al. 2009). Moreover, another study about the antioxidant activity of *A. tuncelianum* showed that aqueous and EtOH extracts (Ağbaşı & al. 2013) and MeOH extract (Şehitoğlu & al. 2018) of bulbs got a significant effect. Other research also found a substantial correlation between antioxidant capacity and total phenolic content (Sytař & al. 2015; Granato & al. 2018) as well.

Previous investigations indicated that the major constituents of essential oils of *A. tuncelianum* bulbs were diallyl tri- sulfides (30.90%) and diallyl disulfide (28.30%) (Takim & al. 2016); diallyl disulfide (in green and red garlic was 67.33% and 72.52%, respectively) (Kiralan & al. 2013).

Nowadays it is known that diallyl disulfide derivatives are chemical agents that modulate various facets of AD (Manral & al. 2015). Also, previous studies showed that some *Allium* species such as *A. sativum* and *A. tuberosum* Rottler ex Spreng. had significant effects on AD (Chauhan 2003; Kim & al. 2007; Ray & al. 2011).

AD is a neurodegenerative disease induced by oxidative stress with a further cholinergic lack in the brain. Particularly, AD is characterised by a reduction in the sum of acetylcholine delivered from cholinergic synapses. A therapy methodology has been stimulated to enhance or maintain the proportion of acetylcholine through inhibiting acetylcholinesterase (Dickson 1997). Essential oils are a miscellaneous family of low molecular weight organic compounds with circumstantial biological activity. Compounds that act as cholinesterase inhibitors still are the only pharmacological therapeutics for AD. Many in vitro examinations showed that some components, in essential oils, may have cholinesterase inhibitory activity (Karakaya & al. 2019).

This paper showed that the EtOAc fraction of *A. tuncelianum* has cholinesterase inhibitory and antioxidant effects. The uses of antioxidants may be beneficial for AD

healing (Gibson & Huang 2005). To the literature surveys, this is the initial exploration of cholinesterase inhibitory activity of extracts, fractions and essential oil from *A. tuncelianum*.

Especially, EtOAc fraction of *A. tuncelianum* bulbs displayed a substantial cholinesterase inhibitory and antioxidant potentials. The studied essential oils, extracts and fractions exhibited radical scavenging capacity (RSC), which were detected to be in correlation to the content of phenolic compounds. The essential oil has been characterized by the presence of diallyl disulfide which was determined to have inhibition towards both cholinesterase inhibitory activities. This paper displays that the EtOAc fraction of *A. tuncelianum* could be a novel potency source of native antioxidant and anticholinesterase components.

ACKNOWLEDGEMENTS

This paper was supported by the Research Fund of Ataturk University (FHD-2019-7227).

REFERENCES

- Ağbaş B., Karakuş D., Adigüzel R., Keser S. & Demir E. 2013. Comparison of Total Antioxidant Properties and Dry Matter Content of Tunceli Garlic (*Allium tuncelianum*) and Normal Garlic (*Allium sativum*). *Bilim ve Gençlik Dergisi* 2: 50–62.
- APGIII 2009. An update of the Angiosperm Phylogeny Group classification for the orders and the families of the flowering plants. *Botanical journal of the Linnean Society* 161: 105–121. <https://doi.org/10.1111/j.1095-8339.2009.00996.x>




Fig. 2. *Allium tuncelianum* (Kollmann) Özhatay, B.Mathew & Şiraneci: **a, e, f, h**, anthesis; **b**, bulb; **c, d**, early stages of the anthesis; **g**, bulbs at a local market. [Photos: a, b, d-f, h, Gülnur Ekşi; c, g, Mehmet Koyuncu].

- Chand R.N., Gopalan R.D. & Christi K. 2018. Evaluation of antioxidant properties in thirteen Fijian medicinal plants used in Alzheimer's disease and related illness. *Free Radicals and Antioxidants* 8 (1): 11–17.
- Chauhan N.B. 2003. Anti-Amyloidogenic effect of *Allium sativum* in Alzheimer's transgenic model Tg2576. *Journal of Herbal Pharmacotherapy* 3: 95–107. https://doi.org/10.1080/J157v03n01_05
- Dickson D.W. 1997. Neuropathological diagnosis of Alzheimer's disease: A perspective from longitudinal clinicopathological studies. *Neurobiology of Aging* 18: 21–26. [https://doi.org/10.1016/S0197-4580\(97\)00065-1](https://doi.org/10.1016/S0197-4580(97)00065-1)
- Duman H., Ekşi G. & Özbek F. 2017. Two new species of *Allium* L. sect. *Allium* (*Amaryllidaceae*) from Turkey. *Plant Systematics and Evolution* 303: 1271–1291. <https://doi.org/10.1007/s00606-017-1437-4>
- Ekşi G., Koyuncu M. & Özkan A.M.G. 2016. *Allium ekimianum*: a new species (*Amaryllidaceae*) from Turkey. *PhytoKeys* 62: 83–93. <https://doi.org/10.3897/phytokeys.62.7796>
- Ellman G.L., Courtney K.D., Andresjr V. & Featherstone R.M. 1961. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology* 7: 88–89. [https://doi.org/10.1016/0006-2952\(61\)90145-9](https://doi.org/10.1016/0006-2952(61)90145-9)
- Fritsch R.M. & Friesen N. 2002. *Evolution, Domestication and Taxonomy*. CABI, Wallingford.
- Gibson G.E. & Huang H.M. 2005. Oxidative stress in Alzheimer's disease. *Neurobiology of Aging* 26: 575–578. <https://doi.org/10.1016/j.neurobiolaging.2004.07.017>
- Granato D., Shahidi F., Wrolstad R., Kilmartin P., Melton L.D., Hidalgo F.J., Miyashita K., Camp J., Alasalvar C., Ismail A.B., Elmore S., Birch G.G., Charalampopoulos D., Astley S.B., Pegg R., Zhou P. & Finglas P. 2018. Antioxidant activity, total phenolics and flavonoids contents: Should we ban in vitro screening methods? *Food Chemistry* 264: 471–475. <https://doi.org/10.1016/j.foodchem.2018.04.012>
- Karakaya S., Göger G., Kılıç C.S. & Demirci B. 2016. Composition of volatile oil of the aerial parts, flowers and roots of *Ferulago blancheana* Post. (*Apiaceae*) growing in Turkey and determination of their antimicrobial activities by bioautography method. *Turkish Journal Of Pharmaceutical Sciences* 13: 173–180.
- Karakaya S., Koca M., Kılıç C.S. & Coskun M. 2018. Antioxidant and anticholinesterase activities of *Ferulago syriaca* Boiss. and *F. isaurica* Peşmen growing in Turkey. *Medical Chemistry Research* 27: 1843–1850. <https://doi.org/10.1007/s00044-018-2196-7>
- Karakaya S., Koca M., Yılmaz S.V., Yıldırım K., Pınar N.M., Demirci B., Brestic M. & Sytar O. 2019. Molecular docking studies of coumarins isolated from extracts and essential oils of *Zosima absinthifolia* Link as potential inhibitors for Alzheimer's Disease. *Molecules* 24 (4), E722: 1–18. <https://doi.org/10.3390/molecules24040722>
- Kim M.J., Choi S., Kim H.K., Kim C.J., Hong B., Kim Y.J. & Shin D.H. 2007. Activation effects of *Allium Tuberosum* Rottl. on choline acetyltransferase. *Bioscience, Biotechnology, and Biochemistry* 71: 226–230. <https://doi.org/10.1271/bbb.60314>
- Kıralan M., Rahimi A., Arslan N. & Bayrak A. 2013. Volatiles in an endemic *Allium* species: *Allium tuncelianum* by headspace solid phase microextraction. *TEOP* 16: 417–420. <https://doi.org/10.1080/0972060X.2013.813283>
- Kollmann F., Özhatay N. & Koyuncu M. 1983. New *Allium* taxa from Turkey. *Notes from the Royal Botanic Garden, Edinburgh* 41 (2): 262.
- López S., Bastida J., Viladomat F. & Codina C. 2002. Acetylcholinesterase inhibitory activity of some *Amaryllidaceae* alkaloids and *Narcissus* extracts. *Life Sciences* 71: 2521–2529. [https://doi.org/10.1016/S0024-3205\(02\)02034-9](https://doi.org/10.1016/S0024-3205(02)02034-9)
- Manral A., Saini V., Meena P. & Tiwari M. 2015. Multifunctional novel diallyl disulfide (DADS) derivatives with b-amyloid-reducing, cholinergic, antioxidant and metal chelating properties for the treatment of Alzheimer's disease. *Bioorganic & Medicinal Chemistry* 23: 6389–6403. <https://doi.org/10.1016/j.bmc.2015.08.024>
- Mathew B. 1996. *A review of Allium sect. Allium*. Royal Botanic Gardens, Kew.
- Mimica N.D., Bozin B., Sokovic M. & Simin N. 2004. Antimicrobial and antioxidant activities of *Melissa officinalis* L. (*Lamiaceae*) essential oil. *Journal of Agricultural and Food Chemistry* 52: 2485–2489. <https://doi.org/10.1021/jf030698a>
- Ng Y.P., Or T.C.T. & Ip N.Y. 2015. Plant alkaloids as drug leads for Alzheimer's disease. *Neurochemistry International* 89: 260 e270. <https://doi.org/10.1016/j.neuint.2015.07.018>
- Özhatay N. & Mathew B. 1995. New taxa and notes on the genus *Allium* (*Alliaceae*) in Turkey and Arabia. *Kew Bulletin* 50 (4): 723–731. <https://doi.org/10.2307/4110233>
- Özkan O., Gül S., Kart A., Çiçek B.A. & Kılıç K. 2013. In vitro antimutagenicity of *Allium tuncelianum* ethanol extract against induction of chromosome aberration by mutagenic agent mitomycin C. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 19: 259–262. <https://doi.org/10.9775/kvfd.2012.7637>
- Ray B., Chauhan B.N. & Lahiri K.D. 2011. The “Aged Garlic Extract” (AGE) and one of its active ingredients S-Allyl-LCysteine (SAC) as potential preventive and therapeutic agents for Alzheimer's disease (AD). *Current Medicinal Chemistry* 18: 3306–3313. <https://doi.org/10.2174/092986711796504664>
- Resetar A., Freytag C., Kalydi F., Gonda S., M-Hamvas M., Ajtay K., Papp L. & Máthé C. 2017. Production and antioxidant capacity of tissue cultures from four *Amaryllidaceae* species. *Acta Societatis Botanicorum Poloniae* 86: 3525. <https://doi.org/10.5586/asbp.3525>
- Şehitoğlu M.H., Karakan F.Y., Kizilkaya B., Öztöpus R.Ö. & Gülçin İ. 2018. Investigation of antioxidant properties and bioactive composition of *Allium tuncelianum* ((Kollman) Ozhatay, Matthew & Siraneci) and *Allium sativum* L. *Journal of the Institute of Science and Technology* 8: 213–221. <https://doi.org/10.21597/jist.427293>
- Sgarbossa A., Giacomazza D. & Carlo M. 2015. Ferulic Acid: A Hope for Alzheimer's Disease Therapy from Plants. *Nutrients* 7: 5764–5782. <https://doi.org/10.3390/nu7075246>
- Sytar O., Bruckova K., Hunkova E., Zivcak M., Kiessoun K. & Brestic M. 2015. The application of multiplex fluorimetric sensor for analysis flavonoids content in the medical herbs family *Asteraceae*, *Lamiaceae*, *Rosaceae*. *Biological Research* 48 (5): 2–9. <https://doi.org/10.1186/0717-6287-48-5>
- Takim K., Kutlu T., İçen S., Demir N. 2016. Characterization of Tunceli mountain garlic essential oil component and determination effect on the antioxidant enzyme levels on rat renal tissue by dmba-induced. International Engineering, Science and Education Conference: 401–418.
- Yerdelen K.O. & Tosun E. 2015. Synthesis, docking and biological evaluation of oxamide and fumaramide analogs as potential AChE and BuChE inhibitors. *Medical Chemistry Research* 24: 588–602. <https://doi.org/10.1007/s00044-014-1152-4>
- Yumrutaş Ö., Demir S., Saygideğer Ö. & Doğan M. 2009. The in vitro antioxidant activity of *Allium tuncelianum*: An endemic. *Journal of Applied Biological Sciences* 3: 61–64.