

# Henry Ford Hospital Medical Journal

---

Volume 26 | Number 2

Article 11

---

6-1978

## Clinical Notes: Bacteremia Associated with Colonoscopy

A. Suarez

B. Schuman

E. Quinn

T. Neblett

Follow this and additional works at: <https://scholarlycommons.henryford.com/hfhmedjournal>

 Part of the [Life Sciences Commons](#), [Medical Specialties Commons](#), and the [Public Health Commons](#)

---

### Recommended Citation

Suarez, A.; Schuman, B.; Quinn, E.; and Neblett, T. (1978) "Clinical Notes: Bacteremia Associated with Colonoscopy," *Henry Ford Hospital Medical Journal* : Vol. 26 : No. 2 , 59-61.

Available at: <https://scholarlycommons.henryford.com/hfhmedjournal/vol26/iss2/11>

This Article is brought to you for free and open access by Henry Ford Health System Scholarly Commons. It has been accepted for inclusion in Henry Ford Hospital Medical Journal by an authorized editor of Henry Ford Health System Scholarly Commons.

## Clinical Notes

### Bacteremia Associated with Colonoscopy

A. Suarez, MD,\* B. Schuman, MD,\* E. Quinn, MD,\* and T. Neblett, MD\*\*

*Twenty-eight patients had a total of 168 blood cultures before and at regular intervals during colonoscopy. No bacteremia was found. Based on our results and other reports, we recommend antibiotic prophylaxis during colonoscopy only for those patients with prosthetic heart valves and for those with valvular heart disease who also have advanced liver disease.*

**T**RANSIENT bacteremia has been reported following several procedures involving trauma to mucosal surfaces, such as dental procedures,<sup>1</sup> nasotracheal intubation,<sup>2</sup> genitourinary tract manipulation,<sup>3</sup> sigmoidoscopy,<sup>4</sup> barium enema,<sup>5</sup> and upper gastrointestinal endoscopy.<sup>6</sup>

In contrast to other endoscopic procedures, colonoscopy involves manipulation of an extensive mucosal surface populated by many microorganisms, and it usually lasts longer than other endoscopic procedures. Thus, it would seem reasonable to anticipate the occurrence of bacteremia with colonoscopy.

#### Material and Methods

The study group was composed of patients who consented to donate samples of blood for culture while undergoing colonoscopy at Henry Ford Hospital for indications listed in the accompanying table. Patients were excluded if they had received antibiotics during the preceding week. The preparation consisted of cleansing enemas the night before and the morning of colonoscopy. Patients also received citrate of magnesium and/or bisacodyl tablets.

Blood for culture was obtained through a #19 butterfly needle placed and secured into a forearm vein. Prepuncture skin preparation was with povidone iodine. Specimens were drawn just before colonoscopy, at 5, 15, 30 and 45 minutes during the procedure, and every 15 minutes thereafter when the procedure lasted longer than 45 minutes. The first nine patients had blood for culture drawn 15 and 30 minutes after colonoscopy. Additional samples were also drawn at the time of biopsy or polypectomies.

In the first nine patients tested, the specimen was a 2 ml blood sample drawn into a heparin-rinsed glass syringe and immediately placed in anaerobic transport vials. Each sample was divided in two halves in the anaerobic chamber. One half was removed and processed aerobically; the other was left in the anaerobic chamber. In both cases the blood was filtered through a 0.45  $\mu$  millipore disc which was then placed on the surface of a reduced blood agar plate and

---

Submitted for publication: April 14, 1978  
Accepted for publication: May 18, 1978

\* Department of Medicine, Henry Ford Hospital  
\*\* Department of Pathology, Henry Ford Hospital

Address reprint requests to Dr. Suarez, Department of Medicine, Henry Ford Hospital, Detroit, MI 48202

incubated at 37°C. Plates were observed for growth daily for seven days.

In the remaining 19 patients, a 5 ml blood specimen was drawn into commercial blood culture bottles (supplemented peptone broth II\*) and incubated anaerobically. Previous, unpublished studies by T. Neblett indicated that this method was effective in detecting aerobic as well as anaerobic enteric organisms. Bottles were examined for turbidity daily, and gram stains were done after 18 hours, 7 and 14 days of incubation. After 7 days of incubation, each bottle was subcultured to two reduced blood agar plates, one of which was incubated aerobically in a CO<sub>2</sub> incubator. The other plate was incubated anaerobically in an environment containing 3% hydrogen, 12% carbon dioxide, and 85% nitrogen. Plates were observed for growth daily for seven days. At the end of 14 days, aliquots from all bottles were examined by gas chromatography.

Colonoscopy in 28 Patients	
Indication	Number
Polypectomy	10
Carcinoma suspected	6
Polyp suspected	4
Unexplained bleeding	4
Inflammatory Bowel Disease	4

### Results

A total of 168 samples were obtained from 28 patients who had colonoscopy. Only two cultures, one in each of two patients, were positive. In both cases *Propionibacterium* species was isolated.

### Discussion

*Propionibacteria* are frequent contaminants of blood cultures obtained through cutaneous puncture. It is more likely that in our patients these isolates represent contamination rather than true bacteremia.

The importance of transient bacteremia in the pathogenesis of bacterial endocarditis is widely recognized. The use of prophylactic antibiotics when patients with valvular or congenital heart disease undergo dental or genitourinary manipulation is generally recommended.<sup>7</sup> Whether colonoscopy would be an indication for prophylactic antibiotics in this circumstance prompted this study.

Several studies have been done to determine the frequency of bacteremia during colonoscopy.<sup>8-14</sup> Results have been

conflicting and, therefore, recommendations for antibiotic prophylaxis in patients at risk for infective endocarditis are also controversial. Our findings agree with those in most other studies; namely, that bacteremia is either nonexistent or very rare during colonoscopy. A possible explanation is the presence of the liver "filter" between the portal and systemic circulation. Indeed, as the report of Dickman et al suggests,<sup>10</sup> patients with severe liver impairment might be more susceptible to bacteremia during colonoscopy.

In view of the available evidence and since cases of endocarditis clearly related to colonoscopy have not been documented, we currently do not recommend prophylactic antibiotics for colonoscopy patients at risk for infective endocarditis. However, we concur with the American Heart Association recommendations<sup>7</sup> for administration of prophylactic antibiotics to patients with prosthetic valves, "since they appear to be at especially high risk." Patients who have advanced liver disease are also candidates for prophylaxis. Prophylaxis might also be indicated during colonoscopy in patients with strongly suspected or proven carcinoma of the colon because of the high carrier rate of *Streptococcus bovis* and the association of infective endocarditis due to this organism in such patients.<sup>15</sup>

### Acknowledgment

We want to thank the Michigan Heart Association, the McGraw Fund and Mr. Herbert Freedland, Howell Industries, Inc., for supporting this study. We also thank Mr. Donald Pohlod and Miss Carol Worek for valuable technical assistance.

\* Beckton Dickinson, Rutherford, NJ



### References

1. Cobe HM: Transitory bacteremia. *Oral Surg* 7:609-615, 1954.
2. Berry FA, Blakenbaker WL, and Ball CG: A comparison of bacteremia occurring with nasotracheal and orotracheal intubation. *Anesth Analg (Cleve)* 52:873-876, 1973.
3. Sullivan NM, Sutter VL, Carter WT, Attebery HR, and Finegold SM: Bacteremia after genitourinary tract manipulation: Bacteriological aspects and evaluation of various blood culture systems. *Appl Microbiol* 23:1101-1106, 1972.
4. LeFrock JL, Ellis CA, Turchik JB, and Weinstein L: Transient bacteremia associated with sigmoidoscopy. *N Engl J Med* 289:467-469, 1973.
5. LeFrock JL, Ellis CA, Klainer AS, and Weinstein L: Transient bacteremia associated with barium enema. *Arch Intern Med* 135:835-837, 1975.
6. Shull HJ, Greene BM, Allen SI, Dunn GD, and Schenkens S: Bacteremia with upper gastrointestinal endoscopy. *Ann Intern Med* 83:212-214, 1975.
7. Kaplan, EL, Anthony BF, Bisno A, Durack D, Houser H, Millard HD, Sanford J, Shulman ST, Stillerman M, Taranta A, and Wenger N: Prevention of bacterial endocarditis. *Circulation* 56:139A-143A, 1977.
8. Raforth RJ, Sorenson RM, and Bond JH: Bacteremia following colonoscopy. *Gastrointest Endosc* 22:32-33, 1975.
9. Craner GE and Ogburn RM: Letter to the editor. *Gastrointest Endosc* 22:105, 1975.
10. Dickman MD, Farrell R, Higgs RH, Wright LE, Humphries TJ, Wojcik JD, and Chappelka R: Colonoscopy associated bacteremia. *Surg Gynecol Obstet* 142:173-176, 1976.
11. Norfleet RG, Mulholland DD, Mitchell PD, Philo J, and Walters EW: Does bacteremia follow colonoscopy? *Gastroenterology* 70:20-21, 1976.
12. Norfleet RG, Mitchell PD, Mulholland DO, and Philo J: Does bacteremia follow colonoscopy? II. Results with blood cultures obtained 5, 10 and 15 minutes after colonoscopy. *Gastrointest Endosc* 23:31-32, 1976.
13. Pelican G, Hentges D, Butt J, Haag T, Rolfe R, and Hutcheson D: Bacteremia during colonoscopy. *Gastrointest Endosc* 23:33-35, 1976.
14. Liebermann TR: Bacteremia and fiberoptic endoscopy. *Gastrointest Endosc* 23:36-37, 1976.
15. Klein RS, Recco RA, Catalano MT, Edberg SC, Casey JI, and Steigbigel NH: Association of *Streptococcus bovis* with carcinoma of the colon. *N Engl J Med* 297:800-802, 1977.