

LE MATEMATICHE

Vol. LXXIV (2019) – Issue II, pp. 363–367

doi: 10.4418/2019.74.2.9

## A SHORT PROOF FOR A DETERMINANTAL FORMULA FOR GENERALIZED FIBONACCI NUMBERS

MILICA ANĐELIĆ - CARLOS M. DA FONSECA

The aim of this note is to provide a short and elegant proof for a recent determinantal formula for generalized Fibonacci numbers. The attractiveness of the proof presented here is its elementary nature.

### 1. Preliminaries

The study of sequences generated by the homogeneous linear second order difference equation with constant coefficients

$$u_{n+1} = au_n + bu_{n-1}, \quad \text{for } n \geq 1, \quad (1)$$

with certain initial conditions, goes back to the beginning of 1960s with the analysis of the algebraic properties of  $(u_n)$  [2, 7, 8]. Many relevant number sequences are obtained from (1), namely the Fibonacci numbers, setting  $a = b = u_1 = 1$  and  $u_0 = 0$ .

It is well-known that (1) can be represented by the determinant of the Jacobi matrix

$$T_n = \begin{pmatrix} a & -1 & & & \\ b & \ddots & \ddots & & \\ & \ddots & \ddots & -1 & \\ & & b & a & \end{pmatrix}_{n \times n} \quad (2)$$

---

Received on July 1, 2019

*AMS 2010 Subject Classification:* 15A18, 65F15, 15B05, 42C05

*Keywords:* Fibonacci numbers, Chebyshev polynomials of second kind, determinant

together with the specialisation of the initial conditions, namely,  $\det T_0 = 1$  and  $\det T_1 = a$ . From the well-established theory of orthogonal polynomials (see, e.g., the standard reference [3]), the determinant of  $T_n$  can be given by (cf. e.g. [6])

$$\det T_n = (-i\sqrt{b})^n U_n \left( \frac{ai}{2\sqrt{b}} \right),$$

where  $\{U_n(x)\}_{n \geq 0}$  are the Chebyshev polynomials of second kind, i.e., the orthogonal polynomials satisfying the three-term recurrence relations

$$U_{n+1}(x) = 2xU_n(x) - U_{n-1}(x), \quad \text{for all } n = 1, 2, \dots,$$

with initial conditions  $U_0(x) = 1$  and  $U_1(x) = 2x$ . The main explicit formula for the Chebyshev polynomials of second kind is

$$U_n(x) = \frac{\sin(n+1)\theta}{\sin \theta}, \quad \text{with } x = \cos \theta \quad (0 \leq \theta < \pi), \tag{3}$$

for all  $n = 0, 1, 2, \dots$ . While (3) is more common to find in the orthogonal polynomials theory, there are other explicit representations and relations for  $U_n(x)$ . Among them, the most frequent to find in number theory are

$$U_n(x) = \frac{(x + \sqrt{x^2 - 1})^{n+1} - (x - \sqrt{x^2 - 1})^{n+1}}{2\sqrt{x^2 - 1}},$$

an immediate consequence of de Moivre's formula, or

$$U_n(x) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k \binom{n-k}{k} (2x)^{n-2k}$$

which can be found for example in [1, (22.3.7)]. As stated in [5, p.187], many of them are paraphrases of trigonometric identities and derivations from (3). Nonetheless, here no explicit formula for  $U_n(x)$  is required for our aims.

Now, the Fibonacci numbers can be obtained directly from (cf. e.g. [4])

$$\det \begin{pmatrix} 1 & -1 & & & \\ 1 & \ddots & \ddots & & \\ & \ddots & \ddots & -1 & \\ & & & 1 & 1 \end{pmatrix}_{n \times n},$$

with  $a = b = 1$ ,  $u_0 = 0$  and  $u_1 = 1$  in (2). This means that the  $n$ th Fibonacci number  $F_n$  can be given by (cf. [2, 7])

$$F_n = (-i)^{n-1} U_{n-1} \left( \frac{i}{2} \right).$$

We observe that the determinant of a tridiagonal matrix is known in the literature as a *continuant* (cf. [10]). The terminology “tridiagonal determinant” is however inaccurate.

**2. A determinantal formula**

Recently in [9], Qi and Guo using intricate techniques proved that

$$u_n = \frac{1}{n!} \begin{vmatrix} \binom{1}{0}a & -1 & & & \\ 2\binom{2}{0}b & \binom{2}{1}a & -1 & & \\ & 2\binom{3}{1}b & \ddots & \ddots & \\ & & \ddots & \ddots & -1 \\ & & & 2\binom{n}{n-2}b & \binom{n}{n-1}a \end{vmatrix}. \tag{4}$$

Using the multilinearity of the determinant, our purpose here is to provide a simple proof for (4). Indeed,

$$\begin{aligned} & \begin{vmatrix} \binom{1}{0}a & -1 & & & \\ 2\binom{2}{0}b & \binom{2}{1}a & -1 & & \\ & 2\binom{3}{1}b & \ddots & \ddots & \\ & & \ddots & \ddots & -1 \\ & & & 2\binom{n}{n-2}b & \binom{n}{n-1}a \end{vmatrix} = \\ & = \begin{vmatrix} 1 \cdot a & -1 & & & \\ 2 \cdot 1 \cdot b & 2 \cdot a & -1 & & \\ & 3 \cdot 2 \cdot b & \ddots & \ddots & \\ & & \ddots & \ddots & -1 \\ & & & n(n-1) \cdot b & n \cdot a \end{vmatrix} \\ & = \begin{vmatrix} 1 \cdot a & -1 & & & \\ 2 \cdot b & 2 \cdot a & -2 & & \\ & 3 \cdot b & \ddots & \ddots & \\ & & \ddots & \ddots & -(n-1) \\ & & & n \cdot b & n \cdot a \end{vmatrix} \\ & = n! \begin{vmatrix} a & -1 & & & \\ b & \ddots & \ddots & & \\ & \ddots & \ddots & -1 & \\ & & b & a & \end{vmatrix}_{n \times n} \end{aligned}$$

$$= n!u_n,$$

for any positive integer  $n$ .

## REFERENCES

- [1] M. Abramowitz - I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, ninth edition, Dover Publications, Inc., New York, 1970.
- [2] R. G. Buschman, *Fibonacci numbers, Chebyshev polynomials generalizations and difference equations*, *Fibonacci Quart.* 1 (1963), 1–8, 19.
- [3] T. S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, New York, 1978.
- [4] L. Elsner - R. M. Redheffer, *Remarks on band matrices*, *Numer. Math.* 10 (1967), 153-161.
- [5] Erdélyi, et al., *Higher Transcendental Functions*, vol. 2, McGraw Hill, New York, 1953.
- [6] C. M. da Fonseca, *Unifying some Pell and Fibonacci identities*, *Appl. Math. Comput.* 236 (2014), 41–42.
- [7] A. F. Horadam, *Basic properties of a certain generalized sequence of numbers*, *Fibonacci Quart.* 3 (1965), 161–176.
- [8] A. F. Horadam, *A generalized Fibonacci sequence*, *Amer. Math. Monthly* 68 (1961), 455–459.
- [9] F. Qi - B.-N. Guo, *Expressing the generalized Fibonacci polynomials in terms of a tridiagonal determinant*, *Le Matematiche* 72(1) (2017), 167–175.
- [10] P. Rózsa, *On periodic continuants*, *Linear Algebra Appl.* 2 (1969), 267–274.

MILICA ANĐELIĆ

Department of Mathematics

Kuwait University

Safat 13060

Kuwait

e-mail: milica@sci.kuniv.edu.kw

*CARLOS M. DA FONSECA*

*Kuwait College of Science and Technology*

*Doha District, Block 4, P.O. Box 27235, Safat 13133*

*Kuwait*

*e-mail: c.dafonseca@kcst.edu.kw*

*University of Primorska*

*FAMNIT*

*Glagoljskaška 8, 6000 Koper*

*Slovenia*

*e-mail: carlos.dafonseca@famnit.upr.si*