



EFFECT OF 6 WEEKS AEROBIC EXERCISES TRAINING PROGRAM TO PHYSICAL FITNESS VARIABLES IN MIDDLE AGED WOMEN

Kiranⁱ

PhD Scholar,

Lakshmibai National Institute of Physical Education,
Gwalior, Madhya Pradesh, India

Abstract:

Purpose of the study was to see the effect of 6 weeks aerobics training program to physical fitness variables in middle aged women, the study was conducted 20 middle aged women, age ranging between 35 to 50 years from Gwalior, were selected as subjects of this study. To see the effect of 6 weeks aerobics training program to physical fitness variables in middle aged women, collected data was analyzed by using paired t-test at 0.05 level of significance. And significant difference was found between means of pre & post test in flexibility & grip strength but there is no significant difference found between means of pre and post test in cardio vascular endurance of middle aged women.

Keywords: aerobics training, middle aged women, physical fitness variables

1. Introduction

Women in the middle ages occupied a number of different social roles. During the middle ages, women held the positions of wife, mother, peasant, artisan, and nun. The very concept of "woman" changed in a number of ways during the middle Ages and several forces influenced women's roles during their period. Aerobic exercise gives a stronger heart, a leaner body, lower cholesterol, improved sleep. You can strengthen your entire cardiovascular system - heart, lungs and blood vessels - through regular aerobic activities. Aerobic exercise (also known as cardio) is physical exercise of low to high intensity that depends primarily on the aerobic energy-generating process. Aerobic literally means "relating to, involving, or requiring free oxygen", and refers to the use of oxygen to adequately meet energy demands during exercise via aerobic metabolism. Generally, light-to-moderate intensity activities that are sufficiently supported by aerobic metabolism can be performed for extended periods of time. Kenneth Cooper was the first person to introduce the concept of aerobic exercise. In the

ⁱ Correspondence: email kiran.chaudhary16@gmail.com

1960s, Cooper started research into preventive medicine. He became intrigued by the belief that exercise can preserve one's health. He sparked millions into becoming active and is now known as the "father of aerobics".

2. Objective of the Study

The purpose of the study was to find out the effect of 6 week aerobic training program to physical fitness variables in middle aged women.

3. Methodology

The study was to see the effect of 6 weeks aerobics training program to physical fitness variables in middle aged women. 20 middle aged women, age ranging between 35 to 50 years from Gwalior, were selected as subjects of this study. To see the effect of 6 weeks aerobics training program to physical fitness variables in middle aged women paired t-test were employed. Cardiovascular endurance were measured by the timing of 800 meter run and walk were recorded in minute, Grip strength was measured by the nearest kilogram of Grip strength (right and left) in Grip dynamometer were recorded & Flexibility measured by the sit and reach equipment (Lafayette) was used for the flexibility test, by measuring the nearest centimetre.

3.1 Statistical Methods

Paired t-test was applied to see the effect of 6 weeks aerobics training program to physical fitness variables in middle aged women. The hypothesis was tested at 0.05 level of significance.

3.1.1 Descriptive statistics for the data on cardiovascular endurance of middle aged women

Table 1: Descriptive statistics for the data
on cardiovascular endurance of middle aged women

		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	CE_PRE	7.109	20	1.623	.36291
	CE_POST	6.909	20	1.664	.37196

Table 1 shows that the mean score of pre and post test, the mean and SD of cardiovascular endurance is 7.109_1.623 & 6.909_1.664 respectively.

Table 2: Paired Samples Test

		Paired Differences					T	Df	Sig. (2-tailed)
		Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
					Lower	Upper			
Pair 1	CE_PRE CE_POST	-.20050	.51980	.11623	-.04277	.44377	1.725	19	.101

According to analysis of data presented in table 2, the calculated t-value (1.725) was found insignificant as its p-value is more than 0.05. The difference can be seen with the help of graphical representation of mean score. (Figure 1)

3.1.2 Descriptive statistics for the data on flexibility of middle aged women

Table 3: Paired Samples Statistics

		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	FLEXIBILITY_PRE	33.2750	20	6.16517	1.37857
	FLEXIBILITY_POST	35.1800	20	6.66038	1.48931

Table 3 Shows that the mean score of pre and post test, the mean and SD of Flexibility 33.2750_6.16517 and 35.1800_6.66038 respectively.

Table 4: Paired Samples Test

		Paired Differences					t	df	Sig. (2-tailed)
		Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
					Lower	Upper			
Pair 1	FLEXIBILITY_PRE FLEXIBILITY_POST	-1.90500	3.54423	.79251	-3.56375	-.24625	-2.404	19	.027

According to analysis of data presented in table 4, the calculated t-value (-2.404) was found significant as its p-value is less than 0.05. The difference can be seen with the help of graphical representation of mean score. (Figure 1)

3.1.3 Descriptive Statistics for the data on grip strength of middle aged women

Table 5: Paired Samples Statistics

		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	GS_PRE	21.1450	20	4.27249	.95536
	GS_POST	22.6450	20	4.68879	1.04844


Table 5 Shows that the mean score of pre and post test, the mean and SD of Grip Strength 21.1450_4.27249 & 22.6450_4.68879 respectively.

Table 6: Paired Samples Test

		Paired Differences					t	df	Sig. (2-tailed)
		Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
					Lower	Upper			
Pair 1	GS_PRE GS_POST	-1.50000	1.65593	.37028	-2.27500	-.72500	-4.051	19	.001

According to analysis of data presented in table 4, the calculated t-value (-4.051) was found significant as its p-value is less than 0.05. The difference can be seen with the help of graphical representation of mean score. (Figure 1)

Figure 1: Graphical representation of mean score of cardiovascular endurance, flexibility and grip strength



4. Findings and Discussion

In the present study, it is showed that there is a significant difference between pre and post test of 2 variables which is flexibility and grip strength; the development in body fat, hypertension, heart disease that creates interest of women to participate in aerobic dance activities, yoga asana's. This study has involved aerobic dance, zomba and yoga sessions which may showed the positive result with significant difference in flexibility and grip strength. There is no significant difference between pre and post test of 1 variable which is cardiovascular endurance. It may be attributed due to the age group, while the evidence is limited, it appears that middle-aged and older women have positive attitudes to exercise but seem unable or unwilling to take action. The habit of not being engaged with physical fitness exercise may not be developing in very short period of time as 6 weeks training. The time duration of this study was insufficient to bring significant difference.

4.1 Discussion of Hypothesis

On the basis of the results of the study, the hypothesis was accepted at 0.05 level of significance in flexibility and grip strength because there was significant difference between pre and post test after six weeks of training program.

The hypothesis was rejected at 0.05 level of significant in cardiovascular endurance.

5. Conclusion

The researcher was able to obtain the following conclusion on the basis of the results obtained after administration of the test.

The result of the statistical application shows that:

1. There was an insignificant difference in cardio vascular endurance after the pre and post test. The reason behind this can be judged because of the age group of the subjects. Another reason for the negative result can be given because of their lack of physical workout during their daily life activities.
2. However, there was a significant difference in flexibility and grip strength of the subjects.

References

1. Blumenthal, J. A., Emery, C. F., Madden, D. J., George, L. K., Coleman, R. E., Riddle, M. W., ... & Williams, R. S. (1989). Cardiovascular and behavioral effects of aerobic exercise training in healthy older men and women. *Journal of gerontology*, 44(5), M147-M157.
2. Blumenthal, J. A., Fredrikson, M., Kuhn, C. M., Ulmer, R. L., Walsh-Riddle, M., & Appelbaum, M. (1990). Aerobic exercise reduces levels of cardiovascular and sympathoadrenal responses to mental stress in subjects without prior evidence of myocardial ischemia. *The American journal of cardiology*, 65(1), 93-98.
3. Bowen, D. J., Fesinmeyer, M. D., Yasui, Y., Tworoger, S., Ulrich, C. M., Irwin, M. L., ... & McTiernan, A. (2006). Randomized trial of exercise in sedentary middle aged women: effects on quality of life. *International Journal of Behavioral Nutrition and Physical Activity*, 3(1), 34.
4. Fleg, J. L., O'connor, F., Gerstenblith, G., Becker, L. C., Clulow, J., Schulman, S. P., & Lakatta, E. G. (1995). Impact of age on the cardiovascular response to dynamic upright exercise in healthy men and women. *Journal of Applied Physiology*, 78(3), 890-900.
5. Gillett, P. A., & Eisenman, P. A. (1987). The effect of intensity controlled aerobic dance exercise on aerobic capacity of middle-aged, overweight women. *Research in nursing & health*, 10(6), 383-390.

6. Gutin, B., Barbeau, P., Owens, S., Lemmon, C. R., Bauman, M., Allison, J., ...& Litaker, M. S. (2002). Effects of exercise intensity on cardiovascular fitness, total body composition, and visceral adiposity of obese adolescents. *The American journal of clinical nutrition*, 75(5), 818-826.

Kiran
EFFECT OF 6 WEEKS AEROBIC EXERCISES TRAINING PROGRAM
TO PHYSICAL FITNESS VARIABLES IN MIDDLE AGED WOMEN

Creative Commons licensing terms

Authors will retain the copyright of their published articles agreeing that a Creative Commons Attribution 4.0 International License (CC BY 4.0) terms will be applied to their work. Under the terms of this license, no permission is required from the author(s) or publisher for members of the community to copy, distribute, transmit or adapt the article content, providing a proper, prominent and unambiguous attribution to the authors in a manner that makes clear that the materials are being reused under permission of a Creative Commons License. Views, opinions and conclusions expressed in this research article are views, opinions and conclusions of the author(s). Open Access Publishing Group and European Journal of Physical Education and Sport Science shall not be responsible or answerable for any loss, damage or liability caused in relation to/arising out of conflict of interests, copyright violations and inappropriate or inaccurate use of any kind content related or integrated on the research work. All the published works are meeting the Open Access Publishing requirements and can be freely accessed, shared, modified, distributed and used in educational, commercial and non-commercial purposes under a [Creative Commons attribution 4.0 International License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/).