



THE EFFECT OF AEROBIC EXERCISE AND WEIGHT-LIFTING PLUS AEROBIC EXERCISE ON BLOOD PRESSURE AND BLOOD PARAMETERS IN SEDENTARY FEMALES

Osman İmamoğlu,

Pelin Akyol, Arif Satici¹

Ondokuz Mayıs University,
Yaşar Doğu Sport Science Faculty,
Turkey

Abstract:

The purpose of this study was determined the effects of 12 weeks aerobic exercise and weight-lifting plus aerobic exercise on blood pressure, blood serum lipids and lipoproteins of sedentary females. Thirty seven sedentary females (Aerobic group mean age: 44.66 years, Weight-Lifting + Aerobic group mean age: 40.80 years) participated in this study. Before initiate the exercise their blood samples, peripheral measurements and thickness of skinfolds were taken. The subjects participated in aerobics group (n = 21) and weight-lifting plus aerobics group (n = 16) 3 times in a week through one hour. The intensity of exercise was increased gradually and their heart rates were raised up to 130-140 rates per-minute and of the tension. Two way ANOVA was used in this study. There were no significant difference between two different type of exercise groups ($p>0.05$) first group (aerobic); HDL-C (High Density Lipoprotein-Cholesterol) 18.12%, second group (weight-lifting plus aerobic) %28.23, there were a significant increase among them ($p<0.01$). Total cholesterol decrease 13.97% in group of aerobic exercise and it decreases 7.13% in group of weight-lifting plus aerobic. Triglyceride decreases 31.74% in group of aerobic exercise; it decreases 19.53% in group of weight-lifting plus aerobic exercise. Level of LDL-C was decreased 28.11% in group of aerobic exercise, 21.46% in group of weight-lifting plus aerobic. When we consider exercise types, the only flats of type is on triglyceride and APO B, ($p<0.05$), there were effects of exercise duration is on HDL-C, LDL-C AND Total cholesterol ($p<0.01$). There were effects of type and duration of exercise on TC, TG, HDL-C, LDL-C, APO A1, and APO B ($P<0.01$). In this study,

¹ Correspondence: email arifsatici@gmail.com

regular aerobic and weight-lifting plus aerobic exercises have been shown to reduce the risk of cardiovascular disease in sedentary women and a significant improvement in systolic and diastolic blood pressures as well as reduction of cholesterol, triglyceride, and of LDL-C. Level of HDL-C was increased in both of exercise groups.

Keywords: aerobic exercise, weight-lifting, blood pressure, blood parameters, sedentary females

1. Introduction

The lipid and lipoprotein abnormalities play a major role in the development and progression of coronary artery disease and participate in many physiological and biochemical events in human body. Benefits include improved serum lipid profiles, blood pressure and inflammatory markers as well as reduced risk of stroke, acute coronary syndrome and overall cardiovascular mortality (Skoumas et al., 2003; Cinar, 2010; Shiraev and Barclay, 2012; Baydil, 2013). Human is dynamic existence and should move to survive. If the movement is performed with some rules and discipline, in order to purpose, ensure our physical and physiological health (Baydil, 2013; Yılmaz, 2016). Recently, number of women who exercise for fitness, except from professional's athletes, are lower than number of women who don't perform any sport activity (Özdal and Bostancı, 2016). Sport activity can improve the total and regional fat mass of women (Özdal et al., 2016). Because of modern life physically motionless effects every aged of people. The problems such as; hypertension, obesity, muscle atrophy, postural disorder and inadequate cardiovascular system caused by lack of exercise and sedentary life style or harmful habits such as smoking. Because of differences in percentage body fat according to sex difference and inadequate possibility or could not find the leisure time for their own life health female population affected seriously from lack of exercise than male counter parts. These results encourage many researchers to make investigation to find out effects of exercise to females physical and philological health. Finding about Coronary Heart disease (CHD) in active people is really; and in older ages the rate of the mortality is less than sedentary people (Fripp et al, 1985; Hessel et al., 1988; Saçaklı, 1992; Mayda, 2016; Pancar, 2017; Özdal et al., 2017).

The importance of regular physical activity is emphasized in order to maintain a healthy life and to have both preventive and healing effects for many diseases and injuries (Hillsdon et al., 2005; Brown et al., 2012; Yılmaz, 2017; Vural, 2017; Biçer et al., 2015).

When literature was reviewed about effects of exercise on plasma lipids, lipoproteins, blood pressure and heart rates, results were indicated that moderate and low intensity exercise, if its performed adequate time period (at least 6-8 weeks) cause decrease in body weight and body fat ratio, additionally decrease serum TC level (Dufaux et al., 1982). In the most studies, it has been noticed that regular exercise affects levels of lipid and lipoprotein positively and it is salutary for the heart. It has been identified that regular exercise reduces total blood cholesterol, serum triglycerides and LDL-C and increases high-density lipoprotein cholesterol HDL-C (Champe at al, 1997; Imamoğlu, 2014).

It is known that cardiovascular risk factors are greatly improved through the cardiovascular changes that occur with regular and long-term exercise (Çicek et al., 2017). Cardiovascular disease (CVD) in women is the leading cause of mortality in the United States, and less than optimal lipid and lipoprotein levels are major risk factors for CVD (George at al, 2004). There is consolidated evidence that physical activity exerts beneficial effects on several chronic conditions and longevity, on the basis of its proposed biological effects, especially on lipid profiles (Lippi et al., 2006). Aerobic exercise is efficacious for increasing High density lipoprotein (HDL-C) and decreasing Total Cholesterol (TC), Low density lipoprotein (LDL-C), and Triglyceride (TG) in women (George at al, 2004). High-intensity aerobic training results in improvement in high-density lipoprotein cholesterol (Tambalis et al., 2008).

At the end of aerobic exercise, hearts rates decrease and (O₂) need of heart muscle and heart activity become economical. These events show increase of HDL-C, decrease of LDL-C and increase protective effects to arteriosclerosis (Imamoğlu et al., 1998). Cardiovascular disease (CVD) is the leading cause of death worldwide (Cutler, 2006). Low blood levels of high-density lipoprotein cholesterol (HDL-C) are an independent risk factor for CVD (Franceschini, 2001; Boden, 2000). Cross-sectional data provide strong evidence that people who are more physically active have higher HDL-C levels (Drygas et al., 2000). Thus, the value of regular aerobic exercise in increasing serum HDL-C level and in reducing the risk of CVD has received widespread acceptance (Rippe et al., 1988). In contrast, results of aerobic exercise studies vary considerably, depending on the exercise program (duration, intensity, or frequency) and characteristics of subjects at baseline (Crouse et al, 1997).

In this study we examined effect of 12 weeks two different types of exercise (aerobic and weight-lifting plus aerobic) on blood pressure and blood serum parameters on middle aged females.

2. Material and Methods

Thirty seven healthy sedentary females (mean age: Aerobic group, 44.66 years; and Weight-Lifting + Aerobic group 40.80 years) participated in this study as a subject. Subjects were divided in to two group; aerobic exercise (n = 21) and weight-lifting plus aerobic exercise (n = 16). Both groups participate the training during 12 weeks period with one hour training session 3 days in a week period. Intensity of training increased slightly and heart rates were 130-140 rates in a minute at the end of exercise.

At the beginning of the study, blood samples were taken from all subjects before breakfast in the morning. HDL-C, LDL-C, TC, TG and polyproteins I (APO A1 and APO B) levels were determined these blood samples. The blood pressure and heart rates were measured by Digital Blood Pressure Meters ALP K2 777. At the end of 12 weeks period, all measurements were repeated.

All statistical data obtained from the study was calculated with licensed SPSS 22.0 packet program. Means and standard deviation were calculated with the program. Test of normality for the data were made by Shapiro-Wilk test. The difference between pre-test, post-test results of group was determined by two way ANOVA. The types of exercises are aerobic and weight-lifting plus aerobic; the duration of exercise is between pre-test and post-test. Statistical significance were accepted as and $p < 0.05$, $p < 0.01$.

3. Results

Table 1 shows physical parameters of pre-test and post-test results sedentary females who participated to our study, and table 2 shows alteration rates of blood parameters, table 3 shows blood pressure and heart rates alteration, and two way ANOVA test results of blood parameters exercise duration and types are determined in Table 4.

Table 1: Physical parameters of females before and after exercise who performed aerobic exercise and weight-lifting plus aerobic exercise

Variables	Aerobic Exercise		Weight-Lifting + Aerobic		Aerobic	Difference Weight-lifting plus aerobic
	Before 3 months mean	After 3 month mean	Before 3 month mean	After 3 month mean		
Age (year)	44.66	-	40.8	-	-	-
Height (cm)	158.45	-	157.3	-	-	-
Body Weight (kg)	72.30	66.43	69.36	63.44	-8.12	-8.54
BMI (kg/m ²)	29.36	26.11	29.04	25.14	-11,07	-13,43

Table 2: Alteration Percentage of Blood Parameters of Sedentary Females

	Aerobic Exercise		Weight-Lifting+ Aerobic		Difference	
	Before 3 months mean	After 3 months mean	Before 3 months means	After 3 months mean	Aerobic	Weight-lifting plus aerobic
Mg/100						
Cholesterol	221.2	190.3	196.9	182.5	-13.97	-7.32
Triglyceride	126.35	86.25	91.89	73.95	-31.74	-19.53
HDL-C	35.76	42.30	34.43	44.15	18.29	28.23
LDL-C	118.15	84.94	111.92	87.91	-28.11	-21.46
APO A1	153.21	137.94	151.08	131.07	-9.97	-13.25
APO B	126.14	109.84	108.81	100.60	-7.93	-7.55

Decrease and increase percentage of blood values of females (Table 2); cholesterol decrease 13.97% in group of aerobic exercise and it decreases 7.13% in group of weight-lifting plus aerobic. Triglyceride decreases 31.74% in group of aerobic exercise; it decreases 19.53% in group of weight-lifting plus aerobic exercise. Level of HDL-C was increased in both of exercise groups. This increase is 18.29% in group of aerobic exercise, 28.23% in group of weight-lifting plus aerobic. Level of LDL-C was decreased 28.11% in group of aerobic exercise, 21.46% in group of weight-lifting plus aerobic. APO A1 was decreased 9.97% in group of aerobic exercise, 13.25% in group of weight-lifting plus aerobic exercise. APO B was decreased 7.93% in group of aerobic exercise, 7.55% in group of weight-lifting plus aerobic exercise.

Table 3: Alteration percentage of the blood pressure and heart rates of sedentary females

Variables	Aerobic Exercise		Weight-Lifting + A		Difference	
	Before Exercise mean	After Exercise mean	Before Exercise mean	After Exercise mean	Aerobic	Weight-lifting plus aerobic
	Systolic (mmHg)	119.05	108.15	113.16	100.29	-9.16
Diastolic (mmHg)	76.34	73.42	74.57	69.64	-3.83	-6.61
Heart rate (bpm)	89.58	77.22	84.20	72.29	-13.8	-14.14

According to Table 3, in generally there is no significant percentage difference between the groups. The sistolic blood pressure was decreased 9.16% in group of aerobic exercise, 11.38% in group of weight-lifting plus aerobic. Diastolic blood pressure was decreased 3.83% in group of weight-lifting plus aerobic, 6.61% in group of aerobic exercise. Heart rate was decrease 13.8% in group of aerobic exercise, 14.14% in group of weight-lifting plus aerobic.

Table 4: Two way ANOVA results of effects of exercise types and duration to the blood parameters

Variables	Source of Variation	Sum of Square	Df	Mean of square	F	Level of significant
Cholesterol	Main effect	16574.67	2	8236.08	10.52	.000**
	Type of Exercise	6010.83	1	6021.83	7.770	.006*
	Period of Exercise	10353.33	1	10451.33	13.13	.001**
Triglyceride	Main effect	17175.23	2	8625.61	6.61	.003*
	Type of Exercise	7229.53	1	7239.53	5.58	.021*
	Period of Exercise	10026.70	1	10036.70	7.69	.007*
HDL-C	Main effect	807.41	2	403.20	10.40	.000**
	Type of Exercise	2.30	1	2.41	.62	.804
	Period of Exercise	805.00	1	805.00	20.69	.000**
LDL-C	Main effect	724.67	2	3643.83	9.49	.000**
	Type of Exercise	1853.62	1	1853.62	7.38	.011*
	Period of Exercise	5432.05	1	5430.05	16.32	.001**
APO A1	Main effect	7195.79	2	3599.39	12.53	.000**
	Type of Exercise	1952.74	1	1952.74	6.71	.012*
	Period of Exercise	5244.05	1	5242.05	17.86	.000**
APO B	Main effect	4271.03	2	214.01	4.09	.023*
	Type of Exercise	2592.770	1	2592.77	4.92	.031*
	Period of Exercise	1678.26	1	1675.26	3.05	.081

**level of significant according to .001 *Level of significant according to .05

Effect of type and period of exercise on blood cholesterol found significant ($p < 0,01$). It is obvious that, more effect of period of exercise than type of exercises was significant ($p < 0.05$ and $p < 0.001$). While type and period of exercise is effecting is effecting HDL-C and LDL-C significant ($p < 0.01$), only Period of exercise can it effect on HDL-C them ($p < 0.05$). Significant of APO A1 is same as significance LDL-C type and period of exercise can be significant, together type of exercise and period of exercise is significant ($p < 0.01$). The only period of exercise is not significant effect on APO B. Because of duration and type of exercise are significant ($p < 0.05$), (Table 4) when they controlled together.

4. Discussion

The changing of plasma lipid can be affected by endogenic sex hormones, meanwhile reducing body fat mass can be affected by physical activities after training (Sacks and Dzau, 1986). Because of this working which is about enough density and strength

exercise, HDL-C level raises, LDL-C level falls down (Massarei and Pyke, 1982). We accept that physical activity changes lipid and lipoprotein profiles (Crigui, 1986; Stein et al., 1990).

Several studies have shown that exercise can cause improving effect on serum HDL-C, LDL-C and TG (Güllü et al., 2013; Boardley et al., 2007) while some other studies didn't found any significant differences after exercises (Stefanick et al., (1998; Thomas et al., 2000).

Çiçek et al (2017) found 16 weeks of step-aerobic and core exercise showed significant changes of inflammatory and lipid markers with cardiac dimensions and had favorable effects on both left ventricular systolic function. Massarei et al (1982), found that long period of exercise increases HDL-C with aerobic capacities increase and reduce heart illness risk. In this study for a month, females who are obese, rode bicycle made usual, special and passive gymnastic. At the end they couldn't found significance, triglyceride, total cholesterol, LDL-C was decreased, HDL-C was increased. But when we think of effect of exercise which are on plasma lipid and lipoprotein we see that exercises which have not low and low density and enough period (6-8 weeks) reduce body mass, body fat percentage and total serum cholesterol (Dufaux et al., 1982). Trainings which have low and not low levels provide real significant fall (Franklin et al., 1984).

Ready et al. (1995), found to reduce; remaining cholesterol of women after menopause with walking program. They compared control group and walking group and they determined increase HDL-C and to decrease in total cholesterol and triglyceride too much.

Yanagibari et al. (1993), investigated 12 weeks walking effect of exercise on serum lipids lipoproteins and apolipoproteins of middle aged female. They compared to part of two females groups where walking. And they determined both groups to increase in HDL-C and to decrease in APO A1, APO B, total cholesterol and LDL-C.

Comparisons between intensities of aerobic exercise programs resulted in favorable effects only for high intensity. The most frequently observed alteration was an increase in the high-density lipoprotein cholesterol, whereas reductions in triglycerides, total cholesterol, and low-density lipoprotein cholesterol appeared less often (Tambalis et al.2008). Imamoğlu et al (1998), investigated to effect short time exercise on serum lipids of sedentary females before and after menopause and they determined to increase in HDL-C (%27.76) and to decrease in LDL-C (20.06%), Total cholesterol (12.96%), triglyceride (27.76%) and apolipoproteins. Koca (2017) in one study was finding on the effects of 12 weeks aerobic exercise 13.9% in cholesterol, 25.3% in Triglyceride, and 22.07% in LDL-C have been registered. In this study, after 3 month exercise present

results show that; Amount of cholesterol decreased in first group (aerobic exercise) from 221.2 to 190.3 (13.97%) in second group (aerobic exercise plus weight-lifting) from 199.9 to 182.5 (7.32%) triglyceride decreased in 1 st group from 126.35 to 86.25 (31.74%) in 2nd group from 91.89 to 73.95 (19.53%), LDL-C decreased in 1 st group from 118.15 to 84.94 (28.11%) in 2nd group from 111.92 to 87.95 (21.46%). Level of HDL-C was increased in both of exercise groups. This increase is 18.29% in group of aerobic exercise, 28.23% in group of weight-lifting plus aerobic.

Coroner heart disease risk decrease %1.5 cause of to decrease %1 of HDL-C so that, in our study, decrease of LDL-C and increases of HDL-C have decreased risk of CHD (Cardiovascular Heart Daises). Our study concordant Yanagibori et al., Imamoğlu et al and Koca studies. In our study, performing aerobic and weight-lifting plus aerobic exercise groups changing percentage of serum lipids and lipoproteins resemble to each other and before the exercise there is no difference, after the exercise all variables decreased significantly.

In present study, at the end of 12 weeks exercise, decrease of cholesterol is normal ($p < 0.01$) type of exercise and time of exercise are meaningful to each effect from statistical direction on the cholesterol ($p < 0.01$). In our study, time of exercise effect to cholesterol, triglyceride, HDL-C, LDL-C, APO A1. In our study, Type of Exercise not effect on the HDL-C. Main effective each effect of type and of exercise is manful on blood parameters.

Most of studies about blood pressure and heart rate, claim that the exercise decreased the blood pressure and heart rate (Nelson, 1986; Parker et al., 1991; Suzuki et al., 1996, Wilmore et al., 1996). Parker et al. (1996), investigated heartrate, systolic blood pressure during submaximal walking on the female subjects. During 16 weeks, 3 times 1 hour study at a week finally average of heart rate decreased from 135-120 rate/minute to 108-104 rate/minute. So that, systolic blood pressure decreased. Wilmore et al. (1996), made same study. They investigated 62 subjects at certain times of day and they found that, the aerobic exercise decreased heart rate and blood pressure. They determined that moderate strength aerobic exercise saved blood pressure in daily stresses of life (Brownley et al., 1996). Koca (2017) found at end of the activity 12 weeks aerobic exercises, decrease effects of blood pressure, heart beats and blood parameters and cardiovascular system in middle-aged sedentary females.

In these study findings, the sistolic blood pressure was decreased 9.16% in group of aerobic exercise, 11.38% in group of weight- lifting plus aerobic. Diastolic blood pressure was decreased 3.83% in group of weight-lifting plus aerobic, 6.61% in group of aerobic exercise. Heart rate was decrease 13.8% in group of aerobic exercise, 14.14% in

group of weight-lifting plus aerobic. This reduction in heart rate variability can be considered in significant.

In this study, regular aerobic and weight-lifting plus aerobic exercises have been shown to reduce the risk of cardiovascular disease in sedentary women and a significant improvement in systolic and diastolic blood pressures as well as reduction of cholesterol, triglyceride, and of LDL-C. Level of HDL-C was increased in both of exercise groups.

References

1. Baydil, B. (2013). Serum macro-micro element responses to acute maximal physical exercise. *World Applied Sciences Journal*, 23(7), 945-949.
2. Baydil, B. (2013). The effect of acute maximal aerobic exercise on humoral immune parameters on healthy young males. *International Journal of Academic Research*, 5(3).
3. Biçer, M., Özdal, M., Akcan, F., Mendes, B., & Patlar, S. (2015). Effect of strength training program with elastic band on strength parameters. *Journal of Biology of Exercise*, 11(2), 111-122.
4. Boardley, D., Fahlman, M., Topp, R., Morgan, A. L., & McNevin, N. (2007). The impact of exercise training on blood lipids in older adults. *The American journal of geriatric cardiology*, 16(1), 30-35.
5. Boden, W. E. (2000). High-density lipoprotein cholesterol as an independent risk factor in cardiovascular disease: assessing the data from Framingham to the Veterans Affairs High-Density Lipoprotein Intervention Trial. *The American journal of cardiology*, 86(12), 19-22.
6. Brown, W. J., McLaughlin, D., Leung, J., McCaul, K. A., Flicker, L., Almeida, O. P., & Dobson, A. J. (2012). Physical activity and all-cause mortality in older women and men. *Br J Sports Med*, bjsports-2011.
7. Brownley, K. A., West, S. G., Hinderliter, A. L., & Light, K. C. (1996). Acute aerobic exercise reduces ambulatory blood pressure in borderline hypertensive men and women. *American journal of hypertension*, 9(3), 200-206.
8. Cicek, G., Imamoglu, O., Gullu, A., Celik, O., Ozcan, O., Gullu, E., & Yamaner, F. (2017). The effect of exercises on left ventricular systolic and diastolic heart function in sedentary women: Step-aerobic vs core exercises. *Journal of Exercise Science & Fitness*, 15(2), 70-75.

9. Cinar, V., Mogulkoc, R., Baltaci, A. K., & Bostanci, O. (2010). Effects of calcium supplementation on glucose and insulin levels of athletes at rest and after exercise. *Biological trace element research*, 133(1), 29-33.
10. Criqui, M. H. (1986). Epidemiology of atherosclerosis: an updated overview. *The American journal of cardiology*, 57(5), C18-C23.
11. Drygas, W., Kostka, T., Jegier, A., & Kuski, H. (2000). Long-term effects of different physical activity levels on coronary heart disease risk factors in middle-aged men. *International journal of sports medicine*, 21(04), 235-241.
12. Dufaux, B., Assmann, G., & Hollmann, W. (1982). Plasma lipoproteins and physical activity: a review. *International journal of sports medicine*, 3(03), 123-136.
13. Franceschini, G. (2001). Epidemiologic evidence for high-density lipoprotein cholesterol as a risk factor for coronary artery disease. *The American journal of cardiology*, 88(12), 9-13.
14. Fripp, R. R., Hodgson, J. L., Kwiterovich, P. O., Werner, J. C., Schuler, H. G., & Whitman, V. (1985). Aerobic capacity, obesity, and atherosclerotic risk factors in male adolescents. *Pediatrics*, 75(5), 813-818.
15. Gullu, E., Gullu, A., Cicek, G., Yamaner, F., Imamoglu, O., & Gumusdag, H. (2013). The effects of aerobic exercises on cardiovascular risk factors of sedentary women. *Int. J. Acad. Res*, 5, 160-167.
16. Hillsdon, M. M., Brunner, E. J., Guralnik, J. M., & Marmot, M. G. (2005). Prospective study of physical activity and physical function in early old age. *American journal of preventive medicine*, 28(3), 245-250.
17. İmamoğlu, O., Altun, P., Çebi, M., & Kışalı, N. F. (2000). Menapoz öncesi ve sonrası sedanter bayanlarda 3 aylık egzersizin vücut kompozisyonu, serum lipidleri, lepoproteinleri ve fiziksel fitness üzerine etkileri. *Journal of Physical Education and Sport Sciences*, 1(1).
18. Kelley, G. A., Kelley, K. S., & Tran, Z. V. (2004). Aerobic exercise and lipids and lipoproteins in women: a meta-analysis of randomized controlled trials. *Journal of women's health*, 13(10), 1148-1164.
19. Koca F. (2017). The effect of 12 weeks aerobic exercise on blood pressure, heart rates and blood parameters in sedentary females, *International Journal of Academic Research, Part A. Applied and Natural Sciences*, 9(1), 5-7.
20. Lippi, G., Schena, F., Salvagno, G. L., Montagnana, M., Ballestrieri, F., & Guidi, G. C. (2006). Comparison of the lipid profile and lipoprotein (a) between sedentary and highly trained subjects. *Clinical Chemical Laboratory Medicine*, 44(3), 322-326.

21. Massarelli J.R.L., Pyke F.S., (1982). Physical fitness and plasma HDL cholesterol male in business executives. *Atherosclerosis*, 42(1):77-83.
22. Mayda, M. H., Karakoç, Ö., & Özdal, M. (2016). The Investigation of Some Physical, Physiological and Anthropometric Parameters of Visually Impaired and Non-impaired a National Male Judoka. *Journal of Education and Training Studies*, 4(6), 192-198.
23. Nelson, L., Esler, M., Jennings, G., & Korner, P. (1986). Effect of changing levels of physical activity on blood-pressure and haemodynamics in essential hypertension. *The lancet*, 328(8505), 473-476.
24. Özdal, M., Bostancı, Ö., Kabadayi, M., & Akcan, F. (2016). Influence of Electromyostimulation Training with Voluntary Muscle contractions on Basal Metabolic Rate of Sedentary Women. *International Journal of Sport Studies*, 6(1), 12-15.
25. Özdal M., & Bostancı, Ö. (2016). Effects of whole-body electromyostimulation with and without voluntary muscular contractions on total and regional fat mass of women. *Archives of Applied Science Research*, 8(3), 75-79.
26. Özdal, M., Pancar, Z., Çınar, V., Bilgiç, M. (2017). Effect of Smoking on Oxygen Saturation in Healthy Sedentary Men and Women. *EC Pulmonology and Respiratory Medicine*, 4(6), 178-182.
27. Pancar, Z., Özdal, M., & Çınar, V. (2017). The effect of 4-weekly low intensity physical activity program in thyroid hormone levels in obese and overweight children. *European Journal of Physical Education and Sport Science*, 3(11), 1-8.
28. Parker, N. D., Hunter, G. R., Treuth, M. S., Kekes-Szabo, T., Kell, S. H., Weinsier, R., & White, M. (1996). Effects of strength training on cardiovascular responses during a submaximal walk and a weight-loaded walking test in older females. *Journal of Cardiopulmonary Rehabilitation and Prevention*, 16(1), 56-62.
29. Ready A.E., Drinkwater D.T., Ducas J., Fitzpatrick D.W. Brereton D.G., Oades S.C., (1995). Walking program reduces elevated cholesterol in women postmenopausal. *Canadian Journal of Cardiology*. 11 (10):905-12.
30. Sacks, F. M., & Dzau, V. J. (1986). Adrenergic effects on plasma lipoprotein metabolism: speculation on mechanisms of action. *The American journal of medicine*, 80(2), 71-81.
31. Skoumas, J., Pitsavos, C., Panagiotakos, D. B., Chrysohoou, C., Zeimbekis, A., Papaioannou, I. ... & Stefanadis, C. (2003). Physical activity, high density lipoprotein cholesterol and other lipids levels, in men and women from the ATTICA study. *Lipids in health and disease*, 2(1), 3.

32. Stefanick, M. L., Mackey, S., Sheehan, M., Ellsworth, N., Haskell, W. L., & Wood, P. D. (1998). Effects of diet and exercise in men and postmenopausal women with low levels of HDL cholesterol and high levels of LDL cholesterol. *N Engl J Med*, 1998 (339), 12-20.
33. Stein, R. A., Michielli, D. W., Glantz, M. D., Sardy, H., Cohen, A., Goldberg, N., & Brown, C. D. (1990). Effects of different exercise training intensities on lipoprotein cholesterol fractions in healthy middle-aged men. *American Heart Journal*, 119(2), 277-283.
34. Tambalis, K., Panagiotakos, D. B., Kavouras, S. A., & Sidossis, L. S. (2009). Responses of blood lipids to aerobic, resistance, and combined aerobic with resistance exercise training: a systematic review of current evidence. *Angiology*, 60(5), 614-632.
35. Thomas, E. L., Brynes, A. E., McCarthy, J., Goldstone, A. P., Hajnal, J. V., Saeed, N., & Bell, J. D. (2000). Preferential loss of visceral fat following aerobic exercise, measured by magnetic resonance imaging. *Lipids*, 35(7), 769-776.
36. Vural, M., Özdal, M., & Öztütüncü, S. (2017). The effect of 4-week two different strength training programs on body composition. *European Journal of Physical Education and Sport Science*, 3(7), 1-10.
37. Wilmore, J. H., Stanforth, P. R., Gagnon, J., Leon, A. S., Rao, D. C., Skinner, J. S., & Bouchard, C. (1996). Endurance exercise training has a minimal effect on resting heart rate: the HERITAGE Study. *Medicine and science in sports and exercise*, 28(7), 829-835.
38. Yılmaz A.K., Kabadayı, M., Mayda, M.H., Birinci, M.C. (2016). Examination of isokinetic strength rates of knee joint H Q in football players. *Journal of Social Sciences Research*, 10(4).
39. Yılmaz, A. K., Kabadayı, M., Mayda M, H., Çavuşoğlu, G., & Taşmektepligil, M. Y. (2017). Analysis of q angle values of female athletes from different branches. *Ovidius University Annals, Series Physical Education & Sport/Science, Movement & Health*, 17(2).

Creative Commons licensing terms

Authors will retain the copyright of their published articles agreeing that a Creative Commons Attribution 4.0 International License (CC BY 4.0) terms will be applied to their work. Under the terms of this license, no permission is required from the author(s) or publisher for members of the community to copy, distribute, transmit or adapt the article content, providing a proper, prominent and unambiguous attribution to the authors in a manner that makes clear that the materials are being reused under permission of a Creative Commons License. Views, opinions and conclusions expressed in this research article are views, opinions and conclusions of the author(s). Open Access Publishing Group and European Journal of Physical Education and Sport Science shall not be responsible or answerable for any loss, damage or liability caused in relation to/arising out of conflict of interests, copyright violations and inappropriate or inaccurate use of any kind content related or integrated on the research work. All the published works are meeting the Open Access Publishing requirements and can be freely accessed, shared, modified, distributed and used in educational, commercial and non-commercial purposes under a [Creative Commons attribution 4.0 International License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/).