brought to you by DCORE

EMERGING SCIENCE JOURNAL

# Emerging Science Journal

Available online at www.ijournalse.org

# **Emerging Science Journal**

Vol. 3, No. 6, December, 2019



# Finite-time Stability, Dissipativity and Passivity Analysis of Discrete-time Neural Networks Time-varying Delays

Porpattama Hammachukiattikul <sup>a\*</sup>

<sup>a</sup> Mathematics Department, Phuket Rajabhat University, Phuket 83000, Thailand

### Abstract

The neural network time-varying delay was described as the dynamic properties of a neural cell, including neural functional and neural delay differential equations. The differential expression explains the derivative term of current and past state. The objective of this paper obtained the neural network time-varying delay. A delay-dependent condition is provided to ensure the considered discrete-time neural networks with time-varying delays to be finite-time stability, dissipativity, and passivity. This paper using a new Lyapunov-Krasovskii functional as well as the free-weighting matrix approach and a linear matrix inequality analysis (LMI) technique constructing to a novel sufficient criterion on finite-time stability, dissipativity, and passivity of the discrete-time neural networks with time-varying delays. An effective LMI approach derives by base the appropriate type of Lyapunov functional. Finally, we present the effectiveness of novel criteria of finite-time stability, dissipativity, and passivity, dissipativity, and passivity condition of discrete-time neural networks with time-varying delays in the form of linear matrix inequality (LMI).

#### **Keywords:**

Finite-time Stability; Dissipativity and Passivity Analysis; Lyapunov-Krasovskii Functional.

#### Article History:

Received:	02	October	2019
Accepted:	22	November	2019
Published:	01	December	2019

# **1-Introduction**

Current years we have been attending in researching delay neural networks (NNs), this is mainly to the major feasible applications in many areas, for example, combinatorial optimization, static image processing, pattern recognition, associative memory and signal processing [1]. On the other hand, the time delay is ineluctable in various applied systems, and it is also the principal cause of poor performance, oscillation, and instability of the systems. Therefore, important interest has been considered to delay-dependent conditions of analysis and combination problems of time-delay NNs [2]. Thus, it is significant to learn the stability of discrete-time neural networks (DNNs) with time-varying delay.

First recommended by Popov [3], the idea of passive systems from the beginning occurs in the conditions of electrical circuit theory. In the preliminary 1970s, Willems [4] developed the concept of dissipative systems, and passive systems, by suggestion the symbols of a supply rate and a storage function. Dissipativity theory provides a substructure for the analysis and design of control systems applying an input-output feature used as a basis energy-related judgment.

Neural networks (NNs) have been successfully practiced in a diversity of fields such as signal processing, pattern recognition, static image processing, associative memory, and combinatorial optimization, and these a applications contingent on seriously on their dynamic department. The dynamical conduct of NNs are the key to the above-said applications, and a necessary step for the practical design of NNs. Up to now, there have been fruitful research results available in the literature about the dynamic deportment of NNs [1-6]. Many results have been investigated by the dynamic behaviour of continuous-time NNs [7-10]. However, compared with continuous-time NNs, discrete-time neural networks (DNNs) equally have a strong engineering application background for the sake of computer-based simulation

<sup>\*</sup> CONTACT: Porpattama@pkru.ac.th

DOI: http://dx.doi.org/10.28991/esj-2019-01198

<sup>© 2019</sup> by the authors. Licensee ESJ, Italy. This is an open access article under the terms and conditions of the Creative Commons Attribution (CC-BY) license (https://creativecommons.org/licenses/by/4.0/).

and the dynamics of continuous-time NNs cannot be preserved by discretization as mentioned in [11]. Hence, it is essential to study the dynamical behaviour of DNNs.

Moreover, in many physical and biological phenomena, the rate of variation in the system state depends on the past states. This characteristic is called a delay (or a time delay) and therefore a system with a time delay is called a timedelay system. Time delay phenomena were first discovered in biological systems and were later found in many engineering systems, such as mechanical transmissions, fluid transmissions, metallurgical processes, and networked control systems. They are often a source of instability, periodic oscillatory, chaos, and poor control performance. Timedelay systems have attracted the attention of many researchers [12-14] because of their importance and widespread occurrence.

It is well known that dissipativeness was initially introduced by "Willems" in terms of an inequality involving the storage function and supply rate. Dissipativity hypothesis needs to assume a discriminating some piece in the dissection Also control outline for straight Also nonlinear systems, particularly to high-order framework [15], since from those useful requisition purpose about the view, a significant number frameworks have to make dissipative for accomplishing viable commotion weakening. [16-19]. That provides a strong connection between Physics, system theory, and control engineering. The dissipated theory has proven to be an essential and very useful tool for control applications like robotics, active vibration damping, electromechanical systems, combustion engines, circuit theory, and for control techniques like adaptive control, and inverse optimal control problems. The dissipative theory being a framework for the design and analysis of control systems using an input-output description based on energy-related consideration is applicable in characterizing important system behaviours, such as passivity, and has close connections with passivity theorem, bounded real lemma, Kalman--Yakubovich lemma, and the circle criterion [20-21]. On the other hand, passivity is part of a broader and general theory of dissipativeness. The main idea of passivity theory is that the passive properties of a system can keep the system internally stable. In recent years, dissipated and passivity results for NNs are established in [22-26].

Since Zhang et al. [27] has pointed out that the relaxed passivity conditions for NNs with time-varying delays. New delay-dependent passive criterion is obtained in terms of linear matrix inequalities, which guarantees that the input and output of the considered NNs. Also, Wei et al. [28] shown the passivity problem by using value-map and suitable for Lyaponov-Krasovskii function. Recently, Zeng et al. [29] derived new passivity conditions for NNs with time-varying delays and norm-bounded parameter uncertainties using with the complete delay-decomposing approach. Also, the problem of robust passivity analysis of uncertain NNs with discrete and distributed time-varying delays has been reported, by constructing an augmented Lyapunov functional and combining a new integral inequality with the reciprocally convex approaches respectively. In addition, Park M.J. et al. [30] and Li et al. [31] developed generalized free-matrix-base integral inequality for enhanced passivity condition and derive in form of linear matrix inequalities (LMIs) as complex value. In [32-34], proposed the global asymptotic stability problem for recurrent NNs with multiple time-varying delays. Using the free-weighting matrix technique and incorporating the interconnected information between the upper bounds of multiple time-varying delays.

Recall the past several years. However, there are fewer works have been done on the dynamics of stability, dissipativity, and passivity analysis of DNNs with time-varying delays. Motivated by these earlier efforts, in this paper, we are concerned with the problem of stability, dissipativity, and passivity analysis of discrete-time neural networks with time-varying delays. Based on the newly established integral inequality, a class of new Lyapunov functional including is proposed, and some less conservative delay range-dependent stability, dissipativity and passivity are derived in terms of LMIs.

This paper is organized as follows. Preliminaries is formulates the problem under consideration. Stability, Dissipativity, and passivity conditions for stability, dissipativity, and passivity analysis of DNNs with time-varying delays are derived in main results. Finally, conclusions are drawn in last section.

#### **1-1-Preliminaries**

The following notation will be used in this paper.  $R^+$  denotes the set of all real non-negative numbers;  $\mathbb{R}^n$  denotes the n-dimensional space with the scalar product x, y or  $x^T y$  of two vectors x, y, and the vector norm  $\|.\|$ ;  $M^{n \times r}$  denotes the space of all matrices of  $(n \times r)$ -dimensions.  $A^T$  denotes the transpose of matrix A; A is symmetric if  $A = A^T$ ; I denotes the identity matrix; Matrix A is called semi-positive definite  $(A \ge 0)$  if  $\langle Ax, x \rangle \ge 0$ , for all  $x \in \mathbb{R}^n$ ; A is positive definite (A > 0) if  $\langle Ax, x \rangle > 0$  for all  $x \neq 0$ ; A > B means A - B > 0. The notation diag{...} stands for a block-diagonal matrix. The symmetric term in a matrix is denoted by \* First point.

**Lemma 1.1** ([38]) For a positive definite matrix R > 0 and any sequence of discrete-time variables  $y: [-h, 0] \cap Z \rightarrow R^n$ , the following inequality holds:

$$\sum_{i=-h+1}^{0} \Delta x(i)^{T} R \Delta x(i) \geq \frac{1}{h} \Theta_{0}^{T} R \Theta_{0} + \frac{3}{h} \frac{h+1}{h-1} \Omega_{1}^{T} R \Omega_{1} + \frac{5(h+1)(h+2)}{(h-2)(h-1)} \Omega_{2}^{T} R \Omega_{2},$$

where  $\Theta_0 = x(0) - x(-h), \Omega_1 = x(0) + x(-h) - \frac{2}{h+1} \sum_{k=-h}^0 x(k), \Omega_2 = x(0) - x(-h) + \frac{6h}{(h+1)(h+2)} \sum_{i=-h}^0 x(i) - \frac{12}{(h+1)(h+2)} \sum_{i=-h+1}^0 \sum_{k=i}^0 x(k).$ 

**Lemma 1.2** ([38]) For a positive definite matrix R > 0 and any sequence of discrete-time variables  $y: [-h, 0] \cap Z \rightarrow R^n$ , the following inequality holds:

$$\sum_{i=-h+1}^{0} \sum_{k=i}^{0} \Delta x(k)^{T} R \Delta x(k) \ge \frac{2(h+1)}{h} \left[ x(0) - \frac{1}{(h+1)} \sum_{i=-h}^{0} x(i) \right]^{T} R \left[ x(0) - \frac{1}{(h+1)} \sum_{i=-h}^{0} x(i) \right] \\ + \frac{4(h+1)(h+2)}{h(h-1)} \Omega_{4}^{T} R \Omega_{4},$$
  
Where  $\Omega_{4} = \left[ x(0) + \frac{2}{(h+1)} \sum_{i=-h}^{0} x(i) - \frac{6}{(h+1)(h+2)} \sum_{i=-h}^{0} \sum_{k=i}^{0} x(k) \right].$ 

**Definition 1.3** ([37]) The neural network (1) is said to be (Q, S, R) –dissipative, if the following dissipation inequality

$$\sum_{k_0}^{\kappa_p} r(u(k), y(k)) \ge 0, \quad \forall k_p \ge 0,$$

Holds under zero initial condition for any nonzero input  $u \in l_2[0, +\infty)$ . Furthermore, if for some scalar  $?\gamma > 0$ , the dissipation inequality

$$\sum_{k_0}^{k_p} r(u(k), y(k)) \ge \gamma \sum_{k_0}^{k_p} u^T(k) u(k), \quad \forall k_p \ge 0,$$

Holds under zero initial condition for any nonzero input  $u \in l_2[0, +\infty)$ , then the neural network (1) is said to be strictly  $(Q, S, R) - \gamma$  -dissipative. In this paper, we define a quadratic supply rate r(u, y) associated with neural network (1) as follows:

$$r(u, y) = y^T Q y + 2y^T S u + u^T R u,$$

Where  $Q \le 0, S$ , and R are real symmetric matrices of appropriate dimensions.

# **2- Results and Discussion**

In this section, we will establish a new criterion on dissipativity analysis of DNNs with time-varying delays;

$$\begin{aligned} x(k+1) &= Cx(k) + D_0 f(x(k)) + D_1 g(x(k-h(k)) + u(k), \quad k \ge 0, \\ y(k) &= f(x(k)), \\ x(k) &= \varphi(k), k \in [-h_2, 0]. \end{aligned}$$
(1)

Where  $x(k) = [x_1(k), x_2(k), ..., x_n(k)]^T \in \mathbb{R}^n$  is the state of the neural,  $u(k) \in \mathbb{R}^n$  is the input belonging to  $l_2, \varphi$  is the initial value, n is the number of neurals,

$$f(x(k)) = [f_1(x_1(k)), f_2(x_2(k)), \dots, f_n(x_n(k))]^T, \quad g(x(k-h(k))) = [g_1(x_1(k-h(k)), g_2(x_2(k-h(k)), \dots, g_n(x_n(k-h(k)))]^T, \text{ are the activation functions;}$$

 $C = \text{diag}(\overline{c_1}, \overline{c_2}, \dots, \overline{c_n}), \overline{a_i} > 0$  represents the self-feedback term,  $D_0, D_1$  denote the connection weights, the discretely delayed connection weights and the distributively delayed connection weight. For every  $k \ge 0$ , the variable delay h(k) is defined to be a positive integer and  $h(k) \in [h_1, h_2], \forall k \ge 0$  for some integers  $h_2 \ge h_1 > 1$ .

Assumption 2.1 ([35]) For any  $s_1, s_2 \in R$ ,  $s_1 \neq s_2$ , the continuous and bounded activation functions;

 $\hat{f}_i(\cdot)$  and  $\hat{g}_i(\cdot)$  satisfy

$$F_i^- \le \frac{\hat{f}_i(s_1) - \hat{f}_i(s_2)}{s_1 - s_2} \le F_i^+,$$
  

$$G_i^- \le \frac{\hat{g}_i(s_1) - \hat{g}_i(s_2)}{s_1 - s_2} \le G_i^+, \quad i = 1, 2, ..., n, \text{ where } F_i^-, F_i^+, G_i^-, \text{ and } G_i^+ \text{ are known constants.}$$

The following notations are needed;

$$h_{12} = h_2 - h_1, e_i = \left[\underbrace{0, 0 \dots, \stackrel{i}{\hat{T}}, \dots 0}_{10}\right]_{10n \times n}^T, i = 1, 2, \dots, 10, y(k) = x(k) - x(k-1),$$
  
$$\xi(k) = [x^T(k), x^T(k-h_1), x^T(k-h(k)), x^T(k-h_2), f(x(k)), g(x(k-h(k)), \frac{1}{h_1+1}\sum_{i=k-h_1}^k x^T(i), x^T(k-h_2), f(x(k)), y(k) = x(k) - x(k-1),$$

$$\begin{split} \frac{1}{h(k)-h_1+1} \sum_{l=k-h(k)}^{k-h_1} x^T(i), \frac{1}{h_2-h(k)+1} \sum_{l=k-h_2}^{k-h(k)} x^T(i), \sum_{l=-h_1+1}^{0} \sum_{j=k+l}^{k} x^T(j), u^T(k)]^T, \\ \alpha(k) &= \left[x^T(k), \sum_{l=k-h_1}^{k-1} x^T(l), \sum_{l=k-h_2}^{k-h_1-1} x^T(l), \sum_{l=-h_1+1}^{0} \sum_{j=k+l}^{k} x^T(j)\right]^T, \\ Z_{10} &= diag \left\{Z_1, \frac{3(h_1+1)}{h_1-1} Z_1, \frac{5(h_1+1)(h_1+2)}{(h_1-2)(h_1-1)} Z_1\right\}, Z_2^* &= \begin{bmatrix}Z_2 & 0\\ 0 & 3Z_2\end{bmatrix}, Z_{20} &= \begin{bmatrix}Z_2^* & X_2^*\\ Z_2^*\end{bmatrix}, \\ \Pi_0 &= \begin{bmatrix}C, 0, 0, 0, D_0, D_1, 0, 0, 0, l\end{bmatrix}^T, \\ \Pi_1 &= \begin{bmatrix}\Pi_0, (h_1+1)e_7 - e_2, (h(k) - h_1+1)e_8 + (h_2 - h(k) + 1)e_9 - e_3 - e_4, e_{10} + h_1\Pi_0 - (h_1+1)e_7 + e_2], \\ \Pi_2 &= \begin{bmatrix}e_1, (h_1+1)e_7 - e_1, (h(k) - h_1 + 1)e_8 + (h_2 - h(k) + 1)e_9 - e_3 - e_2, e_{10}], \\ \Pi_3 &= \begin{bmatrix}C - I, 0, 0, 0, D_0, D_1, 0, 0, 0, I\end{bmatrix}^T, \\ \Pi_4 &= \begin{bmatrix}e_1 - e_2, e_1 + e_2 - 2e_7, e_1 - e_2 + \frac{6h_1}{h_1+2}e_7 - \frac{12}{(h_1+1)(h_1+2)}e_{10}\end{bmatrix}, \\ \Pi_5 &= \begin{bmatrix}e_3 - e_4, e_3 + e_4 - 2e_9, e_2 - e_3, e_2 + e_3 - 2e_8], \\ \Pi_5 &= e_1 + \left(2 - \frac{6}{(h_1+2)}\right)e_7 - \frac{6}{(h_1+1)(h_1+2)}e_{10}, \\ \Xi_1 &= \Pi_1P\Pi_1^T - \Pi_2P\Pi_2^T, \\ \Xi_2 &= e_1Q_1e_1^T - e_2Q_1e_2^T + e_2Q_2e_2^T - e_4Q_2e_4^T, \\ \Xi_3 &= \Pi_3(h_1^2Z_1 + h_{12}^2Z_2)\Pi_3^T - \Pi_4Z_10\Pi_4^T - \Pi_5Z_20\Pi_5^T, \\ \Xi_4 &= \frac{h_1(h_1+1)}{2}\Pi_3Z_3\Pi_3^T - \frac{2(h_1+1)}{h_1}\Pi_6Z_3\Pi_6^T - \frac{4(h_1^2-1)}{h_1(h_1+2)}\Pi_7Z_3\Pi_7^T, \\ \Xi_5 &= -e_3G_1\Lambda_2e_3^T + 2e_3G_2\Lambda_2e_6^T - e_6\Lambda_2e_6^T, \\ \Xi_7 &= -e_5\Psi_1e_5^T - 2e_5\Psi_2e_{10}^T - e_{10}\Psi_3e_{10}^T, \\ \Xi_4 &= \frac{h_2(h_1+1)}{2}H_2(h_2e_3^T + 2e_3G_2\Lambda_2e_6^T - e_6\Lambda_2e_6^T, \\ \Xi_7 &= -e_5\Psi_1e_5^T - 2e_5\Psi_2e_{10}^T - e_{10}\Psi_3e_{10}^T, \\ \Xi_4 &= \frac{h_2(h_1+1)}{2}H_2(h_2e_3^T + 2e_3G_2\Lambda_2e_6^T - e_6\Lambda_2e_6^T, \\ \Xi_7 &= -e_5\Psi_1e_5^T - 2e_5\Psi_2e_{10}^T - e_{10}\Psi_3e_{10}^T, \\ \Xi_4 &= \frac{h_2(h_1+1)}{2}H_2(h_2e_3^T + 2e_3G_2\Lambda_2e_6^T - e_6\Lambda_2e_6^T, \\ \Xi_7 &= -e_5\Psi_1e_5^T - 2e_5\Psi_2e_{10}^T - e_{10}\Psi_3e_{10}^T, \\ \Xi_4 &= \frac{h_2(h_1+1)}{2}H_2(h_2e_3^T + 2e_3G_2\Lambda_2e_6^T - e_6\Lambda_2e_6^T, \\ \Xi_7 &= -e_5\Psi_1e_5^T - 2e_5\Psi_2e_{10}^T - e_{10}\Psi_3e_{10}^T, \\ \Xi_8 &= \frac{h_2(h_2e_3^T - 2e_5\Psi_2e_{10}^T - e_{10}\Psi_3e_{10}^T, \\ \Xi_8 &= \frac{h_2(h_2e_3^T - 2e_5\Psi_2e_6^T - e_6\Lambda_2e_6^T, \\ \Xi_7 &= -e_5\Psi_1e_5^T - 2e_5\Psi$$

**Theorem 2.2** For given integer  $h_1$ ,  $h_2$  satisfying  $1 < h_1 \le h_2$ , matrices  $\Psi_1$ ,  $\Psi_2$ , and  $\Psi_3$  with  $\Psi_1$ ,  $\Psi_2$ , and  $\Psi_3$  being real symmetric, system (1) is stability, passivity, and dissipativity for  $h_1 < h(k) \le h_2$ , if there are positive define matrices  $P \in R^{4n \times 4n}, Z_1 \in R^{n \times n}, Z_2 \in R^{n \times n}, Z_3 \in R^{n \times n}, Q_1 \in R^{n \times n}, Q_2 \in R^{n \times n}$ , and any matrix  $X \in R^{2n \times 2n}$ , diagonal matrices  $\Lambda_1 > 0, \Lambda_2 > 0$  of appropriate dimensions such that the following three LMIs are satisfied.

$$\Xi < 0, \qquad Z_{20} \ge 0.$$
 (2)

Proof. Choose a Lyapunov functional candidate as follows:

$$V(k) = \sum_{j=1}^{4} V_j(k),$$
(3)  
Where  $V_1(k) = \alpha^T(k) P \alpha(k),$ 

Where 
$$V_1(I)$$

$$V_{2}(k) = \sum_{i=k-h_{1}}^{k-1} x^{T}(i)Q_{1}x(i) + \sum_{i=k-h_{2}}^{k-h_{1}-1} x^{T}(i)Q_{2}x(i),$$

$$V_{3}(k) = h_{1}\sum_{i=-h_{1}+1}^{0} \sum_{j=k+i}^{k} y^{T}(i)Z_{1}y(i) + h_{12}\sum_{i=-h_{2}+1}^{-h_{1}} \sum_{j=k+i}^{k} y^{T}(i)Z_{2}y(i),$$

$$V_{4}(k) = \sum_{i=-h_{1}+1}^{0} \sum_{j=i}^{k} \sum_{u=k+j}^{k} y^{T}(u)Z_{3}y(u).$$
(4)

Next, we calculate the difference of V(k). For  $V_1(k)$  and  $V_2(k)$ , we have:

$$\Delta V_1(k) = \xi^T(k) \Xi_1 \xi(k) \tag{5}$$

and

$$\Delta V_2(k) = \xi^T(k) \Xi_2 \xi(k). \tag{6}$$

Calculating  $V_3(k)$  gives;

$$\Delta V_3(k) = h_1^2 y_{k+1}^T Z_1 y_{k+1} + h_{12}^2 y_{k+1}^T Z_2 y_{k+1} - h_1 \sum_{i=k-h_1+1}^k y^T(i) Z_1 y(i) - h_{12} - h_{12} \sum_{i=k-h_2+1}^{k-h_1} y^T(i) Z_2 y(i).$$
(7)

By Lemma1.1, we get;

$$-h_1 \sum_{i=k-h_1+1}^k y^T(i) Z_1 y(i) = -\xi^T(k) \Pi_4 Z_{10} \Pi_4^T \xi(k).$$
(8)

Under the condition of  $Z_{20} > 0$ , by Lemma2.1 and the lower bounded lemma, we get;

$$-h_{12}\sum_{i=k-h_2+1}^{k} y^T(i)Z_2 y(i) = -\xi^T(k)\Pi_5 Z_{20}\Pi_5^T \xi(k).$$
(9)

Then we have;  $\Delta V_3(k) = \xi^T(k) \Xi_3 \xi(k).$ (10) Calculating  $\Delta V_4(k)$  gives;

$$\Delta V_4(k) = \frac{h_1(h_1+1)}{2} y_{k+1}^T Z_3 y_{k+1} - \sum_{i=h_1+1}^0 \sum_{i=k+i}^k y^T(j) Z_3 y(j).$$
(11)

By Lemma1.2, we have;

$$-\sum_{i=h_{1}+1}^{0}\sum_{i=k+i}^{k}y^{T}(j)Z_{3}y(j) \leq \xi^{T}(k)\left(-\frac{2(h_{1}+1)}{h_{1}}\Pi_{6}Z_{3}\Pi_{6}^{T}-\frac{4(h_{1}+1)(h_{2}+2)}{h_{1}(h_{1}+1)}\Pi_{7}Z_{3}\Pi_{7}^{T}\right)\xi(k).$$
(12)

Then we have;

$$\Delta V_4(k) = \xi^T(k) \Xi_4 \xi(k). \tag{13}$$

From Assumption 2.1, we have;

$$\begin{bmatrix} x(k) \\ f(x(k)) \end{bmatrix}^{I} \begin{bmatrix} F_{1}\Lambda_{1} & -F_{2}\Lambda_{1} \\ -F_{2}\Lambda_{1} & \Lambda_{1} \end{bmatrix}^{T} \begin{bmatrix} x(k) \\ f(x(k)) \end{bmatrix} \leq 0,$$
(14)

$$\begin{bmatrix} x(k-h(k)) \\ g(x(k-h(k))) \end{bmatrix}^{T} \begin{bmatrix} G_{1}\Lambda_{1} & -G_{2}\Lambda_{1} \\ -G_{2}\Lambda_{1} & \Lambda_{1} \end{bmatrix}^{T} \begin{bmatrix} x(k-h(k)) \\ g(x(k-h(k))) \end{bmatrix} \leq 0,$$
(15)

Where

$$\begin{split} \Lambda_{1} &= diag\{\lambda_{11}, \lambda_{12}, \dots, \lambda_{1n}\}, \Lambda_{2} = diag\{\lambda_{21}, \lambda_{22}, \dots, \lambda_{2n}\}, \\ F_{1} &= diag\{F_{1}^{-}F_{1}^{+}, F_{2}^{-}F_{2}^{+}, \dots, F_{n}^{-}F_{n}^{+}\},_{2} = diag\{\frac{F_{1}^{-}+F_{1}^{+}}{2}, \frac{F_{2}^{-}+F_{2}^{+}}{2}, \dots, \frac{F_{n}^{-}+F_{n}^{+}}{2}\}, \\ G_{1} &= diag\{G_{1}^{-}G_{1}^{+}, G_{2}^{-}G_{2}^{+}, \dots, G_{n}^{-}G_{n}^{+}\},_{2} = diag\{\frac{G_{1}^{-}+G_{1}^{+}}{2}, \frac{G_{2}^{-}+G_{2}^{+}}{2}, \dots, \frac{G_{n}^{-}+G_{n}^{+}}{2}\}. \end{split}$$

Since  $\Delta x_k = x_{k+1} - x_k$ , by introducing relaxation matrices  $W_1, W_2$  with appropriate dimensions, we obtain the following zero equation;

$$2(\Delta^{T}(k)Z_{1}^{T} + x^{T}(k)Z_{2}^{T})[Cx(k) + D_{0}f(x(k)) + D_{1}g(x(k-h(k)) + u(k) - x(k) - \Delta^{T}(k)] = 0.$$
(16)

Define

$$J(i) = y^{T}(i)\Psi_{1}y(i) + 2y^{T}(i)\Psi_{2}u(i) + u^{T}(i)\Psi_{3}u(i).$$
(17)

Adding the equations from (5) to (17), then we can get the upper bound of  $\Delta V(k) - J(k)$  as

$$\Delta V(k) - J(k) \le \xi^{T}(k) \sum_{i=1}^{7} \Xi_{i} \xi(k) = \xi^{T}(k) \Xi \xi(k).$$
(18)  
If  $\Xi < 0$ , then  $\Delta V(k) - J(k) < 0$ .

This completes the proof of Theorem2.2.

**Remark 2.3** In the above Theorem, we analyzed the passivity and dissipavity for DNN (1). In the following corollary analyze the asymptotic stability for NN:

$$\begin{aligned} x(k+1) &= Cx(k) + D_0 f(x(k)) + D_1 g(x(k-h(k))), \quad k \ge 0, \\ x(k) &= \varphi(k), k \in [-h_2, 0]. \end{aligned}$$
(19)

The NN (19) is a special case of (1) when u(k) = y(k) = 0.

The following notations are needed.

$$\begin{aligned} h_{12} &= h_2 - h_1, e_i = \left[ \underbrace{0, 0 \dots, \stackrel{i}{T}, \dots 0}_{9} \right]_{9n \times n}^T, i = 1, 2, \dots, 9, y(k) = x(k) - x(k-1), \\ \xi(k) &= [x^T(k), x^T(k-h_1), x^T(k-h(k)), x^T(k-h_2), f(x(k)), g(x(k-h(k))), \frac{1}{h_1 + 1} \sum_{i=k-h_1}^k x^T(i), \\ &\frac{1}{h(k) - h_1 + 1} \sum_{i=k-h(k)}^{k-h_1} x^T(i), \frac{1}{h_2 - h(k) + 1} \sum_{i=k-h_2}^{k-h(k)} x^T(i), \sum_{i=-h_1+1}^0 \sum_{j=k+i}^k x^T(j)]^T, \\ \alpha(k) &= \left[ x^T(k), \sum_{i=k-h_1}^{k-1} x^T(i), \sum_{i=k-h_2}^{k-h_1 - 1} x^T(i), \sum_{i=-h_1+1}^0 \sum_{j=k+i}^k x^T(j) \right]^T, \end{aligned}$$

$$\begin{split} Z_{10} &= diag \left\{ Z_1, \frac{3(h_1+1)}{h_1-1} Z_1, \frac{5(h_1+1)(h_1+2)}{(h_1-2)(h_1-1)} Z_1 \right\}, Z_2^* = \begin{bmatrix} Z_2 & 0 \\ 0 & 3Z_2 \end{bmatrix}, Z_{20} = \begin{bmatrix} Z_2^* & X \\ & Z_2^* \end{bmatrix}, \\ \Pi_0 &= \begin{bmatrix} C, 0, 0, 0, D_0, D_1, 0, 0, 0 \end{bmatrix}^T, \\ \Pi_1 &= \begin{bmatrix} \Pi_0, (h_1+1)e_7 - e_2, (h(k) - h_1 + 1)e_8 + (h_2 - h(k) + 1)e_9 - e_3 - e_4, e_{10} + h_1\Pi_0 - (h_1+1)e_7 + e_2 \end{bmatrix}, \\ \Pi_2 &= \begin{bmatrix} e_1, (h_1+1)e_7 - e_1, (h(k) - h_1 + 1)e_8 + (h_2 - h(k) + 1)e_9 - e_3 - e_2, e_{10} \end{bmatrix}, \\ \Pi_3 &= \begin{bmatrix} C - I, 0, 0, 0, D_0, D_1, 0, 0, 0 \end{bmatrix}^T, \\ \Pi_4 &= \begin{bmatrix} e_1 - e_2, e_1 + e_2 - 2e_7, e_1 - e_2 + \frac{6h_1}{h_1+2}e_7 - \frac{12}{(h_1+1)(h_1+2)}e_{10} \end{bmatrix}, \\ \Pi_5 &= \begin{bmatrix} e_3 - e_4, e_3 + e_4 - 2e_9, e_2 - e_3, e_2 + e_3 - 2e_8 \end{bmatrix}, \\ \Pi_6 &= e_1 - e_7, \\ \Pi_7 &= e_1 + \left( 2 - \frac{6}{(h_1+2)} \right)e_7 - \frac{6}{(h_1+1)(h_1+2)}e_{10}, \\ \Xi_1 &= \Pi_1P\Pi_1^T - \Pi_2P\Pi_2^T, \\ \Xi_2 &= e_1Q_1e_1^T - e_2Q_1e_2^T + e_2Q_2e_2^T - e_4Q_2e_4^T, \\ \Xi_3 &= \Pi_3(h_1^2Z_1 + h_{12}^2Z_2)\Pi_3^T - \Pi_4Z_{10}\Pi_4^T - \Pi_5Z_{20}\Pi_5^T, \\ \Xi_4 &= \frac{h_1(h_1+1)}{2}\Pi_3Z_3\Pi_3^T - \frac{2(h_1+1)}{h_1}\Pi_6Z_3\Pi_6^T - \frac{4(h_1^2-1)}{h_1(h_1+2)}\Pi_7Z_3\Pi_7^T, \\ \Xi_5 &= -e_3G_1\Lambda_2e_3^T + 2e_3G_2\Lambda_2e_6^T - e_6\Lambda_2e_6^T, \\ \Theta &= \sum_{i=1}^6 \Xi_i. \end{split}$$

**Corollary 2.4** For given integer  $h_1$ ,  $h_2$  satisfying  $0 < h_1 < h_2$  system (19) is asymptotically stable for  $h_1 < h(k) \le h_2$ , if there are positive define matrices  $P \in R^{4n \times 4n}, Z_1 \in R^{n \times n}, Z_2 \in R^{n \times n}, Z_3 \in R^{n \times n}, Q_1 \in R^{n \times n}, Q_2 \in R^{n \times n}$ , and any matrix  $X \in R^{2n \times 2n}$ , diagonal matrices  $\Lambda_1 > 0, \Lambda_2 > 0$  of appropriate dimensions such that the following three LMIs are satisfied.

$$\Theta < 0, \qquad Z_{20} \ge 0. \tag{20}$$

Summing up the above analysis, some sufficient conditions on finite-time stability for dissipativity and passivity analysis of discrete-time neural networks with time-varying delays (1) with (2), (3) are obtained. In the following, we mainly focus on stabilizing by transforming the sufficient conditions into solvable linear matrix inequalities.

# **3-** Conclusion

In this paper, stability, dissipativity and passivity analysis of discrete-time neural networks with time-varying delays was studied. A delay-dependent condition has been provided to ensure the considered discrete-time neural networks with time-varying delays to be stability, dissipativity and passivity. An effective LMI approach has been proposed to derive the stability, dissipativity and passivity criterion. Based on the appropriate type of Lyapunov functional, a sufficient condition for the solvability of this problem is established for the stability, dissipativity and passivity criterion.

## 4- Acknowledgments

The author would to acknowledge Phuket Rajabhat university for providing the financial support. The author are greatly thankful to Assoc. Prof. Dr. Kreangkri Ratchagit, Maejo University.

# **5- Conflict of Interest**

The author declares that there is no conflict of interests regarding the publication of this manuscript. In addition, the ethical issues, including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, and redundancies have been completely observed by the authors.

# **6- References**

[1] Haykin, S. "Neural Networks: A Comprehensive Foundation, Prentice Hall PTR." Upper Saddle River, NJ, USA (1998).

- [2] Wangli He, and Jinde Cao. "Exponential Synchronization of Hybrid Coupled Networks with Delayed Coupling." IEEE Transactions on Neural Networks 21, no. 4 (April 2010): 571–583. doi:10.1109/tnn.2009.2039803.
- [3] Popov, Vasile Mihai, and Radu Georgescu. "Hyperstability of control systems." Springer-Verlag, 1973.
- [4] Willems, J.C., Dissipative dynamical systems part I: general theory, Arch. Ration. Mech. Anal. 45 (5) (1972) 321-351. doi:10.1109/9780470544334.ch20.
- [5] Zhang, Dan, and Li Yu. "Exponential State Estimation for Markovian Jumping Neural Networks with Time-Varying Discrete and Distributed Delays." Neural Networks 35 (November 2012): 103–111. doi:10.1016/j.neunet.2012.08.005.
- [6] Dan Zhang, Li Yu, Qing-Guo Wang, and Chong-Jin Ong. "Estimator Design for Discrete-Time Switched Neural Networks With Asynchronous Switching and Time-Varying Delay." IEEE Transactions on Neural Networks and Learning Systems 23, no. 5 (May 2012): 827–834. doi:10.1109/tnnls.2012.2186824.

- [7] Chen, Guoliang, Jianwei Xia, Ju H. Park, and Guangming Zhuang. "New Delay-Dependent Global Robust Passivity Analysis for Stochastic Neural Networks with Markovian Jumping Parameters and Interval Time-Varying Delays." Complexity 21, no. 6 (April 8, 2015): 167–179. doi:10.1002/cplx.21677.
- [8] Raja, R., and R. Samidurai. "New Delay Dependent Robust Asymptotic Stability for Uncertain Stochastic Recurrent Neural Networks with Multiple Time Varying Delays." Journal of the Franklin Institute 349, no. 6 (August 2012): 2108–2123. doi:10.1016/j.jfranklin.2012.03.007.
- [9] Shi, Kaibo, Hong Zhu, Shouming Zhong, Yong Zeng, and Yuping Zhang. "New Stability Analysis for Neutral Type Neural Networks with Discrete and Distributed Delays Using a Multiple Integral Approach." Journal of the Franklin Institute 352, no. 1 (January 2015): 155–176. doi:10.1016/j.jfranklin.2014.10.005.
- [10] Zeng, Hong-Bing, Ju H. Park, Chang-Fan Zhang, and Wei Wang. "Stability and Dissipativity Analysis of Static Neural Networks with Interval Time-Varying Delay." Journal of the Franklin Institute 352, no. 3 (March 2015): 1284–1295. doi:10.1016/j.jfranklin.2014.12.023.
- [11] Mohamad, S., and K. Gopalsamy. "Exponential Stability of Continuous-Time and Discrete-Time Cellular Neural Networks with Delays." Applied Mathematics and Computation 135, no. 1 (February 2003): 17–38. doi:10.1016/s0096-3003(01)00299-5.
- [12] Q.C. Zhong, Robust Control of Time-Delay Systems, Springer-Verlag, London, 2006.
- [13] M. Vidyasagar, Nonlinear Systems Analysis, Prentice-Hall, Englewood Cliffs, NJ, 1993.
- [14] J. Sarangapani, Neural Network Control of Nonlinear Discrete-Time Systems, Taylor and Francis Group, Boca Raton, London, New York, 2006.
- [15] J.C. Willems, The Analysis of Feedback Systems, The MIT Press, Cambridge, 1971.
- [16] W.M. Hadda, V.S. Chellaboina, Nonlinear Dynamical Systems and Control: A Lyapunov-based Approach, Princeton University Press, USA, 2008.
- [17] Shen, Hao, Ju H. Park, Lixian Zhang, and Zheng-Guang Wu. "Robust Extended Dissipative Control for Sampled-Data Markov Jump Systems." International Journal of Control 87, no. 8 (February 26, 2014): 1549–1564. doi:10.1080/00207179.2013.878478.
- [18] Shen, Hao, Yanzheng Zhu, Lixian Zhang, and Ju H. Park. "Extended Dissipative State Estimation for Markov Jump Neural Networks With Unreliable Links." IEEE Transactions on Neural Networks and Learning Systems 28, no. 2 (February 2017): 346–358. doi:10.1109/tnnls.2015.2511196.
- [19] D. Hill, P. Moylan, The stability of nonlinear dissipative systems, IEEE Trans. Autom. Control 21 (5) (1976) 708-711.
- [20] Shen, Hao, Zheng-Guang Wu, and Ju H. Park. "Reliable Mixed Passive and H∞ Filtering for Semi-Markov Jump Systems with Randomly Occurring Uncertainties and Sensor Failures." International Journal of Robust and Nonlinear Control 25, no. 17 (October 15, 2014): 3231–3251. doi:10.1002/rnc.3255.
- [21] Shen, Hao, Zheng-Guang Wu, Ju H. Park, and Zhengqiang Zhang. "Extended Dissipativity-Based Synchronization of Uncertain Chaotic Neural Networks with Actuator Failures." Journal of the Franklin Institute 352, no. 4 (April 2015): 1722–1738. doi:10.1016/j.jfranklin.2015.01.026.
- [22] Zheng-Guang Wu, Peng Shi, Hongye Su, and Jian Chu. "Passivity Analysis for Discrete-Time Stochastic Markovian Jump Neural Networks With Mixed Time Delays." IEEE Transactions on Neural Networks 22, no. 10 (October 2011): 1566–1575. doi:10.1109/tnn.2011.2163203.
- [23] Zheng-Guang Wu, Peng Shi, Hongye Su, and Jian Chu. "Dissipativity Analysis for Discrete-Time Stochastic Neural Networks With Time-Varying Delays." IEEE Transactions on Neural Networks and Learning Systems 24, no. 3 (March 2013): 345–355. doi:10.1109/tnnls.2012.2232938.
- [24] Zeng, Hong-Bing, Ju H. Park, and Hao Shen. "Robust Passivity Analysis of Neural Networks with Discrete and Distributed Delays." Neurocomputing 149 (February 2015): 1092–1097. doi:10.1016/j.neucom.2014.07.024.
- [25] Zeng, Hong-Bing, Ju H. Park, Chang-Fan Zhang, and Wei Wang. "Stability and Dissipativity Analysis of Static Neural Networks with Interval Time-Varying Delay." Journal of the Franklin Institute 352, no. 3 (March 2015): 1284–1295. doi:10.1016/j.jfranklin.2014.12.023.
- [26] Zhang, Baoyong, Shengyuan Xu, and James Lam. "Relaxed Passivity Conditions for Neural Networks with Time-Varying Delays." Neurocomputing 142 (October 2014): 299–306. doi:10.1016/j.neucom.2014.04.031.
- [27] Zeng, Hong-Bing, Yong He, Min Wu, and Shen-Ping Xiao. "Passivity Analysis for Neural Networks with a Time-Varying Delay." Neurocomputing 74, no. 5 (February 2011): 730–734. doi:10.1016/j.neucom.2010.09.020.
- [28] Li, Ning, and Wei Xing Zheng. "Passivity Analysis for Quaternion-Valued Memristor-Based Neural Networks with Time-Varying Delay." IEEE Transactions on Neural Networks and Learning Systems (2019): 1–12. doi:10.1109/tnnls.2019.2908755.

- [29] Zeng, Hong-Bing, Yong He, Min Wu, and Hui-Qin Xiao. "Improved Conditions for Passivity of Neural Networks With a Time-Varying Delay." IEEE Transactions on Cybernetics 44, no. 6 (June 2014): 785–792. doi:10.1109/tcyb.2013.2272399.
- [30] Kang, Wei, and Shouming Zhong. "Passivity Analysis of Discrete-Time Memristive Neural Networks with Time-Varying Delay." 2017 9th International Conference on Intelligent Human-Machine Systems and Cybernetics (IHMSC) (August 2017). doi:10.1109/ihmsc.2017.177.
- [31] Park, M.J., O.M. Kwon, and J.H. Ryu. "Passivity and Stability Analysis of Neural Networks with Time-Varying Delays via Extended Free-Weighting Matrices Integral Inequality." Neural Networks 106 (October 2018): 67–78. doi:10.1016/j.neunet.2018.06.010.
- [32] Chen, Yonggang, Shumin Fei, and Yongmin Li. "Improved Delay-Dependent Stability Conditions for Recurrent Neural Networks with Multiple Time-Varying Delays." Nonlinear Dynamics 78, no. 2 (June 24, 2014): 803–812. doi:10.1007/s11071-014-1478-y.
- [33] Liu, Pin-Lin. "Improved Delay-Derivative-Dependent Stability Analysis for Generalized Recurrent Neural Networks with Interval Time-Varying Delays." Neural Processing Letters (August 10, 2019). doi:10.1007/s11063-019-10088-8.
- [34] Wang, Liyan, Yang Wang, and Jianxia Liu. "Delay Dependent Robust Stability of Neural Networks with Time-Varying Delays." 2018 Chinese Control and Decision Conference (CCDC) (June 2018). doi:10.1109/ccdc.2018.8407796.
- [35] Liu, Yurong, Zidong Wang, and Xiaohui Liu. "Global Exponential Stability of Generalized Recurrent Neural Networks with Discrete and Distributed Delays." Neural Networks 19, no. 5 (June 2006): 667–675. doi:10.1016/j.neunet.2005.03.015.
- [36] Zheng-Guang Wu, Peng Shi, Hongye Su, and Jian Chu. "Passivity Analysis for Discrete-Time Stochastic Markovian Jump Neural Networks with Mixed Time Delays." IEEE Transactions on Neural Networks 22, no. 10 (October 2011): 1566–1575. doi:10.1109/tnn.2011.2163203.
- [37] Bo Zhou, and Qiankun Song. "Boundedness and Complete Stability of Complex-Valued Neural Networks with Time Delay." IEEE Transactions on Neural Networks and Learning Systems 24, no. 8 (August 2013): 1227–1238. doi:10.1109/tnnls.2013.2247626.
- [38] Wang, Feng-Xian, Xin-Ge Liu, Mei-Lan Tang, and Yan-Jun Shu. "Stability Analysis of Discrete-Time Systems with Variable Delays via Some New Summation Inequalities." Advances in Difference Equations 2016, no. 1 (April 5, 2016). doi:10.1186/s13662-016-0829-z.