

Caudrelier, V. & Crampe, N. (2008). Symmetries of Spin Calogero Models. *Symmetry, Integrability and Geometry: Methods and Applications (SIGMA)*, 4, pp. 1-16.



**CITY UNIVERSITY
LONDON**

[City Research Online](#)

Original citation: Caudrelier, V. & Crampe, N. (2008). Symmetries of Spin Calogero Models. *Symmetry, Integrability and Geometry: Methods and Applications (SIGMA)*, 4, pp. 1-16.

Permanent City Research Online URL: <http://openaccess.city.ac.uk/161/>

Copyright & reuse

City University London has developed City Research Online so that its users may access the research outputs of City University London's staff. Copyright © and Moral Rights for this paper are retained by the individual author(s) and/ or other copyright holders. Users may download and/ or print one copy of any article(s) in City Research Online to facilitate their private study or for non-commercial research. Users may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain. All material in City Research Online is checked for eligibility for copyright before being made available in the live archive. URLs from City Research Online may be freely distributed and linked to from other web pages.

Versions of research

The version in City Research Online may differ from the final published version. Users are advised to check the Permanent City Research Online URL above for the status of the paper.

Enquiries

If you have any enquiries about any aspect of City Research Online, or if you wish to make contact with the author(s) of this paper, please email the team at publications@city.ac.uk.

Symmetries of Spin Calogero Models^{*}

Vincent CAUDRELIER[†] and Nicolas CRAMPÉ[‡]

[†] *Centre for Mathematical Science, City University,
Northampton Square, London, EC1V 0HB, United Kingdom*
E-mail: v.caudrelier@city.ac.uk

[‡] *International School for Advanced Studies, Via Beirut 2-4, 34014 Trieste, Italy*
E-mail: crampe@sissa.it

Received September 24, 2008, in final form December 17, 2008; Published online December 23, 2008
Original article is available at <http://www.emis.de/journals/SIGMA/2008/090/>

Abstract. We investigate the symmetry algebras of integrable spin Calogero systems constructed from Dunkl operators associated to finite Coxeter groups. Based on two explicit examples, we show that the common view of associating one symmetry algebra to a given Coxeter group W is wrong. More precisely, the symmetry algebra heavily depends on the representation of W on the spins. We prove this by identifying two different symmetry algebras for a B_L spin Calogero model and three for G_2 spin Calogero model. They are all related to the half-loop algebra and its twisted versions. Some of the result are extended to any finite Coxeter group.

Key words: Calogero models; symmetry algebra; twisted half-loop algebra

2000 Mathematics Subject Classification: 70H06; 81R12; 81R50

1 Introduction

Dunkl operators [1] were introduced by C.F. Dunkl as part of a program on polynomials in several variables with reflection symmetries related to finite reflection groups (or equivalently finite Coxeter groups). Besides this important mathematical motivation, they quickly became fundamental objects in the study of integrable quantum mechanical many-body systems introduced by F. Calogero and B. Sutherland [2, 3] and generalized to any Weyl group in [4]. Indeed they can be viewed as generalizations of momentum operators with which they share the crucial property of forming an Abelian algebra. This fact allows one to implement the exchange operator formalism described in [5] producing many-body integrable Hamiltonians. In [6, 7], it was realized that this could be pushed further by representing the Coxeter group on spins¹. This gave L -particle integrable spin Calogero systems related to the group A_L . Then one could play several games such as changing the type of potential and/or the underlying group while maintaining integrability.

Another big step was taken in [8] where the “FRT formalism” [9], so powerful with spin chain type integrable models, was used in connection with the algebraic properties of the Dunkl operators to construct the symmetry algebra. By this, we mean the algebra of the operators commuting with the Hamiltonian of the system (and its hierarchy). In the case of [8], it is a non Abelian infinite dimensional algebra which contains the whole hierarchy of the integrable A_L spin Sutherland model in its centre: the Yangian of \mathfrak{gl}_N [10]. This type of symmetry algebra is a very efficient tool to study and, in particular, to compute explicitly the spectrum, the eigenvectors and the correlation functions of a large class of integrable models: Sutherland model [11, 12,

^{*}This paper is a contribution to the Special Issue on Dunkl Operators and Related Topics. The full collection is available at <http://www.emis.de/journals/SIGMA/DunklOperators.html>

¹Here the spin is to be understood as an internal degree of freedom taking N different values.

13], Haldane–Shastry spin model [14, 15, 16, 17], non-linear Schrödinger equation [18, 19] and Hubbard model [20].

The method developed in [8] was then derived in various ways, again accommodating various groups and potentials, to identify the corresponding symmetry algebras. From there, following several results in the literature, some standard views imposed themselves as granted: for instance, the symmetry algebra of an A_L Sutherland type model [8] should be a Yangian while the Calogero counterpart (without harmonic confinement) should be the corresponding half-loop algebra. Similarly, for B_L Sutherland models, the reflection algebra [21] should be symmetry algebra [22] with the corresponding twisted half-loop algebra as symmetry algebra of the Calogero counterpart (this paper).

Actually, we show in this paper that this is not true and in general, it is not enough to specify the type of potential and the underlying Coxeter group to identify the correct symmetry algebra of the associated integrable model. In addition, one has to give the spin representation of the Coxeter group. This is exemplified in this paper by working with rational Dunkl operators (hence producing Calogero type potentials) based on two particular finite Coxeter groups: B_L and G_2 . In the first three sections, we recall the background on finite reflection groups, half-loop algebras and the link to dynamical spin integrable systems that is necessary for our purposes. In Section 5, we deal with B_L spin Calogero model for which we identify two different symmetry algebras by choosing two different ways of representing B_L on the spin configuration space. In Section 6, we apply the same strategy with three different spin representations of G_2 to obtain three different symmetry algebras. All the symmetry algebras are related to the \mathfrak{gl}_N half-loop algebra and its twisted versions. Various generalizations of our discussion are collected in Section 7. Our conclusions are gathered in the last section.

2 Dunkl operators associated to finite reflection groups

Here we recall the ingredients we need from the theory of finite reflection groups (see e.g. [23]) and Dunkl operators [1]. A reflection in the real Euclidean space \mathbb{R}^L endowed with the scalar product $(e_i, e_j) = \delta_{ij}$, where $\{e_i\}_{i=1, \dots, L}$ are the canonical basis vectors of \mathbb{R}^L , is a linear operator s_α ($\alpha \in \mathbb{R}^L, \alpha \neq 0$) on \mathbb{R}^L defined by

$$\forall \mu \in \mathbb{R}^L, \quad s_\alpha(\mu) = \mu - 2 \frac{(\mu, \alpha)}{(\alpha, \alpha)} \alpha.$$

It sends the vector α to $-\alpha$ and leaves invariant the hyperplane H_α orthogonal to α . A root system Φ is a subset of \mathbb{R}^L satisfying, for all $\alpha \in \Phi$,

$$\begin{aligned} (\lambda \in \mathbb{R}, \alpha \in \Phi \quad \text{and} \quad \lambda\alpha \in \Phi) &\Rightarrow \lambda = \pm 1, \\ s_\alpha \Phi &= \Phi. \end{aligned}$$

Φ is a finite set and the group generated by $\{s_\alpha \mid \alpha \in \Phi\}$ is a finite subgroup W of $O(\mathbb{R}^L)$ called the reflection group associated to Φ . Any root system can be written as $\Phi = \Phi^+ \cup (-\Phi^+)$ (Φ^+ is called the positive root system). A simple system Δ is a subset of Φ such that Δ is a vector space basis of $\text{span}_{\mathbb{R}} \Phi$ and each $\alpha \in \Phi$ is a linear combination of Δ with coefficients all of the same sign.

An important result is that the reflection group W is generated by $\{s_\alpha \mid \alpha \in \Delta\}$ subject only to the relations, for $\alpha, \beta \in \Delta$

$$(s_\alpha s_\beta)^{m(\alpha, \beta)} = 1$$

with $m(\alpha, \alpha) = 1$ and $m(\alpha, \beta) > 1$ ($\alpha \neq \beta$). That is to say, by definition, W is a Coxeter group. The possible values of $m(\alpha, \beta)$ such that the group W is finite provides a classification of the finite reflection groups.

To each Coxeter group W , one can associate Dunkl operators which are differential operators acting on functions $\varphi : \mathbb{R}^L \rightarrow \mathbb{C}$. Let us define the natural action of W on such functions by

$$(\hat{s}\varphi)(\mu) = \varphi(s^{-1}(\mu))$$

with $s \in W$ and $\mu \in \mathbb{R}^L$. The map $s \mapsto \hat{s}$ is a representation of the group W . A function $k : \Phi \rightarrow \mathbb{C}$ is called a multiplicity function if

$$\forall \alpha, \beta \in \Phi, \quad k(\beta) = k(s_\alpha(\beta)).$$

Let k be given. For $\xi \in \mathbb{R}^L$, the Dunkl operator d_ξ acts on $C^1(\mathbb{R}^L)$ and is defined by

$$d_\xi = -i\partial_\xi + i \sum_{\alpha \in \Phi^+} k(\alpha) \frac{(\alpha, \xi)}{(\alpha, x)} \hat{s}_\alpha,$$

where ∂_ξ is the derivative in the direction ξ , $x = (x_1, \dots, x_L)^t$ with x_i the operator multiplication by x_i . Note that our choice of Dunkl operators yields hermitian operators provided k is real-valued. This is important for quantum mechanical applications. The Dunkl operators do not depend on the choice of the positive root system Φ^+ and have the following fundamental properties²

Proposition 1 ([1]). *The Dunkl operators are W -equivariant, i.e. for any $s \in W$, we get*

$$\hat{s}d_\xi \hat{s}^{-1} = d_{s(\xi)}.$$

The Dunkl operators commute, i.e. for any $\xi, \zeta \in \mathbb{R}^L$, we get

$$d_\xi d_\zeta = d_\zeta d_\xi.$$

3 Half-loop algebra and twists

This section is meant to give necessary definitions and notations to handle algebraic structures related to some integrable models.

The \mathfrak{gl}_N half-loop algebra³ is the unital algebra $\mathfrak{gl}_N[z]$ of polynomials in an indeterminate z with coefficients in \mathfrak{gl}_N : it is generated by $\{e_{ij} z^n \mid 1 \leq i, j \leq N, n = 0, 1, 2, \dots\}$ where e_{ij} are the generators of \mathfrak{gl}_N satisfying the commutation relations

$$[e_{ij}, e_{kl}] = \delta_{jk} e_{il} - \delta_{il} e_{kj}.$$

Let E_{ij} be the matrix with 1 on the entry (i, j) and 0 elsewhere. The map $e_{ij} \mapsto E_{ij}$ provides a representation of \mathfrak{gl}_N . In the following, we will use auxiliary space notations to simplify computations. We define the monodromy matrix of the half-loop algebra as follows

$$T(u) = \sum_{i,j=1}^N E_{ij} \otimes \sum_{n=0}^{\infty} \frac{e_{ji} z^n}{u^{n+1}} = \sum_{i,j=1}^N \frac{E_{ij} \otimes e_{ji}}{u - z} \in \text{End}(\mathbb{C}^N) \otimes \mathfrak{gl}_N[z][[u^{-1}]],$$

where u is a formal parameter called the spectral parameter and $\text{End}(\mathbb{C}^N)$ is the auxiliary space used to pack nicely the generators. The defining relations of the half-loop algebra can be written

$$[T_a(u), T_b(v)] = [T_a(u) + T_b(v), r_{ab}(u - v)], \quad (3.1)$$

²Strictly speaking, the proof in [1] is given for slightly different Dunkl operators simply related to ours by the gauge transformation $d_\xi \mapsto \phi^{-1}(x) d_\xi \phi(x)$ where ϕ is a W -invariant function $\phi(s(x)) = \phi(x)$ for all $s \in W$.

³Also sometimes called Gaudin algebra.

where a and b denote two copies of the auxiliary space $\text{End}(\mathbb{C}^N)$, $T_a(u) = T(u) \otimes \mathbb{I}_N$, $T_b(u) = \mathbb{I}_N \otimes T(u)$, \mathbb{I}_N is the identity in $\text{End}(\mathbb{C}^N)$, and

$$r_{ab}(u) = \sum_{i,j=1}^N \frac{E_{ij} \otimes E_{ji}}{u} \equiv \frac{P_{ab}}{u}$$

is a classical r -matrix solution of the classical Yang–Baxter equation [24].

Let $\sigma \in \text{End}(\mathbb{C}^N)$ with $\sigma^n = \mathbb{I}_N$ for some $n \in \mathbb{N}$. The eigenvalues of σ are the powers of the n -th root of unity $\tau = e^{\frac{2i\pi}{n}}$. We denote A the inner automorphism of \mathfrak{gl}_N associated to σ . Its action on the generators of \mathfrak{gl}_N is given by

$$A : e_{ij} \mapsto (\sigma)_{jn} e_{mn} (\sigma^{-1})_{mi}.$$

We extend this action to $\mathfrak{gl}_N[z]$ in the natural way and we define

$$\mathfrak{gl}_N[z]^\sigma = \{M(z) \in \mathfrak{gl}_N[z] \mid AM(z) = M(\tau z)\}.$$

One can see that $\mathfrak{gl}_N[z]^\sigma$ is a subalgebra of $\mathfrak{gl}_N[z]$ which is called *the (inner) twisted half-loop algebra of order n* . Now we proceed to derive the commutation relations of this algebra. For $k \in \mathbb{N}$, we introduce the projectors

$$\mathcal{P}_k = \frac{1}{n} \sum_{j=0}^{n-1} \tau^{-jk} A^j.$$

Note that there are only n such projectors as $\mathcal{P}_{n+k} = \mathcal{P}_k$. This allows us to define a surjective projection map from $\mathfrak{gl}_N[z]$ to $\mathfrak{gl}_N[z]^\sigma$ by $e_{ij}z^k \mapsto \mathcal{P}_k e_{ij}z^k$. In turn, this maps $T(u)$ to

$$B(u) = \frac{1}{n} \sum_{j=0}^{n-1} \tau^j \sigma^j T(\tau^j u) \sigma^{-j} \in \text{End}(\mathbb{C}^N) \otimes \mathfrak{gl}_N[z]^\sigma[[u^{-1}]], \quad (3.2)$$

which contains the generators of $\mathfrak{gl}_N[z]^\sigma$ by construction. Next we have [25]

Proposition 2. *$B(u)$ satisfies the symmetry property*

$$\forall j = 0, \dots, n-1, \quad B(u) = \tau^j \sigma^j B(\tau^j u) \sigma^{-j},$$

and encodes the commutation relations of $\mathfrak{gl}_N[z]^\sigma$ as

$$[B_a(u), B_b(v)] = \frac{1}{n} \sum_{j=0}^{n-1} \left[\tau^j B_a(u) + B_b(v), \frac{(\sigma^j)_b P_{ab}(\sigma^{-j})_b}{u - \tau^j v} \right]. \quad (3.3)$$

For our purposes, the following facts are important. If $b(u) = \text{Tr} B(u)$, where the trace is taken over the auxiliary space $\text{End}(\mathbb{C}^N)$, then it follows from (3.3) that

$$[b(u), B(v)] = 0, \quad [b(u), b(v)] = 0. \quad (3.4)$$

4 Link with integrable dynamical spin models

The Dunkl operators allow us to construct integrable Hamiltonians by implementing the strategy discussed in [5]. Indeed, any polynomial in the Dunkl operators commutes with the independent and two by two commuting operators $\{d_{e_i} \mid i = 1, \dots, L\}$. When it is of order 2, this polynomial can usually be regarded as an Hamiltonian which is then integrable.

The next step is to consider Hamiltonians for particles with N internal degrees of freedom. The construction to introduce these “spins” has been pioneered in [6, 7] and the symmetry of the corresponding models has been shown in [8]. The point is that the Dunkl operators only act on the positions x_i while now the wavefunctions become $\varphi(x|s)$, $x \in \mathbb{R}^L$, $s \in (\mathbb{C}^N)^{\otimes M}$. Let us remark that M may be different from L in general. So one has to come up with a method allowing the Dunkl operators to act on the spins while maintaining their nice properties. This is done with a suitable projector

$$\Lambda = \frac{1}{|W|} \sum_{w \in W} \hat{w} R_w.$$

Note that we require that the wavefunctions satisfy the following generalized statistics⁴ for $w \in W$,

$$\hat{w} R_w \varphi(x|s) = \varphi(x|s),$$

where $R : W \rightarrow \text{End}((\mathbb{C}^N)^{\otimes M})$ is a representation of W .

One then acts with $d_\xi \Lambda$ on the wavefunctions. Unfortunately, usually $[d_\xi \Lambda, d_\zeta \Lambda] \neq 0$ so the idea to use directly polynomials in $d_\xi \Lambda$ to get an integrable hierarchy fails. The idea of [8] is to use the monodromy matrix formalism and the underlying algebraic structures to circumvent this problem. In our context, consider a set of M vectors $\{\xi_1, \dots, \xi_M\}$. Then, the matrix

$$T_a(u) = \sum_{k=1}^M \frac{P_{ak}}{u - d_\xi \xi_k}$$

provides a representation of the half-loop algebra (3.1). At this point, let us emphasize that the auxiliary space has no ‘physical’ meaning (in the sense it does not act on the wavefunctions) but it is a powerful and convenient tool to deal with the infinite algebras we are interested in. Then, the ‘physical’ operators are contained in the entries of the matrix.

One obtains $B(u)$ through (3.2) with the properties (3.4). The crucial part is to show that, for a suitable choice of ξ and of the twist, $B(u)\Lambda$ is also a representation of the algebra (3.3). To prove this statement, it is sufficient to show that $\Lambda B(u)\Lambda = B(u)\Lambda$ which is guaranteed by, for g any generator of W ,

$$\hat{g} R_g B(u)\Lambda = B(u)\Lambda. \tag{4.1}$$

Finally, using properties (3.4) for $B(u)\Lambda$, we conclude that $\text{Tr } B(u)\Lambda$ provides a hierarchy of commuting operators when expanding in powers of u^{-1} . To prove the integrability, it is now sufficient to prove that in this hierarchy there are L independent quantities. Noting that these operators are polynomials in d_ξ times Λ , we may choose one of them as the Hamiltonian acting both on the position and spin degrees of freedom of the wavefunctions. As a by-product, but not the least, we prove that this Hamiltonian has the twisted half-loop algebra for symmetry algebra.

It is very important to realize that it is the choice of R , the representation of W on the spins, together with the requirement (4.1), that imposes the form of $B(u)$ and hence the symmetry algebra. This is the essential message of this paper.

In the rest of this paper, we detail this construction for two Coxeter groups, B_L and G_2 . In particular, we get the surprising result that spin Calogero-type integrable models built from Dunkl operators associated to the same Coxeter group can have different symmetry algebras.

⁴Called this way because in the case $W = A_L$ and with the permutation representation on positions and spins, these conditions amount to consider bosonic wavefunctions.

5 Reflection group B_L

We consider the Coxeter group B_L . It is generated by L elements t_1, \dots, t_{L-1} (generating the Coxeter group A_{L-1}) and r with defining relations

$$\begin{aligned} r^2 &= 1, & t_i^2 &= 1, & i &= 1, \dots, L-1, \\ (t_i t_{i+1})^3 &= 1, & i &= 1, \dots, L-2, & (t_i t_j)^2 &= 1, & |i-j| > 1, \\ (r t_{L-1})^4 &= 1, & (r t_j)^2 &= 1, & j &\neq L-1. \end{aligned}$$

5.1 The usual B_L spin Calogero model

To fix ideas, we start with a known model [26]. We choose the usual positive root system $\Phi^+ = \{e_i \pm e_j, e_k \mid 1 \leq i < j \leq L, 1 \leq k \leq L\}$. Then, the representation of the generators of B_L associated to simple roots are

$$t_i = s_{e_i - e_{i+1}} = \begin{pmatrix} \mathbb{I}_{i-1} & & & \\ & 0 & 1 & \\ & 1 & 0 & \\ & & & \mathbb{I}_{L-i-1} \end{pmatrix} \quad \text{and} \quad r = s_{e_L} = \begin{pmatrix} \mathbb{I}_{L-1} & \\ & -1 \end{pmatrix},$$

where \mathbb{I}_j is the $j \times j$ unit matrix. In this case, the Dunkl operators are given by

$$d_{e_k} \equiv d_k = -i\partial_{e_k} + ik_l \sum_{j \neq k} \left(\frac{1}{x_k - x_j} \hat{s}_{e_k - e_j} + \frac{1}{x_k + x_j} \hat{s}_{e_k + e_j} \right) + \frac{ik_s}{x_k} \hat{s}_{e_k}. \quad (5.1)$$

The constants k_l and k_s are the two arbitrary values that the multiplicity function can take on B_L . The action of B_L on $(\mathbb{C}^N)^{\otimes L}$ is taken as

$$R_{t_i} = P_{ii+1} \quad \text{and} \quad R_r = Q_L,$$

where $Q \in \text{End}(\mathbb{C}^N)$ satisfies $Q^2 = \mathbb{I}_N$. We apply (3.2) to

$$T_a(u) = \sum_{k=1}^L \frac{P_{ak}}{u - d_k}$$

with $\sigma = Q$ (so $\tau = -1$ and $n = 2$). We get

$$B_a(u) = \frac{1}{2} (T_a(u) - Q_a T_a(-u) Q_a) = \frac{1}{2} \sum_{k=1}^L \left(\frac{P_{ak}}{u - d_k} + \frac{Q_a P_{ak} Q_a}{u + d_k} \right).$$

To prove that $B(u)\Lambda$ satisfies (3.3), it is enough to prove

$$\hat{r} R_r B(u) = B(u) \hat{r} R_r \quad \text{and} \quad \hat{t}_i R_{t_i} B(u) = B(u) \hat{t}_i R_{t_i}. \quad (5.2)$$

This is readily seen as

$$\hat{r} R_r B(u) = \frac{1}{2} \sum_{k=1}^{L-1} \left(\frac{P_{ak}}{u - d_k} + \frac{Q_a P_{ak} Q_a}{u + d_k} \right) \hat{r} R_r + \frac{1}{2} \left(\frac{Q_L P_{aL} Q_L}{u - \hat{r} d_L \hat{r}} + \frac{Q_L Q_a P_{aL} Q_a Q_L}{u + \hat{r} d_L \hat{r}} \right) \hat{r} R_r$$

and $\hat{r} d_L \hat{r} = d_{re_L} = d_{-e_L} = -d_L$ (recall also that $Q^2 = \mathbb{I}_N$). Similarly, one looks at the i -th and $i+1$ -th term in the second relation in (5.2) and uses $\hat{t}_i d_i \hat{t}_i = d_{t_i e_i} = d_{e_{i+1}} = d_{i+1}$ together with $P_{i+1} P_{ai} P_{i+1} = P_{ai+1}$.

One can now extract the commuting quantities by expanding $b(u)\Lambda$ in powers of u^{-1} . One gets

$$b(u)\Lambda = \sum_{n=0}^{\infty} \frac{1 + (-1)^n}{2u^{n+1}} \sum_{k=1}^L d_k^n \Lambda. \quad (5.3)$$

We see that only $J_{2n} = \sum_{k=1}^L d_k^{2n} \Lambda$ yield non trivial quantities. In particular, one gets the usual B_L spin Calogero Hamiltonian

$$H = J_2 = - \sum_{j=1}^L \partial_{e_j}^2 + k_l \sum_{m \neq j} \left(\frac{k_l - P_{mj}}{(x_m - x_j)^2} + \frac{k_l - Q_j P_{mj} Q_j}{(x_m + x_j)^2} \right) + k_s \sum_{j=1}^L \frac{k_s - Q_j}{x_j^2}. \quad (5.4)$$

Then, we get the well-known result [26] that this hamiltonian is integrable by proving that J_2, J_4, \dots, J_{2L} are independent. In addition, as explained in Section 4, we prove that this model has for symmetry the twisted half-loop algebra of order 2. This result may be also obtained by considering the symmetry of the B_L Sutherland model proved in [22] and taking the suitable limit as explained in [8] for the A_L case.

5.2 Another B_L spin Calogero model

Here we follow the approach of [27] and use another representation of B_L for the spins while keeping the same on \mathbb{R}^L . Let us fix $\mu \in \mathbb{R}^L$. The orbit of the vector μ under the group B_L is written

$$\{w(\mu) \mid w \in B_L\} \equiv \{\mu_1, \dots, \mu_M\}$$

for some $M \in \mathbb{N}$ and with $\mu_j \in \mathbb{R}^L$. The group B_L acts transitively on this set, i.e. for any μ_j, μ_k there exists $w \in B_L$ such that $w(\mu_j) = \mu_k$. For any $w \in B_L$, we obtain an action on the set $\{1, \dots, M\}$ by defining

$$\check{w}(i) = j \quad \text{iff} \quad w(\mu_i) = \mu_j. \quad (5.5)$$

The map $w \mapsto \check{w}$ is a representation of B_L . We can now define a representation of B_L on $(\mathbb{C}^N)^{\otimes M}$ as the map $w \mapsto R_w$ where

$$R_w v_1 \otimes \dots \otimes v_M = v_{\check{w}^{-1}(1)} \otimes \dots \otimes v_{\check{w}^{-1}(M)} \quad (5.6)$$

with $v_i \in \mathbb{C}^N$.

At this stage, one can see that

$$T_a(u) = \sum_{k=1}^M \frac{P_{ak}}{u - d_{\mu_k}}$$

commutes with the projector Λ in the chosen representation. Indeed, for each $w \in B_L$

$$\hat{w} R_w \sum_{k=1}^M \frac{P_{ak}}{u - d_{\mu_k}} = \sum_{k=1}^M \frac{P_{a\check{w}(k)}}{u - d_{w(\mu_k)}} \hat{w} R_w = \sum_{n=1}^M \frac{P_{an}}{u - d_{\mu_n}} \hat{w} R_w,$$

where the last equality is obtained by relabelling the sum according to (5.5). Thus, $T(u)\Lambda$ satisfies the half-loop algebra relations and $\text{Tr } T(u)\Lambda \equiv t(u)\Lambda$ provides the commuting elements.

So we find that all the B_L spin Calogero Hamiltonians obtained in this way (e.g. in [27]) have the half-loop algebra as symmetry algebra.

As an example, we apply this to $\mu = e_1$, the first vector of the canonical basis of \mathbb{R}^L . Then the orbit contains $2L$ elements $\pm e_j$, $j = 1, \dots, L$ which we order as follows $(e_1, -e_1, \dots, e_L, -e_L) \equiv (\mu_1, \mu_{\bar{1}}, \dots, \mu_L, \mu_{\bar{L}})$. Then, by inspection we get

$$\check{r}(j) = \begin{cases} \bar{L}, & j = L, \\ L, & j = \bar{L}, \\ j, & \text{otherwise} \end{cases} \quad \text{and} \quad \check{t}_i(j) = \begin{cases} i+1, & j = i, \\ i, & j = i+1, \\ \bar{i}+1, & j = \bar{i}, \\ \bar{i}, & j = \bar{i}+1, \\ j, & \text{otherwise.} \end{cases}$$

This gives

$$R_r = P_{L, \bar{L}}, \quad R_{t_i} = P_{i, i+1} P_{\bar{i}, \bar{i}+1}.$$

We get

$$T_a(u) = \sum_{k=1}^L \frac{P_{ak}}{u - d_{\mu_k}} + \frac{P_{a\bar{k}}}{u - d_{\mu_{\bar{k}}}} = \sum_{k=1}^L \frac{P_{ak}}{u - d_k} + \frac{P_{a\bar{k}}}{u + d_k},$$

where the Dunkl operators are given by (5.1). The following B_L spin Calogero Hamiltonian has the half-loop algebra as symmetry algebra

$$H = - \sum_{j=1}^L \partial_{e_j}^2 + k_l \sum_{m \neq j} \left(\frac{k_l - P_{mj} P_{\bar{m}\bar{j}}}{(x_m - x_j)^2} + \frac{k_l - P_{m\bar{j}} P_{\bar{m}j}}{(x_m + x_j)^2} \right) + k_s \sum_{j=1}^L \frac{k_s - P_{j\bar{j}}}{x_j^2}. \quad (5.7)$$

As previously, $\text{Tr } T(u)\Lambda$ provides only the even conserved quantities $J_{2k} = \sum_{j=1}^L d_j^{2k} \Lambda$ but it is sufficient to prove the integrability since J_2, J_4, \dots, J_{2L} are again independent. Comparing (5.4) and (5.7), it is manifest that the only difference lies in the action on the spins, the potentials being identical. Yet, these two systems based on B_L have different symmetry algebras. It is also interesting to remark that (5.7) represents a system of L particles on the line with two spin degrees of freedom attached to each particle.

6 Group G_2

We consider the Coxeter group G_2 (which is the dihedral group $I_2(6)$ of order 12). It is generated by 2 elements t and r subject only to the following relations

$$r^2 = 1, \quad t^2 = 1, \quad (tr)^6 = 1.$$

It may also be generated by 2 elements a and b subject only to the following relations

$$a^6 = 1, \quad b^2 = 1, \quad ba = a^{-1}b.$$

The isomorphism between the two presentations reads

$$a \mapsto tr \quad \text{and} \quad b \mapsto t.$$

6.1 Model with three particles and six spins

The choice in this section for the positive roots of $I_2(6)$ in \mathbb{R}^3 are $\Phi^+ = \{e_1 - e_2, e_3 - e_1, e_3 - e_2\} \cup \{-2e_1 + e_2 + e_3, e_1 - 2e_2 + e_3, -e_1 - e_2 + 2e_3\}$. Then, the the representation of the generators of $I_2(6)$ associated to simple roots are

$$t = s_{e_1 - e_2} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad r = s_{-2e_1 + e_2 + e_3} = \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & 2 & -1 \\ 2 & -1 & 2 \end{pmatrix}. \quad (6.1)$$

To write down the Dunkl operators explicitly, it is useful to express all the reflections in terms of r and t

$$s_{e_1 - 2e_2 + e_3} = trt, \quad s_{e_1 - e_3} = rtr, \quad s_{e_2 - e_3} = trtrt, \quad s_{e_1 + e_2 - 2e_3} = rtrtr.$$

We give only one example as the other ones are computed similarly:

$$d_{e_1} = -i\partial_{e_1} + ik_s \left(\frac{1}{x_1 - x_2} \hat{t} + \frac{1}{x_1 - x_3} \widehat{rtr} \right) + ik_l \left(\frac{1}{x_1 - 2x_2 + x_3} \widehat{trt} - 2 \frac{1}{-2x_1 + x_2 + x_3} \hat{r} + \frac{1}{x_1 + x_2 - 2x_3} \widehat{rtrtr} \right).$$

The action on the spins is defined as in Section 5.2 replacing the group B_L by $I_2(6)$. That is, we define for $\mu \in \mathbb{R}^3$,

$$\{\mu_1, \dots, \mu_M\} = \{w(\mu) \mid w \in I_2(6)\}$$

which provides a representation R_w acting on spins. Thus, we have again that

$$T_a(u) = \sum_{k=1}^M \frac{P_{ak}}{u - d_{\mu_k}}$$

commutes with the projector Λ in the chosen representation (the proof follows the same lines than the one in Section 5.2). We conclude that the half-loop algebra is the symmetry algebra of the integrable hierarchy contained in $\text{Tr } T(u)\Lambda$.

We give an example. We may choose $\mu = e_1$. Then the orbit is

$$\left\{ \mu_1 = e_1, \mu_2 = e_2, \mu_3 = e_3, \mu_{\bar{1}} = \frac{1}{3}(-e_1 + 2e_2 + 2e_3), \right. \\ \left. \mu_{\bar{2}} = \frac{1}{3}(2e_1 - e_2 + 2e_3), \mu_{\bar{3}} = \frac{1}{3}(2e_1 + 2e_2 - e_3) \right\}.$$

So, we get $M = 6$. We deduce the following representation of $I_2(6)$ on $(\mathbb{C}^N)^{\otimes 6}$

$$R_t = P_{12}P_{\bar{1}\bar{2}} \quad \text{and} \quad R_r = P_{\bar{1}\bar{1}}P_{2\bar{3}}P_{3\bar{2}}.$$

The coefficient in front of u^{-2} in the expansion of $t(u)\Lambda$ is proportional to the total momentum $P = -i\partial_1 - i\partial_2 - i\partial_3$ and the one in front of u^{-3} is proportional to the Hamiltonian and reads

$$H = - \sum_{i=1}^3 \partial_{e_i}^2 + k_s \sum_{m \neq j} \frac{k_s - P_{mj}P_{\bar{m}\bar{j}}}{(x_m - x_j)^2} + k_l \sum_{n \neq m \neq j} \frac{k_l - P_{n\bar{n}}P_{j\bar{m}}P_{m\bar{j}}}{(-2x_n + x_m + x_j)^2}.$$

By direct computation, we can show that the coefficients in front of u^{-4} , u^{-5} and u^{-6} are not independent of P and H whereas J_6 , the one in front of u^{-7} , is independent of them. Then P , H and J_6 provides 3 independent conserved quantities which proves the integrability of H . Let us remark that the degree in d_i of the conserved quantities (i.e. 1, 2, 6) are in agreement with the degrees of the invariant polynomials by the Coxeter group $I_2(6)$. In fact, it is not a coincidence since the conserved quantities we constructed may be seen as polynomials in terms of the variables d_i and are, by construction, invariant under $I_2(6)$.

6.2 Model with three particles and three spins

We keep the same positive roots as in Section 6.1. Then, the action of $I_2(6)$ on \mathbb{R}^3 is still the one defined by (6.1) but we take now the following representation of $I_2(6)$ on $(\mathbb{C}^N)^{\otimes 3}$

$$R_r = P_{23}Q_1Q_2Q_3, \quad R_t = P_{12},$$

where $Q^2 = \mathbb{I}_N$. The projector may be written as follows

$$\begin{aligned} \Lambda &= \Lambda_Q \Lambda_P = \Lambda_P \Lambda_Q, \\ \Lambda_Q &= \frac{1}{2}(1 + (\widehat{tr})^3 Q_1Q_2Q_3), \\ \Lambda_P &= \frac{1}{6}(1 + \widehat{t} P_{12} + \widehat{trtrt} P_{23} + \widehat{rtrt} P_{12}P_{23} + \widehat{trtr} P_{23}P_{12} + \widehat{rtr} P_{13}). \end{aligned}$$

The projector Λ_P is the usual symmetriser of the eigenfunctions for bosons and Λ_Q yields

$$\phi(x_1, x_2, x_3 \mid s_1, s_2, s_3) = \phi(-x_1 + 2X, -x_2 + 2X, -x_3 + 2X \mid s_1^*, s_2^*, s_3^*),$$

where $s^* = Qs$ and $X = \frac{1}{3}(x_1 + x_2 + x_3)$ is the center of mass of the three particles.

The monodromy matrix is taken to be (with $d_i \equiv d_{e_i}$),

$$T_a(u) = \frac{P_{a1}}{u - d_1} + \frac{P_{a2}}{u - d_2} + \frac{P_{a3}}{u - d_3}.$$

We apply a slight variant of (3.2) and define

$$\begin{aligned} \hat{B}_a(u) &= T_a(u) - Q_a T_a(-u + D) Q_a \\ &= \frac{P_{a1}}{u - d_1} + \frac{P_{a2}}{u - d_2} + \frac{P_{a3}}{u - d_3} + \frac{Q_1 P_{a1} Q_1}{u + d_1 - D} + \frac{Q_2 P_{a2} Q_2}{u + d_2 - D} + \frac{Q_3 P_{a3} Q_3}{u + d_3 - D}, \end{aligned}$$

where $D = \frac{2}{3}(d_1 + d_2 + d_3)$. $\hat{B}(u)$ satisfies the following commutation relations

$$[\hat{B}_a(u), \hat{B}_b(v)] = \frac{1}{2} \left[\hat{B}_a(u) + \hat{B}_b(v), \frac{P_{ab}}{u - v} \right] + \frac{1}{2} \left[-\hat{B}_a(u) + \hat{B}_b(v), \frac{Q_a P_{ab} Q_a}{u + v - D} \right]. \quad (6.2)$$

This algebra is isomorphic to the twisted half-loop algebra of order 2: $\hat{B}(u) \mapsto B(u + D/2)$. The crucial points now are that $\hat{B}(u)$ commutes with Λ and we still have $[\text{Tr } \hat{B}(u), \text{Tr } \hat{B}(v)] = 0$. Then, the coefficient of u^{-2} in $b(u)\Lambda$ is proportional to the total momentum $P = -i\partial_1 - i\partial_2 - i\partial_3$ and the one of u^{-3} contains the following Hamiltonian, studied in [28],

$$H = d_1^2 + d_2^2 + d_3^2 = - \sum_{i=1}^3 \partial_{e_i}^2 + k_s \sum_{m \neq j} \frac{k_s - P_{mj}}{(x_m - x_j)^2} + k_l \sum_{n \neq m \neq j} \frac{k_l - P_{mj} Q_n Q_m Q_j}{(-2x_n + x_m + x_j)^2}. \quad (6.3)$$

As before, the coefficients of u^{-4} , u^{-5} and u^{-6} are not independent of P and H . It is again J_6 , the coefficient of u^{-7} , which provides the third independent conserved quantity and allows us to prove that the previous Hamiltonian is integrable.

Concerning the symmetry algebra, one has to be careful since the commutation relations (6.2) depend on the operator D which is proportional to the total momentum. We can talk about the symmetry of the model (6.3) only in the sectors where the total momentum P has a given value p . This is not a problem as the states are defined by three quantum numbers, p being one of them and in the sectors of fixed p , the symmetry algebra is isomorphic to the twisted half-loop algebra. For example, in the center of the mass frame ($p = 0$), the symmetry algebra of (6.3) is the usual twisted half-loop algebra of order 2.

6.3 Model with two particles and two spins

Our choice for the simple roots of $I_2(6)$ in \mathbb{R}^2 is $e_1 - e_2$ and $-e_1 + \cot\left(\frac{\pi}{12}\right)e_2$. Then the other 4 positive roots are

$$e_1 - \tan\left(\frac{\pi}{12}\right)e_2, \quad e_1 + \tan\left(\frac{\pi}{12}\right)e_2, \quad e_1 + e_2, \quad e_1 + \cot\left(\frac{\pi}{12}\right)e_2.$$

Then, the action of both generators of $I_2(6)$ on \mathbb{R}^2 is given explicitly by

$$b = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad a = \begin{pmatrix} \cos\left(\frac{\pi}{3}\right) & -\sin\left(\frac{\pi}{3}\right) \\ \sin\left(\frac{\pi}{3}\right) & \cos\left(\frac{\pi}{3}\right) \end{pmatrix}.$$

It is useful to express all the reflections in terms of a and b

$$\begin{aligned} s_{e_1 - e_2} &= b, & s_{e_1 - \tan\left(\frac{\pi}{12}\right)e_2} &= ab, & s_{e_1 + \tan\left(\frac{\pi}{12}\right)e_2} &= a^2b, \\ s_{e_1 + e_2} &= ba^3 = a^3b, & s_{e_1 + \cot\left(\frac{\pi}{12}\right)e_2} &= ba^2, & s_{-e_1 + \cot\left(\frac{\pi}{12}\right)e_2} &= ba. \end{aligned}$$

Then, for example, we get

$$\begin{aligned} d_{e_1} &= -i\partial_{e_1} + ik_s \left(\frac{1}{x_1 - x_2} \hat{b} + \frac{1}{x_1 + (2 - \sqrt{3})x_2} \hat{a}^2 \hat{b} + \frac{1}{x_1 + (2 + \sqrt{3})x_2} \hat{b} \hat{a}^2 \right) \\ &\quad + ik_l \left(\frac{1}{x_1 - (2 - \sqrt{3})x_2} \hat{a} \hat{b} + \frac{1}{x_1 + x_2} \hat{a}^3 b + \frac{1}{x_1 - (2 + \sqrt{3})x_2} \hat{b} \hat{a} \right), \end{aligned}$$

where we have used $\tan\left(\frac{\pi}{12}\right) = 2 - \sqrt{3}$ and $\cot\left(\frac{\pi}{12}\right) = 2 + \sqrt{3}$. We define the action on $(\mathbb{C}^N)^{\otimes 2}$ by

$$R_a = Q_1 Q_2^{-1}, \quad R_b = P_{12},$$

where $Q^6 = \mathbb{I}_N$. The projector Λ may be written as follows

$$\begin{aligned} \Lambda &= \Lambda_Q \Lambda_P = \Lambda_P \Lambda_Q, & (6.4) \\ \Lambda_P &= \frac{1}{2}(1 + \hat{b} P_{12}), \\ \Lambda_Q &= \frac{1}{6} \left(1 + \hat{a} Q_1 Q_2^{-1} + \hat{a}^2 (Q_1 Q_2^{-1})^2 + \hat{a}^3 (Q_2 Q_1^{-1})^3 + \hat{a}^4 (Q_1 Q_2^{-1})^4 + \hat{a}^5 (Q_2 Q_1^{-1})^5 \right). \end{aligned}$$

So, it is easy to see that the eigenfunctions are totally symmetric $\phi(x_1, x_2 | s_1, s_2) = \phi(x_2, x_1 | s_2, s_1)$ and satisfy the additional relation

$$\phi(x_1, x_2 | s_1, s_2) = \phi\left(\frac{x_1 + \sqrt{3}x_2}{2}, \frac{x_2 - \sqrt{3}x_1}{2} \mid Qs_1, Q^{-1}s_2\right).$$

It turns out to be convenient to map \mathbb{R}^2 into \mathbb{C} . Let us define

$$d = d_{e_1} + id_{e_2} \quad \text{and} \quad \bar{d} = d_{e_1} - id_{e_2},$$

and similarly for $\partial, \bar{\partial}$. We write the Dunkl operators in terms of $z = x_1 + ix_2$ and $\bar{z} = x_1 - ix_2$

$$d = -i\partial + 2 \sum_{j=0}^2 \left(\frac{ik_s}{i\tau^{2j}z + \bar{z}} \hat{b} \hat{a}^{2j} + \frac{ik_l}{i\tau^{2j+1}z + \bar{z}} \hat{b} \hat{a}^{2j+1} \right),$$

$$\bar{d} = -i\bar{\partial} + 2 \sum_{j=0}^2 \left(\frac{ik_s}{z - i\tau^{2j}\bar{z}} \hat{a}^{2j}\hat{b} + \frac{ik_l}{z - i\tau^{2j+1}\bar{z}} \hat{a}^{2j+1}\hat{b} \right),$$

where $\tau = \exp(i\pi/3)$. The action of the generators of $I_2(6)$ on these Dunkl operators are

$$\hat{a}d = \tau^{-1}\hat{a}\hat{d}, \quad \hat{a}\bar{d} = \tau\bar{d}\hat{a}, \quad \hat{b}d = i\bar{d}\hat{b} \quad \text{and} \quad \hat{b}\bar{d} = -i\bar{d}\hat{b}.$$

Now, we introduce

$$T_a(u) = \frac{P_{a1}}{u - d} + \frac{P_{a2}}{u - i\bar{d}}$$

and also

$$\hat{B}_a(u) = \sum_{j=0}^5 \tau^j Q_a^{-j} T_a((u - d\bar{d})\tau^j) Q_a^j = \sum_{j=0}^5 \frac{Q_a^{-j} P_{a1} Q_a^j}{u - d\bar{d} - \tau^{-j}d} + \frac{Q_a^{-j} P_{a2} Q_a^j}{u - d\bar{d} - i\tau^{-j}\bar{d}}.$$

This satisfies the commutation relations of the twisted half-loop algebra of order 6 with a shift of $d\bar{d}$ in the spectral parameters. We can show by direct investigation that $\hat{B}(u)$ commutes with the projector defined by (6.4). As usual, the series $\text{Tr} \hat{B}(u)\Lambda$ provides the conserved quantities. We can show that 2 are independent: the coefficients in front of u^{-2} and u^{-7} proportional respectively to

$$H = d\bar{d} \quad \text{and} \quad J_6 = d^6 - \bar{d}^6 + 2(d\bar{d})^6.$$

Explicitly, the Hamiltonian is

$$H = -\partial_z \partial_{\bar{z}} + 4i \sum_{j=0}^2 \left(k_s \tau^{2j} \frac{Q_1^{-2j} P_{12} Q_1^{2j} - k_s}{(z - i\tau^{2j}\bar{z})^2} + k_l \tau^{2j+1} \frac{Q_1^{-(2j+1)} P_{12} Q_1^{2j+1} - k_l}{(z - i\tau^{2j+1}\bar{z})^2} \right).$$

As in Section 6.2, the commutation relations satisfied by $\hat{B}(u)$ depend on one operator of our theory (H). Thus, in each sector of fixed energy, the symmetry algebra of the integrable hierarchy is isomorphic to the twisted half-loop algebra of order 6.

7 Generalizations

Let us remark that the construction of Section 6.3 for $I_2(6)$ extends to $I_2(m)$, a presentation of which is

$$a^m = 1, \quad b^2 = 1, \quad ba = a^{-1}b.$$

The results follow by taking Q with $Q^m = \mathbb{I}_N$ and $\tau = e^{\frac{2i\pi}{m}}$. In the expressions for the Dunkl, one has to modify the sum and the two independent operators are to be found at orders u^3 and u^{m+1} . The relevant symmetry algebra is the twisted half-loop algebra of order m .

Another general result holds: the construction of the spin representation described in Sections 5.2 and 6.1 applies to any finite Coxeter group W as explained in [27]. So fixing μ and denoting $\{\mu_1, \dots, \mu_M\}$ the orbit of μ under W , the representation R as defined in (5.6) is always well-defined on $(\mathbb{C}^N)^{\otimes M}$ and the monodromy matrix

$$T_a(u) = \sum_{k=1}^M \frac{P_{ak}}{u - d_{\mu_k}}$$

commutes with the projector Λ in the chosen representation. So we find that all the W spin Calogero Hamiltonians obtained in this way have the half-loop algebra as symmetry algebra.

At this stage, it is worth mentioning that potentially physically interesting integrable Hamiltonians can always be constructed. Indeed, from the general theory of invariant polynomials under finite Coxeter groups, the polynomial of order two in the variables is always invariant. This means that Hamiltonians of the form

$$-\sum_{j=1}^L \partial_{e_i}^2 + \text{interactions terms}$$

can always be extracted from the hierarchy. We note however that M can be different from L so these systems may represent particles with more than one spin attached to them.

8 Conclusions

We have shown that, as far as symmetry is concerned, speaking of a W spin Calogero model is not enough. Even if the root structure of W provides the form of the interaction term in Hamiltonians constructed from Dunkl operators based on W , the symmetry algebra heavily depends on the choice of representation of W on the spins. Using the method pioneered in [8], we showed this explicitly for the case of the Coxeter groups B_L and $I_2(6)$. For the former, two different symmetry algebras were constructed while we proposed three of them for the latter. We stress that in all cases, the identification of the symmetry algebra is an original result in itself.

It should be noted that one systematic method, based on [27], always yields the same symmetry algebra (here the half-loop algebra for Calogero type models) for all Coxeter groups. However, there seems to be no systematic understanding of how to produce other possibilities such as those of Section 5.1 for B_L and Sections 6.2, 6.3 for G_2 . In physical terms, such an understanding would allow one to control the number of spins M and make it coincide (if desired) with the number of particles L . Indeed, it all amounts to finding a construction of spin representations based on orbits of conveniently chosen subgroups of W . We hope to come back to this question in the future.

Other directions of investigation could involve the so-called Sutherland models, whose traditional study involves the same tools as described in this paper. However, it is expected that novel algebraic structures should be understood first. Indeed, already in the A_L case, the symmetry algebra for spin Sutherland models is related to the Yangian of \mathfrak{gl}_N , a deformation of the half-loop algebra. For the B_L case, the reflection algebra appears (when choosing the same spin representation as in Section 5.1) which is a deformation of the twisted half-loop algebra of order 2. However, for higher order twisting, the corresponding deformations are not known.

Finally, the symmetry algebras of the different spin Calogero models obtained in this paper might be useful to compute their eigenstates. Indeed, in the case of the A_L Sutherland model, the Yangian symmetry was the cornerstone of the explicit construction of the eigenvectors [11, 12]. It is based on representation theory of the Yangian and, in particular, on the construction of the Gelfand–Zetlin bases [29] which uses the knowledge of the maximal Abelian subalgebra containing the center. Let us remark that strangely enough, the situation in the case of the half-loop algebra seems more complicated. Indeed, the maximal Abelian subalgebra of the half-loop algebra has been discovered 10 years after the one of the Yangian [30]. Thus, we hope that the present paper provides one additional motivation to look for the maximal Abelian subalgebra of the twisted cases.

Acknowledgements

N.C. would like to thank the hospitality of the Centre for Mathematical Science, City University, where this work was initiated.

References

- [1] Dunkl C.F., Differential-difference operators associated to reflection groups, *Trans. Amer. Math. Soc.* **311** (1989), 167–183.
- [2] Calogero F., Solution of a three-body problem in one dimension, *J. Math. Phys.* **10** (1969), 2191–2196.
Calogero F., Ground state of a one-dimensional N -body system, *J. Math. Phys.* **10** (1969), 2197–2200.
Calogero F., Solution of the one-dimensional N -body problems with quadratic and/or inversely quadratic pair potentials, *J. Math. Phys.* **12** (1971), 419–436, Erratum, *J. Math. Phys.* **37** (1996), 3646.
- [3] Sutherland B., Quantum many-body problem in one dimension: ground state, *J. Math. Phys.* **12** (1971), 246–250.
Sutherland B., Quantum many-body problem in one dimension: thermodynamics, *J. Math. Phys.* **12** (1971), 251–256.
Sutherland B., Exact results for a quantum many-body problem in one dimension, *Phys. Rev. A* **4** (1971), 2019–2021.
- [4] Olshanetsky M.A., Perelomov A.M., Quantum integrable systems related to Lie algebras, *Phys. Rep.* **94** (1983), 313–404.
- [5] Polychronakos A.P., Exchange operator formalism for integrable systems of particles, *Phys. Rev. Lett.* **69**, 703–705.
- [6] Minahan J.A., Polychronakos A.P., Integrable systems for particles with internal degrees of freedom, *Phys. Lett. B* **302** (1993), 265–270, [hep-th/9206046](#).
- [7] Hikami K., Wadati M., Integrability of Calogero–Moser spin system, *J. Phys. Soc. Japan* **62** (1993), 469–472.
- [8] Bernard D., Gaudin M., Haldane F.D.M., Pasquier V., Yang–Baxter equation in long-range interacting systems, *J. Phys. A: Math. Gen.* **26** (1993), 5219–5236, [hep-th/9301084](#).
- [9] Faddeev L.D., Reshetikhin N.Y., Takhtajan L.A., Quantization of Lie groups and Lie algebras, *Leningrad Math. J.* **1** (1990), 193–225.
- [10] Drinfel’d V.G., Hopf algebras and the quantum Yang–Baxter equation, *Sov. Math. Dokl.* **32** (1985), 254–258.
Drinfel’d V.G., Quantum groups, in Proceedings of the International Congress of Mathematicians, Vols. 1, 2 (Berkeley, Calif., 1986), Amer. Math. Soc., Providence, RI, 1987, 798–820.
- [11] Takemura K., Uglov D., The orthogonal eigenbasis and norms of eigenvectors in the spin Calogero–Sutherland model, *J. Phys. A: Math. Gen.* **30** (1997), 3685–3717, [solv-int/9611006](#).
- [12] Takemura K., The Yangian symmetry in the spin Calogero model and its applications, *J. Phys. A: Math. Gen.* **30** (1997), 6185–6204, [solv-int/9701015](#).
- [13] Uglov D., Yangian Gelfand–Zetlin bases, \mathfrak{gl}_N -Jack polynomials and computation of dynamical correlation functions in the spin Calogero–Sutherland model, *Comm. Math. Phys.* **191** (1998), 663–696, [hep-th/9702020](#).
- [14] Ha Z.N.C., Haldane F.D.M., On models with inverse-square exchange, *Phys. Rev. B* **46** (1992), 9359–9368, [cond-mat/9204017](#).
- [15] Haldane F.D.M., Ha Z.N.C., Talstra J.C., Bernard D., Pasquier V., Yangian symmetry of integrable quantum chains with long-range interactions and a new description of states in conformal field theory, *Phys. Rev. Lett.* **69** (1992), 2021–2025.
- [16] Bernard D., Pasquier V., Serban D., A one dimensional ideal gas of spinons, or some exact results on the XXX spin chain with long range interaction, [hep-th/9311013](#).
- [17] Hikami K., Yangian symmetry and Virasoro character in a lattice spin system with long-range interactions, *Nuclear Phys. B* **441** (1995), 530–548.
- [18] Murakami S., Wadati M., Connection between Yangian symmetry and the quantum inverse scattering method, *J. Phys. A: Math. Gen.* **29** (1996), 7903–7915.
- [19] Mintchev M., Ragoucy E., Sorba P., Zaugg Ph., Yangian symmetry in the nonlinear Schrödinger hierarchy, *J. Phys. A: Math. Gen.* **32** (1999), 5885–5900, [hep-th/9905105](#).

- [20] Uglov D.B., Korepin V.E., The Yangian symmetry of the Hubbard model, *Phys. Lett. A* **190** (1994), 238–242, [hep-th/9310158](#).
- [21] Sklyanin E.K., Boundary conditions for integrable quantum systems, *J. Phys. A: Math. Gen.* **21** (1988), 2375–2389.
- [22] Caudrelier V., Crampé N., Integrable N -particle Hamiltonians with Yangian or reflection algebra symmetry, *J. Phys. A: Math. Gen.* **7** (2004), 6285–6298, [math-ph/0310028](#).
- [23] Humphreys J.H., Reflection groups and Coxeter groups, *Cambridge Studies in Advanced Mathematics*, Vol. 29, Cambridge University Press, Cambridge, 1990.
- [24] Belavin A.A., Drinfel'd V.G., Solutions of the classical Yang–Baxter equation for simple Lie algebras, *Funct. Anal. Appl.* **16** (1982), 159–180.
Belavin A.A., Drinfel'd V.G., Classical Yang–Baxter equation for simple Lie algebras, *Funct. Anal. Appl.* **17** (1983), 220–221.
Belavin A.A., Drinfel'd V.G., Triangle equation and simple Lie algebras, *Soviet Sci. Rev. Sect. C Math. Phys. Rev., Mathematical Physics Reviews*, Vol. 4, Harwood Academic Publ., Chur, 1984, 93–165.
- [25] Crampé N., Young C.A.S., Integrable models from twisted half-loop algebras, *J. Phys. A: Math. Theor.* **40** (2007), 5491–5509, [math-ph/0609057](#).
- [26] Yamamoto T., Multicomponent Calogero model of B_N -type confined in a harmonic potential, *Phys. Lett. A* **208** (1995), 293–302, [cond-mat/9508012](#).
- [27] Inozemtsev V.I., Sasaki R., Universal Lax pairs for spin Calogero–Moser models and spin exchange models, *J. Phys. A: Math. Gen.* **34** (2001), 7621–7632, [hep-th/0106164](#).
- [28] Quesne C., An exactly solvable three-particle problem with three-body interaction, *Phys. Rev. A* **55** (1997), 3931–3934, [hep-th/9612173](#).
- [29] Nazarov M., Tarasov V., Yangians and Gelfand–Zetlin bases, *Publ. Res. Inst. Math. Sci.* **30** (1994), 459–478, [hep-th/9302102](#).
- [30] Talalaev D., Quantization of the Gaudin system, [hep-th/0404153](#).