

Guadecitabine and Carboplatin in Ovarian Cancer

A randomized phase 2 trial of epigenetic priming with guadecitabine and carboplatin in platinum-resistant, recurrent ovarian cancer

Amit Oza, Princess Margaret Cancer Centre, University Health Network, University of Toronto, Toronto Canada

Ursula Matulonis, Dana-Farber Cancer Institute, Boston MA

Angeles Alvarez Secord, Duke Cancer Institute, Durham NC

Lynda Roman, USC Norris Comprehensive Cancer Center, Los Angeles CA

Sarah Blagden, University of Oxford, Oxford UK

Susana Banerjee, Royal Marsden NHS Foundation Trust, London UK

John Nemunaitis, University of Toledo College of Medicine and Life Sciences, Toledo, OH

William McGuire, Virginia Commonwealth University, Richmond VA

Sharad Ghamande, Augusta University (Georgia Regents University), Augusta GA

Michael Birrer, University of Alabama at Birmingham, Birmingham AL

Gini F Fleming, The University of Chicago Medicine, Chicago IL

Merry Jennifer Markham, University of Florida, Gainesville FL

Hal Hirte, Juravinski Cancer Centre, Hamilton, Canada

Diane Provencher, Centre Hospitalier de l'Université de Montréal, Université de Montréal, Montréal, Canada

Bristi Basu, Department of Oncology, University of Cambridge and Cambridge University Hospitals NHS Foundation Trust, Cambridge UK

Rebecca Kristeleit, Cancer Institute University College London, UCL, UK

Deborah Armstrong, Johns Hopkins Kimmel Cancer Center, Baltimore MD

Benjamin Schwartz, Northwell Health Gynecologic Oncology, Bay Shore, NY

Patricia Braly, Women's Cancer Care, Covington, LA

Geoffrey Hall, St James University Hospital, Leeds, UK

Kenneth P. Nephew, Indiana University School of Medicine, IU Simon Cancer Center, Bloomington IN

Simone Jueliger, Astex Pharmaceuticals, Cambridge UK

Aram Oganessian, Astex Pharmaceuticals Inc., Pleasanton CA

Sue Naim, Astex Pharmaceuticals Inc., Pleasanton CA

Yong Hao, Astex Pharmaceuticals Inc., Pleasanton CA

Harold Keer, Astex Pharmaceuticals Inc., Pleasanton CA

Mohammad Azab, Astex Pharmaceuticals Inc., Pleasanton CA

Daniela Matei, Northwestern University Feinberg School of Medicine, Chicago IL

Corresponding author: Daniela Matei, 303 E Superior Street Lurie 4-107, Chicago, IL, 60610. Phone: 312-503-4853 Fax: 312-503-4853 Email: Daniela.matei@northwestern.edu

Running title: Guadecitabine and Carboplatin in Ovarian Cancer

Word count: 3259 words

Clinical Trial Registration: ClinicalTrials.gov NCT01696032

Guadecitabine and Carboplatin in Ovarian Cancer

1 **ABSTRACT**

2 **PURPOSE:**

3 Platinum resistance in ovarian cancer (OC) is associated with epigenetic modifications.

4 Hypomethylating agents (HMAs) have been studied as carboplatin re-sensitizing agents in OC.

5 This randomized, multicenter, open-label, phase 2 trial compared combination guadecitabine, a
6 second generation HMA, and carboplatin (G+C) against second-line chemotherapy in women
7 with measurable or detectable platinum-resistant OC.

8 **PATIENTS AND METHODS:**

9 Patients received either G+C (guadecitabine 30 mg/m² SC once-daily for 5 days and carboplatin)
10 or treatment choice (TC; topotecan IV, pegylated liposomal doxorubicin IV, paclitaxel IV, or
11 gemcitabine IV) in 28-day cycles until progression or unacceptable toxicity. Crossover to G+C
12 was allowed at progression. The primary endpoint was progression-free survival (PFS);
13 secondary endpoints were RECIST v1.1 and CA-125 response rate, 6-month PFS, and overall
14 survival (OS).

15 **RESULTS:**

16 Of 100 patients treated, 51 received G+C and 49 received TC, of which 27 crossed over to G+C.
17 The study did not meet its primary endpoint as the median PFS was not statistically different
18 between arms ($P = 0.0654$). However, the 6-month PFS rate was significantly higher in the G+C
19 group (37% vs 11% in TC group; $P = 0.0027$). The incidence of grade 3 or higher toxicity was
20 similar in G+C and TC groups (51% and 49%, respectively), with neutropenia and leukopenia
21 being more frequent in the G+C group.

Guadecitabine and Carboplatin in Ovarian Cancer

22 CONCLUSIONS:

23 Although this trial did not show superiority for PFS of G+C versus TC, the 6-month PFS
24 increased in G+C treated patients. Further refinement of this strategy should focus on
25 identification of predictive markers for patient selection.

26

27

28 **TRANSLATIONAL RELEVANCE**

29 Although women with ovarian cancer (OC) initially respond to platinum-based chemotherapy,
30 platinum-resistance commonly develops leading to fatal outcomes. We set out to determine if
31 epigenetic priming with a hypomethylating agent (HMA) prior to carboplatin improved
32 progression-free survival (PFS) in platinum-resistant OC when compared with physician's
33 choice chemotherapy in a randomized phase II trial. The median PFS and overall survival were
34 not different, but the 6-month PFS rate was higher in the experimental group. Myelosuppression
35 was the main toxicity observed with the experimental regimen and hypomethylating activity was
36 measurable in PBMCs. The efficacy endpoints favored the experimental regimen over
37 established salvage chemotherapy with comparable safety. Further development of the strategy
38 will require identification of predictive biomarkers for patient selection.

39 **INTRODUCTION**

40 Advanced stage high-grade serous ovarian cancer (HGSOC), which is distinctively associated
41 with a p53 mutated signature, has a poor estimated five-year survival of 50% (1). Although
42 patients with HGSOC usually respond to initial platinum-based chemotherapy, relapses occur in
43 most, leading to the development of platinum-resistance and subsequent death (2-3). Progression
44 of HGSOC to a platinum-resistant state is caused by multiple mechanisms, including aberrant
45 DNA repair responses, alterations in efflux pump proteins, and accumulated genomic and
46 epigenomic modifications which impact the response of cancer cells to DNA damage. Adaptive
47 responses include increased DNA methylation and modifications of histone marks (4-5) which
48 cause transcriptional silencing of tumor suppressor (TSGs) and other genes required for
49 chemotherapy-induced cell death (6-7).

50

51 Given preclinical data demonstrating that targeting DNA methylation to re-sensitize HGSOC to
52 platinum is possible (8-11), we hypothesized this approach would restore platinum sensitivity in
53 HGSOC patients (12,13). With early clinical studies demonstrating feasibility of this strategy
54 (13-16), we set out to determine whether targeting DNA methylation induces clinically
55 meaningful activity in platinum-resistant HGSOC by conducting a randomized phase 2 trial. The
56 objectives were to measure and compare clinical outcomes of a combination regimen of the
57 DNA methyltransferase inhibitor (DNMTI), guadecitabine, and carboplatin, versus FDA-
58 approved physician's choice chemotherapy (liposomal doxorubicin, weekly paclitaxel,
59 topotecan, or gemcitabine). Guadecitabine is a dinucleotide linking decitabine to guanosine via a
60 phosphodiester bond. Guadecitabine is resistant to degradation by cytidine deaminase and has a
61 longer half-life compared to other DNMTIs. In a dose finding phase I trial (17), therapeutic

Guadecitabine and Carboplatin in Ovarian Cancer

62 plasma levels of decitabine persisted beyond 8 hours. This pharmacokinetic profile provides a
63 longer window of exposure to the hypomethylating agent (HMA), potentially exposing more
64 cancer cells undergoing S-phase to the parent drug, decitabine, and promoting hypomethylation.
65 Guadecitabine was shown to exert anti-tumor activity in OC xenografts as a single agent and in
66 combination with carboplatin (11, 18, 19).

67
68 A recently reported phase 1 trial established the tolerable and biologically active dose of
69 guadecitabine in combination with carboplatin (17). Guadecitabine was tolerable at 30 mg/m² SC
70 daily for 5 days prior to carboplatin on Day 8 at an AUC of 4. Each cycle was 28 days and the
71 regimen induced ~20% hypomethylation of long interspersed nuclear elements (LINE-1) in
72 peripheral blood mononuclear cells (PBMCs), indicating biological activity. The phase 1 trial
73 reported three patients with partial response (PR) and six patients with stable disease (SD) longer
74 than 3 months (17), providing the rationale for conducting this randomized trial in women with
75 platinum-resistant HGSOE. Here we report clinical outcomes with G+C as compared to
76 physician choice FDA-approved chemotherapy for OC in this high-need patient population.

77

78 METHODS

79 Trial Design and Patient Population:

80 This was a multicenter, randomized, open-label phase 2 trial conducted at 20 centers in the US,
81 UK, and Canada. Eligible patients were ≥18 years old with platinum-resistant histologically- or
82 cytologically-confirmed recurrent high-grade serous, or grade 2-3 endometrioid, mixed cell or
83 clear cell epithelial OC; primary peritoneal carcinoma (PPC); or fallopian tube (FT) cancer. All

Guadecitabine and Carboplatin in Ovarian Cancer

84 patients were required to have received carboplatin and taxanes. Platinum-resistance was
85 defined as recurrence within 6 months of the last platinum-containing regimen. Patients were
86 required to have either measurable disease according to Response Evaluation Criteria in Solid
87 Tumors (RECIST) v1.1 or detectable disease, defined as baseline values of CA-125 at least twice
88 the upper limit of normal and (i) ascites and/or pleural effusion attributed to tumor or (ii) solid
89 and/or cystic abnormalities on radiographic imaging that do not meet RECIST definitions for
90 target lesions. Tumor biopsies, paracentesis, or thoracentesis were performed to recover tumor
91 cells and were required at baseline and on cycle 2 day 8, if clinically safe and feasible. Eligible
92 patients had acceptable organ function based on laboratory data, Eastern Cooperative Oncology
93 Group performance status (ECOG PS) of 0 or 1 and were ≥ 3 weeks from their last therapy.
94 Exclusion criteria included carboplatin hypersensitivity, prior HMA therapy, progression on
95 platinum treatment, left ventricular ejection fraction $< 50\%$, grade 2 or greater peripheral
96 neuropathy, known brain metastases, other malignancies, active infections, or life-threatening
97 illnesses. The trial was conducted in accordance with the International Council for
98 Harmonisation Good Clinical Practice guidelines and applicable local regulatory requirements
99 according to the Declaration of Helsinki. Local Institutional Review Boards and Independent
100 Ethics Committees reviewed and approved the protocol and the informed consent form. Patients
101 provided written informed consent before enrollment. The trial is registered on
102 ClinicalTrials.gov as NCT01696032. Trial protocol and amendments are available as
103 Supplements 1 and 2, respectively.

104 Randomization, Trial Intervention and Clinical Outcomes:

105 Eligible subjects were randomly assigned (1:1) to receive a 28-day treatment cycle of either a
106 G+C combination treatment (guadecitabine 30 mg/m² SC once-daily on Days 1–5 and

Guadecitabine and Carboplatin in Ovarian Cancer

107 carboplatin IV AUC 4 on Day 8), or treatment choice (TC) of topotecan IV (3.5–4.0 mg/m²/wk
108 administered on Days 1, 8 and 15), pegylated liposomal doxorubicin IV (PLD; 40–50 mg/m²
109 administered on Day 1), paclitaxel IV (60–80 mg/m²/wk administered on Days 1, 8, 15 and 22),
110 or gemcitabine IV (800–1000 mg/m² administered on Days 1, 8 and 15); treatment choice in the
111 TC arm was at the investigator’s discretion. Randomization was stratified by number of prior
112 chemotherapies and by treatment center using an unblinded approach using a centralized web-
113 based system. Concomitant medications and therapies were allowed, as deemed necessary for
114 supportive care and safety of subjects; administration of other anti-cancer agents was not
115 permitted. Treatment in both arms continued until disease progression or unacceptable toxicity.
116 If the investigator decided to stop carboplatin treatment after 4 or more cycles, guadecitabine
117 could be continued until progression or initiation of an alternative anti-cancer treatment.
118 Crossover from the TC arm to the G+C arm was permitted after evidence of disease progression
119 in the standard therapy arm.

120 The primary endpoint was PFS. Secondary efficacy endpoints included objective response rate
121 (ORR: defined as complete response [CR] and partial response [PR] based on both measurable
122 and evaluable disease), PFS at 6 months, clinical benefit rate (CBR: defined as CR+ PR + stable
123 disease for at least 3 months), proportion of patients with CA-125 reduction of at least 50%,
124 duration of response (DOR), and overall survival (OS); in subjects crossing over from the TC to
125 the G+C arm, ORR was measured. Response was assessed using RECIST v1.1 for patients with
126 measurable disease (20), and modified Rustin criteria for patients with detectable disease
127 according to CA-125 criteria (21-22). Tumor measurements were obtained by CT or MRI at
128 screening, after every 2 cycles for the first six cycles, and every three months until progression.

Guadecitabine and Carboplatin in Ovarian Cancer

129 Safety was assessed by subject-reported and investigator-observed adverse event (AE) recording,
130 along with physical examination, 12-lead electrocardiograms, hematology, chemistry, and
131 urinalysis with each cycle. There was a 30-day (+5 day) safety visit after the last treatment. AEs
132 were graded by Common Terminology Criteria for Adverse Events (CTCAE) v4.0. Treatment-
133 emergent AEs (TEAEs) were defined as events that first occurred or worsened after the first dose
134 of trial drug given on the first day of the first treatment cycle until 30 days after the last dose of
135 treatment. Related serious AEs (SAEs) that occurred more than 30 days after the last dose were
136 also considered TEAEs; AEs occurring after the start of an alternative anti-cancer treatment were
137 not considered TEAEs. Patients lost to follow-up were included in statistical analyses to the
138 point of their last evaluation.

139 Exploratory pharmacodynamic endpoints included quantitative analysis of methylation of LINE-
140 1 in PBMCs and tumor DNA, and of selected gene promoters in tumor tissue. Blood samples for
141 methylation assays were collected weekly during cycle 1 and on Day #1 and Day #8 thereafter.
142 Global DNA methylation was evaluated by sodium bisulfite pyrosequencing for LINE-1 CpGs
143 using PyroMark Q24 as previously described (17). Ascites, pleural fluid, or fresh tumor biopsies
144 were obtained at screening and on Day 8 of cycle 2 for assessment of methylation of selected
145 genes listed in the supplementary information (Supplementary Table S1). DNA was extracted
146 from tumor biopsies or ascites using DNeasy Blood & Tissue Kit (Qiagen, Netherlands) and
147 LINE-1 and specific gene pyrosequencing was performed at EpigenDx Inc (Hopkinton, MA).

148 **Statistical Design and Analyses:**

149 It was estimated a sample size of ≥ 96 patients randomized 1:1 into two treatment arms would
150 provide approximately 80% power to detect a difference between the two PFS curves (median

Guadecitabine and Carboplatin in Ovarian Cancer

151 PFS of 15 vs. 28 weeks for the TC and G+C arms) at 5% significance level using a two-sided
152 log-rank test, assuming uniform accrual of subjects over 12 months, a 24-month trial duration
153 and an exponential distribution of the PFS endpoint. PFS, OS and 95% CIs were evaluated using
154 the Kaplan-Meier method. PFS and OS were compared using the log-rank test, while ORR and
155 CBR were compared using Fisher's exact test. Subjects still alive with no progression and those
156 who withdrew were censored on the date of the last adequate tumor, CA-125, or clinical
157 progression assessment. Subjects initiating subsequent anti-cancer therapy, including those who
158 crossed over, were censored accordingly, but prior to the initiation Survival time was censored
159 on the last date the subject was known to be alive or lost to follow-up before reaching the event
160 of death. Efficacy and safety data for subjects who crossed over were tabulated separately once
161 guadecitabine was first administered. All analyses are descriptive and inferential statistical tests
162 and CIs were two-sided with alpha equal to 0.05 unless otherwise specified. The database was
163 locked for analysis on July 7, 2016 with very mature PFS data; 97 of the 100 treated patients
164 progressed or did not survive and all patients discontinued protocol therapy at this time (Figure
165 1). LINE-1 and gene-specific methylation level differences before and after G+C treatment were
166 determined using paired *t*-tests. SAS version 9.3 was used for all statistical analyses.

167 **RESULTS**

168 One hundred and three patients with HGSOC, FT cancer or PPC were enrolled and randomized
169 (52 G+C, 51 TC) and 100 received treatment (51 G+C, 49 TC; Figure 1). Baseline characteristics
170 are summarised in Table 1 and were well balanced between the two arms in terms of age,
171 performance status, prior therapy and ethnicity. More patients randomized to the G+C arm had
172 PPC compared to those randomized to TC (10 vs. 0). Most subjects were white, with a median
173 age of 62 years, and all received prior platinum-based therapy (Table 1). Of the patients

Guadecitabine and Carboplatin in Ovarian Cancer

174 randomized to TC, 11 received weekly paclitaxel, 15 received liposomal doxorubicin, 20
175 received topotecan, and 3 received gemcitabine. Patients in the G+C arm received more
176 treatment cycles than subjects in the TC arm (median of 4.0 vs 2.0 cycles, respectively), with
177 59% of subjects in the G+C arm receiving at least 3 cycles of treatment and 37% receiving at
178 least 6 cycles of treatment versus 47% and 31% of subjects in the TC arm, respectively. Fifty-
179 five percent of patients from the TC arm crossed over to G+C arm following progression (Figure
180 1). Disease progression was the most common reason for discontinuing treatment (~80% of
181 patients in each group; Figure 1). The most common TEAEs occurring in more than 5% of the
182 trial population are reported in Table 2. AE frequencies between the two arms were similar, but
183 neutropenia, diarrhea, nausea and vomiting were more common in the G+C arm (Tables 2 and
184 3).

185 The median duration of PFS in the G+C arm was 16.2 weeks compared to 9.1 weeks in TC arm
186 ($p=0.0654$; Figure 2A and Table 4). The 6-month PFS rate was 37% in the G+C arm (95% CI,
187 [0.24; 0.50]) compared to 11% in the TC arm (95% CI, [0.04; 0.22]; $p=0.0027$) and did not meet
188 the pre-specified criterion for superiority (HR 0.686, 95% CI, [0.456; 1.030]; Figure 2 and Table
189 4). There was no difference between the two arms in OS (43 and 40 weeks in the G+C and TC
190 arms, respectively; Figure 2B and Table 4), OS survival rate at 6 months (0.72 and 0.67 in the
191 G+C and TC arms, respectively; Table 4), overall response rate (ORR; 16% and 8% in the G+C
192 and TC arms, respectively; Table 4), or clinical benefit response by RECIST v1.1 or CA-125
193 (Table 4, Supplementary Table S2). Twenty-seven patients from the TC arm crossed over post-
194 progression into the G+C arm and received a median of 3 cycles (14 subjects received ≥ 3 cycles
195 and 5 subjects received ≥ 6 cycles) with a CA-125 response being confirmed in 6 of 21 evaluable
196 subjects (29%). Patient disposition and outcomes are included in Supplementary Table S3.

Guadecitabine and Carboplatin in Ovarian Cancer

197 To determine the biological activity of the G+C regimen, LINE1 methylation was assessed in
198 PBMCs from 48 patients randomized to the G+C arm. Similar to the first stage of this trial (17),
199 LINE1 hypomethylation approximated 20% (C1D8 vs. C1D1; range +15% to -55%;
200 Supplementary Figure S1A) (17). In 15 patients who continued treatment beyond 2 cycles and
201 for whom PBMCs were available, LINE1 hypomethylation observed during Cycle 1 was
202 maintained or increased during subsequent cycles (Supplementary Figure S1B), indicating that
203 G+C maintains its biological effects throughout treatment. Correlation between clinical response
204 and pharmacodynamic effects as measured by LINE-1 hypomethylation in PBMCs was not
205 observed. Promoter methylation of selected genes representing TSGs (23-24) or tumor antigens
206 known to be methylated in OC (25-26) was measured in bisulfite-converted DNA obtained from
207 paired tumor biopsies on C1D1 and C2D8 (n = 8 paired specimens). Treatment-induced
208 hypomethylation of *MAGE-A2* and *MAGE-A3* promoters in tumor DNA was significant
209 (Supplementary Figure S1C). A non-significant decrease in promoter CpG methylation was also
210 observed for LINE-1 and for the tumor antigens *NY-ESO-1* and *MAGE-A11*, but not for the
211 TSGs *RASSF1A*, *MLH1* and *BRCA1* (data not shown) or for the differentiation associated gene
212 *HOXA11*. Taken together, these results provide evidence that G+C treatment exerts *in vivo*
213 hypomethylating activity detectable in PBMCs and tumors.

214

215 **DISCUSSION**

216 This is the first randomized study comparing a regimen of G+C to standard of care
217 chemotherapy for recurrent platinum-resistant OC. The study did not meet its primary endpoint,
218 but the 6-month PFS in the G+C arm was superior ($P < 0.05$) versus TC in this heavily pre-treated
219 patient population. These results are comparable with previous single arm phase II studies using

Guadecitabine and Carboplatin in Ovarian Cancer

220 an epigenetic priming with decitabine (13-14) or 5-azacitadine (15) prior to carboplatin. Those
221 trials used repetitive low doses of DNMTIs, which is similar to the strategy employed with this
222 class of HMAs in hematological malignancies (27-28). The repetitive administration of the HMA
223 increases exposure of cells undergoing S-phase to the drug and incorporation of the nucleoside
224 analogue into the replicating DNA, trapping DNMTs and inhibiting *de novo* methylation.

225

226 In contrast, a previous trial conducted by the Scottish Gynecological Trials Group that used
227 bolus administration of decitabine on Day 1 prior to administration of carboplatin a week later
228 was prematurely closed due to high hematological toxicity and indicated lower efficacy of the
229 combination regimen compared to carboplatin alone (29). This trial reported reduction in
230 efficacy with the addition of decitabine to patients with partially platinum sensitive recurrence
231 when given in conjunction with carboplatin (29). Whether the difference in administration (bolus
232 vs. low dose repetitive administration) was solely responsible for the differences in levels of
233 clinical activity remains unknown. The clinical efficacy differences with this trial may be
234 attributable to the Scottish trial's inclusion of less heavily pre-treated subjects who retained
235 partial platinum sensitivity. Since increased DNA methylation is observed in advanced bladder,
236 colon cancer, cholangiocarcinoma, and germ cell tumors (30), DNMTI-induced sensitization to
237 platinum or to chemotherapy is also explored in these settings with early promising results
238 having been reported recently in colon cancer (31).

239

240 The G+C regimen had myelosuppression as the main toxicity. Prolonged neutropenia required
241 growth factor support in >80% of the patient population to maintain the intended every 4-week
242 administration of the combination. However, infections were rare and no episodes of neutropenic

Guadecitabine and Carboplatin in Ovarian Cancer

243 sepsis were recorded. Hypersensitivity and other adverse infusion reactions were observed in 9
244 (18%) and 8 (15%) patients in the G+C arm compared with 6% in the TC arm in this trial, which
245 is concordant with similar observations from prior trials of DNMTIs and carboplatin (13, 29).
246 This is most likely due to increased exposure to platinum therapy in the experimental arm, but it
247 is also possible HMA treatment may augment type II allergic reactions.

248

249 The study has few limitations. While all patients in this trial had platinum-resistant disease,
250 platinum-refractory disease was excluded. Given that carboplatin was not included among the
251 potential control regimens, and could conceivably induce clinical benefit in selected patients, this
252 trial cannot exclude the activity of single agent carboplatin in this population. Additionally,
253 topotecan administration in the TC arm followed a weekly administration schedule. While this
254 schedule was favored among treating oncologists due to its favorable toxicity profile and early
255 reports of activity (32), the regimen was subsequently shown to induce a decreased response rate
256 compared to the schedule using daily administration for 5 days, although OS was not affected
257 (33). Chemotherapy with bevacizumab became FDA-approved and an accepted standard for
258 patients with platinum resistant OC after results of Aurelia trial were reported (34), which
259 occurred after the inception of this protocol. Of note is that prior therapy with bevacizumab was
260 not excluded, and 33 patients enrolled in this trial had received bevacizumab. The shorter
261 median PFS observed in the control group of this study (~2 months) compared to the Aurelia
262 trial (3.4 months; 34) reflects a more heavily pre-treated group patients included here (mean of 3-
263 4 prior regimens) for whom limited treatment options currently exist.

264

Guadecitabine and Carboplatin in Ovarian Cancer

265 High quality nucleic acids were extracted from tumor biopsies from 40 subjects at baseline and
266 from 8 patients after two cycles of G+C. Methyloomic and transcriptomic analyses of biopsies
267 from G+C-treated subjects reported previously indicated activation of immune pathways,
268 including interleukin (IL)-8, IL-6, B-cell receptor, leukocyte extravasation, and nuclear factor- κ B
269 activation by viruses (35). Treatment with other DNMTIs in preclinical OC models also
270 increased host anti-tumor immunity by directly reducing the population of myeloid-derived
271 suppressor cells (36). Other preclinical studies in cancer models reported treatment with
272 DNMTIs induced an IFN- γ response, triggered by cellular sensing of double-stranded RNA and
273 retroviral transcripts, which are upregulated in response to global genomic hypomethylation (37).
274 Taken together, these clinical and preclinical observations indicate priming with DNMTIs
275 promotes anti-tumor immunity through a variety of mechanisms and may enhance response to
276 other immune therapies. Clinical trials evaluating this concept are currently ongoing. The
277 precise mechanism by which G+C induces anti-tumor responses remains unknown. Our tissue-
278 and cell-based analyses showed a number of genes and pathways involved in DNA repair and
279 response to chemotherapy (e.g. *DOK2*, *miR193a*, *14-3-3 σ* , *RASSF1A*) are silenced through
280 promoter methylation and re-expressed after guadecitabine treatment. Using overexpression or
281 knock-down strategies, we have shown some of these pathways restore platinum sensitivity in
282 OC cell lines and xenografts (10, 35). It is likely that not one gene, but a more global genomic
283 program is reactivated in response to DNA hypomethylation, allowing tumor cells to undergo
284 apoptosis in response to chemotherapy. Since preclinical models show that guadecitabine
285 selectively eliminates chemotherapy-resistant OC stem cells (11) by inducing a cellular
286 differentiation program, the G+C regimen may exert anti-tumor activity through multiple

Guadecitabine and Carboplatin in Ovarian Cancer

287 mechanisms. The low number of post-treatment biopsies collected in the trial limits the strength
288 of the conclusions we can draw regarding the mechanisms elicited by this HMA *in vivo*.

289

290 This randomized trial demonstrated that epigenetic priming in combination with carboplatin did
291 not increase PFS compared to standard chemotherapy, but improved 6-month PFS in platinum-
292 resistant OC. Although these results do not support development of this strategy for an
293 unselected population, they do indicate a subgroup benefitted from G+C treatment. Future
294 studies should focus on developing predictive markers to enrich a patient population more likely
295 to benefit from the use of HMAs.

296

297

298 **DISCLOSURE OF POTENTIAL CONFLICTS OF INTEREST**

299 Angeles Alvarez Secord reported being paid for consulting or participating in an advisory role
300 for Alexion, Aravive, AstraZeneca, Clovis, Janssen/Johnson & Johnson, Mesano, Myriad,
301 Roche/Genentech, and Tesaro, and received research funding from Amgen, AbbVie, Amgen,
302 Astellas Pharma Inc., Astex Pharmaceuticals Inc., AstraZeneca, Boehringer Ingelheim, Bristol
303 Myers Squibb, Eisai, Exelixis, Endocyte, Roche/Genentech, Incyte, Merck, PharmaMar, Prima
304 Biomed, and Tesaro. Sarah Blagden reported serving in a consultant or advisory role for
305 Novartis, Octimet and Roche, receiving travel, accommodation and expense reimbursement from
306 NuCana, BioMed and Tesaro, receiving research funding from NuCana, BioMed, Sierra
307 Oncology, Incyte, DesigneRx and Tesaro, and holds patents or receives royalties from RNA
308 Guardian Ltd. Susana Banerjee reported receiving honoraria from AstraZeneca and Tesaro,
309 serving in a consultant or advisory role for AstraZeneca, Tesaro, Clovis, Seattle Genetics, and
310 receiving research funding from AstraZeneca. John Nemunaitis disclosed employment with
311 Gradulis, leadership roles with Gradulis and Symvivo. He has stock or other ownership interest
312 to disclose with Gradulis, received honoraria from AstraZeneca, has consulted for AstraZeneca
313 and Symvivo, participated in a speaker's bureau for AstraZeneca, received research funding from
314 Gradulis, holds patents or receives royalties from Gradulis, receives travel, accommodations, or
315 expenses from AstrazZeneca, Symvivo and Gradulis, and been paid to provide expert testimony
316 on behalf of Foundation Medicine. Hal Hirte reported receiving honoraria from AstraZeneca,
317 Merck and Roche. Diane Provencher reported consulting and advising AstraZeneca. Benjamin
318 Schwartz reported receiving honoraria from NOVADAQ. Patricia Braly reported participating in
319 a speakers' bureau for Myriad, Invitae, Tesaro, AstraZeneca, Clovis and Roche, and receiving
320 research funding from Tesaro, AstraZeneca, Merck, Janssen, Pharma Mar and Xenetic. Geoffrey

Guadecitabine and Carboplatin in Ovarian Cancer

321 Hall reported receiving honoraria from and serving in a consultant or advisory role for
322 AstraZeneca and IQVIA. Daniela Matei reported serving in a consulting or advisory role for
323 Genentech, Tesaro, AstraZeneca, and Anydyn, and receiving travel, accommodation and expense
324 reimbursement from Genentech. The following authors are employed by Astex Pharmaceuticals
325 Inc: Aram Ogenesian, Sue Naim, Yong Hao, Harold Keer, Mohammad Azab and Simone
326 Jueliger.

327

328 **AUTHORS' CONTRIBUTIONS:**

329 *Conception and design:* Matei, Nephew, Azab, Ogenesian, Naim, Hao, Keer.

330 *Development of methodology:* Matei, Nephew, Azab, Ogenesian, Naim, Hao, Keer.

331 *Acquisition of data (provided animals, acquired and managed patients, provided*

332 *facilities, etc.):* All authors.

333 *Analysis and interpretation of data (e.g., statistical analysis, biostatistics,*

334 *computational analysis):* Matei, Nephew, Azab, Ogenesian, Naim, Hao, Keer.

335 *Writing, review, and/or revision of the manuscript:* All authors.

336 *Administrative, technical, or material support (i.e., reporting or organizing*

337 *data, constructing databases):* Matei, Oza, Azab, Naim, Hao, Keer.

338 *Study supervision:* Matei, Oza, Azab, Jueliger, Ogenesian, Naim, Hao, Keer.

339

340 **ACKNOWLEDGEMENTS**

341 **Additional Contributions:** This work was funded in part by the National Cancer Institute

342 Award CA182832-01 and the V- Foundation (to DM and KPN). Imperial, UCH and Royal

343 Marsden Hospitals are supported as CRUK and Experimental Cancer Medicine Centres

Guadecitabine and Carboplatin in Ovarian Cancer

344 (ECMCs). This study was supported in part by Astex Pharmaceuticals Inc. Medical writing
345 support was provided by Paul Sobol and Ryan Draker from Six Degrees Medical Consulting and
346 funded by Astex Pharmaceuticals, Inc.

347

348

REFERENCES

1. Society AC. Cancer Facts and Figures 2018. Atlanta, GA, American Cancer Society.
2. Liu CM. Cancer of the ovary. *New Engl J Med*. 2005;352:1268-9; author reply 1268-9.
3. Sandercock J, Parmar MK, Torri V, Wian W. First-line treatment for advanced ovarian cancer: paclitaxel, platinum and the evidence. *Br J Cancer* 2002;87:815-24.
4. Watts GS, Futscher BW, Holtan N, Degeest K, Domann, FE, Rose SL. DNA methylation changes in ovarian cancer are cumulative with disease progression and identify tumor stage. *BMC Med Genomics* 2008;1:47.
5. Barton CA, Hacker NF, Clark SJ, O'Brien PM. DNA methylation changes in ovarian cancer: implications for early diagnosis, prognosis and treatment. *Gynecol Oncol* 2008;109:129-39.
6. Balch C, Fang F, Matei DE, Huang TH, Nephew KP. Minireview: epigenetic changes in ovarian cancer. *Endocrinology* 2009;150:4003-11.
7. Barton CA, Clark SJ, Hacker NF, O'Brien PM. Epigenetic markers of ovarian cancer. *Adv Exp Med Biol* 2008;622:35-51.
8. Plumb JA, Finn PW, Williams RJ, Bandara MJ, Watkins CJ, La Thangue NB, *et al*. Pharmacodynamic response and inhibition of growth of human tumor xenografts by the novel histone deacetylase inhibitor PXD101. *Mol Cancer Ther* 2003;2:721-8.
9. Plumb JA, Strathdee G, Sludden J, Kaye SB, Brown R. Reversal of drug resistance in human tumor xenografts by 2'-deoxy-5-azacytidine-induced demethylation of the hMLH1 gene promoter. *Cancer Res* 2000;60:6039-44.
10. Fang F, Munck J, Tang J, Taverna P, Wang Y, Miller DF, *et al*. The novel, small-molecule DNA methylation inhibitor SGI-110 as an ovarian cancer chemosensitizer. *Clin Cancer Res* 2014;20:6504-16.
11. Wang Y, Cardenas H, Fang F, Condello S, Taverna P, Segar M, *et al*. Epigenetic targeting of ovarian cancer stem cells. *Cancer Res* 2014;74:4922-36.
12. Fang F, Zuo Q, Pilrose J, Wang Y, Shen C, Li M, *et al*. Decitabine reactivated pathways in platinum resistant ovarian cancer. *Oncotarget* 2014;5:3579-89.
13. Matei D, Fang F, Shen C, Schilder J, Arnold A, Zeng Y, *et al*. Epigenetic resensitization to platinum in ovarian cancer. *Cancer Res* 2012;72:2197-205.
14. Fang F, Balch C, Schilder J, Breen T, Zhang S, Shen C, *et al*. A phase 1 and pharmacodynamic study of decitabine in combination with carboplatin in patients with recurrent, platinum-resistant, epithelial ovarian cancer. *Cancer* 2010;116:4043-53.
15. Fu S, Hu W, Iyer R, Kavanagh JJ, Coleman RL, Levenback CF, *et al*. Phase 1b-2a study to reverse platinum resistance through use of a hypomethylating agent, azacitidine, in patients with platinum-resistant or platinum-refractory epithelial ovarian cancer. *Cancer* 2011;117:1661-9.
16. Appleton K, Mackay HJ, Judson I, Plumb JA, McCormick C, Strathdee G, *et al*. Phase I and pharmacodynamic trial of the DNA methyltransferase inhibitor decitabine and carboplatin in solid tumors. *J Clin Oncol* 2007;25:4603-9.
17. Matei D, Ghamande S, Roman L, Alvarez Secord A, Nemunaitis J, Markham MJ, *et al*. A Phase I Clinical Trial of Guadecitabine and Carboplatin in Platinum-Resistant, Recurrent Ovarian Cancer: Clinical, Pharmacokinetic, and Pharmacodynamic Analyses. *Clin Cancer Res* 2018;24:2285-93.

18. Wang Y, Zong X, Mitra S, Mitra AK, Matei D, Nephew KP. IL-6 mediates platinum-induced enrichment of ovarian cancer stem cells. *JCI Insight* 2018; 3:doi:10.1172/jci.insight.122360.
19. Alvero AB, Chen R, Fu HH, Montagna M, Schwartz PE, Rutherford T, *et al.* Molecular phenotyping of human ovarian cancer stem cells unravels the mechanisms for repair and chemoresistance. *Cell Cycle* 2009;8:158-66.
20. Eisenhauer EA, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, *et al.* New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009;45:228-47.
21. Rustin GJ. Use of CA-125 to assess response to new agents in ovarian cancer trials. *J Clin Oncol* 2003;21:187s-193s.
22. Rustin GJ, Bast RC, Jr., Kelloff GJ, Barrett JC, Carter SK, Nisen PD, *et al.* Use of CA-125 in clinical trial evaluation of new therapeutic drugs for ovarian cancer. *Clin Cancer Res* 2004;10:3919-26.
23. Ibanez de Caceres I, Battagli C, Esteller M, Herman JG, Dulaimi E, Edelson MI, *et al.* Tumor cell-specific BRCA1 and RASSF1A hypermethylation in serum, plasma, and peritoneal fluid from ovarian cancer patients. *Cancer Res* 2004;64:6476-81.
24. Imura M, Yamashita S, Cai LY, Furuta JI, Wakabayashi M, Yasugi T, *et al.* Methylation and expression analysis of 15 genes and three normally-methylated genes in 13 Ovarian cancer cell lines. *Cancer Lett* 2006;241:213-20.
25. Odunsi K, Matsuzaki J, James SR, Mhawech-Fauceglia P, Tsuji T, Miller A, *et al.* Epigenetic potentiation of NY-ESO-1 vaccine therapy in human ovarian cancer. *Cancer Immunol Res* 2014;2:37-49.
26. Zhang W, Barger CJ, Link PA, Mhawech-Fauceglia P, Miller A, Akers SN, *et al.* DNA hypomethylation-mediated activation of Cancer/Testis Antigen 45 (CT45) genes is associated with disease progression and reduced survival in epithelial ovarian cancer. *Epigenetics* 2015;10:736-48.
27. Issa JP, Gharibyan V, Cortes J, Jelinek J, Morris G, Verstovsek S, *et al.* Phase II study of low-dose decitabine in patients with chronic myelogenous leukemia resistant to imatinib mesylate. *J Clin Oncol* 2005;23:3948-56.
28. Issa JJ, Roboz G, Rizzieri D, Jabbour E, Stock W, O'Connell C, *et al.* Safety and tolerability of guadecitabine (SGI-110) in patients with myelodysplastic syndrome and acute myeloid leukaemia: a multicentre, randomised, dose-escalation phase 1 study. *Lancet Oncol* 2015;16:1099-110.
29. Glasspool RM, Brown R, Gore ME, Rustin GJ, McNeish IA, Wilson RH, *et al.* A randomised, phase II trial of the DNA-hypomethylating agent 5-aza-2'-deoxycytidine (decitabine) in combination with carboplatin vs carboplatin alone in patients with recurrent, partially platinum-sensitive ovarian cancer. *Br J Cancer* 2014;110:1923-9.
30. Albany C, Hever-Jardine MP, von Herrmann KM, Yim CY, Tam J, Warzecha JM, *et al.* Refractory testicular germ cell tumors are highly sensitive to the second generation DNA methylation inhibitor guadecitabine. *Oncotarget* 2017;8:2949-59.
31. Lee V, Wang JS, Zahurak ML, Gootjes E, Verheul HM, Parkinson R, *et al.* A phase I trial of a guadecitabine (SGI-110) and irinotecan in metastatic colorectal cancer patients previously exposed to irinotecan. *Clin Cancer Res* 2018;24:6160-7.

32. Levy T, Inbar M, Menczer J, Grisaru D, Glezerman M, Safra T. Phase II study of weekly topotecan in patients with recurrent or persistent epithelial ovarian cancer. *Gynecol Oncol* 2004;95:686-90.
33. Sehouli J, Stengel D, Harter P, Kurzeder C, Belau A, Bogenrieder T, *et al.* Topotecan Weekly Versus Conventional 5-Day Schedule in Patients With Platinum-Resistant Ovarian Cancer: a randomized multicenter phase II trial of the North-Eastern German Society of Gynecological Oncology Ovarian Cancer Study Group. *J Clin Oncol* 2011;29:242-8.
34. Pujade-Lauraine E, Hilpert F, Weber B, Reuss A, Poveda A, Kristensen G, *et al.* Bevacizumab combined with chemotherapy for platinum-resistant recurrent ovarian cancer: The AURELIA open-label randomized phase III trial. *J Clin Oncol* 2014;32:1302-8.
35. Fang F, Cardenas H, Huang H, Jiang G, Perkins SM, Zhang C, *et al.* Genomic and Epigenomic Signatures in Ovarian Cancer Associated with Re-sensitization to Platinum Drugs. *Cancer Res* 2018;78: 631-644.
36. Stone ML, Chiappinelli KB, Li H, Murphy LM, Travers ME, Topper MJ, *et al.* Epigenetic therapy activates type I interferon signaling in murine ovarian cancer to reduce immunosuppression and tumor burden. *Proc Natl Acad Sci U S A* 2017;114:E10981-E10990.
37. Chiappinelli KB, Strissel PL, Desrichard A, Li H, Henke C, Akman B, *et al.* Inhibiting DNA Methylation Causes an Interferon Response in Cancer via dsRNA Including Endogenous Retroviruses. *Cell* 2015;162:974-86.

FIGURE LEGENDS

Figure 1. Disposition of subjects in the trial. AUC indicates the target area under the concentration versus time curve.

Figure 2. Survival of subjects assigned to G+C arm versus TC arm. A: Kaplan-Meier estimates of progression-free survival with the G+C treatment and TC regimens. B: Kaplan-Meier estimates of overall survival with the G+C treatment and TC regimens. For subjects in the TC group who crossed over to receive G+C, OS time was censored at the crossover time point.

Figure 1: Oza et al 2018




Figure 2: Oza et al 2018





Table 1. Patient characteristics

	G+C (n=51)		TC (n=49)		TC crossover (TC to G+C) (n=27)	
Demographics and Baseline Characteristics						
Age (y)						
Mean	62.0		62.1		63.7	
SD	9.2		9.6		8.8	
Median	62.1		62.2		64.1	
Min-max	40.6-88.4		38.9-78.5		49.6-78.5	
Race						
Asian	5	(10)	6	(12)	3	(11)
Black or African American	2	(4)	2	(4)	2	(7)
Native Hawaiian or Other Pacific Islander	0		1	(2)	0	
White	43	(84)	40	(82)	22	(81)
Other	1	(2)	1	(2)	1	(4)
Ethnicity						
Hispanic or Latino	2	(4)	3	(6)	3	(11)
Not Hispanic or Latino	49	(96)	46	(94)	24	(89)
Diagnosis, n (%)						
HGSOC	39	(76)	47	(96)		
High-grade serous PPC	10	(20)	0			
High-grade serous FT cancer	2	(4)	2	(4)		
Disease Assessment, n (%)						
Measurable disease (RECIST)	44	(86)	46	(94)		
Detectable disease (GCIC)	7	(14)	3	(6)		
ECOG Performance Status, n (%)						
0	18	(35)	23	(47)	13	(48)
1	33	(65)	25	(51)	14	(52)
2	0		1	(2)	0	
Number of Prior Regimens, n (%)						
1-2	11	(22)	14	(29)	7	(26)
3-4	21	(41)	16	(33)	8	(30)
≥5	19	(37)	19	(39)	12	(44)
Number of subjects with prior bevacizumab	17	(33)	14	(28.6)		
Time since last platinum therapy ^a (days)						
Mean	288		378			
SD	245		290			

^a Includes cisplatin, carboplatin, or oxaliplatin

Table 2. Treatment-related AEs occurring in ≥10% of patients in either arm

AE ^a		G+C (n=51)		TC (n=49)
ANY RELATED EVENT	47	(92)	43	(88)
Neutropenia ^b	36	(71)	16	(33)
Nausea	28	(55)	21	(43)
Fatigue	24	(47)	21	(43)
Injection site reaction ^{b,d}	20	(39)	0	
Vomiting ^c	18	(35)	8	(16)
Anemia	16	(31)	23	(47)
Leukopenia	16	(31)	10	(20)
Hypomagnesaemia	12	(24)	4	(8)
Thrombocytopenia	11	(22)	12	(24)
Constipation	10	(20)	9	(18)
Decreased appetite	10	(20)	8	(16)
Stomatitis	10	(20)	12	(24)
Drug hypersensitivity ^d	9	(18)	3	(6)
Arthralgia	7	(14)	5	(10)
Diarrhea	7	(14)	5	(10)
Headache	6	(12)	1	(2)
Alopecia	5	(10)	7	(14)

^a $P > 0.050$ unless otherwise stated

^b $P < 0.001$

^c $P = 0.040$

^d Due to variability in reporting terms, events of injection site reaction (typically attributed to guadecitabine SQ injection), drug hypersensitivity (typically attributed to carboplatin), anaphylactic reaction, adverse drug reaction, and infusion related reaction were analyzed as a group term and were observed in 32 subjects (33%) who received the G+C treatment.

Table 3. AEs of CTCAE grade 3 or higher occurring in >1 patient in either arm

AE ^a		G+C (n=51)		TC (n=49)
ANY GRADE \geq 3 EVENTS ^b	48	(94)	31	(63)
Neutropenia ^b	34	(67)	9	(18)
Leukopenia ^c	13	(25)	4	(8)
Anaemia	9	(18)	8	(16)
Bowel obstruction	12	(24)	8	(16)
Fatigue	6	(12)	6	(12)
Diarrhea	3	(6)	0	
Thrombocytopenia	3	(6)	4	(8)
Vomiting	3	(6)	4	(8)
Abdominal distension	2	(4)	1	(2)
Abdominal pain	2	(4)	2	(4)
Ascites	2	(4)	2	(4)
Hypertension	2	(4)	2	(4)
Hypokalemia	2	(4)	3	(6)
Nausea	2	(4)	4	(8)
Pyrexia	2	(4)	0	
Decreased appetite	1	(2)	2	(4)
Dehydration	1	(2)	2	(4)
Pulmonary embolism	1	(2)	2	(4)
Pneumonia	0		2	(4)
Sepsis	0		2	(4)

^a $P > 0.050$ unless otherwise stated

^b $P < 0.001$

^c $P = 0.032$

Table 4. Survival and response

	G+C^a (n=51)	TC (n=49)	p-value
Survival			
PFS, median in weeks [95% CI]	16.3 [9, 24.1]	9.1 [7.4, 15]	0.0654 ^b
PFS rate at 6 months, median [95% CI]	0.37 [0.24,0.50]	0.11 [0.04,0.22]	0.0027 ^c
OS, (TC censored) median in weeks [95% CI]	47.3 [33, 59.3]	31.5 [20.7, 53.1]	0.5852 ^b
OS rate at 6 months, (TC censored) median [95% CI]	0.72 [0.58, 0.83]	0.67 [0.47, 0.80]	0.5629 ^c
Response Rate			
ORR (CR/FR+PR), n (%)	8 (16)	4 (8)	0.3580 ^d
[95% CI]	[7.0, 28.6]	[2.3, 19.6]	
CBR (CR/FR+PR+stable disease), n (%)	21 (41)	14 (29)	0.2130 ^d
[95% CI]	[27.6, 55.8]	[16.6, 43.3]	
Duration of Response in Responders			
Number of responders	21	14	
Median duration, weeks [95% CI]	26.6 [21, 34.4]	24.7 [17.3, 38.1]	
CA-125 Response, n			
Number (%) of subjects with ≥50% reduction	15 (36)	13 (32)	
Median best % change from baseline (min, max)	-43 (-98, 154)	-10 (-98, 248)	

AE=adverse event; CBR= clinical benefit rate; CI=confidence interval; CR=complete response; CRc=composite complete response; FR=full response per GCIC criteria; ORR=objective response rate; OS=overall survival; PFS=progression free survival; PR=partial response. Subjects were primarily assessed by RECIST, but in the event that an enrolled subject with measurable disease was not evaluable by RECIST (e.g. inadequate follow up scan) and had evaluable data by GCIC CA-125 criteria, the latter was used. From the G group, there were 5 PR by RECIST of 44 evaluable and 3 PR/FR by GCIC of 7 subjects with detectable disease. From the TC group there were 4 PR by RECIST of 44 evaluable and 0 PR/FR of 3 evaluable by GCIC CA-125 criteria.

^aGuadecitabine 30 mg/m² on Days 1-5 and carboplatin AUC 4 on Day 8 of 28-day treatment cycles.

^bLog-rank test for the overall PFS or OS curve.

^cChi-square test.

^dFisher's exact test.