DOI: <u>https://doi.org/10.1016/j.aquaculture.2019.734677</u> © 2019, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <u>http://</u> creativecommons.org/licenses/by-nc-nd/4.0/

- 1 Enriching Artemia nauplii with selenium from different sources and interactions with essential
- 2 fatty acid incorporation

3

- 4 Authors
- 5 Thomas Cavrois-Rogacki^{1*}, Andrew Rolland¹, Hervé Migaud¹, Andrew Davie^{1*}, Oscar
- 6 Monroig^{1,2}
- 7

8 Addresses

- 9 ¹Institute of Aquaculture, Faculty of Natural Sciences, University of Stirling, Stirling FK9
- 10 4LA, Scotland, UK
- ¹¹ ²Instituto de Acuicultura Torre de la Sal, Consejo Superior de Investigaciones Científicas
- 12 (IATS-CSIC), 12595 Ribera de Cabanes, Castellón, Spain

13

14 ***Corresponding author**

- 15 Andrew Davie
- 16 Institute of Aquaculture, Faculty of Natural Sciences, University of Stirling, Stirling FK9 4LA,
- 17 Scotland, UK
- 18 E-mail : andrew.davie@stir.ac.uk

20 Abstract

The production of high-quality marine fish fry is limited by the low survival observed during 21 22 the larval phase, which is often attributed to dietary deficiencies of the diets at first feeding. 23 Despite progress made with live feed (i.e. rotifers, Artemia), enrichments in essential fatty acids 24 for marine fish larvae, little is known on the micronutrient requirements such as selenium (Se). 25 Se is a critical component of several enzymes maintaining important biological functions such as cellular oxidation, and therefore plays a key role in oxidative and stress status of marine 26 27 larvae. The levels of Se found in the larvae's natural diet (i.e. copepods) is generally higher 28 than those of the enriched live preys used in hatcheries. This study aimed at establishing a 29 protocol to enrich Artemia nauplii with Se using different inorganic (sodium selenite) and 30 organic (selenoyeast). Results indicated that the use of dissolved sodium selenite, an alternative 31 inorganic and cheaper form of Se, did not increase the levels of Se in the nauplii. However, the 32 use of selenoyeast (Sel-Plex) confirmed that it is possible to enrich the nauplii with targeted 33 levels of Se, since this process followed a dose-response pattern with Se enrichment ranging from 1.7 to 12.4 mg kg⁻¹. In addition, the supplementation of Sel-Plex to the regular enrichment 34 35 product did not impact on lipids and fatty acids enrichment irrespective of the dose dispensed. Overall, this study contributes to the refinement of the live prey enrichment protocols that are 36 37 critical to the success of marine finfish larviculture protocols.

Keywords: Artemia; Essential fatty acids; Selenium; Sel-Plex; Sodium selenite.

38

39

- 40
- 41

44 Marine finfish larviculture has remarkably improved during the last three decades but it still remains one of the most challenging research areas and commercial bottleneck in marine fish 45 46 aquaculture. Despite significant advances in the hatcheries' operational procedures, survival 47 rates during the larval phases usually remains low with at best 30 % in well-established species 48 such as gilthead seabream (Sparus aurata) (Atalah et al., 2011) and European seabass (Dicentrachus labrax) (Villamizar et al., 2009), and <10 % in most other species such as 49 50 California yellowtail (Seriola lalandi) (Hawkyard et al., 2016), Atlantic halibut (Hippoglossus hippoglossus) (Bjornsdottir et al., 2009) and ballan wrasse (Labrus bergylta) (Øie et al., 2015). 51 52 Causes accounting for the low survival in marine larviculture are multifactorial but nutrition 53 and feeding have been often regarded as major causal factors (Hamre et al., 2013). While it is 54 fairly well-understood which nutrients are essential in marine fish larvae diets (NRC, 2011), 55 quantitative requirements are largely unknown for most fish species and nutrients. This is 56 mainly due, among other reasons, to the inherent difficulty to manipulate and reliably control 57 dietary content of individual nutrient within live preys (i.e. rotifers, Artemia) to match marine 58 finfish larvae requirements (Støttrup, 2000; Conceição et al., 2010; Dhont et al., 2013). 59 Nutrients can be incorporated into live preys through the so-called "bioencapsulation" or 60 "enrichment" protocols (Sorgeloos et al., 2001), however, targeting a recommended level is 61 challenging due primarily to the metabolic activity of live preys towards the enrichment 62 products (Navarro et al., 1999; Reis et al., 2017) and the inherent variability associated with 63 the live prey enrichment process (Navarro et al., 1999; Hamre, 2016). Furthermore, the 64 nauplii's gut can only contain a defined amount of nutrient thus increasing the level of one 65 nutrient (i.e. lipid) can result in the decrease of another nutrient's level (e.g. micromineral).

The brine shrimp *Artemia* sp. is the most commonly used live feed in marine finfish
hatcheries (Conceição et al., 2010; Dhont et al., 2013). Unlike copepods, the natural preys of

68 marine fish larvae (Hunter, 1980), Artemia have a suboptimal nutritional profile which does not satisfy the requirements of marine fish larvae. It is well-established that Artemia is deficient 69 70 in certain essential lipids for larval stages of fish, particularly phospholipids and long-chain 71 polyunsaturated fatty acids (LC-PUFA) (i.e. arachidonic acid, ARA; eicosapentaenoic acid, EPA; docosahexaenoic acid, DHA) (Navarro et al., 1999; Monroig et al., 2003; Dhont et al., 72 73 2013). Along with essential lipids, successful Artemia enrichment strategies have also been tested and established for other essential nutrients such as methionine (Tonheim et al., 2000), 74 75 vitamins A, C and E (Monroig et al., 2007; Adloo, 2012), minerals including cobalt (Fehér et 76 al., 2013), iodine (Moren et al., 2006), manganese (Nguyen et al., 2008) and selenium (Se) 77 (Hamre et al., 2008a,b; Penglase et al., 2011). Importantly, Se is regarded as an essential trace 78 element for virtually all animal species (Hefnawy and Tórtora-Pérez, 2010) and it is a major 79 component of the glutathione peroxidase (GPx), which is involved in the regulation of the 80 antioxidant status in finfish by reducing hydrogen peroxide and hydroperoxides to their base 81 constituents (Lall, 2003; Pacitti et al., 2015). Moreover, Se has also been shown to play a 82 protective role by reducing oxidative stress caused by heavy metals such as copper (Cu), 83 resulting in enhanced immune response in fish (Lin and Shiau, 2007). Studies have shown that 84 dietary deficiencies of Se can result in reduced growth in channel catfish Ictalurus punctatus (Wang and Lovell, 1997) and increased mortality in rainbow trout Oncorhynchus mykiss 85 86 (Hilton et al., 1980). The NRC (2011) recommendations for Se dietary requirements in fish 87 vary across species but in average, Se dietary requirements for juvenile and adult stages are ~0.35 mg kg⁻¹ (NRC, 2011). Importantly, Se requirements in marine fish larvae are largely 88 89 unknown despite playing a crucial role as an antioxidant and subsequently reducing stress levels in fragile marine larvae (Saleh et al., 2014). Se levels in enriched Artemia nauplii are 90 usually around 2 mg kg⁻¹ (Ribeiro et al., 2012a,b), which is at the lower end of wild copepods 91 Se content (i.e. between 2 to 5 mg kg⁻¹) (Hamre et al., 2008b; Mæhre et al., 2013). This may 92

therefore not satisfy the dietary requirements of marine fish larvae (Solbakken et al., 2002;
NRC, 2011; Hamre et al., 2013). A study performed in Atlantic cod (*Gadus morhua*) showed
that larvae fed rotifers enriched with Se exhibited a higher survival compared to larvae fed the
control rotifers (Penglase et al., 2010). In addition, the inclusion of Se in marine fish larval
diets has been suggested to potentially offset the high oxidation risk from diets boosted in LCPUFA (Saleh et al., 2014).

99 Bioavailability of Se, as per other minerals (NRC, 2011), is greatly dependent upon its 100 form when accumulated in the diet (Pacitti et al., 2015). One of the most common forms of 101 dietary Se is sodium selenite (Na₂SeO₂, hereafter referred to as Na-Se), a highly water-soluble 102 inorganic compound. An alternative Se source is selenoyeast (Se-yeast), an organic source of 103 Se produced by exposing yeast (Saccharomyces cerevisiae) to Na-Se (Suhajda et al., 2000), 104 which results in an accumulation of selenomethionine (Se-Met). The latter (organic) form of 105 Se is regarded to be more bioavailable to organisms, thus explaining why it is broadly used as a livestock feed additive (Wang and Lovell, 1997; Rayman, 2004; Thiry et al., 2012). 106 107 Nevertheless, inappropriate dietary Se levels can induce toxicity effects. Se dietary toxicity levels has been reported in rainbow trout (Onchorynchus mykiss) at 13 mg Se kg⁻¹ when 108 presented as Na-Se (Hilton et al., 1980) and at levels of 20 mg Se kg⁻¹ when presented as Se-109 110 Met in hybrid striped bass (Morone chrysops $\times M$. saxatilis) (Jaramillo et al., 2009). In both 111 studies. Se toxicities resulted in growth impairment and elevated mortality.

Importantly, with a few exceptions (e.g. Penglase et al., 2010), most studies mentioned above were conducted in fish juveniles and information on Se requirements and toxicity levels in fish larvae remain scarce. The present study aimed to compare the effectiveness of different Se enrichment protocols for *Artemia* nauplii using different Se sources and determine how these impact on essential fatty acids enrichment using a commercially available enrichment

- 117 product. The effects of varying doses of Se-yeast were first tested then the efficiency of
- 118 different Se sources were studied on *Artemia* nauplii content in Se and essential fatty acids.

119 **2.** Materials and Methods

120 2.1.Artemia hatching and culture

121 Artemia cysts GSL (EG, Inve, Belgium) were decapsulated and hatched according to Sorgeloos 122 et al. (2001). Newly hatched Artemia nauplii were subsequently used for the enrichment 123 experiments. Enrichments were carried in 1-litre Imhoff cones using glass pipes to strongly aerate the cones from the bottom. The cones were subsequently placed in a 28 °C water bath. 124 125 After the 24 h hatching process, enriched nauplii were collected on a 100 µm sieve, rinsed with freshwater, and distributed in each cone at 300 nauplii ml⁻¹ for further enrichment. Artificial 126 seawater (32 ppt) (Instant Ocean, Virginia, USA), disinfected with Pyceze® (0.05 ml l⁻¹), was 127 128 dispensed in each 1-litre Imhoff cone to provide a final volume of 800 ml after the addition of 129 the enrichment products (see below). The enrichment experiments were run under constant 130 illumination of about 47 klx at the surface of the water.

131 2.2. Experiment 1: Effects of varying doses of Se-yeast on Se content of Artemia nauplii.

Newly hatched Artemia were first enriched with Sel-Plex (SP) (Alltech, Kentucky, USA), 132 during 4 h at different concentrations: 0 mg l⁻¹ (Treatment SP0), 12 mg l⁻¹ (Treatment SP12), 133 24 mg l⁻¹ (Treatment SP24) and 36 mg l⁻¹ (Treatment SP36). Each SP dose was tested in 134 135 triplicated 1-litre Imhoff cones (i.e. 4 treatments x 3 replicates). After 4 h, Larviva Multigain (MG) (BioMar, Denmark) was added to each of the 12 cones at 0.6 g l⁻¹ for a further 24 h. At 136 137 the end of the MG enrichment period, nauplii were collected on a 100 µm sieve, thoroughly 138 rinsed with freshwater to remove the excess of enrichment product, and gently dried on 139 absorbent paper before transferring them into universal sample tubes. The samples were frozen at -20 °C before being freeze-dried and stored at -20 °C for further analysis. 140

141 2.3.Experiment 2: Effects of different sources of Se on Artemia nauplii enrichment 142 efficiency.

143 Experiment 2 investigated the efficiency of two sources of Se to increase the content of Se in Artemia nauplii. Four enrichment diets were tested in triplicate: no addition of Se sources 144 (Treatment SP0); SP at 12 mg l⁻¹ for 4 h (Treatment SP12); sodium selenite Na-Se (Sigma 145 Aldrich, UK) at the Se equivalent dose (i.e. 24 µg l⁻¹) of Treatment SP12 for 4 h (Treatment 146 NS); soya lecithin emulsion (0.6 g l^{-1}) (Optima Health, UK) containing Na-Se (24 μ g l^{-1}) for 4 147 148 h (Treatment SL+NS). Treatments SP12, NS and SL+NS contained the same effective dose of Se $(24 \ \mu g \ l^{-1})$ irrespectively of the form under which Se was presented. For treatment SL+NS, 149 we used a soya lecithin emulsion, potentially creating lipid vesicles (liposomes) that can 150 151 encapsulate dissolved Se, as this has been proven to be a good strategy to deliver water-soluble 152 nutrients in Artemia nauplii (Monroig et al., 2007). The hereby tested sodium selenite was an 153 inorganic, water soluble compound with \geq 95.0 % purity. During the first 4 h, nothing was 154 added to enrichment medium of Treatment SP0, whereas the other treatments were enriched with Se as indicated above. After the 4 h, Larviva Multigain (0.6 g l⁻¹) was added to all 155 156 treatments for a further 24 h. Sample collection and storage was done as described in 157 Experiment 1.

158 2.4.Nutritional analysis

Total lipids (TL) from the enrichment products MG, SL and SP (Table 1), as well as freezedried *Artemia* samples collected from Experiments 1 and 2, were extracted according to Folch et al. (1957), with modifications as described by Monroig et al. (2006). Fatty acid methyl esters (FAME) from TL were prepared, extracted and purified according to Christie (2003). Identification and quantification of FAME were carried out using a gas chromatograph coupled with flame ionisation detection as previously described (Houston et al. 2017). Se concentration was determined after digestion of Sel-Plex, enrichment products and freeze-dried *Artemia* samples in AristAR nitric acid (VWR International, Pennsylvania, USA) in a microwave MARSXpress (CEM, North Carolina, USA) for 40 min (20 min ramping to 120 °C and 20 min holding that temperature). Digests were transferred into a volumetric flask and made up into x 25 dilutions with distilled water. Samples were analysed by Inductively Coupled Plasma Mass Spectrometry (ICP-MS, Thermo Scientific Model X Series 2, Massachusetts, USA) (Smedley et al., 2016).

172 2.5.Statistical analysis

173 All enrichment treatments in both experiments were carried out in triplicate cones (n = 3). 174 Biological and analytical data are expressed as means ± standard deviation (SD). Percentage 175 data were transformed using the arcsine square root function prior to statistical analysis. 176 Difference among treatments for total lipids and fatty acids were analysed by one-way ANOVA 177 followed by a Tukey post-hoc multiple comparison test at a significance level of $P \le 0.05$ (IBM 178 SPSS Statistics 23, New-York (state), USA).

179

180 **3. Results**

181 *3.1.Experiment 1: Effects of varying doses of Se-yeast on Se content of Artemia nauplii.*

The analysis of the unenriched *Artemia* nauplii revealed the presence of Se $(1.83 \pm 0.18 \text{ mg kg}^{-1} \text{ DW})$. In the experimental groups, Se concentration in *Artemia* nauplii increased linearly with increasing levels of selenoyeast Sel-Plex (Fig. 1). The equation of the linear regression was: $[Se]_{Artemia} = 3.79 \times [SelPlex]_{enrichment} - 2.42 (r^2 = 0.97)$ where $[Se]_{Artemia}$ is the Se content in the enriched *Artemia* (mg kg⁻¹ DW) and $[SelPlex]_{enrichment}$

187 is the dose of Sel-Plex used to prepare enrichment (mg l^{-1}). The Se content in *Artemia* nauplii

varied from $1.7 \pm 0.1 \text{ mg kg}^{-1}$ (Treatment SP0), defined as the basal level of Se found in the 188 189 non-enriched nauplii, to $12.4 \pm 1.0 \text{ mg kg}^{-1}$ (Treatment SP36), with significant differences between treatments (P < 0.05) (Fig. 1). Sel-Plex is a yeast-based product that contains lipids 190 (Table 1). However, the dose of Sel-Plex used in Experiment 1 (from 0 to 36 mg l⁻¹) did not 191 affect the TL contents of enriched Artemia nauplii (P > 0.05) (Table 2), which ranged between 192 $209.5 \pm 9.0 \text{ mg g}^{-1}$ (Treatment SP0) and $220.4 \pm 22.5 \text{ mg g}^{-1}$ (Treatment SP12). Similarly, no 193 194 statistical differences were observed in the levels of ARA, EPA and DHA, nor DHA/EPA ratios 195 in Artemia nauplii enriched with varying doses of Se (Table 2).

196 *3.2.Experiment 2: effects of different sources of Se on Artemia nauplii enrichment efficiency*

In Experiment 2, Se levels of *Artemia* from Treatments SP0 and SP12 (1.7 ± 0.1 and 4.3 ± 0.4 mg kg⁻¹, respectively) were consistent with results from Experiment 1 (1.70 and 4.17 mg kg⁻¹, respectively). Moreover, *Artemia* nauplii from NS and SL+NS treatments contained Se levels of 1.7 ± 0.0 and 1.7 ± 0.1 mg kg⁻¹, respectively, very similar to those of the control *Artemia* (Treatment SP0).

202 The different enrichment regimes resulted in variations in the TL content and fatty acid profiles of Artemia nauplii (Table 3). While nauplii from Treatments SP0, SP12 and NS 203 204 showed similar TL contents (P > 0.05), nauplii from Treatment SL+NS exhibited significantly 205 higher TL content (Table 3). In terms of fatty acid profiles, the nauplii enriched with SL+NS 206 showed a significantly higher n-6 fatty acid content (Table 3), largely due to the contribution 207 of 18:2n-6 (16.2 \pm 0.9 %) present in the soybean lecithin (Table 1). Consistently, the n-3/n-6 208 ratio was significantly lower in SL+NS nauplii (1.7 ± 0.1) compared to that of the other 209 treatments (~3.0). ARA levels were significantly lower in the nauplii SL+NS ($1.7 \pm 0.2 \%$) 210 than in the other treatments. The EPA contents of the SPO nauplii were overall the highest (4.9 \pm 0.2 %), but the difference was only significant when compared to the SL+NS nauplii (3.7 \pm 211 212 0.1 %). DHA contents were significantly higher in SP0 and SP12 nauplii (13.8 ± 0.7 and 13.2 213 \pm 0.5 %, respectively) compared to SL+NS (10.8 \pm 0.3 %). DHA/EPA ratios were not 214 significantly different between treatments.

215

4. Discussion

217 Nutritional deficiencies play an important role in explaining elevated mortalities in marine fish 218 during the early larval stages (Hamre et al., 2013), themselves resulting from a knowledge gap 219 in our understanding of the digestive development and physiology of marine larvae (Rønnestad 220 et al., 2013). Se is an essential trace element required for a variety of biological functions 221 throughout the entire fish life-cycle including early larval stages (Hamre et al., 2008a; Ribeiro 222 et al., 2012a). Copepods, natural preys of marine finfish larvae, contain relatively high levels 223 of Se compared to live preys and it is therefore critical to develop Se enrichment protocols to 224 guarantee that live preys, otherwise deficient in Se, provide adequate Se levels to meet larvae 225 requirements. Importantly, Se enrichment must be achieved along with the provision of 226 essential fatty acids, micronutrients which are typically encapsulated into live preys with 227 commercial products with very low or, even non-exiting, levels of Se.

228 The effects of varying doses of selenoyeast Sel-Plex, a commercial Se additive used for 229 animal feed, were first tested on the levels of Se and essential fatty acids of Artemia nauplii. 230 The control treatment in Exp. 1 (Treatment SP0) without Sel-Plex resulted in Artemia 231 containing 1.7 ± 0.1 mg Se kg⁻¹, similar to previously published results (Ribeiro et al., 2012b) 232 despite using a different enrichment protocol (i.e. 3 h instead of 20 h in the current study) and 233 enrichment product (i.e. DHA Selco instead of MG in the current study). In the control 234 treatment, Se concentrations of the nauplii reflected the background level of Se present in 235 unenriched Artemia, mainly provided by the enrichment product MG, which contained 2.2 mg Se kg⁻¹. Interestingly, the provision of selenoyeast Sel-Plex into the enrichment medium 236 237 resulted in increased levels of Se in Artemia nauplii. The relationship between Sel-Plex and 238 Artemia Se was linear thus enabling us to predict Se content in Artemia nauplii when enriched with a given Sel-Plex dose. SP12 nauplii showed Se contents $(4.2 \pm 0.1 \text{ mg kg}^{-1})$ within the 239 upper range of Se concentration reported in wild copepods (i.e. 5 mg kg⁻¹) (Hamre et al., 240 2008b), whereas SP24 and SP36 nauplii, enriched with Sel-Plex doses of 12 and 24 mg l⁻¹, 241 respectively, contained Se above 10 mg kg⁻¹. Thus, SP12 enrichment treatment resulted in a 242 243 diet with Se levels well below the potential dietary toxicity threshold observed in rainbow trout 244 and hybrid striped bass (> 10 mg kg⁻¹; Hilton et al., 1980; Jaramillo et al., 2009). However, SP24 and particularly SP36 treatments resulted in nauplii Se contents that could cause toxicity 245 246 for fish larvae. When compared to other published studies, the efficiency of Se enrichment 247 obtained in Experiment 1 differs from those reported by Ribeiro et al. (2012a). In the latter study, a Sel-Plex dose of 0.6 mg l⁻¹ resulted in a Se content in Artemia of 3.11 ± 0.27 mg kg⁻¹ 248 while in our study, a similar Se content in Artemia was obtained using 12 mg l⁻¹ of Sel-Plex 249 250 (4.17 mg kg⁻¹). While the reasons explaining such discrepancy remain unknown, Ribeiro's 251 study lacks details on the enrichment protocol used (e.g. rinsing Artemia prior sampling, 252 selenium content of Sel-Plex), which could help explain the differences observed in the 253 efficiency of Se incorporation in nauplii. In addition, the lack of data on the fatty acid composition of the enriched nauplii in Ribeiro et al. (2012a) does not allow us to clarify 254 255 whether the high Se incorporation correlated with a concomitant increase in essential fatty acids 256 within Artemia nauplii.

Results from Exp. 1 clearly showed that simultaneous delivery of Se and essential fatty acids is possible under our enrichment protocol. Nauplii from SP12 treatment contained, in addition to 4.2 mg Se kg⁻¹, markedly higher levels of essential fatty acids such as DHA (18.0 \pm 1.1 %). This data is consistent with an *Artemia* enrichment study that showed the high efficiency of Larviva Multigain at supplying DHA, in which levels of 21.8 \pm 0.7 % DHA postenrichment were obtained (Cavrois-Rogacki et al., 2019). Furthermore, none of the treatments with Sel-Plex significantly affected the levels of essential fatty acids (i.e. ARA, EPA and DHA) in the *Artemia* compared to the control, despite the potential dilution effect derived from the inclusion of LC-PUFA free lipids from yeast (Santomartino et al., 2017). Therefore, these results showed that Sel-Plex can be successfully used to enrich *Artemia* in Se while preserving the essential fatty acid contents achieved using commercial enrichment products. Our analyses suggested that Sel-Plex does contain traces of EPA and DHA, although their low levels (<0.2 %) do not appear to have a major contribution to the essential fatty acids of nauplii.

270 Although it is known that organic Se (e.g. selenomethionine) may be more easily 271 absorbed by living animals compared to inorganic forms (e.g. sodium selenite) (Wang and 272 Lovell, 1997; Izquierdo et al., 2017), high cost of the former can constitute a barrier to their 273 use at a large commercial scale. Cheaper inorganic source of Se have previously been tested 274 on fish larvae fed Se-enriched rotifers (Hamre et al., 2008a) and thus represent potential 275 alternatives. This was investigated in Exp. 2, in which NS and SL+NS treatments consisted of a Se dose of 24 μ g l⁻¹ (equivalent to Se contained in SP12 treatment with Sel-Plex) supplied as 276 277 dissolved sodium selenite. Results indicated that neither NS nor SL+NS treatment appeared to 278 be effective ways to enhance Se contents in Artemia nauplii since they did not differ from those 279 of control nauplii. While previous studies reported on the low efficiency of delivering dissolved 280 materials into Artemia (Tonheim et al., 2000; Monroig et al., 2007), it was somewhat 281 unexpected that delivering the same dose of Se encapsulated into phospholipid vesicles 282 (SL+NS) did not result in any increased Se enrichment efficiency despite Artemia being adapted to filtrate discrete particles. The reasons for such a result are unknown but it is 283 284 reasonable to believe that lipid vesicles produced with the soya lecithin source used in the 285 present study were leaky and did not retain the dissolved Se in the inner aqueous phase of the 286 vesicle. Other highly purified sources of phospholipids, particularly when constituted of more saturated fatty acyl chains, have proven to produce relatively stable vesicles with good 287

288 efficiency in delivering water soluble compounds into live preys (Hontoria et al., 1994; 289 Monroig et al., 2003, 2007). Thus, in spite of low Se incorporation into Artemia nauplii, 290 incorporation of soya lecithin was observed as evidenced by the increased levels of linoleic 291 acid (18:2n-6), its most abundant fatty acid, and the corresponding reduced percentage of other fatty acid contents including EPA, DHA and ARA. The importance of essential fatty acids for 292 293 marine fish larvae nutrition has been extensively reviewed (Izquierdo, 1996; Tocher 2010, 294 2015) and it is therefore crucial that the enrichment of live preys with micronutrients is not 295 detrimental to the fatty acids' levels of live feed.

296 One important aspect of live prey enrichment is its reproducibility and predictability, 297 which is essential in commercial hatcheries to ensure a constant daily production of high-298 quality live preys. Producing enriched Artemia with consistent levels of essential nutrients such 299 as LC-PUFA can be challenging as shown in previous studies (Navarro et al., 1999; Monroig 300 et al., 2006). Importantly, the results from the present study suggests that enrichment of 301 Artemia nauplii with selenoyeast Sel-Plex is highly reproducible, according to the consistent 302 levels of Se found in nauplii from the SP12 treatment in two independent experiments (Exp. 1 303 to 2). While this proves the reproducibility of the method on a small-scale enrichment system, 304 further trials are necessary to confirm the reproducibility at commercial scale.

The study showed that it is possible to enrich *Artemia* with targeted levels of Se using selenoyeast Sel-Plex. Enriching *Artemia* nauplii with 12 mg of Sel-Plex per litre for 4 h prior to a 24 h enrichment with LC-PUFA rich commercial diets produces *Artemia* with Se contents similar to those found in the natural preys (wild zooplankton) of marine fish larvae and high levels of essential fatty acids. The use of inorganic Se was not an effective strategy to enrich *Artemia* nauplii even when it was delivered through phospholipid vesicles. In the case of soya lecithin, the use of low-quality liposomes in the experiment could be a potential cause and should be deeper investigated. Ultimately, these results can be implemented to the *Artemia*enrichment protocols of any marine fish species.

314 Acknowledgments

- 315 The project, T. Cavrois Rogacki PhD studentship and A. Rolland MSc studentship were co-
- funded by the Scottish Aquaculture Innovation Centre (SAIC) and University of Stirling.

317

319 **References**

Adloo, M.N., 2012. Effects of Feeding Enriched *Artemia fransiscana* with HUFA, Vitamin C
and E on Growth Performance, Survival and Stress Resistance of Yellowfin Seabream
Larvae. J. Aquac. Res. Dev., 3:157.

- Atalah, E., Hernández-Cruz, C.M., Benítez-Santana, T., Ganga, R., Roo, J., Izquierdo, M.,
 2011. Importance of the relative levels of dietary arachidonic acid and eicosapentaenoic
 acid for culture performance of gilthead seabream (*Sparus aurata*) larvae. Aquac. Res.
 42, 1279–1288.
- Bjornsdottir, R., Johannsdottir, J., Coe, J., Smaradottir, H., Agustsson, T., Sigurgisladottir, S.,
 Gudmundsdottir, B.K., 2009. Survival and quality of halibut larvae (*Hippoglossus hippoglossus* L.) in intensive farming: Possible impact of the intestinal bacterial
 community. Aquaculture 286, 53–63.
- Cavrois-Rogacki, T., Davie, A., Kiné"rg, E., Esnault, S., Migaud, H., Monroig, O., 2019.
 Short-term lecithin enrichments can enhance the phospholipid and DHA contents of the
 polar lipid fraction of *Artemia* nauplii. Aquaculture 510, 122-130.
- Christie, W.W., 2003. Isolation, Separation, Identification and Lipidomic Analysis, The Oily
 Press, Bridgwater, UK. Oily Press, an imprint of PJ Barnes & Associates.
- Conceição, L.E.C., Yúfera, M., Makridis, P., Morais, S., Dinis, M.T., 2010. Live feeds for early
 stages of fish rearing. Aquac. Res. 41, 613–640.
- 338 Dhont, J., Dierckens, K., Støttrup, J., Van Stappen, G., Wille, M., Sorgeloos, P., 2013. Rotifers,
 339 *Artemia* and copepods as live feeds for fish larvae in aquaculture, in: Advances in
 340 Aquaculture Hatchery Technology. Elsevier, pp. 157–202.

341	Fehér, M., Baranyai, E., Simon, E., Bársony, P., Szucs, I., Posta, J., Stündl, L., 2013. The
342	interactive effect of cobalt enrichment in Artemia on the survival and larval growth of
343	barramundi, Lates calcarifer. Aquaculture 414–415, 92–99.

- Folch, J., Lees, M., Sloane Stanley, G.H., 1957. A simple method for the isolation and
 purification of total lipids from animal tissues. J. Biol. Chem 226, 497–509.
- Hamre, K., 2016. Nutrient profiles of rotifers (*Brachionus* sp.) and rotifer diets from four
 different marine fish hatcheries. Aquaculture 450, 136–142.
- Hamre, K., Mollan, T.A., Sæle, Ø., Erstad, B., 2008a. Rotifers enriched with iodine and
 selenium increase survival in Atlantic cod (*Gadus morhua*) larvae. Aquaculture 284, 190–
 195.
- Hamre, K., Srivastava, A., Rønnestad, I., Mangor-Jensen, A., Stoss, J., 2008b. Several
 micronutrients in the rotifer *Brachionus* sp. may not fulfil the nutritional requirements of
 marine fish larvae. Aquac. Nutr. 14, 51–60.
- Hamre, K., Yúfera, M., Rønnestad, I., Boglione, C., Conceição, L.E.C., Izquierdo, M., 2013.
- Fish larval nutrition and feed formulation: Knowledge gaps and bottlenecks for advancesin larval rearing. Rev. Aquac. 5, S26–S58.
- Hawkyard, M., Stuart, K., Langdon, C., Drawbridge, M., 2016. The enrichment of rotifers
 (*Brachionus plicatilis*) and *Artemia franciscana* with taurine liposomes and their
 subsequent effects on the larval development of California yellowtail (*Seriola lalandi*).
 Aquac. Nutr. 22, 911–922.
- Hefnawy, A.E.G., Tórtora-Pérez, J.L., 2010. The importance of selenium and the effects of its
 deficiency in animal health. Small Rumin. Res. 89, 185–192.

- Hilton, J.W., Hodson, P. V, Slinger, S.J., 1980. The requirement and toxicity of selenium in
 rainbow trout (*Salmo gairdneri*). J. Nutr. 110, 2527–2535.
- Hontoria, F., Crowe, J.H., Crowe, L.M., Amat, F., 1994. Potential use of liposomes in
 larviculture as a delivery system through *Artemia* nauplii. Aquaculture 127, 255–264.
- 367 Houston, S.J.S., Karalazos, V., Tinsley, J., Betancor, M.B., Martin, S.A.M., Tocher, D.R.,
- Monroig, O., 2017. The compositional and metabolic responses of gilthead seabream (*Sparus aurata*) to a gradient of dietary fish oil and associated n-3 long-chain PUFA
- 370 content. Br. J. Nutr. 118, 1010–1022.
- Hunter, J., 1980. Feeding Ecobgy and Predation of Marine Fish Larvae, R. Lasker. ed,
 Swfsc.Noaa.Gov.
- 373 Izquierdo, M.S., 1996. Essential fatty acid requirements of cultured marine fish larvae. Aquac.
 374 Nutr. 2, 183–191.
- Izquierdo, M.S., Ghrab, W., Roo, J., Hamre, K., Hernández-Cruz, C.M., Bernardini, G.,
 Terova, G., Saleh, R., 2017. Organic, inorganic and nanoparticles of Se, Zn and Mn in
 early weaning diets for gilthead seabream (*Sparus aurata*; Linnaeus, 1758). Aquac. Res.
 48, 2852–2867.
- Jaramillo, F., Peng, L., Gatlin, D.M., 2009. Selenium nutrition of hybrid striped bass (*Morone chrysops × M. saxatilis*) bioavailability, toxicity and interaction with vitamin e. Aquac.
 Nutr. 15, 160–165.
- Lall, S.P., 2003. The Minerals, in: Fish Nutrition. Elsevier, pp. 259–308.
- 383 Lin, Y.H., Shiau, S.Y., 2007. The effects of dietary selenium on the oxidative stress of grouper,
- *Epinephelus malabaricus*, fed high copper. Aquaculture 267, 38–43.

- Mæhre, H.K., Hamre, K., Elvevoll, E.O., 2013. Nutrient evaluation of rotifers and zooplankton:
 Feed for marine fish larvae. Aquac. Nutr. 19, 301–311.
- Monroig, Ó., Navarro, J.C., Amat, F., González, P., Bermejo, A., Hontoria, F., 2006.
 Enrichment of *Artemia* nauplii in essential fatty acids with different types of liposomes
 and their use in the rearing of gilthead sea bream (*Sparus aurata*) larvae. Aquaculture 251,
 491–508.
- Monroig, Ó., Navarro, J.C., Amat, F., Hontoria, F., 2007. Enrichment of *Artemia* nauplii in
 vitamin A, vitamin C and methionine using liposomes. Aquaculture 269, 504–513.
- Monroig, Ó., Navarro, J.C., Amat, I., González, P., Amat, F., Hontoria, F., 2003. Enrichment
 of *Artemia* nauplii in PUFA, phospholipids, and water-soluble nutrients using liposomes.
 Aquac. Int. 11, 151–161.
- Moren, M., Opstad, I., Van Der Meeren, T., Hamre, K., 2006. Iodine enrichment of *Artemia*and enhanced levels of iodine in Atlantic halibut larvae (*Hippoglossus hippoglossus* L.)
 fed the enriched *Artemia*. Aquac. Nutr. 12, 97–102.
- Navarro, J.C., Henderson, R.J., McEvoy, L.A., Bell, M. V., Amat, F., 1999. Lipid conversions
 during enrichment of *Artemia*. Aquaculture 174, 155–166.
- 401 Nguyen, V.T., Satoh, S., Haga, Y., Fushimi, H., Kotani, T., 2008. Effect of zinc and manganese
- 402 supplementation in *Artemia* on growth and vertebral deformity in red sea bream (*Pagrus*
- 403 *major*) larvae. Aquaculture 285, 184–192.
- 404 NRC, 2011. Nutrient Requirements of Fish and Shrimp, Animal nut. ed. National Academies
 405 Press, Washington, D.C.
- 406 Øie, G., Galloway, T., Sørøy, M., Holmvaag Hansen, M., Norheim, I.A., Halseth, C.K., Almli,

407	M., Berg, M., Gagnat, M.R., Wold, P.A., Attramadal, K., Hagemann, A., Evjemo, J.O.,
408	Kjørsvik, E., 2015. Effect of cultivated copepods (Acartia tonsa) in first-feeding of
409	Atlantic cod (Gadus morhua) and ballan wrasse (Labrus bergylta) larvae. Aquac. Nutr.
410	23, 3–17.

- Pacitti, D., Lawan, M.M., Sweetman, J., Martin, S.A.M., Feldmann, J., Secombes, C.J., 2015.
 Selenium supplementation in fish: A combined chemical and biomolecular study to
 understand Sel-Plex assimilation and impact on selenoproteome expression in rainbow
 trout (*Oncorhynchus mykiss*). PLoS One 10, e0127041.
- 415 Penglase, S., Hamre, K., Sweetman, J.W., Nordgreen, A., 2011. A new method to increase and
 416 maintain the concentration of selenium in rotifers (*Brachionus* spp.). Aquaculture 315,
 417 144–153.
- Penglase, S., Nordgreen, A., van der Meeren, T., Olsvik, P.A., Sæle, Sweetman, J.W.,
 Baeverfjord, G., Helland, S., Hamre, K., 2010. Increasing the level of selenium in rotifers
 (*Brachionus plicatilis* 'Cayman') enhances the mRNA expression and activity of
 glutathione peroxidase in cod (*Gadus morhua* L.) larvae. Aquaculture 306, 259–269.
- Rayman, M.P., 2004. The use of high-selenium yeast to raise selenium status: how does it
 measure up? Br. J. Nutr. 92, 557.
- 424 Reis, D.B., Acosta, N.G., Almansa, E., Navarro, J.C., Tocher, D.R., Andrade, J.P., Sykes, A.
- 425 V., Rodríguez, C., 2017. Comparative study on fatty acid metabolism of early stages of
- 426 two crustacean species: Artemia sp. metanauplii and Grapsus adscensionis zoeae, as live
- 427 prey for marine animals. Comp. Biochem. Physiol. Part B Biochem. Mol. Biol. 204, 53–
 428 60.
- 429 Ribeiro, A.R.A., Ribeiro, L., Sæle, Dinis, M.T., Moren, M., 2012a. Iodine and selenium

- 430 supplementation increased survival and changed thyroid hormone status in Senegalese
 431 sole (*Solea senegalensis*) larvae reared in a recirculation system. Fish Physiol. Biochem.
 432 38, 725–734.
- Ribeiro, A.R.A., Ribeiro, L., Saele, Ø., Hamre, K., Dinis, M.T., Moren, M., 2012b. Selenium
 supplementation changes glutathione peroxidase activity and thyroid hormone production
 in Senegalese sole (*Solea senegalensis*) larvae. Aquac. Nutr. 18, 559–567.
- Rønnestad, I., Yúfera, M., Ueberschär, B., Ribeiro, L., Sæle, Ø., Boglione, C., 2013. Feeding
 behaviour and digestive physiology in larval fish: Current knowledge, and gaps and
 bottlenecks in research. Rev. Aquac. 5.
- Saleh, R., Betancor, M.B., Roo, J., Montero, D., Zamorano, M.J., Izquierdo, M., 2014.
 Selenium levels in early weaning diets for gilthead seabream larvae. Aquaculture 426–
 427, 256–263.
- 442 Santomartino, R., Riego-Ruiz, L., Bianchi, M.M., 2017. Three, two, one yeast fatty acid
 443 desaturases: regulation and function. World J. Microbiol. Biotechnol. 33, 89.
- Smedley, M.A., Clokie, B.G.J., Migaud, H., Campbell, P., Walton, J., Hunter, D., Corrigan,
 D., Taylor, J.F., 2016. Dietary phosphorous and protein supplementation enhances
 seawater growth and reduces severity of vertebral malformation in triploid Atlantic
 salmon (*Salmo salar* L.). Aquaculture 451, 357–368.
- 448 Solbakken, J.S., Berntssen, M.H.G., Norberg, B., Pittman, K., Hamre, K., Berntessen, M.H.G.,
- 449 Norberg, B., Pittman, K., Hamre, K., 2002. Different iodine and thyroid hormone levels
- between Atlantic halibut larvae fed wild zooplankton or *Artemia* from first exogenous
 feeding until post metamorphosis. J. Fish Biol. 61, 1345–1362.
- 452 Sorgeloos, P., Dhert, P., Candreva, P., 2001. Use of the brine shrimp, Artemia spp., in marine

- 453 fish larviculture. Aquaculture 200, 147–159.
- 454 Støttrup, J.G., 2000. The elusive copepods: Their production and suitability in marine 455 aquaculture. Aquac. Res. 31, 703–711.
- 456 Suhajda, Á., Hegóczki, J., Janzsó, B., Pais, I., Vereczkey, G., 2000. Preparation of selenium
- 457 yeasts I. Preparation of selenium-enriched *Saccharomyces cerevisiae*. J. Trace Elem.
 458 Med. Biol. 14, 43–47.
- Thiry, C., Ruttens, A., De Temmerman, L., Schneider, Y.J., Pussemier, L., 2012. Current
 knowledge in species-related bioavailability of selenium in food. Food Chem. 130, 767–
 784.
- 462 Tocher, D.R., 2015. Omega-3 long-chain polyunsaturated fatty acids and aquaculture in
 463 perspective. Aquaculture 449, 94–107.
- 464 Tocher, D.R., 2010. Fatty acid requirements in ontogeny of marine and freshwater fish. Aquac.
 465 Res.
- 466 Tonheim, S.K., Koven, W., Rønnestad, I., 2000. Enrichment of *Artemia* with free methionine.
 467 Aquaculture 190, 223–235.
- Villamizar, N., García-Alcazar, A., Sánchez-Vázquez, F.J., 2009. Effect of light spectrum and
 photoperiod on the growth, development and survival of European sea bass
 (*Dicentrarchus labrax*) larvae. Aquaculture 292, 80–86.
- Wang, C., Lovell, R.T., 1997. Organic selenium sources, selenomethionine and selenoyeast,
 have higher bioavailability than an inorganic selenium source, sodium selenite, in diets
 for channel catfish (*Ictalurus punctatus*). Aquaculture 152, 223–234.

Table 1. Contents of Selenium (Se), total lipids, and selected fatty acids in enrichment products

475 used in Experiments 1 and 2.

	Larviva Multigain	Soya lecithin	Sel-Plex
Se (mg kg ⁻¹)	2.2	ND	2000
Total lipids (mg g ⁻¹)	397.4	808.6	13.4
Fatty acids (% of total)			
14:0	6.1	0.1	0.6
15:0	0.4	ND	0.4
16:0	32.3	19.0	15.3
18:0	0.9	4.2	5.2
Saturates	40.1	24.2	22.2
16:1n-9	0.3	ND	NE
16:1n-7	0.1	0.8	25.1
18:1n-9	1.9	10.2	28.5
18:1n-7	ND	2.0	NE
Monounsaturates	2.3	13.4	54.3
18:2n-6	2.2	54.1	14.7
18:3n-6	0.2	ND	0.1
20:4n-6	1.2	ND	0.1
22:5n-6	14.4	ND	NE
Total n-6	18.4	54.3	15.0
18:3n-3	0.3	6.9	3.9
18:4n-3	0.3	0.1	0.2
20:3n-3	0.1	ND	0.1
20:4n-3	0.7	ND	NE
20:5n-3	0.8	ND	0.1
22:5n-3	0.3	ND	NE
22:6n-3	36.6	ND	0.2
Total n-3	39.0	7.0	4.3
DW, dry weight; ND, not	detected.		

Table 2. Total lipids and selected fatty acids of *Artemia* nauplii from Experiment 1 enriched for 4 h with different doses of selenoyeast Sel-Plex (SP0: 0 mg l⁻¹; SP12: 12 mg l⁻¹; SP24: 24 mg l⁻¹; SP36: 36 mg l⁻¹) followed by a 24 h enrichment with Larviva Multigain (0.6 g l⁻¹). Data are expressed as means \pm standard deviations (n = 3). Differences among treatments were analysed by a one-way ANOVA followed by a Tukey post-hoc test ($P \le 0.05$).

Treatment	SP0	SP12	SP24	SP36
Total lipids (mg g ⁻¹ DW)	209.5 ± 9.0	220.4 ± 22.5	212.5 ± 8.7	211.9 ± 8.9
Fatty acids (% of total)				
14:0	1.6 ± 0.1	1.5 ± 0.0	1.7 ± 0.3	1.7 ± 0.3
15:0	0.2 ± 0.0	0.2 ± 0.0	0.2 ± 0.0	0.2 ± 0.0
16:0	14.6 ± 0.9	14.2 ± 0.3	16 ± 1.9	16.8 ± 3.0
18:0	4.4 ± 0.1	4.3 ± 0.3	4.6 ± 0.3	5.1 ± 1.2
Saturates	21.2 ± 1.1	20.6 ± 0.6	22.8 ± 2.5	24.3 ± 4.6
16:1n-9	0.5 ± 0.0	0.4 ± 0.1	0.5 ± 0.0	0.5 ± 0.2
16:1n-7	1.3 ± 0.1	1.2 ± 0.1	1.5 ± 0.0	1.5 ± 0.6
18:1n-9	13.4 ± 0.4	12.7 ± 0.9	14.6 ± 0.1	15.7 ± 4.8
18:1n-7	4.0 ± 0.1	3.9 ± 0.2	4.3 ± 0.0	4.7 ± 1.3
Monounsaturates	19.7 ± 0.6	18.8 ± 1.4	21.3 ± 0.1	23.0 ± 7.0
18:2n-6	4.7 ± 0.2	4.6 ± 0.1	4.8 ± 0.3	4.8 ± 0.3
18:3n-6	0.4 ± 0.0	0.4 ± 0.0	0.4 ± 0.0	0.4 ± 0.0
20:4n-6	2.6 ± 0.1	2.8 ± 0.3	2.1 ± 0.2	2.1 ± 1.5
22:5n-6	6.7 ± 0.2	7.0 ± 0.6	5.6 ± 0.5	4.9 ± 3.0
Total n-6	14.8 ± 0.3	15.4 ± 0.9	13.3 ± 0.5	12.7 ± 4.3
18:3n-3	16.9 ± 0.6	16.7 ± 0.3	18.5 ± 2.8	18.6 ± 2.2
18:4n-3	2.5 ± 0.1	2.4 ± 0.1	2.9 ± 0.6	2.8 ± 0.7
20:3n-3	0.8 ± 0.0	0.8 ± 0.0	0.8 ± 0.1	0.8 ± 0.0
20:4n-3	0.9 ± 0.0	0.9 ± 0.0	0.9 ± 0.1	0.8 ± 0.2
20:5n-3	5.4 ± 0.2	5.9 ± 0.6	4.4 ± 0.4	4.3 ± 2.9
22:5n-3	0.4 ± 0.0	0.3 ± 0.2	0.2 ± 0.2	0.2 ± 0.3
22:6n-3	17.3 ± 0.4	18.0 ± 1.1	14.5 ± 1.3	12.2 ± 6.9
Total n-3	44.1 ± 0.8	45.0 ± 1.2	42.2 ± 2.5	39.7 ± 7.4
n-3/n-6	3.0 ± 0.1	2.9 ± 0.1	3.2 ± 0.2	3.2 ± 0.5
DHA/EPA	3.2 ± 0.0	3.1 ± 0.2	3.3 ± 0.1	3.0 ± 0.4
Total FA (mg g ⁻¹ DW)	137.6 ± 11.0	136.0 ± 7.4	129.1 ± 2.8	116.5 ± 25.2

DW: dry weight; FA: fatty acids.

487

488

490	Table 3. Total lipids and selected fatty acids in the Artemia nauplii from Experiment 2 treated
491	with different enrichment diets (MG: Larviva Multigain; SP: Sel-Plex; NS: sodium selenite;
492	SL: soya lecithin emulsion). Data are expressed as means \pm standard deviations ($n = 3$).
493	Differences in fatty acid contents among treatments were analysed by a one-way ANOVA
494	followed by a Tukey post-hoc test ($P \le 0.05$). Variables that do not share the same superscript
495	letter within a row are significantly different from each other.

			a	
Treatment	SP0	SP12	NS	SL+NS
Total lipids (mg g ⁻¹ DW)	177.1 ± 8.6^{a}	180.1 ± 2.4^{a}	184.7 ± 2.9^{a}	$213.9\pm7.8^{\rm b}$
Fatty acids (% of total)				
14:0	1.3 ± 0.0^{ab}	1.3 ± 0.1^{ab}	$1.5\pm0.0^{\mathrm{b}}$	$1.2\pm0.1^{\rm a}$
15:0	0.2 ± 0.0	0.2 ± 0.0	0.2 ± 0.0	0.2 ± 0.0
16:0	13.1 ± 0.5	13.0 ± 0.2	14.8 ± 2.1	13.0 ± 0.3
18:0	4.3 ± 0.2	4.2 ± 0.1	4.7 ± 0.6	4.2 ± 0.1
Saturates	19.3 ± 0.7	18.9 ± 0.3	21.6 ± 2.8	19.0 ± 0.3
16:1n-9	0.6 ± 0.0	0.6 ± 0.0	0.4 ± 0.2	0.5 ± 0.0
16:1n-7	1.1 ± 0.9	1.6 ± 0.0	1.5 ± 0.1	1.3 ± 0.1
18:1n-9	14.8 ± 0.2^{ab}	$15.1\pm0.4^{\text{b}}$	14.9 ± 0.5^{ab}	$13.7\pm0.6^{\rm a}$
18:1n-7	4.4 ± 0.1^{b}	$4.5\pm0.1^{\rm b}$	$4.5\pm0.2^{\rm b}$	$3.8\pm0.1^{\text{a}}$
Monounsaturates	$21.3\pm0.6^{\text{b}}$	$22.1\pm0.4^{\text{b}}$	21.7 ± 0.7^{b}	$19.7\pm0.7^{\rm a}$
18:2n-6	$5.2\pm0.2^{\rm a}$	$5.3\pm0.1^{\rm a}$	$5.2\pm0.1^{\rm a}$	$16.2\pm0.9^{\text{b}}$
18:3n-6	0.4 ± 0.1	0.4 ± 0.0	0.4 ± 0.0	0.4 ± 0.0
20:4n-6	$2.3\pm0.2^{\text{b}}$	$2.2\pm0.1^{\text{b}}$	$2.3\pm0.1^{\text{b}}$	1.7 ± 0.2^{a}
22:5n-6	$5.3\pm0.1^{\text{b}}$	$4.9\pm0.2^{\text{b}}$	4.8 ± 0.4^{ab}	$4.3\pm0.1^{\text{a}}$
Total n-6	$13.7\pm0.0^{\rm a}$	$13.2\pm0.2^{\rm a}$	$13.1\pm0.4^{\rm a}$	23.0 ± 0.8^{b}
18:3n-3	21.4 ± 0.4^{b}	$22.1\pm0.5^{\text{b}}$	21.4 ± 1.1^{b}	$19.3\pm0.6^{\rm a}$
18:4n-3	$3.3\pm0.1^{\text{b}}$	$3.4\pm0.1^{\text{b}}$	3.3 ± 0.1^{b}	$2.7\pm0.2^{\rm a}$
20:3n-3	$0.8\pm0.0^{\rm b}$	$0.8\pm0.0^{\rm b}$	$0.8\pm0.0^{\rm b}$	$0.7\pm0.0^{\mathrm{a}}$
20:4n-3	$0.9\pm0.0^{\rm b}$	$0.9\pm0.0^{\rm b}$	$0.9\pm0.0^{\rm b}$	$0.7\pm0.0^{\mathrm{a}}$
20:5n-3	$4.9\pm0.2^{\text{b}}$	4.7 ± 0.1^{ab}	4.5 ± 0.7^{ab}	$3.7\pm0.1^{\rm a}$
22:5n-3	0.2 ± 0.2	0.2 ± 0.2	0.3 ± 0.0	0.1 ± 0.1
22:6n-3	13.8 ± 0.7^{b}	$13.2\pm0.5^{\text{b}}$	12.0 ± 1.4^{ab}	10.8 ± 0.3^{a}
Total n-3	$45.4 \pm 1.0^{\text{b}}$	$45.3\pm0.1^{\text{b}}$	$43.1\pm2.6^{\text{b}}$	$38.0\pm0.7^{\rm a}$
n-3/n-6	$3.3\pm0.1^{\text{b}}$	$3.4\pm0.1^{\text{b}}$	$3.3\pm0.2^{\rm b}$	$1.7\pm0.1^{\rm a}$
DHA/EPA	2.8 ± 0.3	2.8 ± 0.1	2.7 ± 0.2	2.9 ± 0.1
Total FA (mg g ⁻¹ DW)	$122.1\pm7.0^{\rm a}$	$122.7\pm1.3^{\rm a}$	121.4 ± 2.4^a	155.8 ± 12.1^{b}

DW: dry weight; FA: fatty acids.

Figure 1. Selenium (Se) concentration of *Artemia* nauplii (μ g Se g⁻¹ dry weight, "DW") from Experiment 1 enriched 4 h with different dose of selenoyeast Sel-Plex (SP0: 0 mg l⁻¹; SP12: 12 mg l⁻¹; SP24: 24 mg l⁻¹; SP36: 36 mg l⁻¹) followed by a 24 h enrichment with Larviva Multigain (0.6 g l⁻¹). Data are expressed as means ± standard deviations (*n* = 3). Differences in Se contents among treatments were analysed by a one-way ANOVA followed by a Tukey post-hoc test (*P* ≤ 0.05). Treatments with different superscripts are significantly different from each other.

Figure 2. Selenium (Se) concentration of *Artemia* nauplii (µg Se g⁻¹ dry weight, "DW") from 504 Experiment 2 enriched with different treatments. SP0: SP (0 mg l⁻¹, 4 h); SP12: SP (12 mg l⁻¹, 505 4 h); NS: NS (24 μ g l⁻¹, 4 h); SL+NS: SL emulsion (0.6 mg l⁻¹) + NS (24 μ g l⁻¹ emulsion). All 506 treatments were followed by a second enrichment with MG (0.6 g l⁻¹, 24 h) Data are expressed 507 508 as means \pm standard deviations (n = 3). Differences in fatty acid contents among treatments 509 were analysed by a one-way ANOVA followed by a Tukey post-hoc test ($P \le 0.05$). Variables 510 that do not share the same superscript letter within a row are significantly different from each 511 other.

Figure 1



