



Effect of music listening on hypertonia in neurologically impaired patients—systematic review

Tamaya Van Crieking¹, Kristiaan D’Août², Jonathon O’Brien³ and Eduardo Coutinho⁴

¹ Department of Rehabilitation Sciences and Physiotherapy, REVAKI/MOVANT, Universiteit Antwerpen, Antwerp, Belgium

² Department of Musculoskeletal Biology, University of Liverpool, Liverpool, United Kingdom

³ School of Health Sciences, University of Liverpool, Liverpool, United Kingdom

⁴ Applied Music Research Lab, Department of Music, University of Liverpool, Liverpool, United Kingdom

ABSTRACT

Background. As music listening is able to induce self-perceived and physiological signs of relaxation, it might be an interesting tool to induce muscle relaxation in patients with hypertonia. To this date effective non-pharmacological rehabilitation strategies to treat hypertonia in neurologically impaired patients are lacking. Therefore the aim is to investigate the effectiveness of music listening on muscle activity and relaxation.

Methodology. The search strategy was performed by the PRISMA guidelines and registered in the PROSPERO database (no. 42019128511). Seven databases were systematically searched until March 2019. Six of the 1,684 studies met the eligibility criteria and were included in this review. Risk of bias was assessed by the PEDro scale. In total 171 patients with a variety of neurological conditions were included assessing hypertonia with both clinical and biomechanical measures.

Results. The analysis showed that there was a large treatment effect of music listening on muscle performance (SMD 0.96, 95% CI [0.29–1.63], $I^2 = 10\%$, $Z = 2.82$, $p = 0.005$). Music can be used as either background music during rehabilitation (dual-task) or during rest (single-task) and musical preferences seem to play a major role in the observed treatment effect.

Conclusions. Although music listening is able to induce muscle relaxation, several gaps in the available literature were acknowledged. Future research is in need of an accurate and objective assessment of hypertonia.

Submitted 23 July 2019

Accepted 18 November 2019

Published 19 December 2019

Corresponding author

Kristiaan D’Août,
Kristiaan.daout@liverpool.ac.uk

Academic editor

Jafri Abdullah

Additional Information and
Declarations can be found on
page 15

DOI 10.7717/peerj.8228

 Copyright
2019 Van Crieking et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Neuroscience, Kinesiology, Neurology

Keywords Hypertonia, Spasticity, Music, Electromyography, Relaxation, Neurology

INTRODUCTION

A common symptom of upper motor neuron syndromes is hypertonia which is defined as an increased muscle tone, while lower motor neuron syndromes are related to hypotonia or decreased muscle tone. Two types of hypertonia can be distinguished: pyramidal or extrapyramidal hypertonia. First, pyramidal hypertonia also known as spasticity is defined as “a velocity-dependent increase in muscle tone as a result hyper excitability of the stretch reflex” (*Lance, 1980*) and is mostly pronounced during voluntary movements.

Second, extrapyramidal hypertonia or rigidity, is not speed-dependent and is therefore also present during slow passive movement (*Singer et al., 2010*). As inhibition of supraspinal inputs is impaired, antagonistic muscles become less effective in responding to agonist muscle contractions (*Gracies, 2005a; Gracies, 2005b; Singer et al., 2010*). This can either lead to prolonged shortening of a muscle as the joint is fixed in a certain position (a.k.a. contracture) (*Gracies, 2005b*) or it can lead to simultaneous co-contraction of agonist and antagonist muscles which results in an immediate resistance of the movement (*Singer et al., 2010*). Contractions are the result of decreased muscle mass and sarcomeres and an accumulation of connective tissue and fat (*Gracies, 2005a; Gracies, 2005b*).

Spasticity-related interventions mainly consist of pharmacological and/or physio- and occupational therapeutic approaches (*Katz, 1988; Nair & Marsden, 2014; Rekand, 2010; Thompson et al., 2005*). Pharmacological treatment consists of oral anti-spasticity drugs, botulinum toxin injections or intrathecal baclofen pump. Physio- and occupational therapeutic management consists of stretching/splinting and positioning as preventive care for contractures and to preserve comfort. Subsequently, exercise therapy mainly consists of reducing abnormal sensory inputs and providing adequate muscle length. However, high-quality evidence for supporting the effectiveness of these rehabilitation techniques is lacking (*Amatya et al., 2013; Khan et al., 2017; Nair & Marsden, 2014*). A low quality for evidence was found for rehabilitation programs, electrical stimulation, physical activity programs, vibration therapy, stretching and passive movement in neurological patients (*Khan et al., 2017*). There is a clear need for effective non-pharmacological rehabilitation strategies to treat spasticity. In addition, there is only a limited amount of evidence regarding the treatment of rigidity and this relates mainly to the general management of Parkinson's disease (*Diamond & Jankovic, 2006; Guttman, Kish & Furukawa, 2003*).

Music interventions are gaining popularity in rehabilitation programs and are able to induce self-perceived and physiological outcomes of relaxation in healthy adults (*Largo-Wight, O'Hara & Chen, 2016; Madson & Silverman, 2010; Staum & Brotons, 2000; Stratton & Zalanowski, 1984*), hospitalized (*Krout, 2001; Sand-Jecklin & Emerson, 2010*), burn (*Tan et al., 2010*), psychiatric (*Thaut, 1989*), oncology (*Cooper & Foster, 2008*) and neurological patients (*Sihvonen et al., 2017*). Music as an intervention is mostly empowered by its patient-centered character, it is a pleasurable intervention which has the secondary benefit of eliciting arousal and emotional responses. Listening to music activates cortical and paralimbic areas related to neural systems of reward and emotion (*Blood & Zatorre, 2001*), which makes music an intervention which can be rewarding and motivating and at the same time regulate emotions, arousal and cognitive functions (*Sarkamo, 2018*). So music interventions might be a good multi-modal treatment option for inducing muscle relaxation in neurological patients with hypertonia, and might lead to a better therapy compliance by its enjoyable character.

Music interventions can be categorized into two types of interventions—passive or receptive (i.e., listening to music) compared to active (e.g., producing music or have an active role during therapy) (*Tang & Vezeau, 2010*). This study is mainly focused on the effects of passive interventions as active music interventions (e.g., playing an instrument) are not able to distinguish the cause of the observed changes in motor behavior, which can

either be due to the music heard or other behaviors associated with the therapy received. Music listening does not include sound-based interventions which are merely based on rhythmical sound sequences or cues, also known as Rhythmic Auditory Stimulation (RAS) or auditory cueing. This treatment does not elicit similar changes in performance and secondary beneficial effects on mood and arousal (*Styns et al., 2007; Thaut, Rathbun & Miller, 1997; Wittwer, Webster & Hill, 2013*).

The aim of this study is to investigate if music listening interventions (MLI) are an effective tool to decrease muscle tension in neurologically impaired patients suffering from hypertonia. As listening to music is able to induce general relaxation in several patient populations, we hypothesize that relaxation on muscular level will be present.

SURVEY METHODOLOGY

Protocol and registration

This review was conducted according to the Preferred Reporting Items for Systematic Review and Meta-Analysis Statement (PRISMA). The checklist can be found as [Supplementary Material 1](#) (*Moher et al., 2009*). The study was registered in the PROSPERO database (CRD: no. 42019128511).

Eligibility criteria

Studies were included if they met the following criteria:

- (1) The study population included patients suffering from a neurological disorder which could result in hypertonia e.g., stroke, cerebral palsy, Parkinson's disease, spinal cord injury, multiple sclerosis, etc.
- (2) Interventions had to include MLI
- (3) Muscle tone or activity had to be assessed after intervention. Both clinical and biomechanical analysis of muscle performance were included, e.g., modified Ashworth scale, motor assessment scale, electromyography (EMG), etc.
- (4) Studies had to be written in English, Dutch, German, French or Spanish.
- (5) Study design was a (randomized) clinical trial.

Studies were excluded using the following criteria:

- (1) Music listening involving other sound-based interventions, e.g., RAS or auditory cueing;
- (2) Studies pertaining to active music interventions (i.e., singing, playing rhythms on musical instruments).

Information sources

A systematic search strategy was conducted using the electronic databases of PubMed, Web of Science, Cochrane Library, Science Direct, Scopus, ResearchGate and the Physiotherapy Evidence Database (PEDro). A combination of free text words (spastic, spasticity, muscle tone, hypertonicity, hypertonus, hypertonic, hypertonia and muscle activity) and Medical Subject Headings were used (music, muscle spasticity, muscle hypertonia and electromyography) after careful consideration from all authors. Discrepancies were discussed by all authors and resolved when a majority agreed. The details of the final search strategy, performed in March 2019, can be found as [Supplementary Material 2](#).

Study screening

The screening procedure was performed by three independent researchers (Van Criekinge, O'Brien, and D'Août). To collect potentially relevant studies, eligibility was screened based on title and abstract. Full texts were retrieved and evaluated based on the a-priori provided inclusion and exclusion criteria. Afterwards full texts were gathered and evaluated on the previously set inclusion criteria. Reference lists were manually screened to identify additional relevant studies. Discrepancies were discussed with a third independent person (either O'Brien, D'Août, depending on the allocated studies).

Assessment of quality

The risk of bias was assessed by two independent reviewers (Van Criekinge and Coutinho) by using the PEDro scale for randomized controlled trials (Maher *et al.*, 2003). In case of uncertainty at any point during the scoring process, consensus was sought by a third reviewer (O'Brien). The PEDro scale assesses eleven items such as random allocation of the subjects, concealed allocation and blinding of therapists and assessors (Maher *et al.*, 2003). The total PEDro score can be divided into three sections; high quality = PEDro score 6–10, fair quality = PEDro score 4–5 and poor quality = PEDro score ≤ 3 .

Data extraction and analysis

Extracted data consisted of subject characteristics (age, gender, pathology), outcome measures, type of music or apparatus, intervention and results (see Table 1). Results were mostly described as a difference between groups based on intervention.

To facilitate the quantitative interpretation of the randomized controlled trials, forest plots were created using RevMan 5.3 (Review Manager (RevMan), 2014. Version 5.3. Copenhagen: The Nordic Cochrane Centre). Since a great amount of heterogeneity was present in the outcome measures and assessed muscles resulting in difficulties presenting each measure and muscle separately, a generalized outcome measure “muscle performance” was created. Multiple outcome measures concerning muscle tone within one study were collectively estimated as one outcome measure to ensure an accurate and general image of muscle performance. Either the mean of different muscles or the mean of several muscle tone outcome measures were used to calculate muscle performance. Combining outcome measures should only be allowed when similar responsiveness has been reported (Puhan *et al.*, 2006), which is the case in this study since the combined outcome measurement were assessed with the same assessment tool. The recalculated mean differences and standard deviations were inserted in the RevMan 5.3 template. When the necessary data was not available, authors were contacted to complete the data form. If authors did not respond, missing data were manually calculated using the RevMan 5.3 calculator, if possible and if necessary. To calculate pooled effect sizes, inverse variance was used as statistical method, a random effects model was used as analysis model and standardized mean differences (SMD) were calculated as the effect measure. Heterogeneity between the studies was assessed using I^2 statistics (Higgins *et al.*, 2019; Higgins & Thompson, 2002). Cochrane guidelines were used to interpret the heterogeneity: 0–40% (might not be important), 30–60% (may represent moderate heterogeneity), 50–90% (may represent substantial

Table 1 Data-extraction of the included studies.

Author (Year)	Participants		Music	Outcome measure	Intervention		Results	
	Condition (n, m/w)	Age (years, SD)			Experimental	Control	Experimental	Control
<i>Ben-Pazi et al. (2018)</i>	CP (n = 18, 13/5)	7.5 (4.1)	4–5 audio tracks: music/nature sounds-headphones	CCHQ (/7)	Standardized sound freq/amp: gamma range (n = 9)	Only sound/music listening (n = 9)	Pre: 3.6 Post: 2.1	Pre: 3.26 Post: 3.23
					Receptive listening to relaxing music prior to conventional therapy (n = 10)	30 min, 4x/week, 4 weeks Only conventional therapy: inhibitory/facilitatory techniques, weight bearing, stretch, rhythmic rotation (n = 9)	Shoulder abduction (p = 0.22) Pre: 1 (#2), 1+ (#7), 2 (#1) Post: 1 (#6), 1+ (#2), 2 (#2) Elbow flexors (p = 0.02) Pre: 1 (#4), 1+ (#2), 2 (#3), 3 (#1) Post: 0 (#1), 1 (#5), 1+ (#3), 2 (#1) Forearm pronators (p = 0.018) Pre: 1 (#2), 1+ (#3), 2 (#5) Post: 0 (#1), 1 (#6), 1+ (#3) Hip adductors (p = 0.04) Pre: 0 (#1), 1 (#3), 1+ (#6) Post: 0 (#2), 1 (#7), 1+ (#1) Knee extensors (p = 0.04) Pre: 0 (#1), 1 (#7), 1+ (#1), 3 (#1) Post: 0 (#4), 1 (#5), 2 (#1) Ankle plantar flexors (p = 0.01) Pre: 1 (#1), 2 (#1), 3 (#6), 4 (#1) Post: 1 (#2), 2 (#5), 3 (#2), 4 (#1)	Shoulder abduction (p = 0.02) Pre: 1 (#2), 1+ (#3), 2 (#4) Post: 1 (#5), 1+ (#) Elbow flexors (p = 0.02) Pre: 1 (#3), 1+ (#4), 2 (#2) Post: 0 (#1), 1 (#7), 2 (#1) Forearm pronators (p = 0.04) Pre: 1 (#3), 1+ (#1), 2 (#4), 3 (#1) Post: 0 (#1), 1 (#4), 1+ (#3), 2 (#1) Hip adductors (p = 0.05) Pre: 0 (#1), 1 (#3), 1+ (#1), 2 (#3), 3 (#1) Post: 0 (#2), 1 (#3), 1+ (#3), 1 (#1) Knee extensors (p = 0.01) Pre: 0 (#1), 1 (#3), 1+ (#3), 2 (#2) Post: 0 (#5), 1 (#2), 1+ (#2) Ankle plantar flexors (p = 0.10) Pre: 2 (#2), 3 (#3), 4 (#4) Post: 1+ (#1), 2 (#2), 3 (#3), 4 (#3)
<i>Jeba & Joshi (2016)</i>	Stroke (n = 19)	53.30 (9.18) 52.11 (9.98)	Eastern classical instrumental piece—headphones	MAS (/5)	30 min music + 30 to 45 min conventional therapy, 1x/day, 3 days		Shoulder abduction: p = 0.87 Elbow flexors: p = 0.39 Forearm pronators: p = 0.56 Hip adductors: p = 0.32 Knee extensors: p = 0.74 Ankle plantar flexors: p = 0.13	
<i>Kvam (1997)</i>	CP (n = 12, 9/3)	40.5	Music chair	NWMT	Vibroacoustic therapy (n = 6)	Only music (n = 6)	No significant differences based on treatment (no statistical results) None was regarded worse after treatment. No tendency was observed for different muscle groups.	

(continued on next page)

Table 1 (continued)

Author (Year)	Participants		Music	Outcome measure	Intervention		Results	
	Condition (n, m/w)	Age (years, SD)			Experimental	Control	Experimental	Control
					Exp 1: Music (n = 30) Exp 2: Messages (n = 26)	Silence (n = 20)	Music Relaxation: 4:25 Tension: 12:75 No alteration: 0:0 Messages Relaxation: 3:23 Tension: 10:76.9 No alteration: 0:0	Silence Relaxation: 7:70 Tension: 2:20 No alteration: 1:10
<i>Puggina & Da Silva (2015)</i>	Coma, VS or sedated (n = 76,	42.5 (19)	Music between 60 and 70 dB, selected by family members, patient's preferences	ENG (ratio)		2 to 4 min, 2 sessions		<i>p</i> = 0.019 (only data of participants in coma/vegetative state)
<i>Scartelli (1982)</i>	CP (n = 6, 3/3)	23–33	Music 45–60 dB, -stereo cassette unit	EMG (μV)		Sedative music-assisted EMG biofeedback relaxation training (n = 3) 20 min, 3x/week, 5 weeks	Only EMG biofeedback relaxation training (n = 3)	Decrease of 65% in finger extensor muscle activity
								Decrease of 32.5% in finger extensor muscle activity

(continued on next page)

Table 1 (continued)

Author (Year)	Participants		Music	Outcome measure	Intervention		Results	
	Condition (n, m/w)	Age (years, SD)			Experimental	Control	Experimental	Control
					Mozart K.448 (n = 20)	Smooth, soft relaxing music (n = 20)	Left elbow Pre: 1.85 Post: 1.35 Right elbow Pre: 1.65 Post: 1.05 Left knee: Pre: 1.75 Post: 1.30 Right knee: Pre: 1.75 Post: 1.25	Left elbow Pre: 1.87 Post: 1.70 Right elbow Pre: 1.75 Post: 1.17 Left knee: Pre: 2.25 Post: 1.95 Right knee: Pre: 2.15 Post: 1.72
<i>Wong, Mak & Mok (2018)</i>	Stroke, CP and TBI (n = 40, 21/19)	40 (15.9) 37 (16.3)	2 types of relaxing music—portable music player	MAS (/5)	8 min, 3x/week, 8 weeks		Left elbow: $p = 0.161$ Right elbow: $p = 0.260$ Left knee: $p = 0.740$ Right knee: $p = 0.917$	

Notes.

n, number of; m, men; w, woman; SD, standard deviation; CP, cerebral palsy; VS, vegetative state; TBI, traumatic brain injury; dB, decibels; CCHQ, comfort and care hypertonicity questionnaire; MAS, modified Ashworth scale; NWMT, Nic Waals muscle test; ENG, electroneuromyography; EMG, electromyography; μ V, microvolts; x, times; exp, experimental; pre, pre-treatment; post, post-treatment; #, amount of; freq, frequency; amp, amplitude.




Figure 1 PRISMA flow chart of the included studies.

Full-size DOI: 10.7717/peerj.8228/fig-1

heterogeneity), and 75–100% (considerable heterogeneity) (Higgins *et al.*, 2019). Effect sizes were categorized as a standard mean effect size of 0 which represented no change, 0.2 representing a small effect, 0.5 representing a medium effect and 0.8 representing a large effect (Cohen, 1988). Based on the standardized mean differences extracted from the meta-analysis, a Spearman’s correlation analysis was performed with the amount of therapy time. Confidence intervals (CI) were set to 95%.

RESULTS

Study selection

Of the 1995 studies obtained from all databases, 6 studies met all inclusion criteria. The study selection process is depicted in Fig. 1. Concerning the quality assessment, a median score of 6.5 was observed with a maximum of eight and minimum of three (see Table 2). In total, four studies had a high methodological quality, while the other two had a fair and poor quality. Most studies did not meet the criteria of blinding of participants and therapist as this does not seem possible with respect to treatment. Listening to music is quite apparent for both patient and therapist compared to a placebo or no therapy. In addition, only two studies reported variability measures which makes quantitative analysis difficult.

Table 2 Risk of bias assessment with PEDro scale.

Author (Year)	Eligibility	Random allocation	Concealed allocation	Baseline comparability	Blinding subjects	Blinding therapist	Blinding assessors	Adequate follow-up	Intention-to-treat analysis	Between-group statistical comparison	Point measures and measures of variability	Score
<i>Ben-Pazi et al. (2018)</i>	Yes	Yes	No	Yes	Yes	No	Yes	No	Yes	Yes	No	6
<i>Jeba & Joshi (2016)</i>	Yes	Yes	Yes	Yes	No	No	No	Yes	Yes	Yes	Yes	7
<i>Kvam (1997)</i>	No	Yes	No	No	No	No	Yes	Yes	No	Yes	No	4
<i>Puggina & Da Silva (2015)</i>	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	No	Yes	Yes	8
<i>Scartelli (1982)</i>	No	Yes	No	No	No	No	No	Yes	Yes	No	No	3
<i>Wong, Mak & Mok (2018)</i>	Yes	Yes	Yes	Yes	Yes	No	No	No	No	Yes	No	5

Study characteristics

In total, data from 171 patients (50 females, 102 males, 19 unknown), of which 30 were diagnosed with stroke (*Jeba & Joshi, 2016; Wong, Mak & Mok, 2018*), 61 with cerebral palsy (*Ben-Pazi et al., 2018; Kvam, 1997; Scartelli, 1982; Wong, Mak & Mok, 2018*), 4 with traumatic brain injury (*Wong, Mak & Mok, 2018*) and 76 patients in a decreased conscious state (*Puggina & Da Silva, 2015*) were included in this study (Table 1). One hundred and four participants were allocated to the experimental group receiving MLI, while 67 patients were in the control group. The age of the examined subjects ranged between 7 and 53 years.

There was a great variety in outcomes measures used to assess hypertonia. Two studies assessed muscle tone clinically by means of the Modified Ashworth Scale (MAS) (*Jeba & Joshi, 2016; Wong, Mak & Mok, 2018*), while the other studies used the Care and Comfort Hypertonicity Questionnaire (*Ben-Pazi et al., 2018*) or Nic Waals muscle test (*Kvam, 1997*). Hypertonia and muscle activity was biomechanically assessed by either electroneuromyography (ENG) (*Puggina & Da Silva, 2015*) or EMG (*Scartelli, 1982*).

Three studies used MLI as an additional technique during rehabilitation—vibroacoustic (*Kvam, 1997*), biofeedback relaxation training (*Scartelli, 1982*) and conventional neurorehabilitation (*Jeba & Joshi, 2016*). Vibroacoustic therapy was executed with a relaxation chair with built-in loudspeakers, the loudspeakers are able to produce low frequency acoustic vibrations and music simultaneously. Biofeedback relaxation training was executed with a Cyborg EMG biofeedback unit (Model #EMG J33), feedback of the muscular activity was seen on screen and by a clicking sound to keep patients aware of the acceptable level of muscle activity. Conventional neurorehabilitation consisted of inhibitory techniques for hypertonia, facilitatory techniques for antagonistic muscles, weight bearing exercises, stretching and rhythmic rotations. Other studies used music listening in comparison to no music (*Puggina & Da Silva, 2015*) or different music sounds (*Ben-Pazi et al., 2018; Wong, Mak & Mok, 2018*). The amount of interventions ranged from two to 24 sessions, while therapy time ranged from two minutes to 30 min. Four studies (*Ben-Pazi et al., 2018; Kvam, 1997; Scartelli, 1982; Wong, Mak & Mok, 2018*) investigated

long-term effects, ≥ 4 weeks of therapy, while two studies (*Jeba & Joshi, 2016; Puggina & Da Silva, 2015*) examined short-term effects, < 1 week of therapy.

The type of music was either predetermined by the study protocol (*Jeba & Joshi, 2016; Kvam, 1997; Scartelli, 1982; Wong, Mak & Mok, 2018*) or based on the participant's preferences (*Ben-Pazi et al., 2018; Puggina & Da Silva, 2015*). Predetermined music was selected by the researchers and consisted of the following musical pieces:

1. *Scartelli (1982)*: “The Gift” and “Grandfather’s Story” in *The Red Pony* by Aaron Copland (Columbia Y31016); “Aspen” in *Captured Angel* by Dan Fogelberg (Epic PE 33499); “Lullabye” in *Children of Sanchez* by Chuck Mangione (A & M SP 6700); and “Trois Gymnopedies” in *Blood, Sweat, and Tears* by Eric Satie (Columbia CS 9720);
2. *Jeba & Joshi (2016)*: Eastern classical instrumental pieces which were described as relaxing;
3. *Wong, Mak & Mok (2018)* compared to musical pieces: The Mozart K.448 musical piece was selected for the experimental group compared to a relaxing melody which was randomly searched on the internet using keywords “relaxing music, 8 min”. The chosen piece was composed by Michael in 2011 and was played on an electronic piano (<https://www.youtube.com/watch?v=MAjXiyKPBU8>).

Some studies did not mention the music selection process or music pieces used (*Kvam, 1997*). The two studies who allowed music based on the patient's preferences selected the music as follows:

1. *Puggina & Da Silva (2015)*: Family members were asked to choose the patient's (with decreased conscious state) favorite music style from a predetermined list with 98 songs. Eight different musical styles were represented in this list, i.e., Brazilian country music, samba, international music, movie themes, new age, classical, and gospel.
2. *Ben-Pazi et al. (2018)* compared two musical pieces: The control group received 4 to 5 audio tracks including music or nature sounds according to the child's or parent's preferences, while the experimental group also received music or nature sound to the child's or parent's preferences with fixed sound frequencies embedded in the musical piece.

Synthesis of results

Half of the included studies concluded that MLI is effective in influencing muscle tone (*Ben-Pazi et al., 2018; Puggina & Da Silva, 2015; Scartelli, 1982*), while the others found no significant between-group differences between experimental and control therapy (*Jeba & Joshi, 2016; Kvam, 1997; Wong, Mak & Mok, 2018*). Detailed results of the individual studies can be found in [Table 1](#) and as [Supplementary Material 3](#). Although the studies did not find any significant between-group differences, the findings support the use of MLI in reducing hypertonia. First, *Wong, Mak & Mok (2018)* concluded that both Mozart K.448 and general relaxing music were able to reduce the spasticity levels after eight weeks, no differences were found between-groups (two types of music). Second, *Jeba & Joshi (2016)* did not find any significant between-group differences (conventional therapy vs. conventional therapy with music), yet concluded that the clinical impressions suggested supportive effects of MLI on spasticity. They found decreased spasticity levels after only




Figure 2 Effectiveness of music listening on muscle performance: forest plot.

Full-size [DOI: 10.7717/peerj.8228/fig-2](https://doi.org/10.7717/peerj.8228/fig-2)

three sessions within one week in the ankle plantar flexors ($p = 0.01$) and hip adductors ($p = 0.04$) which were not found with conventional therapy alone. *Kvam (1997)* found no significant differences between music plus vibroacoustic therapy compared to music alone, yet both groups significantly improved after eight weeks.

In total, five of the six studies and 133 of the 171 patients were included in the quantitative analysis since *Kvam (1997)* did not provide any data or statistical analysis (only descriptive). The parameter muscle performance was used as a combined outcome measures for different muscles in the study of *Jeba & Joshi (2016)* and *Wong, Mak & Mok (2018)*, and muscle tension assessments in the study of *Puggina & Da Silva (2015)*. The entire population dataset of *Scartelli (1982)* and *Ben-Pazi et al. (2018)* used only one outcome measure or muscle, which allowed calculation of mean (and SD) muscle tone between participants. The analysis shows that there was a large treatment effect of MLI on muscle performance (SMD 0.96, 95% CI [0.29–1.63]) as depicted in [Fig. 2](#). The level of heterogeneity was considered not important ($I^2 = 10\%$). No correlation between the effectiveness of MLI and amount of therapy was found ($r = 0.30$, $p = 0.62$).

DISCUSSION

Synthesis of results

The aim of this study was to investigate the effectiveness of MLI on hypertonia in neurological patients. We reviewed a total of six randomized controlled trials that provided information of 171 neurologically impaired patients after MLI. The overall risk of bias of the included studies was moderate to low, three studies had a high methodological quality, while two had a fair and one a poor quality. Although not all studies reported between-group differences, all reported improvement in muscle tone over time. Although it is difficult to differentiate these results from natural recovery as no study provided follow-up data after the treatment period, it is reasonable to assume that spasticity tends to get worse if left untreated. A general conclusion of the quantitative analysis suggests that the treatment effect after MLI on hypertonia in neurologically impaired patients was large (SMD 0.96, 95% CI [0.29–1.63]), with a comparatively low level of heterogeneity to declare ($I^2 = 10\%$). Yet, no correlation was found with the amount of therapy which was probably due to the low amount of studies included in the analysis.

Pathology-dependent effectiveness

Four different pathologies were included in this review, patients with cerebral palsy, stroke, traumatic brain injury and decreased conscious state. No clear distinction in treatment effect could be made between the different pathologies. The pathophysiology of these disorders are related to upper motor neuron involvement and all of them had signs of pyramidal hypertonia. None of the included participants had extrapyramidal hypertonia or rigidity, which is a specific characteristic of Parkinson's disease. Nevertheless, sound-based interventions are a major part in the rehabilitation of patients with Parkinson's disease. The pathophysiology of these patients is related to degeneration of the substantia nigra, located in the basal ganglia (*Galvan & Wichmann, 2008*). Sound-based interventions such as auditory stimulation or cueing is used as a technique to bypass the involvement of the basal ganglia which enables the control of automatic movements in the premotor cortex, and does therefore not necessary influence muscle tone (*Sarma et al., 2012*). Current treatment options for extrapyramidal hypertonia are mainly pharmaceuticals or mobilization techniques. This led us to believe that maybe not all neurologically impaired patients benefit from MLI for reducing hypertonia and that the type of hypertonia is probably extremely important in obtaining the desired treatment effects. However, these results bring to light a clear gap in the literature—no studies, to our knowledge, have investigated the effect of MLI on extrapyramidal hypertonia or rigidity.

Assessment of spasticity

Assessment of pyramidal hypertonia or spasticity is mainly performed based on clinical outcome measures such as the (Modified) Ashworth scale, (Modified) Tardieu scale or spasm frequency scales (*Biering-Sorensen, Nielsen & Klinge, 2006; Rekind, 2010*). However, since spasticity scales are dependent on the therapist's experience and judgement, they are deemed invalid and unreliable (*Fleuren et al., 2010; Kumar, Pandyan & Sharma, 2006*). Therefore, the authors suggest to stop using these subjective outcome measures and include more promising techniques such as surface EMG (*Fleuren et al., 2010; Kumar, Pandyan & Sharma, 2006*) or considering the impact on functioning and activities of daily living (*Rekind, 2010*). Although there is a definite need for simple clinical outcome tools, accurate and reliable assessment of spasticity is mainly performed by combining biomechanical with electrophysiological techniques (*Biering-Sorensen, Nielsen & Klinge, 2006; Kim et al., 2005; Kumar, Pandyan & Sharma, 2006*). However, in this current review the majority of studies used clinical outcome measures, while only two used electrophysiological assessment tools. ENG analysis was performed in patients with a decreased conscious state and EMG was assessed in patients with cerebral palsy. The limited amount of research in these two population groups and the lack of evidence in others e.g., stroke or traumatic brain injury represents the need for further research to adequately quantify the effectiveness of MLI on spasticity.

Music as an intervention

In this review music was either supportive of another rehabilitation technique (*Ben-Pazi et al., 2018; Jeba & Joshi, 2016; Kvam, 1997; Scartelli, 1982*) or a stand-alone treatment

(Puggina & Da Silva, 2015; Wong, Mak & Mok, 2018). Treatment effects were of greater magnitude for the latter (SMD 0.91; 1.13) compared to executed as supportive treatment (SMD 0.78; 0.14), except for Ben-Pazi et al., who found the greatest effects (SMD 1.86) comparing gamma stimulation to only music suggesting the importance of specific frequency ranges. However, the general tendency of the results is not so surprising, since treatment effects in music as a supportive therapy are probably generated by the primary therapy. Therefore differences in treatment effect between both groups will be smaller as to comparing treatment effects which are solely generated by music. During these type of listening sessions the focus on the music is much higher than during exercises. The main focus of the patient during conventional therapy is accurate execution of the performed task which might result in decreased attention being directed to the music. On the other hand, if the patient is actively listening to the music, they are executing a dual-task. Deficits in divided attention and dual-tasks have been widely reported in neurologically impaired patients (Mathias & Wheaton, 2007; Yang et al., 2007; Yogeve et al., 2005). It might be that, although it is important to train dual-tasks, decreased treatment effects were a result of divided attention in comparison to the single-task where patients only had to focus on the music. Although divided attention between motor task and music generates smaller treatment effects, they are still present. Clinically, this would suggest that MLI can be used either as background music during rehabilitation (dual-task) or during rest (single-task) to induce muscle relaxation.

Characteristics of music

Self-selected music seems to affect physiological parameters such as muscle activity, heart and respiration rate in a different way than not self-selected music (Blood & Zatorre, 2001). Different types of music, such as jazz, rock, country or blues, affect specific brain regions and result in various effects in EEG waves (Sun et al., 2013). These music preferences are generally discriminated in the frontal cortex of the brain, which seems highly dependent on frequency (gamma) bands (Pan et al., 2013; Sammler et al., 2007). As spasticity is a result of an upper motor neuron lesion, located in the cortex or brain stem, inducing changes in brain activity could therefore result in differences in muscle tone. In this review, there were two studies using patient's preferred music, in children with cerebral palsy and adults with decreased conscious state (Ben-Pazi et al., 2018; Puggina & Da Silva, 2015). These studies reported the largest treatment effects of all the included studies. One study compared two different forms of relaxation music (fixed protocol frequency embedded in preferred music compared only preferred music) in children with CP and found a significant decrease on the Care and Comfort Hypertonicity Questionnaire (SMD 1.86, 95% CI [0.71–3.01]) (Ben-Pazi et al., 2018). Listening to music in patients with a decreased conscious state resulted in greater muscle tension ($p = 0.019$) and non-significant improvements in muscle performance (SMD 1.13, 95% CI [−0.81 to 3.06]) (Puggina & Da Silva, 2015). It seems that music preferences are of major importance when selecting songs when aiming to influence muscle tone. Studies have shown that music preferences, inherent to musical genre, are dependent on three music attributes (Fricke et al., 2018; Greenberg et al., 2016): (1) arousal: tense and strong; (2) valence: amusing but not depressing; and (3)

depth: complex, deep and intelligent. These music dimensions are, in its turn, related to personality traits, social connotation, race, cognitive abilities and self-views ([Marshall & Naumann, 2018](#); [Rentfrow, Goldberg & Levitin, 2011](#); [Rentfrow & Gosling, 2003](#))

Music listening in neurologically impaired patients

The majority of these studies investigating brain activity during music listening are performed on healthy subjects. Normal music listening involves several cortical areas extending well beyond the primary auditory cortex, such as the corpus callosum, superior-temporal plane and middle/inferior temporal gyri ([Stewart et al., 2006](#)). Yet, neurologically impaired patients have acquired lesions in the brain which could influence music listening. Damage of these areas can lead to deficits in the processing of pitch, timbre, timing and emotional response ([Stewart et al., 2006](#)). Since the majority of these areas is located in the right hemisphere, patients with left hemiplegia are more predisposed for musical deficits. Moreover, it seems possible to use pathology-specific music, by considering favorable frequencies of several brain structures ([Miller et al., 2017](#)). For example, thalamic lesion would benefit most from musical frequencies at 2,300, 2,349 and 2,400 Hz in note D and in the 7th octave ([Miller et al., 2017](#)). This bio-guided model of brain music was also adopted by [Miller \(2011\)](#), who found that altering several music parameters can help in specifically targeting stress, ADHD, aging and pain ([Miller, 2011](#)). Therefore it might be interesting not to only incorporate a patient's preferences when selecting music, targeting the lesion locations may be of additional value during recovery. However, more research regarding this bio-guided model for MLI is necessary.

Directions for future research and clinical practice

Although results of this study suggest that MLI is able to induce muscle relaxation, the limited number of studies included prevents the implementation in clinical practice. The authors suggest that new studies with larger populations are needed to verify these findings. In addition, future studies should include objective measurement tools when assessing hypertonia as a primary outcome, while also examining the musical characteristics and functional tasks leading to the best outcomes.

Limitations

There are a few limitations of this review that should be acknowledged. First, during the systematic literature search, only studies written in Dutch, English, German, French or Spanish were included. It is therefore possible relevant studies and important information was missed during the search process. Second, although the search strategy was elaborate, some caution for these proposed recommendations is required since conclusions were based on the results of only five studies (included in meta-analysis). Third, due to lack of standard deviations or statistical data provided in the articles or after contacting authors, we decided to create the outcome measures "muscle performance" which was the mean muscle tone of several individual muscles or various tension outcome measures. Although combining outcome measures with similar responsiveness has been described to be an effective method, including the raw data would be more statistically accurate. However, by

adopting this new outcome measure, quantitative analysis could be performed which was otherwise not possible.

CONCLUSIONS

Quantitative analysis of the results in this review suggest that MLI are able to increased muscle relaxation in neurologically impaired patients, although one study reported increased muscle activity. MLI can be used as either background music during rehabilitation (dual-task) or during rest (single-task). In addition, musical preferences seem to play a major role in the observed treatment effect. We therefore advice using patient's preferred music when selecting songs. However, several gaps were found in the literature which necessitates further research. First of all, effectiveness of music listening was only examined in pyramidal hypertonia (spasticity) and no research was found on extrapyramidal hypertonia (rigidity). Second, a great amount of variety was present in the use of spasticity assessment tools. Only a limited amount of research has been performed with adequately quantifiable spasticity measurements in neurological patients. In conclusion, music listening and processing requires several cortical brain areas which might be affected after diagnosis. It might therefore be of interest to further explore a bio-guided model in MLI for these patients.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The authors received no funding for this work.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Tamaya Van Criekinge conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Kristiaan D'Août, Jonathon O'Brien and Eduardo Coutinho conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the paper, approved the final draft.

Data Availability

The following information was supplied regarding data availability:

Most of the raw data is available in [Table 1](#). The remaining data is available as [Supplemental File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8228#supplemental-information>.

REFERENCES

- Amatya B, Khan F, La Mantia L, Demetrios M, Wade DT. 2013.** Non pharmacological interventions for spasticity in multiple sclerosis. *Cochrane Database of Systematic Reviews* 2013(2):CD009974 DOI 10.1002/14651858.CD009974.pub2.
- Ben-Pazi H, Aran A, Pandyan A, Gelkop N, Ginsberg G, Pollak Y, Elnatan D. 2018.** Auditory stimulation improves motor function and caretaker burden in children with cerebral palsy—a randomized double blind study. *PLOS ONE* 13(12): e0208792 DOI 10.1371/journal.pone.0208792.
- Biering-Sorensen F, Nielsen JB, Klinge K. 2006.** Spasticity-assessment: a review. *Spinal Cord* 44(12):708–722 DOI 10.1038/sj.sc.3101928.
- Blood AJ, Zatorre RJ. 2001.** Intensely pleasurable responses to music correlate with activity in brain regions implicated in reward and emotion. *Proceedings of the National Academy of Sciences of the United States of America* 98(20):11818–11823 DOI 10.1073/pnas.191355898.
- Cohen J. 1988.** *Statistical power analysis for the behavioral sciences*. Second Edition. Hillsdale: Erlbaum Associates.
- Cooper L, Foster I. 2008.** The use of music to aid patients' relaxation in a radiotherapy waiting room. *Radiography* 14(3):184–188 DOI 10.1016/j.radi.2007.02.001.
- Diamond A, Jankovic J. 2006.** Treatment of advanced Parkinson's disease. *Expert Review of Neurotherapeutics* 6(8):1181–1197 DOI 10.1586/14737175.6.8.1181.
- Fleuren JF, Voerman GE, Erren-Wolters CV, Snoek GJ, Rietman JS, Hermens HJ, Nene AV. 2010.** Stop using the Ashworth Scale for the assessment of spasticity. *Journal of Neurology, Neurosurgery and Psychiatry* 81(1):46–52 DOI 10.1136/jnnp.2009.177071.
- Fricke K, Greenberg D, Rentfrow P, Herzberg P. 2018.** Computer-based music feature analysis mirrors human perception and can be used to measure individual music preference. *Journal of Research in Personality* 75:94–102 DOI 10.1016/j.jrp.2018.06.004.
- Galvan A, Wichmann T. 2008.** Pathophysiology of parkinsonism. *Clinical Neurophysiology* 119(7):1459–1474 DOI 10.1016/j.clinph.2008.03.017.
- Gracies JM. 2005a.** Pathophysiology of spastic paresis. I: paresis and soft tissue changes. *Muscle and Nerve* 31(5):535–551 DOI 10.1002/mus.20284.
- Gracies JM. 2005b.** Pathophysiology of spastic paresis. II: emergence of muscle overactivity. *Muscle and Nerve* 31(5):552–571 DOI 10.1002/mus.20285.
- Greenberg D, Kosinski M, Stillwell D, Monteiro B, Levitin D, Rentfrow P. 2016.** The song is you: preferences for musical attribute dimensions reflect personality. *Social Psychological and Personality Science* 7(6):597–605 DOI 10.1177/1948550616641473.
- Guttman M, Kish SJ, Furukawa Y. 2003.** Current concepts in the diagnosis and management of Parkinson's disease. *Canadian Medical Association Journal* 168(3):293–301.
- Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (eds.) 2019.** *Cochrane handbook for systematic reviews of interventions: the cochrane collaboration*. 2nd Edition. Chichester: John Wiley & Sons.

- Higgins JP, Thompson SG. 2002.** Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* **21**(11):1539–1558 DOI [10.1002/sim.1186](https://doi.org/10.1002/sim.1186).
- Jeba C, Joshi D. 2016.** Influence of music therapy on spasticity, functional independence and quality of life in subjects with hemiplegia: a randomized controlled trial. *International Journal of Neurorehabilitation* **3**:219 DOI [10.4172/2376-0281.1000219](https://doi.org/10.4172/2376-0281.1000219).
- Katz RT. 1988.** Management of spasticity. *American Journal of Physical Medicine & Rehabilitation* **67**(3):108–116 DOI [10.1097/00002060-198806000-00004](https://doi.org/10.1097/00002060-198806000-00004).
- Khan F, Amatya B, Bensmail D, Yelnik A. 2019.** Non-pharmacological interventions for spasticity in adults: an overview of systematic reviews. *Annals of Physical and Rehabilitation Medicine* **62**(4):265–273 DOI [10.1016/j.rehab.2017.10.001](https://doi.org/10.1016/j.rehab.2017.10.001).
- Kim DY, Park CI, Chon JS, Ohn SH, Park TH, Bang IK. 2005.** Biomechanical assessment with electromyography of post-stroke ankle plantar flexor spasticity. *Yonsei Medical Journal* **46**(4):546–554 DOI [10.3349/ymj.2005.46.4.546](https://doi.org/10.3349/ymj.2005.46.4.546).
- Krout RE. 2001.** The effects of single-session music therapy interventions on the observed and self-reported levels of pain control, physical comfort, and relaxation of hospice patients. *American Journal of Hospice and Palliative Care* **18**(6):383–390 DOI [10.1177/104990910101800607](https://doi.org/10.1177/104990910101800607).
- Kumar RT, Pandyan AD, Sharma AK. 2006.** Biomechanical measurement of post-stroke spasticity. *Age and Ageing* **35**(4):371–375 DOI [10.1093/ageing/afj084](https://doi.org/10.1093/ageing/afj084).
- Kvam MH. 1997.** The effect of vibroacoustic therapy. *Physiotherapy* **83**(6):290–295 DOI [10.1016/S0031-9406\(05\)66176-7](https://doi.org/10.1016/S0031-9406(05)66176-7).
- Lance JW. 1980.** The control of muscle tone, reflexes, and movement: Robert Wartenberg Lecture. *Neurology* **30**(12):1303–1313 DOI [10.1212/WNL.30.12.1303](https://doi.org/10.1212/WNL.30.12.1303).
- Largo-Wight E, O’Hara BK, Chen WW. 2016.** The efficacy of a brief nature sound intervention on muscle tension, pulse rate, and self-reported stress: nature contact micro-break in an office or waiting room. *Health Environments Research and Design Journal* **10**(1):45–51 DOI [10.1177/1937586715619741](https://doi.org/10.1177/1937586715619741).
- Madson AT, Silverman MJ. 2010.** The effect of music therapy on relaxation, anxiety, pain perception, and nausea in adult solid organ transplant patients. *Journal of Music Therapy* **47**(3):220–232 DOI [10.1093/jmt/47.3.220](https://doi.org/10.1093/jmt/47.3.220).
- Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. 2003.** Reliability of the PEDro scale for rating quality of randomized controlled trials. *Physical Therapy* **83**(8):713–721.
- Marshall S, Naumann L. 2018.** What’s your favorite music? Music preferences cue racial identity. *Journal of Research in Personality* **76**:74–91 DOI [10.1016/j.jrp.2018.07.008](https://doi.org/10.1016/j.jrp.2018.07.008).
- Mathias JL, Wheaton P. 2007.** Changes in attention and information-processing speed following severe traumatic brain injury: a meta-analytic review. *Neuropsychology* **21**(2):212–223 DOI [10.1037/0894-4105.21.2.212](https://doi.org/10.1037/0894-4105.21.2.212).
- Miller EB. 2011.** *Bio-guided music therapy: a practitioner’s guide to the clinical integration of music and biofeedback*. London: Jessica Kingsley.
- Miller E, Miller L, Turner RP, Evans JR. 2017.** Chapter 6—the use of music for neuromodulation. In: Evans JR, Turner RP, eds. *Rhythmic stimulation procedures in neuromodulation*. London: Academic Press, 159–192.

- Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. 2009.** Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLOS Medicine* 6(7):e1000097 DOI 10.1371/journal.pmed.1000097.
- Nair KP, Marsden J. 2014.** The management of spasticity in adults. *BMJ* 349:g4737 DOI 10.1136/bmj.g4737.
- Pan YZ, Guan CT, Yu JH, Ang KK, Chan TE. 2013.** Common frequency pattern for music preference identification using frontal EEG. In: *2013 6th international IEEE/Embs conference on neural engineering (Ner)*. 505–508.
- Puggina AC, Da Silva MJ. 2015.** Patients with disorders of consciousness: vital, facial and muscular responses to music or messages. *Revista Brasileira de Enfermagem* 68(1):94–102 DOI 10.1590/0034-7167.2015680114.
- Puhan MA, Soesilo I, Guyatt GH, Schunemann HJ. 2006.** Combining scores from different patient reported outcome measures in meta-analyses: when is it justified? *Health and Quality of Life Outcomes* 4:94 DOI 10.1186/1477-7525-4-94.
- Rekand T. 2010.** Clinical assessment and management of spasticity: a review. *Acta Neurologica Scandinavica. Supplementum* (190):62–66 DOI 10.1111/j.1600-0404.2010.01378.
- Rentfrow PJ, Goldberg LR, Levitin DJ. 2011.** The structure of musical preferences: a five-factor model. *Journal of Personality and Social Psychology* 100(6):1139–1157 DOI 10.1037/a0022406.
- Rentfrow PJ, Gosling SD. 2003.** The do re mi's of everyday life: the structure and personality correlates of music preferences. *Journal of Personality and Social Psychology* 84(6):1236–1256 DOI 10.1037/0022-3514.84.6.1236.
- Review Manager (RevMan) [Computer program]. 2014.** Version 5.3. Copenhagen: The Nordic Cochrane Centre, T.C.C.
- Sammler D, Grigutsch M, Fritz T, Koelsch S. 2007.** Music and emotion: electrophysiological correlates of the processing of pleasant and unpleasant music. *Psychophysiology* 44(2):293–304 DOI 10.1111/j.1469-8986.2007.00497.x.
- Sand-Jecklin K, Emerson H. 2010.** The impact of a live therapeutic music intervention on patients' experience of pain, anxiety, and muscle tension. *Holistic Nursing Practice* 24(1):7–15 DOI 10.1097/HNP.0b013e3181c8e435.
- Sarkamo T. 2018.** Music for the ageing brain: cognitive, emotional, social, and neural benefits of musical leisure activities in stroke and dementia. *Dementia* 17(6):670–685 DOI 10.1177/1471301217729237.
- Sarma SV, Cheng ML, Eden U, Williams Z, Brown EN, Eskandar E. 2012.** The effects of cues on neurons in the basal ganglia in Parkinson's disease. *Frontiers in Integrative Neuroscience* 6(40) DOI 10.3389/fnint.2012.00040.
- Scartelli J. 1982.** The effect of sedative music on electro-myographic biofeedback assisted relaxation training of spastic cerebral-palsied adults. *Journal of Music Therapy* 19(4):210–218 DOI 10.1093/jmt/19.4.210.
- Sihvonen AJ, Sarkamo T, Leo V, Tervaniemi M, Altenmuller E, Soynila S. 2017.** Music-based interventions in neurological rehabilitation. *Lancet Neurology* 16(8):648–660 DOI 10.1016/S1474-4422(17)30168-0.

- Singer HS, Mink JW, Gilbert DL, Jankovic J. 2010.** 17—Cerebral palsy. In: Singer HS, Mink JW, Gilbert DL, Jankovic J, eds. *Movement disorders in childhood*. Philadelphia: W.B. Saunders, 219–230.
- Staum MJ, Brotons M. 2000.** The effect of music amplitude on the relaxation response. *Journal of Music Therapy* **37**(1):22–39 DOI [10.1093/jmt/37.1.22](https://doi.org/10.1093/jmt/37.1.22).
- Stewart L, Von Kriegstein K, Warren JD, Griffiths TD. 2006.** Music and the brain: disorders of musical listening. *Brain* **129**(Pt 10):2533–2553 DOI [10.1093/brain/awl171](https://doi.org/10.1093/brain/awl171).
- Stratton VN, Zalanowski AH. 1984.** The relationship between music, degree of liking, and self-reported relaxation. *Journal of Music Therapy* **21**(4):184–192 DOI [10.1093/jmt/21.4.184](https://doi.org/10.1093/jmt/21.4.184).
- Styns F, Van Noorden L, Moelants D, Leman M. 2007.** Walking on music. *Human Movement Science* **26**(5):769–785 DOI [10.1016/j.humov.2007.07.007](https://doi.org/10.1016/j.humov.2007.07.007).
- Sun CB, Bao YM, Xu JT, Kong DQ, Zhou H, Wang Q, Shang H, Wang W, Jin M, Wang X, Duan Y. 2013.** The effects of different types of music on electroencephalogram. In: *2013 IEEE international conference on bioinformatics and biomedicine (Bibm)*.
- Tan X, Yowler CJ, Super DM, Fratianne RB. 2010.** The efficacy of music therapy protocols for decreasing pain, anxiety, and muscle tension levels during burn dressing changes: a prospective randomized crossover trial. *Journal of Burn Care and Research* **31**(4):590–597 DOI [10.1097/BCR.0b013e3181e4d71](https://doi.org/10.1097/BCR.0b013e3181e4d71).
- Tang HY, Vezeau T. 2010.** The use of music intervention in healthcare research: a narrative review of the literature. *Journal of Nursing Research* **18**(3):174–190 DOI [10.1097/JNR.0b013e3181efeb1b1](https://doi.org/10.1097/JNR.0b013e3181efeb1b1).
- Thaut MH. 1989.** The influence of music-therapy interventions on self-rated changes in relaxation, affect, and thought in psychiatric prisoner-patients. *Journal of Music Therapy* **26**(3):155–166 DOI [10.1093/jmt/26.3.155](https://doi.org/10.1093/jmt/26.3.155).
- Thaut MH, Rathbun JA, Miller RA. 1997.** Music versus metronome timekeeper in a rhythmic motor task. *International Journal of Arts Medicine* **5**(1):4–12.
- Thompson AJ, Jarrett L, Lockley L, Marsden J, Stevenson VL. 2005.** Clinical management of spasticity. *Journal of Neurology, Neurosurgery and Psychiatry* **76**(4):459–463 DOI [10.1136/jnnp.2004.035972](https://doi.org/10.1136/jnnp.2004.035972).
- Wittwer JE, Webster KE, Hill K. 2013.** Music and metronome cues produce different effects on gait spatiotemporal measures but not gait variability in healthy older adults. *Gait Posture* **37**(2):219–222 DOI [10.1016/j.gaitpost.2012.07.006](https://doi.org/10.1016/j.gaitpost.2012.07.006).
- Wong A, Mak L, Mok V. 2018.** The effect of listening to Mozart K448 music for individuals with spasticity—RCT. *International Journal of Neurorehabilitation* **5**:329 DOI [10.4172/2376-0281.1000329](https://doi.org/10.4172/2376-0281.1000329).
- Yang YR, Chen YC, Lee CS, Cheng SJ, Wang RY. 2007.** Dual-task-related gait changes in individuals with stroke. *Gait Posture* **25**(2):185–190 DOI [10.1016/j.gaitpost.2006.03.007](https://doi.org/10.1016/j.gaitpost.2006.03.007).
- Yogev G, Giladi N, Peretz C, Springer S, Simon ES, Hausdorff JM. 2005.** Dual tasking, gait rhythmicity, and Parkinson’s disease: which aspects of gait are attention demanding? *European Journal of Neuroscience* **22**(5):1248–1256 DOI [10.1111/j.1460-9568.2005.04298](https://doi.org/10.1111/j.1460-9568.2005.04298).