

<https://doi.org/10.31896/k.23.4>

Original scientific paper

Accepted 6. 11. 2019.

RUŽICA KOLAR-ŠUPER

Bouvaist Cubic of a Triangle in an Isotropic Plane

Bouvaist Cubic of a Triangle in an Isotropic Plane

ABSTRACT

The cubic in an isotropic plane which passes through the intersections of the sides of an orthic triangle with the sides of a complementary triangle of a given triangle, and through the point which is complementary to the Steiner point of triangle is studied in this paper. It is proved that its non-isotropic asymptote is parallel to Lemoine line of a given triangle.

Key words: isotropic plane, Bouvaist cubic, point complementary to the Steiner point

MSC2010: 51N25

Bouvaistova kubika trokuta u izotropnoj ravnini

SAŽETAK

U članku se proučava kubika koja prolazi kroz sjecišta stranica ortotrokuta i komplementarnog trokuta danog trokuta i kroz točku komplementarnu Steinerovoj točki tog trokuta. Dokazuje se da je neizotropna asimptota kubike paralelna s Lemoineovim pravcem danog trokuta.

Ključne riječi: izotropna ravnina, Bouvaistova kubika, komplementarna točka Steinerovoj točki

In [1], Bouvaist showed the existence of a cubic in Euclidean geometry, which passes through all nine intersections of the sides of an orthic triangle and a complementary triangle of a given triangle and through a point complementary to the Steiner point of that triangle. He proved that this cubic is circular and its real asymptote is parallel to the Lemoine line of a given triangle.

It will be shown in this paper that some analogous statement holds in the isotropic plane as well.

The isotropic (or Galilean) plane is a projective-metric plane, where the absolute consists of one line, i.e., the absolute line ω , and one point on that line, i.e., the absolute point Ω . The lines through the point Ω are isotropic lines, and the points on the line ω are isotropic points (the points at infinity). Two points $P_1 = (x_1, y_1)$, $P_2 = (x_2, y_2)$ with $x_1 = x_2$ are said to be *parallel*, and we shall say they are on the same *isotropic* line. Any isotropic line is perpendicular to any non-isotropic line.

A triangle is said to be *allowable* if none of its sides is isotropic. Each allowable triangle ABC can be set by a suitable choice of the coordinate system in the *standard position*, in which its circumscribed circle has the equation $y = x^2$, and its vertices are the points $A = (a, a^2)$, $B = (b, b^2)$, $C = (c, c^2)$, where $a + b + c = 0$. We shall say then that ABC is a *standard triangle*. To prove geometric facts for each allowable triangle it is sufficient to give a proof for the standard triangle (see [3]).

With the labels

$$p = abc \quad \text{and} \quad q = bc + ca + ab$$

a number of useful equalities are proved in [3], as e.g.

$$\begin{aligned} a^2 &= bc - q, \\ (b - c)^2 &= -(q + 3bc), \\ (c - a)(a - b) &= 2q - 3bc. \end{aligned}$$

In [3], it is proved that the sides $B_h C_h$ and $B_m C_m$ of the orthic triangle $A_h B_h C_h$ and the complementary triangle

$A_m B_m C_m$ of the standard triangle have the equations:

$$y - 2ax + q - 2bc = 0,$$

$$y + ax + q - \frac{bc}{2} = 0,$$

and the equations of their other sides are obtained by a cyclic permutation $a \rightarrow b \rightarrow c \rightarrow a$. That is why every cubic through all nine intersections of the sides of these two triangles has the equation of the form:

$$\prod (y - 2ax + q - 2bc) - \lambda \prod \left(y + ax + q - \frac{bc}{2} \right) = 0, \quad (1)$$

where \prod denotes the product of three factors, the first of which is written, and the other two arise from the first one by cyclic permutations $a \rightarrow b \rightarrow c \rightarrow a$.

In [4], it is shown that the point

$$S = \left(\frac{3p}{2q}, -\frac{9p^2}{2q^2} - q \right)$$

is complementary to the Steiner point of the standard triangle ABC . For that point we obtain

$$\begin{aligned} y - 2ax + q - 2bc &= -\frac{9p^2}{2q^2} - \frac{3ap}{q} - 2bc \\ &= -\frac{bc}{2q^2} (9a^2bc + 6a^2q + 4q^2) \\ &= -\frac{bc}{2q^2} [9bc(bc - q) + 6q(bc - q) + 4q^2] \\ &= \frac{bc}{2q^2} (2q^2 + 3bcq - 9b^2c^2) \\ &= \frac{bc}{2q^2} (q + 3bc)(2q - 3bc) \\ &= -\frac{bc}{2q^2} (b - c)^2 (c - a)(a - b), \end{aligned}$$

$$\begin{aligned} y + ax + q - \frac{bc}{2} &= -\frac{9p^2}{2q^2} + \frac{3ap}{2q} - \frac{bc}{2} \\ &= -\frac{bc}{2q^2} (9a^2bc - 3a^2q + q^2) \\ &= -\frac{bc}{2q^2} [9bc(bc - q) - 3q(bc - q) + q^2] \\ &= -\frac{bc}{2q^2} (4q^2 - 12bcq + 9b^2c^2), \\ &= -\frac{bc}{2q^2} (2q - 3bc)^2 \\ &= -\frac{bc}{2q^2} (c - a)^2 (a - b)^2 \end{aligned}$$

and then

$$\prod (y - 2ax + q - 2bc) = -\frac{a^2b^2c^2}{8q^6} (b - c)^4 (c - a)^4 (a - b)^4,$$

$$\prod \left(y + ax + q - \frac{bc}{2} \right) = -\frac{a^2b^2c^2}{8q^6} (b - c)^4 (c - a)^4 (a - b)^4.$$

Thus, the cubic of the pencil of the cubics with equation (1) passes through the point S if one takes $\lambda = 1$ (Figure 1).

If that cubic of the allowable triangle ABC , which passes through the intersections of the sides of its orthic triangle with the sides of its complementary triangle, and through the point S complementary to the Steiner point of the triangle ABC (Figure 1), is called the *Bouvaist cubic* of that triangle, then we have:

Theorem 1 *The Bouvaist cubic \mathcal{B} of the standard triangle ABC has the equation:*

$$\begin{aligned} &(y - 2ax + q - 2bc)(y - 2bx + q - 2ca) \\ &(y - 2cx + q - 2ab) - \left(y + ax + q - \frac{bc}{2} \right) \\ &\left(y + bx + q - \frac{ca}{2} \right) \left(y + cx + q - \frac{ab}{2} \right) = 0. \end{aligned} \quad (2)$$

Let us now find the intersection points of the cubic (2) and the absolute line. We have to solve the equation

$$(y - 2ax)(y - 2bx)(y - 2cx) - (y + ax)(y + bx)(y + cx) = 0,$$

which can also be written in the following form:

$$-3(a + b + c)xy^2 + 3(bc + ca + ab)x^2y - 9abcx^3 = 0,$$

and finally as $3qx^2y - 9px^3 = 0$. We have the double solution $x = 0$ and the solution $y = \frac{3p}{q}x$, which means that

the cubic has an asymptote with a slope $\frac{3p}{q}$, which is by [2] a slope of the Lemoine line \mathcal{L} of the triangle ABC . We obtained:

Theorem 2 *The non-isotropic asymptote of Bouvaist cubic of an allowable triangle is parallel to the Lemoine line of a given triangle. Absolute point is an intersection point of the Bouvaist cubic and absolute line with intersection multiplicity 2.*




Figure 1: *Bouvaist cubic of a triangle ABC in isotropic plane*

References

- [1] R. BOUVAIST, *Généralization de la question 2811*, *Mathesis* **48** (1934), 451–452.
- [2] Z. KOLAR-BEGOVIĆ, R. KOLAR-ŠUPER, J. BEBAN-BRKIĆ, V. VOLENEC, Symmedians and the symmedian centre of the triangle in an isotropic plane, *Math. Pannon.* **17** (2006), 287–301.
- [3] R. KOLAR-ŠUPER, Z. KOLAR-BEGOVIĆ, V. VOLENEC, J. BEBAN-BRKIĆ, Metrical relationships in a standard triangle in an isotropic plane, *Math. Commun.* **10** (2005), 149–157.
- [4] R. KOLAR-ŠUPER, Z. KOLAR-BEGOVIĆ, V. VOLENEC, Steiner point of a triangle in an isotropic plane, *Rad Hrvat. Akad. Znan. Umjet. Mat. Znan.* **20** 528 (2016), 83–95.

Ružica Kolar-Šuper

orcid.org/0000-0002-8945-2745

e-mail: rkolar@foozos.hr

Faculty of Education, University of Osijek
 Cara Hadrijana 10, 31 000 Osijek, Croatia