

Gratkowski, Maciej; Schmidt, Silvio; Güllmar, Daniel; Witte, Otto W.; Haueisen, Jens:

## Decomposition of somatosensory evoked potentials in rats

---

**Zuerst erschienen in:** Biomedical Engineering = Biomedizinische Technik. - Berlin [u.a.] : de Gruyter. - 57 (2012), Suppl. 1, Track-F, p. 177.

**Erstveröffentlichung:** 2012-08-30

**ISSN (online):** 1862-278X

**ISSN (print):** 0013-5585

**DOI:** [10.1515/bmt-2012-4447](https://doi.org/10.1515/bmt-2012-4447)

**[Zuletzt gesehen:** 2019-08-12]

„Im Rahmen der hochschulweiten Open-Access-Strategie für die Zweitveröffentlichung identifiziert durch die Universitätsbibliothek Ilmenau.“

“Within the academic Open Access Strategy identified for deposition by Ilmenau University Library.”

„Dieser Beitrag ist mit Zustimmung des Rechteinhabers aufgrund einer (DFG-geförderten) Allianz- bzw. Nationallizenz frei zugänglich.“

„This publication is with permission of the rights owner freely accessible due to an Alliance licence and a national licence (funded by the DFG, German Research Foundation) respectively.“



# Decomposition of Somatosensory Evoked Potentials in Rats

M. Gratkowski, Institute of Biomedical Engineering and Informatics, Ilmenau University of Technology, Ilmenau, Germany, maciej.gratkowski@tu-ilmenau.de

S. Schmidt, Department of Neurology, Friedrich-Schiller-University, Jena, Germany

D. Güllmar, Institute of Diagnostic and Interventional Radiology, Friedrich-Schiller-University, Jena, Germany

O.W. Witte, Department of Neurology, Friedrich-Schiller-University, Jena, Germany

J. Haueisen, Institute of Biomedical Engineering and Informatics, Ilmenau University of Technology, Ilmenau, Germany

## Introduction

Somatosensory evoked potentials (SEP) contain several components that are, due to their low amplitude and overlap in time, difficult to distinguish and quantify. In this paper we present a methodology of component extraction and a description of the SEPs recorded from rat.

## Methods

Silicon strips containing 4x4 recording electrodes were positioned on both hemispheres over the forelimb sensomotory areas. Around 500 stimulations were applied both on the left and right N. medianus and the responses on each hemisphere were averaged for each subject. The Topographic Matching Pursuit (TMP) method was used to decompose the averaged SEPs. This method is able to decompose multichannel data into spatio-temporal atoms defined by only a few parameters. Regions of the time-frequency (TF) space with high activation were identified in the averaged TF representation and chosen as possible stable SEP components. Atoms, whose temporal and frequency parameters were located inside these regions, were assigned to the respective components for each subject. This process can be seen as a kind of parametric filtering of data.

## Results

In the present study five distinct components could be identified in the SEP data. The most prominent components had their main frequency of  $51.6 \pm 5.33$  Hz and  $144 \pm 16.8$  Hz by young and  $50.7 \pm 3.99$  Hz and  $152 \pm 17.2$  Hz by aged subjects. Furthermore, additional high frequency components could be found at  $289 \pm 35.5$  Hz,  $456 \pm 21.2$  Hz and  $626 \pm 37.6$  Hz by young subjects and  $346 \pm 21.5$  Hz,  $458 \pm 21.5$  Hz and  $621 \pm 24$  Hz by aged subjects. Moreover, significant differences of the amplitudes, temporal localizations and spatial distributions of the components were found between young and aged subjects.

## Conclusion

The components are described by only a few parameters in the time, frequency and spatial domains, which is convenient and can be used for investigation of group differences of SEP components.