

Strohmeier, Daniel; Kowalski, Matthieu; Haueisen, Jens; Hämäläinen, Matti S.; Gramford, Alexandre:

MEG/EEG single trial source estimates based on inter-trial priors in the time-frequency domain for source connectivity analysis

Zuerst erschienen in: Biomedical Engineering = Biomedizinische Technik. - Berlin [u.a.] : de Gruyter. - 57 (2012), Suppl. 1, Track-F, p. 391.

Erstveröffentlichung: 2012-08-30

ISSN (online): 1862-278X

ISSN (print): 0013-5585

DOI: [10.1515/bmt-2012-4461](https://doi.org/10.1515/bmt-2012-4461)

[Zuletzt gesehen: 2019-08-12]

„Im Rahmen der hochschulweiten Open-Access-Strategie für die Zweitveröffentlichung identifiziert durch die Universitätsbibliothek Ilmenau.“

“Within the academic Open Access Strategy identified for deposition by Ilmenau University Library.”

„Dieser Beitrag ist mit Zustimmung des Rechteinhabers aufgrund einer (DFG-geförderten) Allianz- bzw. Nationallizenz frei zugänglich.“

„This publication is with permission of the rights owner freely accessible due to an Alliance licence and a national licence (funded by the DFG, German Research Foundation) respectively.“



MEG/EEG single trial source estimates based on inter-trial priors in the time-frequency domain for source connectivity analysis

Strohmeier D., Institute of Biomedical Engineering and Informatics, Ilmenau University of Technology, Ilmenau, Germany, daniel.strohmeier@tu-ilmenau.de

Kowalski M., Laboratoire des Signaux et Systèmes (CNRS-SUPELEC-Univ. Paris-Sud)
SUPELEC, Plateau de Moulon, Gif-sur-Yvette cedex, France, matthieu.kowalski@lss.supelec.fr

Haueisen J., Institute of Biomedical Engineering and Informatics, Ilmenau University of Technology, Ilmenau, Germany, jens.haueisen@tu-ilmenau.de

Hämäläinen M., Martinos Center for Biomedical Imaging, Department of Radiology, Massachusetts General Hospital, Boston, MA, msh@nmr.mgh.harvard.edu

Gramfort A., Parietal Project Team, INRIA Saclay Ile-de-France, CEA - NeuroSpin, Gif-sur-Yvette cedex, France, alexandre.gramfort@inria.fr

Introduction

Analysing functional connectivity of neuronal sources based on MEG/EEG source estimates is a non-invasive approach to examine the brain functioning. In order to apply functional connectivity measures such as coherency and phase locking value (PLV) to evoked responses, which are highly non-stationary over time, the source activations and the respective complex-valued time-frequency distributions have to be estimated on a single-trial basis. However, due to the low signal-to-noise ratio of unaveraged data, the applicability of present source reconstruction methods to the study of single-trial MEG/EEG data is limited.

Methods

We apply the multi-trial time-frequency mixed-norm estimate (MT-TF-MxNE) for reconstructing single-trial source estimates, which is based on a composite multidimensional mixed norm prior in the time-frequency domain combining a sparse intra-trial prior and a smooth inter-trial prior. By thresholding the Gabor transform of the source signals, the intra-trial prior promotes spatial sparsity and temporal smoothness of the neuronal activity per trial. The inter-trial prior retains consistency of the respective sparsity structure in the time-frequency domain to impose stationarity over trials. Amplitudes and phases of the active time-frequency coefficients are allowed to vary in order to account for trial-to-trial variations. Thus, the MT-TF-MxNE computes single-trial estimates while regarding multiple trials simultaneously to increase the information content. The resulting sparse regression problem is a convex optimization problem, which is solved using the fast iterative shrinkage-thresholding algorithm. Subsequently, for analysing source connectivity, coherency, coherence and imaginary coherence, and the PLV can be computed using the reconstructed single-trial time-frequency distributions.

Results

The MT-TF-MxNE is applied to simulated MEG/EEG data showing its ability to reconstruct location, activation and time-frequency distribution of evoked neuronal activity for functional source connectivity analysis.

Conclusion

The results show that the MT-TF-MxNE is a promising source reconstruction approach for analysing evoked MEG/EEG data. Applying the MT-TF-MxNE to real MEG/EEG data is part of the future work.