

**HHS PUBLIC ACCESS**

Author manuscript

*Genet Epidemiol.* Author manuscript; available in PMC 2019 September 01.

Published in final edited form as:

*Genet Epidemiol.* 2018 September ; 42(6): 571–586. doi:10.1002/gepi.22137.**Hierarchical modeling of melanocortin 1 receptor variants with skin cancer risk****Amit D. Joshi<sup>1,2,3,\*</sup>, Xin Li<sup>4,\*</sup>, Peter Kraft<sup>1,2</sup>, and Jiali Han<sup>4,5</sup>**<sup>1</sup>Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA 02115<sup>2</sup>Program in Statistical Genetics and Genetic Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA 02115<sup>3</sup>Clinical and Translational Epidemiology Unit, Department of Medicine, Massachusetts General Hospital, Boston, MA 02114<sup>4</sup>Department of Epidemiology, Richard M. Fairbanks School of Public Health, Indiana University, Indianapolis, IN 46202<sup>5</sup>Indiana University Melvin and Bren Simon Cancer Center, Indianapolis, IN 46202**Abstract**

The human *MC1R* gene is highly polymorphic among lightly pigmented populations, and several variants in the *MC1R* gene have been associated with increased risk of both melanoma and non-melanoma skin cancers. The functional consequences of *MC1R* gene variants have been studied *in vitro* and *in vivo* in postulated causal pathways, such as G-protein-coupled signaling transduction, pigmentation, immune response, inflammatory response, cell proliferation, and extracellular matrix adhesion. In a case-control study nested within the Nurses' Health Study, we utilized hierarchical modeling approaches, incorporating the quantitative information from these functional studies, to examine the association between particular *MC1R* alleles and the risk of skin cancers. Different prior matrices were constructed according to the phenotypic associations in controls, cell surface expression, and enzymatic kinetics. Our results showed the parameter variance estimates of each SNP were smaller when using a hierarchical modeling approach compared to standard multivariable regression. Estimates of second-level parameters gave information about the relative importance of *MC1R* effects on different pathways, and odds ratio estimates changed depending on prior models (for example, the change ranged from –21% to 7% for melanoma risk assessment). In addition, the estimates of prior model hyper-parameters in the hierarchical modeling approach allow us to determine the relevance of individual pathways on the risk of each of the skin cancer types. In conclusion, hierarchical modeling provides a useful analytic approach in addition to the widely used conventional models in genetic association studies that can incorporate measures of allelic function.

\*Authors contributed equally to the manuscript. **Corresponding author:** Jiali Han, Ph.D, Department of Epidemiology, Richard M. Fairbanks School of Public Health, Indiana University, Address: 1050 Wishard Boulevard, RG 5112, Indianapolis, IN 46202-2872, Tel: 317-278-0370, [jjalhan@iu.edu](mailto:jjalhan@iu.edu).

Conflict of interest

The authors declare no conflict of interest.

## Keywords

hierarchical modeling; MC1R; functional variants; pigmentation; skin cancer

---

## Introduction

Skin cancer is the most common form of cancer in the United States and accounts for more than 1 million new cases per year, including approximately 87,000 cases of cutaneous malignant melanoma (hereafter referred to simply as melanoma) [2017; Howe, et al. 2001; Siegel, et al. 2017]. There are three major types of skin cancer, with melanoma being the most fatal; the most common types of non-melanoma skin cancer are basal cell carcinoma (BCC) followed by squamous cell carcinoma (SCC). Ultra-violet (UV) radiation exposure is an important risk factor for skin cancer, and dark pigmentation is an important inherited factor protecting against UV-induced skin cancer. Pigmentary melanin is synthesized in melanocytes and secreted into keratinocytes. Brown/black eumelanin absorbs UV and neutralizes free radicals to protect skin from UV damage, whereas yellow/red pheomelanin generates free radicals in response to UV. Human pigmentation is an inherited trait partially determined by the melanocortin 1 receptor (*MC1R*) gene, located at 16q24.3 [Gantz, et al. 1994]. It encodes a 317-amino acid seven-pass-transmembrane G protein-coupled receptor. When activated by  $\alpha$ -melanocyte-stimulating hormone ( $\alpha$ -MSH) or adrenocorticotrophic hormone (ACTH), MC1R activates adenylate cyclase, thereby elevating intracellular cyclic adenosine monophosphate (cAMP). MC1R was shown to be the limiting factor controlling the output of the cAMP signaling pathway in heterologous cells expressing the wild-type *MC1R* gene [Mas, et al. 2003]. This signaling induces the maturation of the phenomelanosome to eumelanosome, resulting in darker pigmentation [Rees 2003; Sturm, et al. 2001].

The human *MC1R* gene is highly polymorphic among lightly pigmented populations [Harding, et al. 2000; Sturm 2002]. Among more than 80 nonsynonymous variants identified to date, Arg151Cys, Arg160Trp, and Asp294His were associated with red-hair phenotype [Beaumont, et al. 2005; Box, et al. 1997; Valverde, et al. 1995] and are known as “red hair color” (RHC) variants; other variants with weaker association are referred to as “non-red hair color” (NRHC) variants. Several variants in the *MC1R* gene have been associated with increased risks for melanoma and non-melanoma skin cancers after pigmentation phenotype was taken into account [Bastiaens, et al. 2001; Box, et al. 2001; Kennedy, et al. 2001; Landi, et al. 2005; Matichard, et al. 2004; Palmer, et al. 2000]. Previous studies grouped RHC alleles and NRHC alleles in tests of genetic associations between the MC1R alleles and risk of skin cancer [Han, et al. 2006; Landi, et al. 2005; Matichard, et al. 2004]. Distinct functions of specific MC1R variants in relation to pigmentation, immune response, inflammatory response, cell proliferation, and extracellular matrix adhesion pathways have been examined using data on pigmentation phenotypes or quantitative *in vitro* assays.

In this study, we utilized this knowledge regarding the role of MC1R alleles in biological processes potentially relevant to skin cancers to improve the accuracy of the estimated effects of MC1R variants on predisposition to skin cancer. Using a nested case-control

design within the Nurses' Health Study, we explored hierarchical modeling analyses, incorporating quantitative information on the function of particular MC1R alleles [Aragaki, et al. 1997; Brenner, et al. 2013; Greenland 2000; Hung, et al. 2004; Witte 1997]. We evaluated the associations of seven common MC1R variants (Val60Leu, Val92Met, Arg151Cys, Ile155Thr, Arg160Trp, Arg163Gln, and Asp294His) with three major types of skin cancer (melanoma, SCC, and BCC) in women of European ancestry. Hierarchical regression approach allows the integration of biological data, reduces the overall estimation error by shrinking the estimated coefficients to their mean, and at the same time protects against over-parameterization of the model.

## Methods

### Study Population

The Nurses' Health Study was established in 1976, when 121,700 female registered nurses between the ages of 30 and 55 completed a self-administered questionnaire on their medical histories and baseline health-related exposures. Updated information has been obtained by questionnaires every 2 years since then. Between 1989 and 1990, blood samples were collected from 32,826 of the cohort members. Detailed methods of this nested case-control study were described previously [Han, et al. 2006]. The distributions of risk factors for skin cancer in the subcohort of those who donated blood samples were very similar to those in the overall cohort. Briefly, eligible cases in this study consisted of women with incident skin cancer from the subcohort who had given a blood specimen, including cases of SCC and BCC with a diagnosis any time after blood collection up to June 1, 1998, and cases of melanoma up to June 1, 2000, with no previously diagnosed skin cancer. A common control series was randomly selected from participants who gave a blood sample and were free of diagnosed skin cancer up to and including the questionnaire cycle in which the case was diagnosed. One or two controls were matched to each case by year of birth ( $\pm 1$  year) and self-reported race (Caucasian/missing or others). Fewer than 5% of cases and controls had missing or other race/ethnicity. The nested case-control study consisted of 219 melanoma cases, 286 SCC cases, 300 BCC cases, and 874 matched controls. We restricted analysis to 197 melanoma cases, 264 SCC cases, 263 BCC cases, and 791 controls with complete genotype information. The study protocol was approved by the Committee on Use of Human Subjects of Brigham and Women's Hospital, Boston, MA.

Information regarding skin cancer risk factors was obtained from the prospective biennial questionnaires and the retrospective supplementary questionnaire. Information on natural hair color and childhood and adolescent tendency to tan was solicited in the 1982 prospective questionnaire. To avoid potential population stratification, we excluded one Asian melanoma case and one control, one Hispanic SCC case and two controls.

### Single Nucleotide Polymorphism (SNP) Identification

The distribution and frequency of MC1R variants in 179 Caucasian controls from the US were determined by Kanetsky et al. using a direct-sequencing method [Kanetsky, et al. 2004]. Seven non-synonymous polymorphisms with allele frequency  $>1\%$  were identified (Val60Leu, Val92Met, Arg151Cys, Ile155Thr, Arg160Trp, Arg163Gln, and Asp294His) in

the coding region. We genotyped these variants in our case-control study. There are comprehensive and quantitative *in vitro* assays on the function of these particular variants, which we incorporated into hierarchical modeling analyses. We did not genotype the three variants (86insA, Asp84Glu, and Arg142His) with allele frequency 1% [Han, et al. 2004].

### Laboratory Assays

The Arg160Trp polymorphism was genotyped by restriction fragment length polymorphism (RFLP), and the other polymorphisms were genotyped by the 5'-nuclease assay (TaqMan®) in 384-well format, using the ABI PRISM 7900HT Sequence Detection System (Applied Biosystems, Foster City, CA). TaqMan® primers and probes were designed using the Primer Express® Oligo Design software v2.0 (ABI PRISM). Laboratory personnel were blinded to case-control status, and blinded quality-control samples were inserted to validate genotyping procedures; concordance for the blinded samples was 100%. Primers, probes, and conditions for genotyping assays are available upon request.

### Statistical Analyses

**Hierarchical modeling outline**—We utilized a two-level hierarchical modeling approach that incorporates functional information in a second-level (prior) model for regression parameters. Specifically, it assumes the log-odds ratio parameters  $\beta$  from the first-level logistic regression of disease risk on MC1R variants are themselves independent random variables with means  $Z\pi$  and variance  $\tau^2$ . The first level of the regression models the log odds of disease on the seven MC1R SNPs,  $X$ , and the vector of covariates,  $W$ .

$$\log\left(\frac{p}{1-p}\right) = \alpha + X\beta + W\gamma$$

The second level is a linear model on which the first-level regression parameter  $\beta$  is regressed on a matrix obtained using functional information,  $Z$ , and this helps improve the estimation of  $\beta$ .

$$\beta = Z\pi + \delta$$

$Z$  is a user-specified prior design matrix, and the dimension of the second-level parameter vector,  $\pi$ , can be much smaller than the dimension of  $\beta$ . For example, if  $\beta$  is a vector of seven SNP log odds ratios,  $Z$  might be a vector of seven ones, indicating that the average effect for each of the modeled SNPs is the same (namely  $\pi$ ). By treating  $\beta$  as a random variable, this approach incorporates the uncertainty in this prior structure (e.g., prediction error in the second-level covariates in  $Z$ ). The residuals in the second-level regression,  $\delta$ , are assumed to have a normal distribution, with a mean 0 and variance  $\tau^2$ .

We coded each variant using a dominant model and fit all variants simultaneously. We used dominant models because the MC1R variants being studied follow a dominant inheritance pattern. Hierarchical analyses can be implemented using the %glimmix SAS macro [Witte, et al. 1998; Witte, et al. 2000], in which the two levels of the hierarchical regression are

combined into a mixed model with fixed coefficients ( $\boldsymbol{\pi}$ , and  $\boldsymbol{\gamma}$ ) and random coefficient ( $\boldsymbol{\delta}$ ). By substituting the value of  $\boldsymbol{\beta}$  from the second-level model into the first-level model, we get the following mixed-model equation:

$$\log\left(\frac{p}{1-p}\right) = \alpha + XZ\pi + X\delta + W\gamma$$

Therefore, in addition to the SNPs  $X$  and the covariates  $W$ , we also included the product of  $X$  and  $Z$  matrices ( $XZ$ ) in the mixed model, and their fixed effect coefficients can be interpreted as  $\boldsymbol{\pi}$ .

In this analysis, we used a Semi-Bayes approach and pre-specified the common variance of the random coefficient vector  $\boldsymbol{\delta}$ , for effect sizes to be consistent with those found in previous studies of genetic variants. Rather than allowing  $\boldsymbol{\tau}$  to be estimated empirically, we fixed the value of  $\boldsymbol{\tau}^2$  at 0.1 [Greenland 1994; Witte, et al. 1994]. At this value of  $\boldsymbol{\tau}$ , we assume with 95% confidence that the variation in the true odds ratios for SNPs, after adjusting for the effects of the attributes considered in the  $Z$  matrix, will fall within a 1.5-fold range, i.e., ( $\exp(0.1 \times 3.92) \approx 1.5$ ).

**Construction of prior design matrix**—The prior design matrix  $Z$  is a critical component of the hierarchical approach. We considered several prior design matrices corresponding to different working hypotheses about the causal role of different *MC1R* variants. First, we simply assumed that all variants were exchangeable, i.e., they have the same average effect. This corresponds to the  $Z$  matrix described in the previous section. We then defined several prior matrices according to the functional parameters generated from phenotypic correlations and *in vitro* assays (Table 1). The functional parameters for model 4–9 in Table 2 were obtained from published literature [Beaumont, et al. 2005; Sanchez-Mas, et al. 2004] and not based on our samples in the NHS.

We performed multiple linear regression analyses among controls to evaluate the associations between the phenotypes and the *MC1R* variants (Table 1, models 2–4). We treated hair color (red, blond, light brown, dark brown or black), skin color (fair, medium, olive), and tanning ability (deep tan, average tan, light tan, practically none) as ordinal outcome variables. The three RHC alleles (Arg151Cys, Arg160Trp, and Asp294His) had a stronger association with skin color and hair color compared with other variant alleles. For tanning ability, the three RHC alleles and 155Thr allele showed strong associations. The variant-specific estimated regression coefficients  $\mathbf{z}_i$  from these analyses were then used in the second-level model, so that the mean log-odds ratio for variant  $i$  was  $\boldsymbol{\pi}_0 + \boldsymbol{\pi}_1 \mathbf{z}_i$ .

*MC1R* polymorphisms are also associated with a change in receptor localization. cAMP production depends on the number of surface *MC1R* sites per cell, and a lower receptor number would result in decreased production of cAMP and downstream signaling [Sanchez-Mas, et al. 2004]. Beaumont et al. [Beaumont, et al. 2005] quantified cell-surface *MC1R* antibody binding detected by immuno-fluorescence in transient cell expression of variant alleles. We used the log-transformed fold decrease relative to the wildtype allele in construction of  $Z$  matrix (Table 1, model 5). Cell-surface expression of *MC1R* variant alleles

in relation to the wildtype was also quantified by binding of the radio-labeled MC1R ligand 125I-NDP-MSH [Beaumont, et al. 2005]. We used the log-transformed fold decrease in maximum binding values of the variant alleles relative to the wildtype allele (Table 1, model 6).

Reduced MC1R coupling activity is also an important factor contributing to the genetic association between the MC1R variants and the RHC phenotype. Pharmacological characterization of common MC1R variants was conducted previously [Ringholm, et al. 2004; Schioth, et al. 1999]. The log-transformed dissociation constants from competition studies ( $K_i$ ) using  $\alpha$ -MSH and NDP-MSH are listed in Table 1 (models 7–8). The  $K_i$  is the concentration of the competing ligand that binds to half of the binding sites at equilibrium in the absence of radioligand or other competitors. If the  $K_i$  is low, the affinity of the receptor for the inhibitor is high. The differential binding to MC1R between the  $\alpha$ -MSH and NDP-MSH is related to the general binding characteristics of these peptides. The ability of the variant alleles to increase the production of cAMP after stimulation with  $\alpha$ -MSH was measured [Ringholm, et al. 2004; Schioth, et al. 1999]. We listed the log-transformed fold decrease in the maximum production by the MC1R variant alleles relative to that of the wildtype in transfection assays (Table 1, model 9). In this signaling transduction process, the downstream signaling from cAMP production influences the ratio of pheomelanosome to eumelanosome. The cAMP production is most reduced for three RHC alleles, Arg151Cys, Arg160Trp, and Asp294His, which is in good agreement with their association with lighter skin color and hair color.

Finally, we considered a multivariable second-level model. We selected three representative measures, one for each “pathway”: hair color as pigmentary phenotype, cell surface MSH ligand binding as cell surface expression, and kinetics ( $K_i$ ) using  $\alpha$ -MSH as MC1R coupling activity. In this case, the mean log odds ratio for variant  $i$  is  $\frac{\pi_1 + \pi_2 z_{2i} + \pi_3 z_{3i} + \pi_4 z_{4i}}{z_{4i}}$ , where  $z_{hi}$  are the variant-specific coefficients for functional measure  $h$ .

## Results

The genotype distributions of the seven non-synonymous polymorphisms did not show departure from Hardy-Weinberg equilibrium among controls. The minor allele frequency of the seven polymorphisms among the controls was 12.7% for Val60Leu, 9.8% for Val92Met, 7.0% for Arg151Cys, 1.3% for Ile155Thr, 7.8% for Arg160Trp, 4.0% for Arg163Gln, and 1.6% for Asp294His, which were compatible with the previous report on 179 US Caucasian controls [Han, et al. 2004]. Overall, based upon the seven polymorphisms genotyped, 31% of the controls were homozygous for the consensus allele. Half of the controls carried one variant allele; 32% carried one NRHC allele and 18% carried one RHC allele. 20% of the controls carried two variant alleles; 9% carried two NRHC alleles, 8% carried one NRHC allele and one RHC allele, and 3% carried two RHC alleles.

The fixed-effect parameter ( $\pi_0$  and  $\pi_1$ ) estimates are listed in Table 2. The results of the genetic main effect of each variant on melanoma risk are presented in Table 3 and Figures 1–6, which compare the conventional and hierarchical analyses. In the conventional analysis, five of the seven common variants included in the analysis (except 155Thr and 163Gln) were

associated with a significantly increased risk of melanoma. All genetic factors were first assumed to be exchangeable in the hierarchical model (model 1). We then pre-specified the prior second-level covariates according to regression coefficients with phenotypes (models 2–4), MC1R cell-surface expression (models 5–6), and parameters of *in vitro* pharmacologic assays (models 7–9). We observed that the CIs from hierarchical modeling were narrower than those from conventional estimates, i.e., the precision of the risk estimates was enhanced. The change of ORs ranged from –21% to 7%. As was observed by Hung et al. [Hung, et al. 2004], the shrinkage of estimates was not always towards the null, but toward the prior means.

Alterations in the prior matrix influenced the risk estimates (Table 3). The results for the RHC variants -- Arg151Cys, Arg160Trp, and Asp294His -- were robust for melanoma risk; all three variants were significantly associated with melanoma risk in all the models. Similarly, the Val60Leu and Val92Met variants were significantly associated with melanoma risk in all models. The association of Ile155Thr variant with melanoma risk became non-significant in model 6 and borderline significant in model 9. In the models incorporating phenotypic parameters, the association between the Arg163Gln variant and melanoma risk was significant in some models, but not in others (model 3,4 and 9).

For SCC, the Val60Leu, Arg151Cys and Arg160Trp variants were significantly associated with risk in all models. The Asp294His variant was not associated with SCC risk in the conventional model but was significantly associated with SCC risk in three hierarchical models, incorporating hair color, skin color or generation of cAMP in response to MSH. The association of Ile155Thr variant with SCC was not significant in any model except in the hierarchical one based on cell surface MC1R antibody binding. The Val92Met and Arg163Gln variants were not associated with SCC risk in any model. (Table 3)

For BCC, Arg151Cys, Arg160Trp and Arg163Gln were significantly associated with increased risk in all models, whereas Asp294His, Val92Met and Ile155Thr were not associated with BCC risk in any model. The association of Val60Leu variant with BCC risk was not significant only in one of the hierarchical models (model 8).

We constructed a multivariable second-level model using three selected representative measures: hair color as pigmented phenotype, cell surface MSH ligand binding as cell surface expression, and kinetics (Ki) using  $\alpha$ -MSH as MC1R coupling activity. The parameter estimates are listed in Table 4. By comparing the  $\pi$  estimates, we found that cell-surface MSH ligand binding and Ki using  $\alpha$ -MSH parameters showed relevance for BCC but not melanoma and SCC, while pigmented hair color parameters did not strongly affect SNP parameters for any of the three outcomes. However the precision of these parameter estimates is low perhaps due to a small sample size. The results of the genetic main effect of each variant on each type of skin cancer are presented in Table 5. We found that the Val60Leu, Arg151Cys and Arg160Trp variants remained significant for the risk of each type of skin cancer. The Asp163Gln variant was associated with the risk of melanoma and BCC, but not SCC. The Asp294His and Val92Met variants were associated only with melanoma risk. The Ile155Thr variants were not associated with the risk of any type of skin cancer.

## Discussion

Hierarchical modeling has previously been used in genetic association studies. Aragaki et al. [Aragaki, et al. 1997] evaluated the NAT2 genotype-specific dietary effects on adenomatous polyps by constructing a Z matrix of conversion rates calculated for NAT2 genotype-specific enzymatic activity and dietary item-heterocyclic amine combinations. Hung et al. [Hung, et al. 2004] evaluated multiple polymorphisms in different pathways in relation to bladder cancer risk by constructing a Z matrix according to the involvement of each polymorphism in specific pathways. The genetic variants in the *MC1R* gene have been evaluated in relation to pigimentary phenotypes and the risk of melanoma and non-melanoma skin cancer, and characterized quantitatively in *in vitro* phenotypic assays. To our knowledge, this is the first report incorporating biological functional relevance in evaluation of associations between MC1R genetic variants and skin cancer risk. We incorporated the quantitative phenotypic parameters into the Z matrix construction. Consistent with previous reports, our data indicate that hierarchical modeling increases the precision of the risk estimates by tightening the confidence intervals.

Defects in MC1R function are attributable to either decreased affinity for the ligand  $\alpha$ -MSH or altered G-protein coupling. These alleles have important structural functions and in turn influence the binding to the ligand and/or signaling process. In this study, we used some phenotypic parameters of the MC1R variants in the construction of the Z matrix such as correlations with constitutional phenotypes (hair color, skin color, and tanning ability), cell surface expression, and enzymatic kinetic parameters (dissociation constant  $K_i$  and maximum cAMP generation in response to MSH). The MC1R variants were characterized quantitatively in other assays, such as measurements of hair color [Naysmith, et al. 2004] and experimentally induced erythematous response to UV [Flanagan, et al. 2001]. However, only some, but not all, of the seven common variants were assayed in these studies. Hence, we were not able to use these data to evaluate the seven variants simultaneously.

We emphasize that the construction of the Z matrix, i.e., the specification of the prior, is a key component of this approach. The MC1R gene is involved in G-protein-coupled signaling transduction, which in turn is thought to be involved in multiple pathways in addition to pigmentation, such as immune response, inflammatory response, cell proliferation, and extracellular matrix adhesion [Kalden, et al. 1999; Luger, et al. 2003; Murata, et al. 1997; Naysmith, et al. 2004; Robinson and Healy 2002; Smalley and Eisen 2000]. In the construction of the Z matrix in this study, we used three series of parameters: the phenotypic associations in controls (models 2–4), cell surface expression (models 5 and 6), and enzymatic kinetics (models 7–9). The first of these parameters pertained primarily to the pigmentation pathway; the second and third may represent signaling transduction in multiple pathways.

In the multivariable second-level model using three selected representative measures, the two RHC variants, Arg151Cys and Arg160Trp, showed robust association with the risk of each type of skin cancer. This was consistent with the results from the conventional multivariate model. The other RHC variant, Asp294His, was associated only with melanoma risk in the conventional model. After taking into account the three functional parameters in



the hierarchical model, this variant became associated with the risk of SCC as well. This suggests that Asp294His may be involved in the etiology of SCC through these potential mechanisms. Neither the conventional model nor the hierarchical model indicated that Asp294His was associated with the risk of BCC. In addition, the Val60Leu variant was associated with the risk of BCC in most of the hierarchical models, but not in the conventional model. These results suggest that the hierarchical models may provide additional information on risk assessment by incorporating biological functional relevance.

We can also utilize the estimates of the second-level parameter vector, to make inferences about relevant mechanisms that may play a role in carcinogenesis for different skin cancer types. For example, altered affinity for the ligand  $\alpha$ -MSH may play a role in basal cell carcinoma – as evidenced by higher estimates for cell-surface MSH ligand binding and  $\alpha$ -MSH Ki parameters for BCC risk. In contrast, higher estimates of prior-model hyper-parameters suggest that variation in pathways relevant to hair color phenotype may play a bigger role in melanoma and SCC.

For this application several experimentally derived quantitative measures of variant effects on different etiologic pathways were available. However, this is not always the case. In the absence of such data, *in silico* estimates of variant effects could be used. For example, the effects of non-synonymous polymorphisms could be regressed on their SIFT score. The SIFT (Sorting Intolerant From Tolerant) algorithm predicts whether an amino acid substitution will have an impact on protein function based on the alignment of highly similar proteins by evaluating its identity and physicochemical characteristics [Ng and Henikoff 2001; Ng and Henikoff 2002; Ng and Henikoff 2003]. The predictions rely on the evolutionary conservation of amino acids among the protein's family members, which can suggest their importance for the function/structure of the protein. For example, Kanetsky et al. [Kanetsky, et al. 2004] used SIFT analysis to identify MC1R positions that were predicted to be intolerant of amino acid substitutions; the predicted intolerant variants were Asp84Glu, Arg142His, Arg151Cys, Ile155Thr, Arg160Trp, and Asp294His, which is in good agreement with previous publications on their reduced function.

One limitation of the study is our use of self-reported phenotypes such as hair color, skin color, and tanning ability. Self-report has been shown to be an appropriate and widely used method of assessing risk factors for skin cancer. Test-retest reliability of collecting information on phenotypic risk factors, including skin color, tanning/burning tendency, and sunburn history, from questionnaires is moderate to substantial [Branstrom, et al. 2002; Glanz, et al. 2003; Westerdahl, et al. 1996]. Several previous studies have evaluated the validity of self-reported melanoma risk factors. Kang et al. reported 85% agreement between self-reported skin phototypes and those determined by a dermatologist, and these phototypes had a positive correlation with the minimal erythemal dose [Kang, et al. 1992]. Self-reported skin characteristics predict melanoma risk reasonably well. In our study, we observed that the OR of self-reported skin color (fair vs. medium/olive) for melanoma risk was 2.24 (95% CI, 1.59–3.16), which is in good agreement with two previous reports. In a case-control study of 511 cases and 511 age- and gender-matched controls in Western Australia, reflectance-measured skin color was associated with melanoma risk (fair vs. medium/olive: OR, 2.54; 95% CI, 1.94–3.31) [Holman and Armstrong 1984]. In a meta-analysis of 31 case-

control studies, fair skin color was associated with a two-fold increased risk of melanoma (fair vs. medium/olive: OR, 2.06; 95% CI, 1.68–2.52) [Gandini, et al. 2005]. For other constitutional risk factors, such as hair color, sunburn history, and mole counts, the relative risks in our study were also similar to previous reports [Cho, et al. 2005].

In summary, we performed hierarchical modeling analyses, incorporating quantitative information on the function of particular MC1R alleles in an association study with skin cancer risk. Hierarchical modeling stabilizes estimates and provides an alternative analytic approach in genetic association studies.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgements

We would like to thank Dr. Richard A. Sturm at University of Queensland for providing us with the absolute values corresponding to the bar graphs in his original publications. We also thank Dr. Helgi B. Schiöth at Uppsala University for his helpful discussions. We are grateful to the Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School for providing us the resources of the Nurses' Health Study. We are also indebted to the participants in the Nurses' Health Study for their dedication and commitment. We thank the following state cancer registries for their help: Alabama, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Nebraska, New Hampshire, New Jersey, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Virginia, Washington, and Wyoming. The authors assume full responsibility for analyses and interpretation of these data. This work is supported by NIH R01 CA49449, P01 CA87969, UM1 CA186107, UM1 CA167552 and K01DK110267 (ADJ).

## References

- 2017 Cancer facts and figures, American Cancer Society Atlanta.
- Aragaki CC, Greenland S, Probst-Hensch N, Haile RW. 1997 Hierarchical modeling of gene-environment interactions: estimating NAT2 genotype-specific dietary effects on adenomatous polyps. *Cancer Epidemiol Biomarkers Prev* 6(5):307–14. [PubMed: 9149889]
- Bastiaens MT, ter Huurne JA, Kielich C, Gruis NA, Westendorp RG, Vermeer BJ, Bavinck JN. 2001 Melanocortin-1 receptor gene variants determine the risk of nonmelanoma skin cancer independently of fair skin and red hair. *Am J Hum Genet* 68(4):884–94. [PubMed: 11254446]
- Beaumont KA, Newton RA, Smit DJ, Leonard JH, Stow JL, Sturm RA. 2005 Altered cell surface expression of human MC1R variant receptor alleles associated with red hair and skin cancer risk. *Hum Mol Genet* 14(15):2145–54. [PubMed: 15972726]
- Box NF, Duffy DL, Irving RE, Russell A, Chen W, Griffyths LR, Parsons PG, Green AC, Sturm RA. 2001 Melanocortin-1 receptor genotype is a risk factor for basal and squamous cell carcinoma. *J Invest Dermatol* 116(2):224–9. [PubMed: 11179997]
- Box NF, Wyeth JR, O'Gorman LE, Martin NG, Sturm RA. 1997 Characterization of melanocyte stimulating hormone receptor variant alleles in twins with red hair. *Hum Mol Genet* 6(11):1891–7. [PubMed: 9302268]
- Branstrom R, Kristjansson S, Ullen H, Brandberg Y. 2002 Stability of questionnaire items measuring behaviours, attitudes and stages of change related to sun exposure. *Melanoma Res* 12(5):513–9. [PubMed: 12394194]
- Brenner DR, Brennan P, Boffetta P, Amos CI, Spitz MR, Chen C, Goodman G, Heinrich J, Bickeboller H, Rosenberger A and others. 2013 Hierarchical modeling identifies novel lung cancer susceptibility variants in inflammation pathways among 10,140 cases and 11,012 controls. *Hum Genet* 132(5): 579–89. [PubMed: 23370545]

- Cho E, Rosner BA, Feskanich D, Colditz GA. 2005 Risk factors and individual probabilities of melanoma for whites. *J Clin Oncol* 23(12):2669–75. [PubMed: 15837981]
- Flanagan N, Ray AJ, Todd C, Birch-Machin MA, Rees JL. 2001 The relation between melanocortin 1 receptor genotype and experimentally assessed ultraviolet radiation sensitivity. *J Invest Dermatol* 117(5):1314–7. [PubMed: 11710950]
- Gandini S, Sera F, Cattaruzza MS, Pasquini P, Zanetti R, Masini C, Boyle P, Melchi CF. 2005 Meta-analysis of risk factors for cutaneous melanoma: III. Family history, actinic damage and phenotypic factors. *Eur J Cancer* 41(14):2040–59. [PubMed: 16125929]
- Gantz I, Yamada T, Tashiro T, Konda Y, Shimoto Y, Miwa H, Trent JM. 1994 Mapping of the gene encoding the melanocortin-1 (alpha-melanocyte stimulating hormone) receptor (MC1R) to human chromosome 16q24.3 by Fluorescence in situ hybridization. *Genomics* 19(2):394–5. [PubMed: 8188275]
- Glanz K, Schoenfeld E, Weinstock MA, Layi G, Kidd J, Shigaki DM. 2003 Development and reliability of a brief skin cancer risk assessment tool. *Cancer Detect Prev* 27(4):311–5. [PubMed: 12893080]
- Greenland S 1994 Hierarchical regression for epidemiologic analyses of multiple exposures. *Environ Health Perspect* 102 Suppl 8:33–9.
- Greenland S 2000 When should epidemiologic regressions use random coefficients? *Biometrics* 56(3): 915–21. [PubMed: 10985237]
- Han J, Colditz GA, Samson LD, Hunter DJ. 2004 Polymorphisms in DNA double-strand break repair genes and skin cancer risk. *Cancer Res* 64(9):3009–13. [PubMed: 15126335]
- Han J, Kraft P, Colditz GA, Wong J, Hunter DJ. 2006 Melanocortin 1 receptor variants and skin cancer risk. *Int J Cancer* 119(8):1976–84. [PubMed: 16721784]
- Harding RM, Healy E, Ray AJ, Ellis NS, Flanagan N, Todd C, Dixon C, Sajantila A, Jackson IJ, Birch-Machin MA and others. 2000 Evidence for variable selective pressures at MC1R. *Am J Hum Genet* 66(4):1351–61. [PubMed: 10733465]
- Holman CD, Armstrong BK. 1984 Pigmentary traits, ethnic origin, benign nevi, and family history as risk factors for cutaneous malignant melanoma. *J Natl Cancer Inst* 72(2):257–66. [PubMed: 6582314]
- Howe HL, Wingo PA, Thun MJ, Ries LA, Rosenberg HM, Feigal EG, Edwards BK. 2001 Annual report to the nation on the status of cancer (1973 through 1998), featuring cancers with recent increasing trends. *J Natl Cancer Inst* 93(11):824–42. [PubMed: 11390532]
- Hung RJ, Brennan P, Malaveille C, Porru S, Donato F, Boffetta P, Witte JS. 2004 Using hierarchical modeling in genetic association studies with multiple markers: application to a case-control study of bladder cancer. *Cancer Epidemiol Biomarkers Prev* 13(6):1013–21. [PubMed: 15184258]
- Kalden DH, Scholzen T, Brzoska T, Luger TA. 1999 Mechanisms of the antiinflammatory effects of alpha-MSH. Role of transcription factor NF-kappa B and adhesion molecule expression. *Ann N Y Acad Sci* 885:254–61. [PubMed: 10816658]
- Kanetsky PA, Ge F, Najarian D, Swoyer J, Panossian S, Schuchter L, Holmes R, Guerry D, Rebbeck TR. 2004 Assessment of polymorphic variants in the melanocortin-1 receptor gene with cutaneous pigmentation using an evolutionary approach. *Cancer Epidemiol Biomarkers Prev* 13(5):808–19. [PubMed: 15159314]
- Kang S, Fitzpatrick TB, Youn JI. 1992 Identification of population-at-risk for melanoma and epithelial skin cancer using skin phototyping. *Book of Abstracts of 18th World Congress of Dermatology*.
- Kennedy C, ter Huurne J, Berkhout M, Gruis N, Bastiaens M, Bergman W, Willemze R, Bavinck JN. 2001 Melanocortin 1 receptor (MC1R) gene variants are associated with an increased risk for cutaneous melanoma which is largely independent of skin type and hair color. *J Invest Dermatol* 117(2):294–300. [PubMed: 11511307]
- Landi MT, Kanetsky PA, Tsang S, Gold B, Munroe D, Rebbeck T, Swoyer J, Ter-Minassian M, Hedayati M, Grossman L and others. 2005 MC1R, ASIP, and DNA repair in sporadic and familial melanoma in a Mediterranean population. *J Natl Cancer Inst* 97(13):998–1007. [PubMed: 15998953]
- Luger TA, Scholzen TE, Brzoska T, Bohm M. 2003 New insights into the functions of alpha-MSH and related peptides in the immune system. *Ann N Y Acad Sci* 994:133–40. [PubMed: 12851308]

- Mas JS, Gerritsen I, Hahmann C, Jimenez-Cervantes C, Garcia-Borrón JC. 2003 Rate limiting factors in melanocortin 1 receptor signalling through the cAMP pathway. *Pigment Cell Res* 16(5):540–7. [PubMed: 12950734]
- Matichard E, Verpillat P, Mezziani R, Gerard B, Descamps V, Legroux E, Burnouf M, Bertrand G, Bouscarat F, Archimbaud A and others. 2004 Melanocortin 1 receptor (MC1R) gene variants may increase the risk of melanoma in France independently of clinical risk factors and UV exposure. *J Med Genet* 41(2):e13. [PubMed: 14757863]
- Murata J, Ayukawa K, Ogasawara M, Fujii H, Saiki I. 1997 Alpha-melanocyte-stimulating hormone blocks invasion of reconstituted basement membrane (Matrigel) by murine B16 melanoma cells. *Invasion Metastasis* 17(2):82–93. [PubMed: 9561027]
- Naysmith L, Waterston K, Ha T, Flanagan N, Bisset Y, Ray A, Wakamatsu K, Ito S, Rees JL. 2004 Quantitative measures of the effect of the melanocortin 1 receptor on human pigmentation status. *J Invest Dermatol* 122(2):423–8. [PubMed: 15009725]
- Ng PC, Henikoff S. 2001 Predicting deleterious amino acid substitutions. *Genome Res* 11(5):863–74. [PubMed: 11337480]
- Ng PC, Henikoff S. 2002 Accounting for human polymorphisms predicted to affect protein function. *Genome Res* 12(3):436–46. [PubMed: 11875032]
- Ng PC, Henikoff S. 2003 SIFT: Predicting amino acid changes that affect protein function. *Nucleic Acids Res* 31(13):3812–4. [PubMed: 12824425]
- Palmer JS, Duffy DL, Box NF, Aitken JF, O’Gorman LE, Green AC, Hayward NK, Martin NG, Sturm RA. 2000 Melanocortin-1 receptor polymorphisms and risk of melanoma: is the association explained solely by pigmentation phenotype? *Am J Hum Genet* 66(1):176–86. [PubMed: 10631149]
- Rees JL. 2003 Genetics of hair and skin color. *Annu Rev Genet* 37:67–90. [PubMed: 14616056]
- Ringholm A, Klovins J, Rudzish R, Phillips S, Rees JL, Schiöth HB. 2004 Pharmacological characterization of loss of function mutations of the human melanocortin 1 receptor that are associated with red hair. *J Invest Dermatol* 123(5):917–23. [PubMed: 15482480]
- Robinson SJ, Healy E. 2002 Human melanocortin 1 receptor (MC1R) gene variants alter melanoma cell growth and adhesion to extracellular matrix. *Oncogene* 21(52):8037–46. [PubMed: 12439754]
- Sanchez-Mas J, Hahmann C, Gerritsen I, Garcia-Borrón JC, Jimenez-Cervantes C. 2004 Agonist-independent, high constitutive activity of the human melanocortin 1 receptor. *Pigment Cell Res* 17(4):386–95. [PubMed: 15250941]
- Schiöth HB, Phillips SR, Rudzish R, Birch-Machin MA, Wikberg JE, Rees JL. 1999 Loss of function mutations of the human melanocortin 1 receptor are common and are associated with red hair. *Biochem Biophys Res Commun* 260(2):488–91. [PubMed: 10403794]
- Siegel RL, Miller KD, Jemal A. 2017 Cancer statistics, 2017. *CA: A Cancer Journal for Clinicians* 67(1):7–30. [PubMed: 28055103]
- Smalley K, Eisen T. 2000 The involvement of p38 mitogen-activated protein kinase in the alpha-melanocyte stimulating hormone (alpha-MSH)-induced melanogenic and anti-proliferative effects in B16 murine melanoma cells. *FEBS Lett* 476(3):198–202. [PubMed: 10913613]
- Sturm RA. 2002 Skin colour and skin cancer - MC1R, the genetic link. *Melanoma Res* 12(5):405–16. [PubMed: 12394181]
- Sturm RA, Teasdale RD, Box NF. 2001 Human pigmentation genes: identification, structure and consequences of polymorphic variation. *Gene* 277(1–2):49–62. [PubMed: 11602344]
- Valverde P, Healy E, Jackson I, Rees JL, Thody AJ. 1995 Variants of the melanocyte-stimulating hormone receptor gene are associated with red hair and fair skin in humans. *Nat Genet* 11(3):328–30. [PubMed: 7581459]
- Westerdahl J, Anderson H, Olsson H, Ingvar C. 1996 Reproducibility of a self-administered questionnaire for assessment of melanoma risk. *Int J Epidemiol* 25(2):245–51. [PubMed: 9119548]
- Witte JS. 1997 Genetic analysis with hierarchical models. *Genet Epidemiol* 14(6):1137–42. [PubMed: 9433637]
- Witte JS, Greenland S, Haile RW, Bird CL. 1994 Hierarchical regression analysis applied to a study of multiple dietary exposures and breast cancer. *Epidemiology* 5(6):612–21. [PubMed: 7841243]


- Witte JS, Greenland S, Kim LL. 1998 Software for hierarchical modeling of epidemiologic data. *Epidemiology* 9(5):563–6. [PubMed: 9730038]
- Witte JS, Greenland S, Kim LL, Arab L. 2000 Multilevel modeling in epidemiology with GLIMMIX. *Epidemiology* 11(6):684–8. [PubMed: 11055630]

Author Manuscript


Author Manuscript

Author Manuscript


Author Manuscript



**Figure 1.** Beta estimates from the conventional, prior, and hierarchical models for association between MC1R variants and risk of melanoma.




**Figure 2.** Beta estimates from the conventional, prior, and hierarchical models for association between MC1R variants and risk of SCC.




**Figure 3.** Beta estimates from the conventional, prior, and hierarchical models for association between MC1R variants and risk of BCC.






**Figure 4.** Standard error (se) of the beta estimates from the conventional and hierarchical models for the association between MC1R variants and risk of melanoma.



**Figure 5.** Standard error (se) of the beta estimates from the conventional and hierarchical models for the association between MC1R variants and risk of SCC.



**Figure 6.** Standard error (se) of the beta estimates from the conventional and hierarchical models for the association between MC1R variants and risk of BCC.

**Table 1.**

## Hierarchical modeling of MC1R variants

Model		Wildtype	Val60Leu	Val92Met	Arg151Cys	Ile155Thr	Arg160Trp	Arg163Gln	Asp294His
1	Assumes that all variants have the same effects	0	1	1	1	1	1	1	1
2	Regression coefficient with hair color in controls	0	-0.020	-0.092	0.368	0.195	0.539	0.028	0.605
3	Regression coefficient with skin color in controls	0	0.149	0.083	0.310	0.140	0.250	0.026	0.428
4	Regression coefficient with tanning ability in controls	0	0.317	0.259	0.625	0.477	0.453	0.136	0.899
5	Cell surface MC1R antibody binding	0	0.772	0.004	1.505	1.911	1.537	0.411	-0.378
6	Cell surface MSH ligand binding	0	1.234	0.126	1.966	3.808	2.137	0.648	-0.466
7	Kinetics, $\alpha$ -MSH Ki	0	0.276	3.191	-0.034	1,878	0.259	1.563	1.887
8	Kinetics, NDP Ki	0	-0.084	0.069	0.808	0.313	0.425	0.468	-0.166
9	Maximum cAMP generation in response to MSH	0	1.006	0.211	1.634	0.483	1.516	-0.401	2.088

Parameters for models 5–9 were log transformed.

**Table 2.**

Estimates of prior-model hyper-parameters: univariate second-level models

Model	Melanoma		SCC		BCC	
	$\pi_0$ est (SD)	$\pi_1$ est (SD)	$\pi_0$ est (SD)	$\pi_1$ est (SD)	$\pi_0$ est (SD)	$\pi_1$ est (SD)
1 Fixed mean	0.63 (0.18)	–	0.46 (0.17)	–	0.38 (0.17)	–
2 Hair color	0.52 (0.22)	0.56 (0.58)	0.33 (0.21)	0.61 (0.58)	0.30 (0.21)	0.38 (0.58)
3 Skin color	0.37 (0.30)	1.34 (1.24)	0.22 (0.29)	1.23 (1.25)	0.26 (0.29)	0.61 (1.22)
4 Tanning ability	0.29 (0.35)	0.79 (0.69)	0.20 (0.34)	0.59 (0.70)	0.34 (0.33)	0.10 (0.68)
5 Cell surface MC1R antibody binding	0.62 (0.24)	0.01 (0.21)	0.29 (0.24)	0.21 (0.20)	0.36 (0.23)	0.02 (0.20)
6 Cell surface MSH ligand binding	0.64 (0.24)	–0.003 (0.13)	0.32 (0.24)	0.11 (0.13)	0.45 (0.24)	–0.06 (0.14)
7 Kinetics, $\alpha$ -MSH Ki	0.67 (0.24)	–0.03 (0.13)	0.61 (0.23)	–0.13 (0.13)	0.57 (0.23)	–0.16 (0.13)
8 Kinetics, NDP Ki	0.58 (0.22)	0.22 (0.47)	0.34 (0.21)	0.41 (0.46)	0.23 (0.21)	0.53 (0.46)
9 Maximum cAMP generation	0.48 (0.26)	0.16 (0.19)	0.27 (0.25)	0.20 (0.19)	0.31 (0.24)	0.08 (0.19)

**Table 3.**

Hierarchical modeling of associations between MC1R variants and skin cancer risk

	Val60Leu	Val92Met	Arg151Cys	Ile155Thr	Arg160Trp	Arg163Gln	Asp294His	
<b>Melanoma</b>	<b>Conventional model</b>	1.52 (1.04 – 2.24)	1.64 (1.10 – 2.45)	2.47 (1.63 – 3.75)	1.87 (0.80 – 4.40)	1.79 (1.15 – 2.80)	1.71 (0.96 – 3.04)	2.67 (1.35 – 5.31)
	<b>Model 1</b>	1.60 (1.13 – 2.28)	1.69 (1.18 – 2.44)	2.30 (1.58 – 3.36)	1.88 (1.07 – 3.31)	1.82 (1.22 – 2.70)	1.78 (1.12 – 2.84)	2.22 (1.32 – 3.72)
	<b>Model 2</b>	1.55 (1.09 – 2.23)	1.63 (1.12 – 2.37)	2.35 (1.61 – 3.44)	1.87 (1.06 – 3.44)	1.92 (1.27 – 2.88)	1.71 (1.06 – 2.75)	2.50 (1.42 – 4.41)
	<b>Model 3</b>	1.58 (1.11 – 2.25)	1.63 (1.13 – 2.37)	2.39 (1.63 – 3.51)	1.79 (1.01 – 3.18)	1.86 (1.25 – 2.76)	1.62 (0.98 – 2.68)	2.63 (1.45 – 4.77)
	<b>Model 4</b>	1.57 (1.10 – 2.23)	1.64 (1.13 – 2.37)	2.39 (1.63 – 3.50)	1.92 (1.09 – 3.38)	1.82 (1.23 – 2.71)	1.61 (0.97 – 2.66)	2.71 (1.47 – 4.98)
	<b>Model 5</b>	1.60 (1.13 – 2.28)	1.69 (1.17 – 2.45)	2.31 (1.57 – 3.40)	1.90 (1.01 – 3.58)	1.82 (1.22 – 2.73)	1.78 (1.12 – 2.85)	2.20 (1.24 – 3.92)
	<b>Model 6</b>	1.60 (1.13 – 2.28)	1.70 (1.17 – 2.45)	2.30 (1.57 – 3.37)	1.87 (0.92 – 3.80)	1.81 (1.21 – 2.71)	1.79 (1.12 – 2.85)	2.22 (1.26 – 3.93)
	<b>Model 7</b>	1.61 (1.13 – 2.30)	1.67 (1.14 – 2.45)	2.33 (1.58 – 3.43)	1.86 (1.05 – 3.30)	1.83 (1.23 – 2.74)	1.78 (1.12 – 2.84)	2.19 (1.30 – 3.71)
	<b>Model 8</b>	1.58 (1.10 – 2.26)	1.68 (1.16 – 2.42)	2.38 (1.59 – 3.56)	1.90 (1.08 – 3.33)	1.84 (1.23 – 2.73)	1.82 (1.14 – 2.92)	2.10 (1.20 – 3.70)
	<b>Model 9</b>	1.60 (1.13 – 2.28)	1.65 (1.14 – 2.39)	2.37 (1.61 – 3.47)	1.79 (1.00 – 3.19)	1.86 (1.25 – 2.78)	1.62 (0.96 – 2.73)	2.46 (1.40 – 4.32)
<b>SCC</b>	<b>Conventional model</b>	1.43 (1.02 – 2.01)	1.25 (0.86 – 1.81)	2.13 (1.47 – 3.09)	1.61 (0.72 – 3.59)	1.97 (1.34 – 2.89)	1.27 (0.74 – 2.17)	1.55 (0.75 – 3.23)
	<b>Model 1</b>	1.46 (1.06 – 2.00)	1.32 (0.94 – 1.85)	1.98 (1.41 – 2.79)	1.59 (0.92 – 2.73)	1.87 (1.31 – 2.66)	1.38 (0.89 – 2.13)	1.56 (0.93 – 2.62)
	<b>Model 2</b>	1.42 (1.03 – 1.96)	1.26 (0.89 – 1.79)	2.03 (1.43 – 2.86)	1.58 (0.91 – 2.72)	1.96 (1.37 – 2.82)	1.32 (0.84 – 2.06)	1.79 (1.01 – 3.18)
	<b>Model 3</b>	1.44 (1.05 – 1.99)	1.28 (0.90 – 1.81)	2.05 (1.44 – 2.90)	1.52 (0.88 – 2.65)	1.90 (1.33 – 2.71)	1.27 (0.79 – 2.03)	1.84 (1.00 – 3.89)
	<b>Model 4</b>	1.44 (1.05 – 1.98)	1.29 (0.91 – 1.82)	2.03 (1.44 – 2.88)	1.61 (0.93 – 2.78)	1.88 (1.32 – 2.67)	1.28 (0.80 – 2.05)	1.83 (0.98 – 3.43)
	<b>Model 5</b>	1.46 (1.06 – 2.00)	1.27 (0.90 – 1.80)	2.06 (1.45 – 2.92)	1.83 (1.00 – 3.33)	1.94 (1.35 – 2.78)	1.35 (0.87 – 2.09)	1.36 (0.75 – 2.45)
	<b>Model 6</b>	1.46 (1.06 – 2.00)	1.28 (0.91 – 1.82)	2.02 (1.43 – 2.86)	1.87 (0.96 – 3.64)	1.91 (1.34 – 2.73)	1.35 (0.87 – 2.10)	1.40 (0.79 – 2.51)
	<b>Model 7</b>	1.49 (1.08 – 2.06)	1.24 (0.86 – 1.78)	2.06 (1.45 – 2.93)	1.50 (0.86 – 1.78)	1.92 (1.35 – 2.75)	1.36 (0.88 – 2.11)	1.48 (0.87 – 2.52)
	<b>Model 8</b>	1.42 (1.03 – 1.96)	1.29 (0.92 – 1.82)	2.08 (1.45 – 2.99)	1.61 (0.93 – 2.77)	1.90 (1.33 – 2.70)	1.43 (0.92 – 2.22)	1.40 (0.79 – 2.49)
	<b>Model 9</b>	1.46 (1.06 – 2.01)	1.28 (0.90 – 1.80)	2.05 (1.44 – 2.90)	1.50 (0.86 – 2.62)	1.92 (1.35 – 2.74)	1.23 (0.76 – 2.01)	1.79 (1.01 – 3.17)
<b>BCC</b>	<b>Conventional model</b>	1.39 (0.99 – 1.95)	1.02 (0.70 – 1.50)	2.06 (1.41 – 2.99)	0.51 (0.15 – 1.73)	1.60 (1.07 – 2.37)	2.10 (1.32 – 3.35)	1.46 (0.71 – 3.04)
	<b>Model 1</b>	1.40 (1.02 – 1.92)	1.13 (0.80 – 1.59)	1.90 (1.34 – 2.68)	1.11 (0.62 – 1.98)	1.57 (1.09 – 2.25)	1.85 (1.23 – 2.78)	1.47 (0.87 – 2.47)
	<b>Model 2</b>	1.38 (1.00 – 1.90)	1.10 (0.77 – 1.56)	1.92 (1.36 – 2.72)	1.11 (0.62 – 1.98)	1.62 (1.12 – 2.34)	1.81 (1.20 – 2.74)	1.60 (0.90 – 2.85)
	<b>Model 3</b>	1.40 (1.02 – 1.92)	1.11 (0.78 – 1.57)	1.93 (1.36 – 2.74)	1.09 (0.60 – 1.96)	1.58 (1.10 – 2.27)	1.79 (1.17 – 2.75)	1.60 (0.87 – 2.94)
	<b>Model 4</b>	1.40 (1.02 – 1.92)	1.12 (0.79 – 1.59)	1.91 (1.34 – 2.70)	1.11 (0.62 – 1.99)	1.57 (1.10 – 2.25)	1.84 (1.20 – 2.81)	1.51 (0.80 – 2.85)

	Val60Leu	Val92Met	Arg151Cys	Ile155Thr	Arg160Trp	Arg163Gln	Asp294His
<b>Model 5</b>	1.40 (1.02 – 1.92)	1.12 (0.79 – 1.60)	1.90 (1.34 – 2.70)	1.13 (0.58 – 2.17)	1.57 (1.09 – 2.27)	1.85 (1.23 – 2.78)	1.45 (0.81 – 2.60)
<b>Model 6</b>	1.40 (1.02 – 1.92)	1.14 (0.80 – 1.62)	1.88 (1.32 – 2.66)	1.00 (0.47 – 2.15)	1.55 (1.07 – 2.23)	1.87 (1.24 – 2.80)	1.55 (0.87 – 2.74)
<b>Model 7</b>	1.45 (1.05 – 1.99)	1.04 (0.72 – 1.51)	1.99 (1.40 – 2.82)	1.03 (0.56 – 1.87)	1.62 (1.13 – 2.33)	1.83 (1.22 – 2.75)	1.38 (0.81 – 2.35)
<b>Model 8</b>	1.36 (0.98 – 1.87)	1.09 (0.77 – 1.55)	2.03 (1.41 – 2.91)	1.13 (0.63 – 2.01)	1.59 (1.11 – 2.89)	1.93 (1.28 – 2.91)	1.28 (0.72 – 2.28)
<b>Model 9</b>	1.40 (1.02 – 1.93)	1.11 (0.78 – 1.58)	1.92 (1.35 – 2.72)	1.08 (0.60 – 1.96)	1.58 (1.10 – 2.28)	1.79 (1.15 – 2.78)	1.55 (0.87 – 2.74)

Each model includes seven variants (carrier versus non-carrier) simultaneously and adjusts for age.

Conventional model is the logistical regression model. Models 1–9 utilize hierarchical modeling approach.

Model 1 assumes that all variants have the same effects.

Model 2 is based on the associations with hair color among controls.

Model 3 is based on the associations with skin color among controls.

Model 4 is based on the associations with tanning ability among controls.

Model 5 is based on cell surface MC1R antibody binding.

Model 6 is based on cell surface MSH ligand binding.

Model 7 kinetics, Ki-a-MSH.

Model 8 kinetics, Ki-NDP.

Model 9 is based on maximum generation of cAMP in response to MSH.

**Table 4.**

Estimates of prior-model hyperparameters: multivariate second-level model

	<b>Melanoma</b>	<b>SCC</b>	<b>BCC</b>
<b>Variable</b>	<b><math>\pi</math> est (SD)</b>	<b><math>\pi</math> est (SD)</b>	<b><math>\pi</math> est (SD)</b>
<b>Mean</b>	0.52 (0.43)	0.37 (0.41)	0.97 (0.45)
<b>Cell surface MSH ligand binding</b>	-0.03 (0.15)	0.05 (0.15)	-0.24 (0.18)
<b>Kinetics, <math>\alpha</math>-MSH Ki</b>	0.02 (0.16)	-0.06 (0.16)	-0.28 (0.18)
<b>Hair color</b>	0.61 (0.65)	0.44 (0.65)	0.09 (0.64)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Table 5.**

Hierarchical modeling of MC1R variants using multivariate second-level model

	<b>Val60Leu</b>	<b>Val92Met</b>	<b>Arg151Cys</b>	<b>Ile155Thr</b>	<b>Arg160Trp</b>	<b>Arg163Gln</b>	<b>Asp294His</b>
<b>Melanoma</b>	1.55 (1.07–2.24)	1.64 (1.11–2.42)	2.3 (1.58–3.44)	1.80 (0.80–4.05)	1.91 (1.26–2.88)	1.71 (1.06–2.77)	2.60 (1.37–4.95)
<b>SCC</b>	1.45 (1.04–2.02)	1.23 (0.85–1.77)	2.07 (1.46–2.94)	1.67 (0.78–3.57)	1.98 (1.38–2.86)	1.31 (0.84–2.06)	1.60 (0.82–3.14)
<b>BCC</b>	1.46 (1.05–2.03)	1.03 (0.71–1.50)	1.97 (1.39–2.81)	0.61 (0.22–1.97)	1.59 (1.09–2.31)	1.86 (1.23–2.81)	1.66 (0.85–3.24)

The model includes seven variants (carrier versus non-carrier) simultaneously and adjusts for age. We use multivariate second-level model incorporate three representative measures: cell surface MSH ligand binding, kinetics,  $\alpha$ -MSH Ki, and hair color.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript