



# EPA PUBLIC ACCESS

Author manuscript

*Neurotoxicology*. Author manuscript; available in PMC 2019 September 24.

About author manuscripts

Submit a manuscript

Published in final edited form as:

*Neurotoxicology*. 2017 March ; 59: 155–163. doi:10.1016/j.neuro.2016.10.012.

## Atypical Microglial Response to Biodiesel Exhaust in Healthy and Hypertensive Rats

Christen L. Mumaw<sup>1</sup>, Michael Surace<sup>2</sup>, Shannon Levesque<sup>3</sup>, Urmila P. Kodavanti<sup>4</sup>, Prasada Rao S. Kodavanti<sup>5</sup>, Joyce E. Royland<sup>6</sup>, Michelle L. Block<sup>1,\*</sup>

<sup>1</sup>Department of Anatomy and Cell Biology, The Stark Neuroscience Research Institute, Indiana University School of Medicine, Indianapolis, IN 46202, USA

<sup>2</sup>Department of Biochemistry & Molecular Biology, Virginia Commonwealth University Medical Campus, Richmond, VA 23298, USA

<sup>3</sup>Department of Anatomy and Neurobiology, Virginia Commonwealth University Medical Campus, Richmond, VA 23298, USA

<sup>4</sup>Environmental Public Health Division, National Health and Environmental Effects Research Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina 27711, USA

<sup>5</sup>Neurotoxicology Branch, NHEERL, ORD, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina 27711, USA;

<sup>6</sup>Genetic and Cellular Toxicology Branch, NHEERL, ORD, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina 27711, USA.

### Abstract

Accumulating evidence suggests a deleterious role for urban air pollution in central nervous system (CNS) diseases and neurodevelopmental disorders. Microglia, the resident innate immune cells and sentinels in the brain, are a common source of neuroinflammation and are implicated in how air pollution may exert CNS effects. While renewable energy, such as soy-based biofuel, is of increasing public interest, there is little information on how soy biofuel may affect the brain. To address this, male spontaneously hypertensive rats (SHR) and normotensive Wistar Kyoto (WKY) rats were exposed to 100% Soy Biodiesel Exhaust (100SBDE; 0, 50, 150 and 500  $\mu\text{g}/\text{m}^3$ ) by inhalation for 4 h/day for 4 weeks (5 days/week). IBA-1 staining of microglia in the substantia nigra revealed significant changes in morphology with 100SBDE exposure in rats from both genotypes, where the SHR were less sensitive. Further analysis failed to show consistent changes in pro-inflammatory cytokine expression, nitrated protein, and arginase1 expression in brain tissue from either rat strain exposed to 100SBDE. CX3CR1 and fractalkine mRNA expression were

\*Address correspondence to: Michelle L. Block, Ph.D., Associate Professor, Indiana University School of Medicine, Department of Anatomy & Cell Biology, The Stark Neuroscience Research Institute, Indianapolis, IN 46202, Phone: (317) 274-7180, Fax: (317) 278-2040, mlblock@iupui.edu.

#### CONFLICT OF INTEREST STATEMENT

The authors declare no financial interests or potential conflicts of interest.

#### TRANSPARENCY DOCUMENT

The transparency document associated with this article can be found in the online version.

lower in the striatum of all 100SBDE exposed rats, but greater SBDE exposure was required for loss of fractalkine expression in the SHR. Together, these data support that month-long 100SBDE exposure impacts the basal ganglia with changes in microglia morphology, an impaired fractalkine axis, and an atypical activation response without traditional markers of M1 or M2 activation, where the SHR may be less sensitive to these effects.

## Keywords

Biodiesel; Air Pollution; Brain; Microglia; Atypical Activation

---

## 1. INTRODUCTION

Renewable energy, such as biofuel, has received increasing public attention. However, current understanding about the health effects of biofuel is limited (Larcombe et al. , 2015), particularly regarding how it might impact the brain. Derived from plant or animal oil through the transesterification of triglycerides with alcohol, biodiesel may be used in place of traditional petroleum diesel fuel sources (Larcombe, Kicic, 2015, Swanson et al. , 2007). Importantly, depending on the source and composition, the exhaust emissions of biofuel pollutants (particulate matter, adsorbed chemicals, and gases) can be lower than traditional mineral fuel diesel exhaust (Mutlu et al. , 2015, Prokopowicz et al. , 2015, Shojaeefard et al. , 2013). While this suggests the potential for health benefits, the effects of each exhaust might be unique to the organ system examined. For example, biodiesel has been shown to result in acute pulmonary inflammation similar to petroleum diesel (Mehus et al. , 2015), but also elevated acute vascular effects (Bass et al. , 2015), significant increased liver and lung tissue damage (Shvedova et al. , 2013), elevated pulmonary and liver oxidative stress (Shvedova, Yanamala, 2013), and more pronounced pro-inflammatory pulmonary responses (Shvedova, Yanamala, 2013). At present, the impact of biofuel in the brain is largely unknown.

Urban air pollution is a complex chemical mixture derived from multiple sources, including engine emissions, coal combustion, biomass burning, and secondary photochemical products. Diesel exhaust is a major constituent of near-road and urban air pollution and is commonly used as a surrogate model of air pollution in health effects studies (Hesterberg et al. , 2010, Ma and Ma, 2002). In the U.S. alone, millions of people are exposed to air pollution levels above safety standards (Mauderly et al. , 2010). Importantly, evidence points to a link between high levels of urban air pollution and several CNS conditions including, elevated stroke incidence (Bedada et al. , 2012, Wellenius et al. , 2012b), increased Autism risk (Becerra et al. , 2013, Roberts et al. , 2013, Volk et al. , 2011, Volk et al. , 2013), decreased cognitive function in the elderly (Power et al. , 2011, Power et al. , 2013, Wellenius et al. , 2012a), elevated Alzheimer's disease (AD)-risk (Jung et al. , 2015), increased Parkinson's disease (PD) risk (Kirrane et al. , 2015), and accelerated neurodegenerative disease progression (Kioumourtoglou et al. , 2015). This is consistent with case studies demonstrating an association of living in highly polluted areas with elevated AD-like neuropathology (Calderon-Garciduenas et al. , 2012, Calderon-Garciduenas et al. , 2004, Calderon-Garciduenas et al. , 2008) and Parkinson's-disease like

neuropathology in humans (Calderon-Garciduenas et al. , 2011, Calderon-Garciduenas et al. , 2013). As such, it is of pressing public health interest to understand the impact of different sources of air pollution on the brain and their effect on unique genetic backgrounds that might confer vulnerability.

Microglia are the resident innate immune cells in the brain, mandatory for healthy CNS function, and constantly active, both as sentinels surveying the CNS environment (Hickman et al. , 2013) and as “electricians” regulating synaptic communication (Sun et al. , 2014). As sentinels, microglia detect and respond to cellular damage, pathogens, and environmental exposures (Block et al. , 2007), resulting in a shift in phenotype commonly referred to as “activation”. The deleterious response associated with neuron damage and CNS disease is predicted to occur when the microglial pro-inflammatory response is excessive, unregulated, and fails to resolve (Block, Zecca, 2007, Heneka et al. , 2014). Microglia have long been implicated in the neuronal damage occurring in many CNS diseases and conditions, including those impacted by air pollution such as AD (Heppner et al. , 2015), PD (Block, Zecca, 2007), and Autism (Takano, 2015). Thus, microglia and neuroinflammation have been implicated as a common underlying mechanisms through which air pollution may affect diverse CNS conditions (Block and Calderon-Garciduenas, 2009).

Increasing evidence indicates that ambient air pollutants are a prevalent environmental source of chronic neuroinflammation and microglial activation (Costa et al. , 2015). Urban particulate matter (Allen et al. , 2014, Campbell et al. , 2009, Campbell et al. , 2005, Morgan et al. , 2011), O<sub>3</sub> (Santiago-Lopez et al. , 2010), diesel exhaust (Bolton et al. , 2012, Gerlofs-Nijland et al. , 2010, Levesque et al. , 2011a, Levesque et al. , 2011b), and manganese (Antonini et al. , 2009, Elder et al. , 2006) are examples of inhaled pollutants that elevate CNS cytokines and oxidative stress in animal models. Human reports confirm microglia activation and neuroinflammation in individuals living in highly polluted cities (Calderon-Garciduenas, Solt, 2008). Additionally, *in vitro* studies indicate that the individual components of air pollution, especially particulate matter, may directly activate microglia (Block et al. , 2004, Campbell et al. , 2014, Levesque, Taetzsch, 2011b, Morgan, Davis, 2011). Thus, while the specific mechanisms underlying how microglia detect and respond to air pollution remain unresolved, there appears to be a prototypical pro-inflammatory microglial activation phenotype and neuroinflammation linked to several forms of air pollution, particularly diesel exhaust.

In an effort to begin to understand how pollutants from biodiesel exhaust may affect the brain, we addressed the effects of month-long exposure to soy biodiesel exhaust on the neuroinflammation phenotype in the brains of normal and hypertensive rats, including morphology, M1 markers, and M2 markers. The spontaneously hypertensive rats have been previously reported to be more vulnerable to the extra-pulmonary effects of air pollution (Saxena et al. , 2009) and were originally predicted to be more sensitive to the CNS effects of biodiesel exposure.

## 2. METHODS

### 2.1. Reagents

The polyclonal antibody against the Ionized Calcium-binding adapter molecule-1 (IBA-1) microglial marker was purchased from Wako (Richmond, VA). The Biotinylated goat anti-rabbit secondary antibody was purchased from Vector Laboratories (Burlingame, CA). ELISA kits were purchased from R&D Systems (Minneapolis, MN). All other reagents were procured from Sigma-Aldrich (St. Louis, MO).

### 2.2. Animals

Twelve, fourteen week old male Spontaneously Hypertensive rats (SHR) and normotensive Wistar Kyoto (WKY) rats were purchased from Charles River Laboratories (Raleigh, NC). SHR are a model of cardiovascular disease frequently employed to discern the underlying mechanisms of the cardiopulmonary health effects of PM, where SHR are more vulnerable to these extra-pulmonary effects of air pollution (Saxena, Gilmour, 2009). Animals were acclimated to the housing facility for 1 week before studies began. Rats were housed in an AAALAC-accredited housing facility and maintained at 20–24°C and on a 12h light: dark schedule. All animal experiments were approved by the US EPA IACUC and the National Institutes of Health guidelines were followed. All animals were treated humanely and with regard for alleviation of suffering.

### 2.3. Biodiesel Exhaust Exposure

A 100% soy biodiesel fuel (100SBDE) was purchased from Piedmont Biofuels (Pittsboro, NC). This fuel was used to power a Yanmar L70 diesel engine (Adairville, GA) and Pramac E3750 generator (Marietta, GA). This single cylinder, 320-cm<sup>3</sup> engine ran at 5.8 hp (4.3 kW) with a continuous load, 3600 rpm. The engine exhaust (~ 85 L/min) was diluted with HEPA and charcoal filtered room air (~7:1 dilution) and was directed to three 984 L Hazelton model 1000 inhalation exposure chambers located in an adjacent inhalation exposure laboratory. Additional HEPA filtered room air was introduced for secondary dilution to the desired exposure concentrations. The aerosol concentration in each chamber was continuously monitored using tapered element oscillating microbalances (TEOM, Rupprecht and Patashnick Co., series 1400, Albany, NY). Particle-size distributions were measured using a scanning mobility particle sizer (SMPS, TSI Inc., model 3080, St. Paul, MN) for all chambers. Target exposure concentrations of 50, 150, and 500 µg/m<sup>3</sup> were achieved concurrently in the three chambers by manually adjusting the secondary dilution according to the TEOM determined mass concentration. The fourth Hazelton chamber was used as a control, receiving only HEPA/charcoal filtered room air. Inhalation chamber temperatures, pressures, and relative humidity were maintained hourly during exposure. The flow rate of 283 L/min (10 ft<sup>3</sup>/min) for all chambers was maintained, which is sufficient for approximately 17 air changes per hour. Also, all chambers were continuously monitored for oxygen (O<sub>2</sub>, California Analytical Instruments, model 200, Orange, CA.), carbon monoxide (CO, Thermo Electron Instruments, model 48, Fitchburg, WI.), sulfur dioxide (SO<sub>2</sub>, Thermo Environmental Instruments, model 43C, Franklin, MA.), nitric oxide and total nitrogen oxides (NO and NO<sub>x</sub>, Thermo Scientific, model 42i-HL, West Palm Beach, FL). Nitrogen

dioxide (NO<sub>2</sub>) levels were obtained by subtracting NO values from total NO<sub>x</sub>. Further details are provided in previous publications (Mutlu et al.,2015; Bass et al., 2015).

In brief, Rats were exposed for 4 h/day for 4 weeks (5 days/week) to 100% Soy Biodiesel Exhaust (100SBDE) with particulate matter concentrations of 0, 50, 150 and 500 µg/m<sup>3</sup>, which are levels higher than typically encountered in ambient air, but may be achieved during heavy traffic or occupational situations. The exhaust characterization for 100SBDE was previously reported in detail and compared to values from exhaust obtained from petroleum diesel fuel (Mutlu, Nash, 2015), where 100SBDE PM concentrations were 30% lower than traditional petroleum diesel fuel exhaust. When emissions were diluted with air to control equivalent PM mass concentrations, 100SBDE exhibited lower CO and slightly higher NO concentrations when compared to the traditional petroleum diesel fuel exhaust (Mutlu, Nash, 2015). Further, the organic/elemental carbon ratios and oxygenated methyl esters and organic acids were higher for 100SBDE and it exhibited unimodal-accumulation mode particle-size distributions, with mildly higher concentrations of slightly larger particles when compared to exhaust composition from traditional petroleum diesel fuel (Mutlu, Nash, 2015). Only rats exposed to filtered air and 100SBDE were used in this study, where one day following the last day of exposure, rats were anesthetized with an overdose of sodium pentobarbital (Virbac AH, Inc., Fort Worth, TX; 50–100 mg/kg, ip), and the left hemisphere of the brain was immediately dissected (striatum and remaining brain) for protein and mRNA analyses and the right hemisphere was fixed for histology. The striatum was targeted due to our previous reports demonstrating the midbrain (Levesque, Surace, 2011a, Levesque, Taetzsch, 2011b), and potentially the basal ganglia, may be more sensitive to the pro-inflammatory effects of diesel exhaust.

#### 2.4. Quantitative Reverse Transcriptase Polymerase Chain Reaction

Total RNA was extracted from the rat brain striatum using Trizol (Invitrogen Life Technologies, Grand Island, NY) according to manufacturer's instructions. The RNA was treated with Ambion DNase I (Invitrogen Life Technologies, Grand Island, NY), which was subsequently removed with Qiagen RNeasy RNA cleanup kit (Qiagen, Germantown, MD). Reverse transcription of RNA (0.3 – 1.0 µg/sample) was performed with iScript Reverse Transcription Supermix (BioRad, Hercules, CA) according to manufacturer's instructions. SsoFast Evagreen Supermix (BioRad, Hercules, CA) and 500 nM forward and reverse primers were used to carry out quantitative PCR on a CFX96 (BioRad, Hercules, CA) real-time PCR detection system, per manufacturer's instructions. Cycling parameters were 1 cycle at 95°C for 5 min, 40 cycles of 95°C (5 sec.) and 56°C (5 sec.) followed by a melt curve measurement consisting of 5 second 0.5°C incremental increases from 65°C to 95°C. The primer sequences are listed in Table S1.

#### 2.5. Protein Isolation: Brain Homogenate

Remaining brain tissue was homogenized as general brain homogenate. Protein was isolated by suspending frozen tissue in 10 volumes of lysis buffer (Cytobuster Protein Extraction Reagent; EMD Chemicals; Darmstadt, Germany) with 10 µL/mL HALT protease inhibitor and 10 µL/mL EDTA. Samples were homogenized using a motorized pellet mixer and then

centrifuged for 5 minutes at 5000 g. The protein concentration of the resulting supernatant was determined using a BCA protein assay (ThermoScientific; Rockford, IL).

## 2.6. Immunohistochemistry

The right hemisphere of the brain was fixed in 4% paraformaldehyde for 2 days and cryoprotected in 30% sucrose. Coronal sections (40  $\mu$ m) of the midbrain region containing the substantia nigra pars compacta (SNpc) were collected with a freezing stage microtome (Microm HM 450, Thermo Scientific, Waltham, MA). Dopaminergic neurons were stained to confirm that sectioned samples across rat brain were in the same frame of the SNpc, and were recognized with an anti-tyrosine hydroxylase (TH, 1:1000) rabbit polyclonal antibody (Millipore Billerica, MA) and microglia were stained with a polyclonal rabbit anti-ionized calcium-binding adaptor molecule-1 (IBA-1, 1:1000) antibody (Wako, Richmond, VA). Free-floating brain slices were treated with 1% H<sub>2</sub>O<sub>2</sub>, washed twice for 10 min with phosphate buffered saline (PBS), incubated 20 min with a blocking solution (PBS containing 1% bovine serum albumin, 4% goat serum, 0.4% Triton X-100) and incubated overnight at 4°C with primary antibody diluted 1:1000 in DAKO antibody diluent. Slices were then washed twice in PBS, incubated with biotinylated anti-rabbit antibody (Vector Laboratories, Burlingame, CA) for 1 h, washed twice in PBS, and incubated with Vectastain ABC Kit (Vector Laboratories, Burlingame, CA) reagents according to manufacturer's instructions. Staining was visualized using 3,3'-diaminobenzidine and urea-hydrogen peroxide tablets (Sigma-Aldrich, St. Louis, MO). Images were captured with an Olympus BX51 microscope (Olympus America, Center Valley, PA).

## 2.7. Immunoblotting

Ten micrograms of protein from each sample was electrophoresed on a 12% SDS-PAGE gel. Samples were transferred to nitrocellulose membranes by semi-dry transfer, blocked with 5% nonfat milk for 1 hr. at 24°C, followed by incubation overnight with the anti-GAPDH (1:1000), anti-P38 (1:1000) and anti-phosphorylated P38 (1:1000) antibodies at 4°C (Cell Signaling Technologies, Danvers, MA), where all antibodies have been reported to cross-react with the rat proteins. Blots were then incubated with horseradish peroxidase-linked mouse anti-rabbit (1:5000) or goat anti-mouse (1:5000) for 1 h (24°C) and ECL+Plus reagents (Amersham Biosciences Inc., Piscataway, NJ) were used as a detection system. Band density was quantitated with ImageJ (Abramoff et al. , 2004) and analyzed as a ratio of total P38 and phosphorylated P38 bands. Results are reported as a percent change from control.

## 2.8. Cytokine ELISAs

Fractalkine and TNF $\alpha$  protein concentrations in 100 $\mu$ g/well of brain homogenate protein was measured using a commercial enzyme-linked immunosorbent assay (ELISA) kit from R&D Systems (Minneapolis, MN).

## 2.9. Nitrotyrosine ELISA

The amount of nitrated proteins was measured using 100 µg/well of brain homogenate with a commercial enzyme-linked immunosorbent assay (ELISA) kit from Millipore (Temecula, CA), per the manufacturer protocol.

## 2.10. Statistical Analysis

Data are expressed as representative images, raw values, fold change, and the percentage of control, where control values were set to either 100% or 1 accordingly. The treatment groups are expressed as the mean and SEM, where statistical significance was assessed with a one-way or two-way ANOVA, followed by Bonferroni's post hoc analysis (GraphPad Prism v 6.0). The accepted level of statistical significance was  $P < 0.05$ .

# 3. RESULTS

## 3.1. Microglia in the Substantia Nigra Change Morphology in Response to Biodiesel Exposure

Previous reports indicate that microglia in the substantia nigra are activated and change morphology in response to month-long inhalation to petroleum-based diesel exhaust (Levesque, Taetzsch, 2011b). Using a similar histopathological approach, here we revealed that microglia also demonstrate activated morphology in the substantia nigra in response to 100SBDE (Figure 1). Interestingly, the representative images in Figure 1 suggest that higher concentrations of 100SBDE may be required for changes in morphology to occur in the SHR compared to WKY rats. Activated morphology in response to 100SBDE is denoted by larger cell bodies, fewer processes, and thicker remaining processes (Figure 1). Thus, microglia in the substantia nigra are clearly able to detect and respond to month-long biodiesel exhaust exposure.

## 3.2. Biodiesel Exposure Fails to Consistently Impact Markers of Traditional M1/M2 Activation in the Brain

Microglial activation and healthy neuroinflammation typically exists on a spectrum of pro-inflammatory (M1) or alternative (M2) responses (Boche et al. , 2013, Durafourt et al. , 2012, Jang et al. , 2013). What is frequently characterized as M1 activation is associated with the upregulation of pro-inflammatory mediators (ex. TNF $\alpha$  and IL-6), the production of reactive oxygen species (ROS, ex. H<sub>2</sub>O<sub>2</sub> and ONOO<sup>-</sup>) (Block, Zecca, 2007), and initiating signaling pathways associated with neuroinflammation (ex. p38 MAP Kinase phosphorylation) (Bachstetter and Van Eldik, 2010). In healthy activation responses, the initial M1 response is typically followed by a secondary M2 activation that is important for wound healing and resolving inflammation, which is marked by the expression of factors such as Arginase1 (ARG-1) (Boche, Perry, 2013, Durafourt, Moore, 2012, Mills, 2012). In an effort to discern how microglia were responding to the 100SBDE exposure and to characterize the brain pro-inflammatory milieu, several M1/M2 markers were assessed. In contrast to our previous reports of generalized whole brain neuroinflammation with petroleum-based diesel exhaust exposure (Levesque, Taetzsch, 2011b), 100SBDE exposure resulted in no significant changes in p38 MAP Kinase phosphorylation (Table S3,  $P > 0.05$ )

or TNF $\alpha$  protein (Table S4, P>0.05) in brain homogenates, regardless of genotype. Exposure to 100SBDE did elevate brain nitrotyrosine levels only in WKY rats, but this was not a dose-dependent response (Table S2 P<0.05). Analysis of mRNA from the striatum revealed no consistent dose response for TNF $\alpha$  (Table S5, P>0.05), IL-6 (Table S6, P>0.05), or ARG-1 (Table S7, P>0.05) gene expression due to 100SBDE exposure, regardless of genotype. While IBA-1 mRNA showed no significant gene expression changes in the striatum (Table S8, P>0.05), the microglia-specific receptor CX3CR1 showed significant downregulation, regardless of genotype (Figure 2, P<0.05). Interestingly, there was a genotype by environment interaction in striatum fractalkine mRNA levels with 100SBDE exposure, where SHR rats appear to be less sensitive to 100SBDE-induced reduction of fractalkine expression (Figure 3, P<0.05). Thus, the data indicate the microglial activation phenotype associated with the changes in morphology is independent of genotype and is atypical, where neuroinflammation is absent and there is no discernable effect on the traditional M1/M2 markers measured. However, unique to this activation phenotype, it appears that CX3CR1 may decrease in microglia in the striatum without affecting other myeloid specific markers. In addition, fractalkine expression (the ligand for CX3CR1) is decreased in the striatum in both strains, where the WKY rats are the most sensitive to these effects, suggesting a potential impairment of fractalkine signaling with biodiesel exposure.

#### 4. DISCUSSION

As sentinels, microglia detect and respond to diverse stimuli that reach the brain, which includes responding to several sources of urban air pollution. Our previous reports document a pro-inflammatory microglial response linked to generalized neuroinflammation in response to diesel exhaust from traditional petroleum-based fuel (Levesque, Taetzsch, 2011b). Here, we demonstrate that microglia have a unique response to diesel exhaust exposure from a different source. More specifically, our current findings indicate that microglia detect biodiesel exhaust and change morphology in the substantia nigra (Figure 1) without consistent evidence of generalized neuroinflammation, oxidative stress, nor changes in M1 or M2 marker gene expression (Tables S2–8). Importantly, there was also no evidence to support changes in microglia number, which is supported by a lack of any effect of biodiesel exhaust on striatum IBA-1 gene expression (Table S8). However, CX3CR1 was downregulated in all strains, to further differentiate this atypical activation phenotype (Figure 2).

Together, these findings indicate that the CNS response to 100SBDE breaks from the prototypical pro-inflammatory response we have observed with exposure to urban pollution from several sources. With regards to the microglial activation phenotype, it is important to note that the presence of M1 or M2 markers may reveal little about the beneficial or deleterious nature of microglial function when assessed without knowledge of the appropriate kinetic response (Taetzsch et al. , 2015), and the healthy vs. pathological microglial response to urban air pollution is poorly understood. Notably, fractalkine expression was reduced in the striatum of both the SHR and WKY rats, where it took less biodiesel to cause this effect in WKY rats (Figure 3). As such, this suggests biodiesel may perturb the fractalkine axis in the striatum. At present, it is premature to characterize these CNS responses to biodiesel as toxic or benign, as the exposure was short term and the



analyses were not all inclusive. To begin to address this issue, further inquiry will require assessment of chronic effects, including those occurring across a lifetime, which is more similar to human exposure to urban air pollution.

The reduction of CX3CR1 in the brain (Figure 2) without the modification of other myeloid cell markers, like IBA-1 (Table S8), implicates cellular downregulation of this key receptor in microglia. CX3CR1 is a myeloid specific receptor involved in microglial chemotaxis, general microglial function, and is the receptor for fractalkine, a factor produced by neurons for neuron-microglia communication in health and injury (Limatola and Ransohoff, 2014). The biodiesel-induced CNS reduction of CX3CR1 paired with pronounced changes in microglia morphology (Figure 1) were initially surprising, as deficits in CX3CR1 have been previously associated with early impaired changes in microglia morphology in response to an activator/stressor (Hellwig et al. , 2015). However, the consequences of CX3CR1 reduction for CNS function and health is unclear due to conflicting reports in the literature. For instance reduced CX3CR1 has been associated with disease such as schizophrenia (Bergon et al. , 2015) and impairment of cognitive function (Rogers et al. , 2011) as well as neuroprotection (Merino et al. , 2015), increased beneficial microglial phagocytic activity (Merino, Muneton-Gomez, 2015), and other protective effects from reducing the deleterious microglial response to fractalkine (Merino, Muneton-Gomez, 2015). However, recent findings indicate that downregulation of CX3CR1 may have a more ominous consequence in chronic damage, as has been reported for neurotrauma (Zanier et al. , 2015). As such, ongoing research will be focused on understanding the role of fractalkine and CX3CR1 on the effects of air pollution in the brain.

In summary, these data reveal an atypical activated microglial response to biodiesel exhaust, where hypertensive rats appear to be less sensitive to these effects. At present, it is unclear whether this microglial/CNS response is benign, beneficial, or deleterious. The underlying mechanisms and the consequences of this unusual response remains of significant scientific interest in the ongoing effort to understand the potential impact of urban air pollution on CNS health.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## ACKNOWLEDGEMENTS

This research was supported by the National Institute of Environmental Health Sciences/the National Institute of Health ES01695. We sincerely appreciate the assistance of Jesus Ramirez in the sectioning and TH-IR staining of the brain samples. We are also grateful to Mr. Charley King and Mr. Daniel Janek (Arcadis G&M Inc., Highlands Ranch, CO) for their help with the operation of the engine and exposure facilities, and to Dr. Ian Gilmour, and Dr. Seung-Hyun Cho, Dr. William Linak and Mr. Todd Krantz (USEPA) for their help with the diesel exposures and its characterization. The authors thank Dr. John Hong of NIEHS, Research Triangle Park, NC and Dr. William K. Boyes of NHEERL of USEPA, Research Triangle Park, NC for their helpful comments on an earlier version of this manuscript. The research described in this article has been reviewed by the National Health and Environmental Effects Research Laboratory, US Environmental Protection Agency and approved for publication. Approval does not signify that the contents necessarily reflect the views and the policies of the Agency nor does mention of trade names or commercial products constitute endorsement or recommendation for use.


## REFERENCES

- Abramoff M, Magelhaes PJ, Ram S. Image processing with ImageJ. *Biophotonics Int.* 2004;11:36–42.
- Allen JL, Liu X, Weston D, Prince L, Oberdorster G, Finkelstein JN, et al. Developmental exposure to concentrated ambient ultrafine particulate matter air pollution in mice results in persistent and sex-dependent behavioral neurotoxicity and glial activation. *Toxicol Sci.* 2014;140:160–78. [PubMed: 24690596]
- Antonini JM, Sriram K, Benkovic SA, Roberts JR, Stone S, Chen BT, et al. Mild steel welding fume causes manganese accumulation and subtle neuroinflammatory changes but not overt neuronal damage in discrete brain regions of rats after short-term inhalation exposure. *Neurotoxicology.* 2009;30:915–25. [PubMed: 19782702]
- Bachstetter AD, Van Eldik LJ. The p38 MAP Kinase Family as Regulators of Proinflammatory Cytokine Production in Degenerative Diseases of the CNS. *Aging Dis.* 2010;1:199–211. [PubMed: 22720195]
- Bass VL, Schladweiler MC, Nyska A, Thomas RF, Miller DB, Krantz T, et al. Comparative cardiopulmonary toxicity of exhausts from soy-based biofuels and diesel in healthy and hypertensive rats. *Inhal Toxicol.* 2015;27:545–56. [PubMed: 26514782]
- Becerra TA, Wilhelm M, Olsen J, Cockburn M, Ritz B. Ambient air pollution and autism in Los Angeles county, California. *Environ Health Perspect.* 2013;121:380–6. [PubMed: 23249813]
- Bedada GB, Smith CJ, Tyrrell PJ, Hirst AA, Agius R. Short-term effects of ambient particulates and gaseous pollutants on the incidence of transient ischaemic attack and minor stroke: a case-crossover study. *Environ Health.* 2012;11:77. [PubMed: 23067103]
- Bergon A, Belzeaux R, Comte M, Pelletier F, Herve M, Gardiner EJ, et al. CX3CR1 is dysregulated in blood and brain from schizophrenia patients. *Schizophr Res.* 2015;168:434–43. [PubMed: 26285829]
- Block ML, Calderon-Garciduenas L. Air pollution: mechanisms of neuroinflammation and CNS disease. *Trends Neurosci.* 2009;32:506–16. [PubMed: 19716187]
- Block ML, Wu X, Pei Z, Li G, Wang T, Qin L, et al. Nanometer size diesel exhaust particles are selectively toxic to dopaminergic neurons: the role of microglia, phagocytosis, and NADPH oxidase. *FASEB J.* 2004;18:1618–20. [PubMed: 15319363]
- Block ML, Zecca L, Hong JS. Microglia-mediated neurotoxicity: uncovering the molecular mechanisms. *Nat Rev Neurosci.* 2007;8:57–69. [PubMed: 17180163]
- Boche D, Perry VH, Nicoll JA. Review: activation patterns of microglia and their identification in the human brain. *Neuropathol Appl Neurobiol.* 2013;39:3–18. [PubMed: 23252647]
- Bolton JL, Smith SH, Huff NC, Gilmour MI, Foster WM, Auten RL, et al. Prenatal air pollution exposure induces neuroinflammation and predisposes offspring to weight gain in adulthood in a sex-specific manner. *FASEB J.* 2012;26:4743–54. [PubMed: 22815382]
- Calderon-Garciduenas L, D'Angiulli A, Kulesza RJ, Torres-Jardon R, Osnaya N, Romero L, et al. Air pollution is associated with brainstem auditory nuclei pathology and delayed brainstem auditory evoked potentials. *Int J Dev Neurosci.* 2011;29:365–75. [PubMed: 21458557]
- Calderon-Garciduenas L, Franco-Lira M, Mora-Tiscareno A, Medina-Cortina H, Torres-Jardon R, Kavanaugh M. Early Alzheimer's and Parkinson's disease pathology in urban children: Friend versus Foe responses--it is time to face the evidence. *Biomed Res Int.* 2013;2013:161687. [PubMed: 23509683]
- Calderon-Garciduenas L, Kavanaugh M, Block M, D'Angiulli A, Delgado-Chavez R, Torres-Jardon R, et al. Neuroinflammation, hyperphosphorylated tau, diffuse amyloid plaques, and down-regulation of the cellular prion protein in air pollution exposed children and young adults. *J Alzheimers Dis.* 2012;28:93–107. [PubMed: 21955814]
- Calderon-Garciduenas L, Reed W, Maronpot RR, Henriquez-Roldan C, Delgado-Chavez R, Calderon-Garciduenas A, et al. Brain inflammation and Alzheimer's-like pathology in individuals exposed to severe air pollution. *Toxicologic pathology.* 2004;32:650–8. [PubMed: 15513908]
- Calderon-Garciduenas L, Solt AC, Henriquez-Roldan C, Torres-Jardon R, Nuse B, Herritt L, et al. Long-term air pollution exposure is associated with neuroinflammation, an altered innate immune response, disruption of the blood-brain barrier, ultrafine particulate deposition, and accumulation

- of amyloid beta-42 and alpha-synuclein in children and young adults. *Toxicol Pathol.* 2008;36:289–310. [PubMed: 18349428]
- Campbell A, Araujo JA, Li H, Sioutas C, Kleinman M. Particulate matter induced enhancement of inflammatory markers in the brains of apolipoprotein E knockout mice. *J Nanosci Nanotechnol.* 2009;9:5099–104. [PubMed: 19928188]
- Campbell A, Daher N, Solaimani P, Mendoza K, Sioutas C. Human brain derived cells respond in a type-specific manner after exposure to urban particulate matter (PM). *Toxicol In Vitro.* 2014;28:1290–5. [PubMed: 24999231]
- Campbell A, Oldham M, Becaria A, Bondy SC, Meacher D, Sioutas C, et al. Particulate matter in polluted air may increase biomarkers of inflammation in mouse brain. *Neurotoxicology.* 2005;26:133–40. [PubMed: 15527881]
- Costa LG, Cole TB, Coburn J, Chang YC, Dao K, Roque PJ. Neurotoxicity of traffic-related air pollution. *Neurotoxicology.* 2015.
- Durafourt BA, Moore CS, Zammit DA, Johnson TA, Zaguia F, Guiot MC, et al. Comparison of polarization properties of human adult microglia and blood-derived macrophages. *Glia.* 2012;60:717–27. [PubMed: 22290798]
- Elder A, Gelein R, Silva V, Feikert T, Opanashuk L, Carter J, et al. Translocation of inhaled ultrafine manganese oxide particles to the central nervous system. *Environ Health Perspect.* 2006;114:1172–8. [PubMed: 16882521]
- Gerlofs-Nijland ME, van Berlo D, Cassee FR, Schins RP, Wang K, Campbell A. Effect of prolonged exposure to diesel engine exhaust on proinflammatory markers in different regions of the rat brain. *Part Fibre Toxicol.* 2010;7:12. [PubMed: 20478040]
- Hellwig S, Brioschi S, Dieni S, Frings L, Masuch A, Blank T, et al. Altered microglia morphology and higher resilience to stress-induced depression-like behavior in CX3CR1-deficient mice. *Brain Behav Immun.* 2015.
- Heneka MT, Kummer MP, Latz E. Innate immune activation in neurodegenerative disease. *Nature reviews Immunology.* 2014;14:463–77.
- Heppner FL, Ransohoff RM, Becher B. Immune attack: the role of inflammation in Alzheimer disease. *Nat Rev Neurosci.* 2015;16:358–72. [PubMed: 25991443]
- Hesterberg TW, Long CM, Lapin CA, Hamade AK, Valberg PA. Diesel exhaust particulate (DEP) and nanoparticle exposures: what do DEP human clinical studies tell us about potential human health hazards of nanoparticles? *Inhal Toxicol.* 2010;22:679–94. [PubMed: 20462394]
- Hickman SE, Kingery ND, Ohsumi TK, Borowsky ML, Wang LC, Means TK, et al. The microglial sensome revealed by direct RNA sequencing. *Nature neuroscience.* 2013;16:1896–905. [PubMed: 24162652]
- Jang E, Lee S, Kim JH, Seo JW, Lee WH, Mori K, et al. Secreted protein lipocalin-2 promotes microglial M1 polarization. *FASEB J.* 2013;27:1176–90. [PubMed: 23207546]
- Jung CR, Lin YT, Hwang BF. Ozone, particulate matter, and newly diagnosed Alzheimer's disease: a population-based cohort study in taiwan. *J Alzheimers Dis.* 2015;44:573–84. [PubMed: 25310992]
- Kioumourtzoglou MA, Schwartz JD, Weisskopf MG, Melly SJ, Wang Y, Dominici F, et al. Long-term PM Exposure and Neurological Hospital Admissions in the Northeastern United States. *Environ Health Perspect.* 2015.
- Kirrane EF, Bowman C, Davis JA, Hoppin JA, Blair A, Chen H, et al. Associations of Ozone and PM2.5 Concentrations With Parkinson's Disease Among Participants in the Agricultural Health Study. *J Occup Environ Med.* 2015;57:509–17. [PubMed: 25951420]
- Larcombe AN, Kicic A, Mullins BJ, Knothe G. Biodiesel exhaust: the need for a systematic approach to health effects research. *Respirology.* 2015;20:1034–45. [PubMed: 26179557]
- Levesque S, Surace MJ, McDonald J, Block ML. Air pollution and the brain: Subchronic diesel exhaust exposure causes neuroinflammation and elevates early markers of neurodegenerative disease. *J Neuroinflammation.* 2011a;8:105. [PubMed: 21864400]
- Levesque S, Taetzsch T, Lull ME, Kodavanti U, Stadler K, Wagner A, et al. Diesel exhaust activates and primes microglia: air pollution, neuroinflammation, and regulation of dopaminergic neurotoxicity. *Environ Health Perspect.* 2011b;119:1149–55. [PubMed: 21561831]


- Limatola C, Ransohoff RM. Modulating neurotoxicity through CX3CL1/CX3CR1 signaling. *Front Cell Neurosci.* 2014;8:229. [PubMed: 25152714]
- Ma JY, Ma JK. The dual effect of the particulate and organic components of diesel exhaust particles on the alteration of pulmonary immune/inflammatory responses and metabolic enzymes. *J Environ Sci Health Part C Environ Carcinog Ecotoxicol Rev.* 2002;20:117–47.
- Mauderly JL, Burnett RT, Castillejos M, Ozkaynak H, Samet JM, Stieb DM, et al. Is the air pollution health research community prepared to support a multipollutant air quality management framework? *Inhal Toxicol.* 2010;22 Suppl 1:1–19.
- Mehus AA, Reed RJ, Lee VS, Littau SR, Hu C, Lutz EA, et al. Comparison of Acute Health Effects From Exposures to Diesel and Biodiesel Fuel Emissions. *J Occup Environ Med.* 2015;57:705–12. [PubMed: 26147538]
- Merino JJ, Muneton-Gomez V, Alvarez MI, Toledano-Diaz A. Effects of CX3CR1 and fractalkine chemokines in amyloid beta clearance and p-Tau accumulation in Alzheimer's disease (AD) rodent models: Is fractalkine a systemic biomarker for AD? *Curr Alzheimer Res.* 2015.
- Mills CD. M1 and M2 macrophages: oracles of health and disease. *Crit Rev Immunol.* 2012;32:463–88. [PubMed: 23428224]
- Morgan TE, Davis DA, Iwata N, Tanner JA, Snyder D, Ning Z, et al. Glutamatergic neurons in rodent models respond to nanoscale particulate urban air pollutants in vivo and in vitro. *Environ Health Perspect.* 2011;119:1003–9. [PubMed: 21724521]
- Mutlu E, Nash DG, King C, Krantz TQ, Preston WT, Kooter IM, et al. Generation and characterization of diesel engine combustion emissions from petroleum diesel and soybean biodiesel fuels and application for inhalation exposure studies. *Inhal Toxicol.* 2015;27:515–32. [PubMed: 26514780]
- Power MC, Weisskopf MG, Alexeeff SE, Coull BA, Spiro A, Schwartz J 3rd. Traffic-related air pollution and cognitive function in a cohort of older men. *Environ Health Perspect.* 2011;119:682–7. [PubMed: 21172758]
- Power MC, Weisskopf MG, Alexeeff SE, Wright RO, Coull BA, Spiro A 3rd, et al. Modification by hemochromatosis gene polymorphisms of the association between traffic-related air pollution and cognition in older men: a cohort study. *Environ Health.* 2013;12:16. [PubMed: 23413885]
- Prokopowicz A, Zaciera M, Sobczak A, Bielaczyc P, Woodburn J. The effects of neat biodiesel and biodiesel and HVO blends in diesel fuel on exhaust emissions from a light duty vehicle with a diesel engine. *Environ Sci Technol.* 2015;49:7473–82. [PubMed: 25993509]
- Roberts AL, Lyall K, Hart JE, Laden F, Just AC, Bobb JF, et al. Perinatal air pollutant exposures and autism spectrum disorder in the children of nurses' health study II participants. *Environ Health Perspect.* 2013;121:978–84. [PubMed: 23816781]
- Rogers JT, Morganti JM, Bachstetter AD, Hudson CE, Peters MM, Grimmig BA, et al. CX3CR1 deficiency leads to impairment of hippocampal cognitive function and synaptic plasticity. *J Neurosci.* 2011;31:16241–50. [PubMed: 22072675]
- Santiago-Lopez D, Bautista-Martinez JA, Reyes-Hernandez CI, Aguilar-Martinez M, Rivas-Arancibia S. Oxidative stress, progressive damage in the substantia nigra and plasma dopamine oxidation, in rats chronically exposed to ozone. *Toxicol Lett.* 2010;197:193–200. [PubMed: 20541596]
- Saxena RK, Gilmour MI, Schladweiler MC, McClure M, Hays M, Kodavanti UP. Differential pulmonary retention of diesel exhaust particles in Wistar Kyoto and spontaneously hypertensive rats. *Toxicol Sci.* 2009;111:392–401. [PubMed: 19635756]
- Shojaeefard MH, Etaghni MM, Meisami F, Barari A. Experimental investigation on performance and exhaust emissions of castor oil biodiesel from a diesel engine. *Environ Technol.* 2013;34:2019–26. [PubMed: 24350455]
- Shvedova AA, Yanamala N, Murray AR, Kisin ER, Khaliullin T, Hatfield MK, et al. Oxidative stress, inflammatory biomarkers, and toxicity in mouse lung and liver after inhalation exposure to 100% biodiesel or petroleum diesel emissions. *J Toxicol Environ Health A.* 2013;76:907–21. [PubMed: 24156694]
- Sun Q, Wu W, Hu YC, Li H, Zhang D, Li S, et al. Early release of high-mobility group box 1 (HMGB1) from neurons in experimental subarachnoid hemorrhage in vivo and in vitro. *J Neuroinflammation.* 2014;11:106. [PubMed: 24924349]

- Swanson KJ, Madden MC, Ghio AJ. Biodiesel exhaust: the need for health effects research. *Environ Health Perspect*. 2007;115:496–9. [PubMed: 17450214]
- Taetzsch T, Levesque S, McGraw C, Brookins S, Luqa R, Bonini MG, et al. Redox regulation of NF-kappaB p50 and M1 polarization in microglia. *Glia*. 2015;63:423–40. [PubMed: 25331559]
- Takano T Role of Microglia in Autism: Recent Advances. *Dev Neurosci*. 2015.
- Volk HE, Hertz-Picciotto I, Delwiche L, Lurmann F, McConnell R. Residential proximity to freeways and autism in the CHARGE study. *Environ Health Perspect*. 2011;119:873–7. [PubMed: 21156395]
- Volk HE, Lurmann F, Penfold B, Hertz-Picciotto I, McConnell R. Traffic-related air pollution, particulate matter, and autism. *JAMA Psychiatry*. 2013;70:71–7. [PubMed: 23404082]
- Wellenius GA, Boyle LD, Coull BA, Milberg WP, Gryparis A, Schwartz J, et al. Residential proximity to nearest major roadway and cognitive function in community-dwelling seniors: results from the MOBILIZE Boston Study. *J Am Geriatr Soc*. 2012a;60:2075–80. [PubMed: 23126566]
- Wellenius GA, Burger MR, Coull BA, Schwartz J, Suh HH, Koutrakis P, et al. Ambient air pollution and the risk of acute ischemic stroke. *Arch Intern Med*. 2012b;172:229–34. [PubMed: 22332153]
- Zanier ER, Marchesi F, Ortolano F, Perego C, Arabian M, Zoerle T, et al. Fractalkine Receptor Deficiency Is Associated with Early Protection but Late Worsening of Outcome following Brain Trauma in Mice. *J Neurotrauma*. 2015.




**Figure 1. Changes in Microglia Morphology in Response to Soy Biodiesel Exhaust in SHR and WKY Rats.**

Young adult male Spontaneously Hypertensive rats (SHR) and normotensive Wistar Kyoto (WKY) rats were exposed for 4 h/day for 4 weeks (5 days/week) to 100% Soy Biodiesel Exhaust (100SBDE; 0, 50, 150 and 500  $\mu\text{g}/\text{m}^3$ ). Three coronal sections (40  $\mu\text{m}$ ) from the substantia nigra per animal were stained with the IBA-1 antibody. Representative images of the substantia nigra were taken at 40X, the scale bar depicts 50  $\mu\text{m}$ , and the red arrows depict activated microglia morphology (N=3).



**Figure 2. Striatum Changes in Fractalkine in Response to Soy Biodiesel Exhaust in SHR and WKY Rats.**

Young adult male Spontaneously Hypertensive rats (SHR) and normotensive Wistar Kyoto (WKY) rats were exposed to 100% Soy Biodiesel Exhaust (100SBDE; 0, 50, 150 and 500  $\mu\text{g}/\text{m}^3$ ) for 4 h/day for 4 weeks (5 days/week). Striatum fractalkine mRNA expression was evaluated by quantitative RT-PCR, values were normalized to GAPDH using the  $2^{-\Delta\Delta\text{CT}}$  method. An asterisk indicates significant decrease ( $P < 0.05$ ) from 0.0  $\mu\text{g}/\text{m}^3$  control and an “†” indicates a significant difference between the rat strains ( $P < 0.05$ ,  $n = 3$ ).



**Figure 3. Striatum Changes in Fractalkine Receptor (CX3CR1) in Response to Soy Biodiesel Exhaust in SHR and WKY Rats.**

Young adult male Spontaneously Hypertensive rats (SHR) and normotensive Wistar Kyoto (WKY) rats were exposed to 100% Soy Biodiesel Exhaust (100SBDE; 0, 50, 150 and 500  $\mu\text{g}/\text{m}^3$ ) for 4 h/day for 4 weeks (5 days/week). Striatum CX3CR1 mRNA expression was evaluated by quantitative RT-PCR, values were normalized to GAPDH using the  $2^{-\Delta\Delta\text{CT}}$  method. An asterisk indicates significant decrease ( $P < 0.05$ ) from 0.0  $\mu\text{g}/\text{m}^3$  control in both the **A.**WKY rats and **B.** SHR rats.