



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

22nd International Grassland Congress

Effects of Two Intake Levels of *Leucaena leucocephala* on Rumen Function of Sheep

Marcos Barros-Rodríguez
Universidad Autónoma de Yucatán, México

Javier Solorio-Sánchez
Universidad Autónoma de Yucatán, México

Carlos Sandoval-Castro
Universidad Autónoma de Yucatán, México

Athol V. Klieve
University of Queensland, Australia

Eduardo Briceño-Poot
Universidad Autónoma de Yucatán, México

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1-2/31>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

Marcos Barros-Rodríguez, Javier Solorio-Sánchez, Carlos Sandoval-Castro, Athol V. Klieve, Eduardo Briceño-Poot, Luis Ramírez-Avilés, and Rafael Rojas-Herrera

Effects of two intake levels of *Leucaena leucocephala* on rumen function of sheep

Marcos Barros-Rodríguez^A, Javier Solorio-Sánchez^A, Carlos Sandoval-Castro^A, Athol V Klieve^B, Eduardo Briceño-Poot^A, Luis Ramírez-Avilés^A and Rafael Rojas-Herrera^C

^A Departamento de Nutrición Animal, Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma de Yucatán, Mérida, Yucatán, Carretera Mérida-Xmatkuil km 15.5. Apdo. 4-116 Itzimná, C.P. 97100, Mérida, Yucatán, México

^B School of Animal Studies, University of Queensland, Gatton Campus, Gatton, Queensland, Australia

^C Facultad de Ingeniería Química, Universidad Autónoma de Yucatán, Mexico

Contact email: ma_barrosr@yahoo.es

Keywords: Rumen function, sheep, *Leucaena*, *Pennisetum*.

Introduction

Ruminant production systems based on grass pastures often produce poor animal performance as expressed by growth or reproductive rates. The nutrient imbalance affecting rumen function is due to their low energy and protein intake. The incorporation of leguminous forages such as *Leucaena leucocephala*, in the diet of ruminants, can stimulate rumen function by providing protein-rich forage (Barros-Rodríguez *et al.* 2012). This increases the availability of compounds such as ammonia, amino acids and peptides as well as branched short-chain fatty acids, which are produced as a result of degradation of proteins. These substances promote fibre breakdown by acting as ruminal growth activators for rumen bacteria, especially cellulolytic bacteria (Hoover and Stokes 1991). This study aimed to evaluate the effects of 2 intake levels of *L. leucocephala* on rumen function of sheep fed *Pennisetum purpureum*.

Materials and Methods

The experiment was conducted over 60 days at the Faculty of Veterinary Medicine, University of Yucatán, Mérida, using 18 Pelibuey sheep, cannulated in the rumen and with an average weight of 33.3 ± 6.13 kg. A completely randomised design was used with 3 replicates of the following 3 treatments: T1- 100% grass, T2- 20% leucaena + 80% grass, T3- 40% leucaena + 60% grass.

Dry matter (DM) degradation was determined using the technique described by Ørskov *et al.* (1980), incubating at 4, 8, 12, 24, 36, 48, 72, 84 and 96 hours. Rumen constants of DM were analysed by Graph Pad Prism (GraphPad 2003) that fits the data of the exponential equation $Y = a + b(1 - e^{-ct})$ (Ørskov and McDonald 1979). Concentration of NH₃ and pH were determined on rumen liquid samples, collected 6 hours after feeding. The samples were filtered through cotton gauze and 10 ml of rumen liquor from each sample was stored in 0.5% HCL (1:1) for later analysis of NH₃. The pH was measured immediately after collection with a portable potentiometer (Hanna pHep family).

Apparent digestibility of DM and voluntary intake were estimated directly in metabolic cages. Data were analysed by ANOVA and means were compared by the Tukey test using SAS statistical software (SAS 2002).

Results and discussion

Degradation rate of forage was higher on Treatments T2 and T3 than on the pure grass diet (Table 1).

Voluntary intake and apparent digestibility of DM of the diets containing legume were higher than those of the pure grass diet. Ammonia concentration in rumen liquor was also higher for the diets containing leucaena and rumen pH level declined as the proportion of legume in the diet increased.

Crude protein concentration of grass was apparently a limiting factor for ruminal digestion, restricting microbial function and limiting energy intake as indicated by the low rumen ammonia levels on the pure grass diet. The additional protein provided by the leucaena would have increased availability of ammonia for rumen microflora, stimulating microbial growth and increasing rate of breakdown of the forage, which increased rate of passage of the diet and consequently voluntary intake. Kakengi *et al.* (2001) obtained similar trends with leucaena supplementation in the diet of cattle. The reduction in pH with increasing level of leucaena in the diet indicates increased production of volatile fatty acids from the increased intake and breakdown of the forage. These findings support the results reported by Osakwe and Steingass (2006) that, as leucaena proportion in the diet increases, rumen function improves.

For all parameters evaluated, there were no differences between T2 and T3 ($P > 0.05$). The lack of a major response in T3 can probably be attributed to a crude protein v digestible energy imbalance caused by higher leucaena intake in T3 (data not shown), leading to low efficiency of synthesis of microbial protein in the rumen (Calsamiglia *et al.* 2010). These results are consistent with those reported by Barros-Rodríguez *et al.* (2012).

Table 1. In situ degradation of DM, apparent digestibility of DM, voluntary intake, ruminal pH and ammonia concentration. Means within rows with different superscripts differ significantly ($P < 0.05$). a: soluble fraction (in percentage), b: insoluble but potentially degradable fraction (in percentage), c: degradation rate (in percentage per hour), a + b: potential degradation, r^2 : r square, VI: voluntary intake, LW: live weight, T0: time zero.

DM degradation (%)	Treatments			Sig. level
	T1	T2	T3	
<i>Pennisetum purpureum</i>				
T0	17.68	17.68	17.68	
a	19.02±1.52	19.27±1.27	17.88±1.36	>0.05
b	47.49±1.47	46.59±1.16	47.58±1.24	>0.05
c	0.038±0.003 b	0.050±0.003 a	0.052±0.003a	<0.05
a+b	66.51	65.86	65.46	>0.05
r^2	0.95	0.97	0.97	
<i>Leucaena leucocephala</i>				
T0	34.27	34.27	34.27	>0.05
a	32.53±1.42	31.99±1.54	32.33±1.63	>0.05
b	52.81±1.70	49.17±1.41	49.02±1.49	<0.05
c	0.029±0.003 b	0.061±0.004 a	0.059±0.004 a	>0.05-
a+b	85.34	81.16	81.35	
r^2	0.96	0.97	0.97	
Apparent DM digestibility (%)	52.52±2.17 b	56.5±2.28 a	57.6±2 a	0.002
VI (g/kg LW ^{0.75})	67.58±0.053 b	86.94±6.09 a	83.72±2.31 a	0.0001
pH	6.86±0.03 a	6.62±0.02 b	6.56±0.03 c	0.0001
NH ₃ (mg/l)	25.33±3.78 b	36.06±6.31 a	33.38 ±4.56 a	0.005

Conclusions

Providing supplements of leucaena in the diet of sheep can improve rumen function by increasing the rate of degradation of forages and stimulating voluntary intake. This increases availability of nutrients for metabolic processes resulting in improved growth rates, wool growth and reproductive performance. Our results indicate that providing leucaena at more than 20% of the diet would not produce additional benefits. It would seem preferable to supplement twice as many sheep with this lower level. Further studies may be warranted to determine how effective lower levels of supplementation might be.

Acknowledgments

The senior author acknowledges the Consejo Nacional de Ciencia y Tecnología (CONACyT) of Mexico for support to his living expenses, and MSc Assem Safwat for his help with English editing.

References

Barros-Rodríguez M, Solorio-Sánchez J, Ku-Vera J, Ayala-Burgos A, Sandoval-Castro C, Solís-Pérez G (2012). Productive performance and urinary excretion of mimosine metabolites by hair sheep grazing in a silvopastoral system With high densities of *Leucaena leucocephala*. *Tropical*

Animal Health and Production **44**, 1873-1878.

Calsamiglia S, Ferret A, Reynolds C, Kristensen N, van Vuuren A (2010). Strategies for optimizing nitrogen use by ruminants. *Animal* **4**, 1184-1196.

GraphPad (2003). Graphpad Prism Version 4. 1994-2003. (GraphPad Software, Inc.: San Diego, CA, USA)

Hoover WH, Stokes S (1991). Balancing carbohydrates and proteins for optimum rumen microbial yield. *Journal of Dairy Science* **74**, 3630.

Kakengi A, Shem M, Mtengeti E, Otsyina R (2001). *Leucaena leucocephala* leaf meal as supplement to diet of grazing dairy cattle in semiarid Western Tanzania. *Agroforestry Systems* **52**, 73-82.

Ørskov E, McDonald Y (1979). The estimation of protein degradability in the rumen from determining the digestibility of feeds in the rumen. *Journal of Agricultural Science, Cambridge* **92**, 499-503.

Ørskov E, Hovell F, Mauld F (1980). Uso de la técnica de la bolsa de nylon para la evaluación de los alimentos. (Use of the nylon bag to evaluate feeds). *Tropical Animal Production* **5**, 213-233.

Osakwe I, Steingass H (2006). Ruminant fermentation and nutrient digestion in West African Dwarf (WAD) sheep fed *Leucaena leucocephala* supplemental diets. *Agroforestry Systems* **67**, 129-133.

SAS (SAS Institute Inc.) (2002). SAS/STAT Software, Version 9.00. (SAS Institute Inc.: Cary, NC, USA)