



## Vegetative Propagation of *Stylosanthes scabra*

Alexandre C. L. de Mello  
*Federal Rural University of Pernambuco, Brazil*

Adeneide C. Galdino  
*Federal Rural University of Pernambuco, Brazil*

José C. B. Dubeux Jr.  
*Federal Rural University of Pernambuco, Brazil*

Mércia V. F. dos Santos  
*Federal Rural University of Pernambuco, Brazil*

Mário de A. Lira Jr.  
*Federal Rural University of Pernambuco, Brazil*

*See next page for additional authors*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1-2/32>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

### **Presenter Information**

Alexandre C. L. de Mello, Adeneide C. Galdino, José C. B. Dubeux Jr., Mércia V. F. dos Santos, Mário de A. Lira Jr., Felipe M. Saraiva, Mário de A. Lira, and Márcio V. da Cunha

## Vegetative propagation of *Stylosanthes scabra*

Alexandre C L de Mello<sup>A,B</sup>, Adeneide C Galdino<sup>A</sup>, José Carlos B Dubeux Jr<sup>A,B</sup>, Mércia Virginia F dos Santos<sup>A,B</sup>, Mario de A Lira Jr<sup>A,B</sup>, Felipe M Saraiva<sup>A</sup>, Mario de A Lira<sup>B,C</sup> and Márcio V da Cunha<sup>A,B</sup>

<sup>A</sup> Federal Rural University of Pernambuco, Recife, PE, Brazil

<sup>B</sup> Research Fellow of the National Scientific and Technological Research Council, Brazil

<sup>C</sup> Agronomic Institute of Pernambuco (IPA), Recife, PE, Brazil

Contact email: [mello@dz.ufrpe.br](mailto:mello@dz.ufrpe.br)

**Keywords:** Root development, indolebutyric acid, auxins, hormones, gibberellins, cytokinins.

### Introduction

One way to multiply genetically desirable plants in a more uniform manner is to use vegetative propagation, such as using plant cuttings. This technique multiplies a single plant into several, in an easy low-cost system (Fachinello *et al.* 2005), so it is useful in plant breeding. This technique may be enhanced through the use of hormonal regulators, with indolebutyric acid (IBA) being most frequently used, because of its stability in hydroalcoholic solution, low sensitivity to biological degradation and good establishment of adventitious roots (Fachinello *et al.* 1995). In spite of its large-scale use for fruit production, the technique is not commonly used for forage species, such as *Stylosanthes*. We evaluated different concentrations of IBA for root establishment and initial development of *Stylosanthes scabra* cuttings.


### Materials and Methods

The experiment was conducted at the Federal Rural

University of Pernambuco, Recife, PE. Treatments were 4 IBA concentrations (1, 2, 3 and 4 g/L) applied to *S. scabra* plant cuttings (20 cm long, 2.5 mm diameter), selected from a collection obtained in Bom Jardim, PE. The experiment was fully randomised with 3 replicates, and the experimental unit was a set of 15 x 500 mL plastic recipients, each with one cutting, with a 1:3 sand:vermiculite mixture. The cuttings had flat cuts at the top, and inclined ones at the bottom, and were inserted into the respective IBA solutions to 2.5 cm from the bottom for 10 seconds, and transferred to the experimental units. At the end of the experiment (60 days), percentage of cuttings with roots, root length and root and shoot dry masses were recorded and subjected to regression analysis using ASSISTAT.

### Results and Discussion

All parameters measured responded to IBA concentration according to an exponential relationship (Fig. 1). Highest root length was determined at 421 cm at 1.90 g/L IBA



**Figure 1.** Effects of concentration of indolebutyric acid (IBA) on: (A) root length; (B) percentage of rooted cuttings; (C) shoot dry mass; and (D) root dry mass of *Stylosanthes scabra* cuttings.

concentration (Fig. 1A), which supports the finding of Schuster *et al.* (2011) that root length in *Arachis pintoii* cuttings peaked at IBA concentrations about 2 g/L. Percentage of rooted cuttings was highest at IBA concentrations of 1 and 2 g/L (Fig. 1B), while highest shoot dry mass (1.67 g) occurred at 2.28 g/L IBA (Fig. 1C), and highest root dry mass (246 mg) at 1.71 g/L IBA (Fig. 1D). Pasqual *et al.* (2001) also reported negative effects of higher IBA concentrations on fruit plant cuttings, possibly due to hormonal imbalances among auxins, gibberellins and cytokinins as reported by Cunha *et al.* (2012).

### Conclusion

The 2 g/L IBA concentration seems the most promising for use in stimulating root development on cuttings of *Stylosanthes scabra* in breeding programs. Use of this technique should enhance the success rate in striking cuttings and accelerate the selection process without risks

of cross-pollination contaminating selection lines.

### References

- Cunha CSM, Maia SSS, Coelho MFB (2012) Estaquia de *Croton zehntneri* Pax et Hoffm. com diferentes concentrações de ácido indol butírico. *Ciência Rural* **42**, 621-626.
- Fachinello JC, Hoffmann A, Nachtigal JC (1995) 'Propagação de plantas frutíferas de clima temperado.' (Universidade Federal de Pelotas: Pelotas, RS)
- Fachinello JC, Hoffmann A, Nachtigal JC (2005) 'Propagação de plantas frutíferas.' (Embrapa Informação Tecnológica: Brasília, DF)
- Pasqual M, Chalfun NNJ, Ramos JD (2001) 'Fruticultura Comercial: Propagação de plantas frutíferas.' (Universidade Federal de Lavras/FAEPE: Lavras, MG)
- Schuster MZ, Szymczak LS, Lustosa SBC (2011) Enraizamento de estacas de amendoim forrageiro tratadas com AIB. *Pesquisa Aplicada & Agrotecnologia* **4**, 122-136.