



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

22nd International Grassland Congress

Morphological Divergence among Progeny of *Macroptilium lathyroides* Accessions from the Semi-Arid Region of Pernambuco, Brazil

Mércia V. F. dos Santos
Federal Rural University of Pernambuco, Brazil

Gabriella P. de Albuquerque
Federal Rural University of Pernambuco, Brazil

Márcio V. da Cunha
Federal Rural University of Pernambuco, Brazil

Mário de A. Lira Jr.
Federal Rural University of Pernambuco, Brazil

José C. B. Dubeux Jr.
Federal Rural University of Pernambuco, Brazil

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/1-2/20>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

Mércia V. F. dos Santos, Gabriella P. de Albuquerque, Márcio V. da Cunha, Mário de A. Lira Jr., José C. B. Dubeux Jr., Marta G. S. da Silva, Mário de A. Lira, and Alexandre C. L. de Mello

Morphological divergence among progeny of *Macroptilium lathyroides* accessions from the semi-arid region of Pernambuco, Brazil

Mércia V F dos Santos^A, Gabriella P de Albuquerque^B, Márcio V da Cunha^A, Mario de A Lira Jr^A, José C B Dubeux Jr^A, Marta G S da Silva^C, Mario de A Lira^D and A C L. de Mello^A

^A Federal Rural University of Pernambuco (UFRPE), Recife, PE, Brazil

^B Graduate Program in Animal Science, UFRPE

^C PNPD-Capes scholar

^D Instituto Agronômico de Pernambuco (IPA), Recife, PE, Brazil

Contact email: mercia@dz.ufrpe.br

Keywords: Tropical legumes, population, selection, genetic diversity, heritability.

Introduction

Macroptilium is a legume genus with approximately 20 species, usually annual or biennial, herbaceous and distributed mainly in the Americas. It is widely used as a forage resource in grasslands and usually fixes atmospheric N. Martins *et al.* (2001) indicated that half-sib family selection with progeny testing is the most common plant breeding method used in Brazil. In the scientific literature, however, there are few studies dedicated to *Macroptilium* spp.

This study evaluated morphological divergence among *Macroptilium* spp. progeny from accessions collected in 4 counties located in the semi-arid region of Pernambuco State, NE Brazil.

Methods

The experiment was carried out at the Animal Science Department of the Federal Rural University of Pernambuco (UFRPE), NE Brazil. Accessions of *Macroptilium lathyroides* were collected in Bom Jardim, Caetés, Sertânia and Tupanatinga counties, in the semi-arid region of Pernambuco State, Brazil. Seeds originating from 15 progeny of these accessions were planted and morphological characteristics of the seedlings assessed from October 2012 to January 2013.

Seeds were planted in 20-L pots filled with soil (Ultisol) collected from the 0-20 cm soil layer in Arcoverde-PE. The soil presented the following chemical characteristics: pH (H₂O) = 6.6; Mehlich-I P = 57 mg/dm³; K⁺ = 0.41 cmol_c/dm³; Na⁺ = 0.25 cmol_c/dm³; Ca²⁺ + Mg²⁺ = 3.65 cmol_c/dm³; Ca²⁺ = 2.9 cmol_c/dm³; Al³⁺ = 0.0 cmol_c/dm³; H⁺+Al³⁺ = 2.28 cmol_c/dm³; organic C = 6.11 g/kg; soil organic matter = 10.53 g/kg. No lime was added. Before planting, seeds were scarified with sulphuric acid for 10 min.

A complete randomised design was used with 3 replications. Treatments were different accessions of *Macroptilium lathyroides*. The following response variables were assessed: average plant height (from ground level to the tallest leaf); leaf and leaflet dimensions (length and width; longitudinal and latitudinal measurements using

paquímetro); leaf number per plant; and general aspect of the plant (desirability) using a grading scale ranging from 1 (low) to 5 (high). These evaluations were performed every 45 days. Variance analyses were performed using the software GENES (Cruz 2001). The Scott-Knott test was used (P<0.05) for comparing means.

Results

Only plant height, leaf number and desirability showed any variation between accessions (P<0.05). Four accessions (#2 from Bom Jardim; #12 from Tupanatinga; #14 and #15 from Caetés) showed highest values. In an evaluation of the legume genera *Calopogonium*, *Centrosema*, *Macrotyloma* and *Macroptilium*, Veasey *et al.* (1999) observed genetic variability for morphological characteristics within the *Macroptilium* genus. Garcia *et al.* (2003) assessed the genetic variability of *Macroptilium erythroloma* in Rio Grande do Sul, Brazil, and used a protein polymorphism test to reveal greater similarity among individuals from the same population than between populations, indicating populations with genetic activity.

Progeny presented heritability estimates ranging from 6.47 to 59.6% for leaflet width and plant height, respectively (Table 1). These heritability values are of low to medium magnitude, indicating that these plant responses may be highly affected by the environment.

Conclusions

The variation in plant height, leaf number and desirability among these accessions of *Macroptilium lathyroides* indicates that there is merit in collecting further germplasm of this species and determining if these characteristics can be reflected in increased dry matter production and forage quality.

Acknowledgments

Funding of this research by CNPq is gratefully acknowledged.

References

Cruz CD (2001) Programa Genes: aplicativo computacional em genética e estatística. (Universidade Federal de Viçosa: Viçosa, MG)

Table 1. Morphological traits and their estimated heritability of 15 *Macroptilium lathyroides* accessions collected in the semi-arid region of Pernambuco State, Brazil. Means followed by the same letter within columns do not differ by Scott-Knott test ($P>0.05$).

| Accessions | Plant height (cm) | Leaf Length (cm) | Leaf Width (cm) | Leaflet Length (cm) | Leaflet Width (cm) | Leaf Number | Desirability (1=low, 5=high) |
|-------------------------|-------------------|------------------|-----------------|---------------------|--------------------|-------------|------------------------------|
| 1(Belo Jardim county) | 64.98 b | 5.67 a | 5.02 a | 3.16 a | 2.04 a | 9.85 b | 2.62 b |
| 2 (Belo Jardim county) | 105.83 a | 6.09 a | 5.83 a | 3.47 a | 2.30 a | 19.88 a | 3.94 a |
| 3 (Belo Jardim county) | 66.55 b | 4.88 a | 4.18 a | 2.78 a | 1.92 a | 6.85 b | 2.25 b |
| 4 (Belo Jardim county) | 79.97 a | 5.68 a | 4.94 a | 3.34 a | 2.15 a | 12.36 b | 2.94 b |
| 5 (Belo Jardim county) | 85.44 a | 4.92 a | 4.15 a | 3.00 a | 2.24 a | 10.19 b | 2.94 b |
| 6 (Belo Jardim county) | 81.92 a | 4.87 a | 4.48 a | 2.86 a | 1.76 a | 10.66 b | 3.00 b |
| 7 (Belo Jardim county) | 76.61 b | 4.26 a | 4.33 a | 2.62 a | 1.92 a | 10.07 b | 2.92 b |
| 8 (Belo Jardim county) | 66.01 b | 6.11 a | 5.34 a | 3.42 a | 2.39 a | 9.59 b | 2.95 b |
| 9 (Belo Jardim county) | 84.5 a | 5.79 a | 4.59 a | 3.11 a | 2.11 a | 8.94 b | 2.83 b |
| 10 (Belo Jardim county) | 65.58 b | 5.15 a | 4.87 a | 2.83 a | 2.12 a | 10.94 b | 2.69 b |
| 11 (Belo Jardim county) | 65.92 b | 5.66 a | 5.15 a | 2.87 a | 2.00 a | 11.91 b | 3.03 b |
| 12 (Tupanatinga county) | 97.33 a | 5.86 a | 4.69 a | 3.58 a | 1.92 a | 20.58 a | 3.66 a |
| 13 (Caetés county) | 66.33 b | 3.18 a | 3.05 a | 2.02 a | 1.47 a | 12.41 b | 2.66 b |
| 14 (Caetés county) | 95.00 a | 5.63 a | 5.26 a | 3.22 a | 2.11 a | 16.66 a | 3.50 a |
| 15 (Sertânia county) | 97.25 a | 4.87 a | 5.01 a | 2.96 a | 2.33 a | 17.14 a | 3.55 a |
| Mean | 78.90 | 5.22 | 4.75 | 3.01 | 2.06 | 12.20 | 3.01 |
| CV(%) | 18.23 | 16.36 | 15.87 | 15.36 | 17.36 | 35.47 | 17.28 |
| h^2 | 59.60 | 52.26 | 45.80 | 39.74 | 6.47 | 55.35 | 48.67 |

Garcia AG, Battistin A, Viégas J, Silva JHS (2003) Polimorfismo e similaridade de proteínas totais de seis populações de *Macroptilium erythroloma*, pelo método eletroforético SDS-PAGE. *Bioikos* **17**, 49-56.

Martins IS, Martins RCC, Correia HS (2001) Comparação entre seleção combinada e seleção direta em *Eucalyptus grandis*, sob diferentes intensidades de seleção. *Revista Floresta e Ambiente* **8**, 36-43.

Veasey EA, Werner JC, Colozza MT, Freitas JCT, Lucena MAC, Beisman DA, Gerdes L (1999) Avaliação de caracteres morfológicos, fenológicos e agrônômicos em leguminosas forrageiras tropicais visando a produção de sementes. *Boletim Indústria Animal* **56**, 109-125. <http://www.iz.sp.gov.br/pdfs/bia/1232121592.pdf>. Accessed 15 March 2013.