



Effects of Melatonin Supplementation on Hormonal, Inflammatory, Genetic, and Oxidative Stress Parameters in Women With Polycystic Ovary Syndrome

Mehri Jamilian¹, Fatemeh Foroozanfard², Naghmeh Mirhosseini³, Elham Kavossian², Esmat Aghadavod⁴, Fereshteh Bahmani⁴, Vahidreza Ostadmohammadi⁴, Mersedeh Kia⁵, Tahereh Eftekhar^{6*}, Elnaz Ayati^{6,7}, Mostafa Mahdavinia⁸ and Zatollah Asemi^{4*}

¹ Traditional and Complementary Medicine Research Center, Arak University of Medical Sciences, Arak, Iran,

² Gametogenesis Research Center, Kashan University of Medical Sciences, Kashan, Iran, ³ School of Public Health, University of Saskatchewan, Saskatoon, SK, Canada, ⁴ Research Center for Biochemistry and Nutrition in Metabolic Diseases, Kashan University of Medical Sciences, Kashan, Iran, ⁵ Department of Midwifery, Gorgan Branch, Islamic Azad University, Gorgan, Iran, ⁶ Reproductive Health Research Center, Tehran University of Medical Science, Tehran, Iran, ⁷ Department of Gynecology and Obstetrics, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran, ⁸ Department of Dermatology, Razi Hospital, Tehran University of Medical Sciences, Tehran, Iran

OPEN ACCESS

Edited by:

Claus Yding Andersen,
University of Copenhagen, Denmark

Reviewed by:

Dana Manuela Savulescu,
National Institute of Communicable
Diseases (NICD), South Africa
Alessandro Conforti,
University of Naples Federico II, Italy

*Correspondence:

Tahereh Eftekhar
teftekhaar@gmail.com
Zatollah Asemi
asemi_r@yahoo.com

Specialty section:

This article was submitted to
Reproduction,
a section of the journal
Frontiers in Endocrinology

Received: 22 January 2019

Accepted: 15 April 2019

Published: 14 May 2019

Citation:

Jamilian M, Foroozanfard F, Mirhosseini N, Kavossian E, Aghadavod E, Bahmani F, Ostadmohammadi V, Kia M, Eftekhar T, Ayati E, Mahdavinia M and Asemi Z (2019) Effects of Melatonin Supplementation on Hormonal, Inflammatory, Genetic, and Oxidative Stress Parameters in Women With Polycystic Ovary Syndrome. *Front. Endocrinol.* 10:273. doi: 10.3389/fendo.2019.00273

Purpose: The aim of the current study was to evaluate the effect of melatonin administration on clinical, hormonal, inflammatory, and genetic parameters in women with polycystic ovarian syndrome (PCOS).

Methods: The present randomized, double-blinded, placebo-controlled clinical trial was conducted among 56 patients with PCOS, aged 18–40 years old. Subjects were randomly allocated to take either 5 mg melatonin supplements ($n = 28$) or placebo ($n = 28$) twice a day for 12 weeks.

Results: Melatonin administration significantly reduced hirsutism ($\beta -0.47$; 95% CI, $-0.86, -0.09$; $P = 0.01$), serum total testosterone ($\beta -0.11$ ng/mL; 95% CI, $-0.21, -0.02$; $P = 0.01$), high-sensitivity C-reactive protein (hs-CRP) ($\beta -0.61$ mg/L; 95% CI, $-0.95, -0.26$; $P = 0.001$), and plasma malondialdehyde (MDA) levels ($\beta -0.25$ μ mol/L; 95% CI, $-0.38, -0.11$; $P < 0.001$), and significantly increased plasma total antioxidant capacity (TAC) levels ($\beta 106.07$ mmol/L; 95% CI, 62.87, 149.28; $P < 0.001$) and total glutathione (GSH) ($\beta 81.05$ μ mol/L; 95% CI, 36.08, 126.03; $P = 0.001$) compared with the placebo. Moreover, melatonin supplementation downregulated gene expression of interleukin-1 (IL-1) ($P = 0.03$) and tumor necrosis factor alpha (TNF- α) ($P = 0.01$) compared with the placebo.

Conclusions: Overall, melatonin administration for 12 weeks to women with PCOS significantly reduced hirsutism, total testosterone, hs-CRP, and MDA, while increasing TAC and GSH levels. In addition, melatonin administration reduced gene expression of IL-1 and TNF- α .

Clinical Trial Registration: www.irct.ir, identifier IRCT2017082733941N9, Available online at: <https://www.irct.ir/trial/26051>

Keywords: melatonin, hormonal profiles, inflammatory markers, polycystic ovary syndrome, oxidative stress

INTRODUCTION

Polycystic ovarian syndrome (PCOS) is a highly prevalent endocrinopathy affecting up to 10% of premenopausal women and essentially characterized by hyperandrogenism, reproductive abnormalities, and ovulatory dysfunction (1). Although the exact pathophysiology of PCOS is not yet fully recognized, evidence suggests that hyperandrogenism plays a central role in this syndrome (2, 3). Hyperandrogenism might occur due to the inflammatory response of the abnormal ovarian theca cells to free oxygen radicals (4). Furthermore, prior studies have indicated that increased oxidative stress and elevated inflammatory markers such as C-reactive protein (CRP) in women with PCOS are correlated with obesity, insulin resistance, and an increased risk of cardiovascular disease (CVD) (5, 6). The majority of women with PCOS suffer from hirsutism, acne, oligomenorrhea, and androgenic alopecia, which impair the quality of life of these women (7).

Melatonin (N-acetyl-5-methoxytryptamine) is secreted by the pineal gland (8). The circadian pattern of its secretion is regulated by the biological clock within the hypothalamic suprachiasmatic nucleus (9). Some of the major functions of rhythmic secretion of melatonin in blood include regulation of sleep, modulation of circadian rhythms (10), and involvement in the body's immune response (11). Melatonin is also a powerful endogenous antioxidant (12), and its supplementation has radical-scavenging and anti-inflammatory actions (13, 14). Women with PCOS are often sub-fertile secondary to ovulatory dysfunction, impaired quality of oocyte, and low endometrial receptivity (15). *In vitro* fertilization (IVF) is recommended as an assisted reproduction technology for infertile patients with PCOS when patients do not respond to ovulation induction agents (16). Current evidence suggests the involvement of melatonin in ovarian physiology including ovulation, follicular development, luteal function, and oocyte maturation. Furthermore, melatonin deficiency seems to be involved in the pathophysiology of PCOS (17). It has been shown that melatonin is an efficient predictor of oocyte quality and IVF outcomes: indeed, its high concentration correlates positively with a satisfactory quality of oocytes (18). On the other hand, intra-follicular melatonin level is significantly reduced in patients with PCOS (18). In addition, Jain et al. (19) demonstrated that nocturnal melatonin levels in women with PCOS are higher than in control subjects. Elevated nighttime melatonin levels in PCOS have been explained by a feedback mechanism due to its decreased concentration in the ovarian follicles (20). So, women with PCOS who have high oxidative stress are synthesizing more melatonin, probably in an effort to eliminate extra free radicals (21). High melatonin level in the ovarian follicle fluid is pivotal for ovulation, follicular growth, and quality of oocyte, whereas reduced melatonin level in follicular fluid may be responsible for decreased oocyte quality and anovulation in these women (22). In a recent study, melatonin and myo-inositol co-supplementation significantly improved embryo and oocyte quality in women with PCOS who were candidates for IVF, when compared with myo-inositol alone (23). In addition, treatment with melatonin and myo-inositol significantly improved ovarian stimulation protocols

and pregnancy outcomes in women who failed to conceive in previous IVF cycles because of poor quality of the oocyte (24).

There is promising evidence showing the importance of melatonin supplementation in patients with PCOS who are candidates for IVF. However, whether melatonin has direct benefits on hirsutism, hormonal profiles, and biomarkers of inflammation/oxidative stress in these patients has not been assessed yet. The present study was, therefore, carried out to evaluate the effects of melatonin administration on clinical status, hormonal profiles, oxidative stress biomarkers, inflammatory factors, and gene expression related to inflammation in women diagnosed with PCOS who were candidates for IVF.

MATERIALS AND METHODS

Trial Design and Participants' Characteristics

This randomized, double-blinded, placebo-controlled trial was registered in the Iranian website for registration of clinical trials (<http://www.irct.ir>: IRCT2017082733941N9) and followed the Declaration of Helsinki and Good Clinical Practice guidelines. This trial was conducted among 56 women aged 18–40 years old diagnosed with PCOS, based on the Rotterdam criteria (25), who had been referred to the Kosar Clinic in Arak, Iran, between August and December 2017. The study protocol was approved by the ethics committee of Arak University of Medical Sciences (AUMS). Written informed consent was obtained from all participants prior to the intervention. Exclusion criteria were as follows: pregnancy, breastfeeding, sleeping disorders, adrenal hyperplasia, androgen-secreting tumors, hyperprolactinemia, thyroid dysfunction, and/or diabetes at enrollment.

Supplementation

Initially, patients were randomly divided into two groups to take either melatonin capsules (2×5 mg/day) ($n = 28$) or placebo ($n = 28$) for 12 weeks. Melatonin and the placebo were manufactured by Zahravi Pharmaceutical Company (Tabriz, Iran) and Barij Essence Pharmaceutical Company (Kashan, Iran), respectively. The placebo capsules were matched in color, shape, size, packaging, smell, and taste with the melatonin capsules. Due to a lack of evidence regarding the appropriate dosage of melatonin for patients with PCOS, we used the following dose of melatonin based on a previous published study in diabetic patients with coronary heart disease (CHD) (26). All individuals were taking metformin tablets at an initial dose of 500 mg, which was increased in a stepwise manner during the first 3 weeks to a total of 1,500 mg/day (27). In addition, all individuals were taking 10 mg/day medroxyprogesterone tablet from the 15th until the 25th day of the menstrual cycle. Throughout the study, the consumption of melatonin supplements and placebos was assessed by asking subjects to return the medication containers as well as sending a brief daily cell phone reminder to take the supplements. Patients were instructed to have their regular physical activity and not take any extra nutritional supplements during the 12-week trial. All patients completed a 3-day food record (a weekend and 2 weekdays) at the baseline of the study, week 3, week 6, week 9, and the end of the intervention.

Daily macro- and micronutrient intakes were calculated by analyzing 3-day food records, using nutritionist IV software (First Databank, San Bruno, CA).

Assessment of Outcomes

Malondialdehyde (MDA) change was considered as the primary outcome. Serum total testosterone, sex hormone-binding globulin (SHBG), high-sensitivity C-reactive protein (hs-CRP), plasma nitric oxide (NO), total antioxidant capacity (TAC), total glutathione (GSH), hirsutism, and gene expression related to inflammation were recognized as the secondary outcomes.

Clinical Measures

Hirsutism was assessed using a modified Ferriman–Gallwey (mFG) scoring system as 9 body areas, including the upper lip, chin, chest, upper abdomen, lower abdomen, thighs, back, arm, and buttocks, were investigated for hair, from 0 (no hair) to 4 (frankly virile) (28, 29).

Biochemical Assessment

Fasting blood samples (15 ml) were collected at baseline and the end of the intervention at the Arak reference laboratory. Blood samples were immediately centrifuged (Hettich D-78532, Tuttlingen, Germany) at 3,500 rpm for 10 min to separate serum. Then, the samples were stored at -80°C at the AUMS reference laboratory until further analysis. Whole blood samples were used to check gene expression; serum samples to evaluate total testosterone, SHG, and hs-CRP; and plasma samples to measure NO, TAC, GSH, and MDA. Serum total testosterone and SHBG were measured using commercial validated kits (DiaMetra, Milano, Italy) with inter- and intra-assay coefficient variances (CVs) below 6.5%. Serum hs-CRP concentrations were quantified using a commercial ELISA kit (LDN, Nordhorn, Germany) with inter- and intra-assay CVs below 7%. The plasma NO was measured using the Griess method (30), TAC concentrations using the Benzie and Strain method (31), GSH using the Beutler method (32), and MDA concentrations of thiobarbituric acid-reactive substances using a spectrophotometric test with CVs below 5%. Measurements of hormonal profiles and biomarkers of inflammation and oxidative stress were performed in a blinded fashion, in duplicate, in pairs (pre/post-intervention) at the same time, in the same analytical run, and in random order to reduce systematic error and inter-assay variability.

Isolation of Lymphocyte, RNA Extraction, and cDNA Synthesis

Whole blood samples were collected in anticoagulant EDTA tubes. Lymphocytes were isolated using 50% Percoll solution (Sigma-Aldrich, Dorset, UK) gradient by centrifugation for 20 min at 3,000 rpm at 4°C (33). Total RNA was extracted using the acid guanidinium-phenol-chloroform procedure and the application of RNXTM-plus reagent (Cinnacolon, Tehran, Iran) following manufacturer's instructions. RNAs were treated with DNase I (Fermentas, Lithuania) for the elimination of any genomic DNA contamination. Concentration, integration, and purity of RNA samples were determined by spectrometry and gel electrophoresis. Three micrograms of total RNA

was used for cDNA synthesis with random hexamer and oligo(dT)18 primers through RevertAid™ Reverse Transcriptase (Fermentase, Canada) in a total of 20- μl reaction mixture (33).

Real-Time PCR Analysis

Appropriate primers were designed for interleukin-1 (IL-1), IL-8, tumor necrosis factor- α (TNF- α), transforming growth factor beta (TGF- β), and vascular endothelial growth factor (VEGF), and glyceraldehyde-3 phosphate dehydrogenase as the internal control (housekeeping gene; **Table 1**). Quantitative real-time PCR was performed using LightCycler® 96 sequence detection systems (Roche Diagnostics, Rotkreuz, Switzerland) with the application of 4 μl of 5 \times EVA GREEN I master mix (Salise Biodyne, Japan), 10 ng cDNA, and 200 nM each of forward and reverse primers in a final volume of 20 μl . The PCR was performed through the following instruction: an initial denaturation at 95°C for 10 min, followed by 40 cycles of denaturation at 95°C for 10 s, annealing at $54\text{--}62.1^{\circ}\text{C}$ for 15 s, and extension at 72°C for 30 s. The specificity of PCR products was evaluated by 1.5% agarose gel electrophoresis and melting curve analysis. All experiments were performed at least in triplicate.

Sample Size

To determine sample size, we used a randomized clinical trial (RCT) sample-size formula where type I (α) and type II errors (β) were 0.05 and 0.20 (power = 80%). Based on a previous RCT (26), we used a standard deviation of 0.30 $\mu\text{mol/L}$ and a difference in mean (d) of 0.24 $\mu\text{mol/L}$, considering MDA as the key variable. The calculation indicated that 25 participants were required in each group. Assuming a dropout of five patients per group, the final sample size was determined to be 30 participants per group.


Randomization

Study participants were randomized using computer-generated random numbers. Randomization and allocation were concealed

TABLE 1 | Specific primers used for real-time quantitative PCR.

Gene	Primer	Product size (bp)	Annealing temperature (C)
GAPDH	F: AAGCTCATTCTCTGGTATGACAACG	126	61.3
	R: TCTTCCTCTTGTGCTCTTGTCTGG		
IL-1	F: GCTTCTCTCTGGTCTTGG	174	56
	R: AGGGCAGGGTAGAGAAGAG		
IL-8	F: GCAGAGGGTTGTGGAGAAGT	150	56
	R: ACCCTACAACAGACCCACAC		
TNF- α	F: GTCAACCTCTCTCTGCCAT	188	52
	R: CCAAAGTAGACCTGCCAGAG		
TGF- β	F: TTGAGACTTTTCCGTTGCC	227	56
	R: CGAGGTCTGGGGAAAAGTCT		
VEGF	F: CTTCTGAGTTGCCAGGAGA	216	54
	R: CTCACACACACACACCAGG		

GAPDH, glyceraldehyde-3-phosphate dehydrogenase; *IL-1*, interleukin-1; *IL-8*, interleukin-8; *TNF- α* , tumor necrosis factor alpha; *TGF- β* , transforming growth factor beta; *VEGF*, vascular endothelial growth factor.



from both researchers and patients until the completion of final analyses. The randomized allocation sequence, enrollment of participants, and allocation to interventions were conducted by a trained staff at the clinic. Another person, who was not involved in the trial and not aware of random sequences, assigned the subjects to the numbered bottles of capsules.

Statistical Analyses

The Kolmogorov–Smirnov test was done to determine the normality of data. Differences in dietary intakes and gene expression were detected using independent-sample *t*-tests between treatment groups. Multiple linear regression models were used to assess the treatment effects on study outcomes, after adjusting for confounding parameters including age and BMI. The significance of the treatment effects was presented as the mean differences with 95% confidence interval. *P*-values < 0.05 were considered statistically significant. All statistical analyses were done using the Statistical Package for Social Science version 18 (SPSS Inc., Chicago, Illinois, USA).

RESULTS

During the study period, four participants dropped out of the study due to personal reasons (two participants in each group;

TABLE 2 | General characteristics of study participants^a.

	Placebo group (<i>n</i> = 28)	Melatonin group (<i>n</i> = 28)	<i>P</i> ^b
Age (y)	28.3 ± 2.3	28.7 ± 2.1	0.76
Height (cm)	160.0 ± 2.9	161.0 ± 2.5	0.30
Weight at study baseline (kg)	74.8 ± 8.9	75.1 ± 11.0	0.90
Weight at end of trial (kg)	74.5 ± 8.9	74.7 ± 10.9	0.92
Weight change (kg)	−0.3 ± 0.9	−0.4 ± 1.1	0.79
BMI at study baseline (kg/m ²)	29.2 ± 3.5	29.1 ± 4.6	0.92
BMI at end of trial (kg/m ²)	29.1 ± 3.4	29.0 ± 4.6	0.91
BMI change (kg/m ²)	−0.1 ± 0.3	−0.1 ± 0.4	0.84

^aData are means ± SDs.

^bObtained from independent *t*-test.

Figure 1). Fifty-six participants [placebo (*n* = 28) and melatonin (*n* = 28)] completed the trial.

Anthropometric measures were not statistically different between intervention groups (Table 2).

There was no statistically significant difference in dietary macro- and micro-nutrient intakes between the melatonin and placebo groups (Table 3).

Melatonin administration significantly reduced hirsutism (β −0.47; 95% CI, −0.86, −0.09; *P* = 0.01), serum total testosterone

TABLE 3 | Mean dietary intakes of study participants at baseline, week 3, week 6, and week 9, and the end of the intervention.

	Placebo group (n = 28)	Melatonin group (n = 28)	P ^a
Energy (kcal/d)	2,228 ± 311	2,273 ± 300	0.58
Carbohydrates (g/d)	293.7 ± 70.1	311.7 ± 62.3	0.31
Protein (g/d)	89.1 ± 14.1	85.2 ± 12.2	0.28
Fat (g/d)	80.2 ± 11.1	79.8 ± 16.0	0.90
SFAs (g/d)	25.0 ± 5.7	24.9 ± 5.4	0.96
PUFAs (g/d)	25.4 ± 5.1	24.7 ± 6.6	0.63
MUFAs (g/d)	22.0 ± 5.0	22.4 ± 6.1	0.76
Cholesterol (mg/d)	250.6 ± 117.3	215.7 ± 101.7	0.24
TDF (g/d)	17.2 ± 4.6	18.9 ± 4.9	0.19

Data are means ± SDs.

^aObtained from independent t-test.

MUFAs, monounsaturated fatty acids; PUFAs, polyunsaturated fatty acids; SFAs, saturated fatty acids; TDF, total dietary fiber.

(β -0.11 ng/mL; 95% CI, -0.21 , -0.02 ; $P = 0.01$), hs-CRP (β -0.61 mg/L; 95% CI, -0.95 , -0.26 ; $P = 0.001$), and plasma MDA levels (β -0.25 μ mol/L; 95% CI, -0.38 , -0.11 ; $P < 0.001$), and significantly increased plasma TAC levels (β 106.07 mmol/L; 95% CI, 62.87 , 149.28 ; $P < 0.001$) and GSH (β 81.05 μ mol/L; 95% CI, 36.08 , 126.03 ; $P = 0.001$) compared with the placebo (Table 4). Melatonin supplementation did not affect plasma NO and serum SHBG levels.

Melatonin supplementation downregulated gene expression of IL-1 ($P = 0.03$) and TNF- α ($P = 0.01$) compared with the placebo (Figure 2). Melatonin supplementation did not influence the gene expression of IL-8, TGF- β , and VEGF.

DISCUSSION

In this investigation, we evaluated the impact of melatonin administration for 12 weeks on clinical status, hormonal profiles, oxidative stress biomarkers, inflammatory factors, and gene expression related to inflammation in women with PCOS who were candidates for IVF. Our findings depicted that melatonin administration for 12 weeks to women with PCOS significantly reduced hirsutism, total testosterone, hs-CRP, and MDA, while increasing TAC and GSH levels. In addition, melatonin administration reduced gene expression of IL-1 and TNF- α . Melatonin, in humans, is produced in the pineal gland, cutaneous tissue, ovary, retina, and gastrointestinal system, though its rhythmic synthesis occurs only in the pineal gland (34). Melatonin is detectable in everybody's compartment, such as in the pre-ovulatory follicular fluid, where its level is remarkably higher than that in the serum (35). Although it has been considered that the melatonin in human pre-ovulatory follicular fluid is derived from the body circulation, it may also be produced in the ovary (granulosa and oocyte cells) (36). There is current evidence demonstrating elevated melatonin concentrations in the luteal phase compared with the follicular phase of the human menstrual cycle (37). Melatonin directly stimulates the production of progesterone by granulosa cells or

luteal cells and might act at the ovary level to regulate luteal function (18).

Effects on Clinical Status and Hormonal Profiles

We found that melatonin consumption for 12 weeks in women with PCOS significantly decreased hirsutism and total testosterone levels yet did not influence SHBG values. Hyperandrogenism, the hallmark of PCOS, is a consequence of excessive androgen production by the abnormal ovarian theca cells and the augmented activity and expression of various enzymes involved in the production of steroids (2, 38). Furthermore, adrenal glands contribute to hyperandrogenism, as evidenced by raised levels of dehydroepiandrosterone sulfate (DHEAS) (39). Existing evidence suggests that decreased androgen levels are associated with an improvement in ovulatory functions, reducing hirsutism, and enhancing health-related quality of life (40, 41). It has been shown that the ovary (oocyte and granulosa cells) can produce melatonin and that melatonin receptors are detected in the membrane of human luteal and granulosa cells (20). Women with PCOS have significantly elevated melatonin concentrations and hyperandrogenemia along with an elevated number of atretic follicles (19). After 6 months of supplementation, melatonin significantly improved menstrual irregularities and biochemical hyperandrogenism in women with PCOS through a direct, insulin-independent effect on the ovary (42). In another study, melatonin administration to a PCOS mouse model potentially induced oocyte nuclear maturation and guaranteed fertilization potential (43). Melatonin is favored toward the treatment of PCOS via its influence on the production of steroids, thereby modulating ovulation and reducing insulin resistance and dyslipidemia (44). Overall, there are different parameters, including participants' characteristics, pancreatic beta-cell function, higher dose of melatonin, or longer period of intervention, that can provide appropriate circulating levels of melatonin necessary for improving SHBG levels. However, we were not able to assess circulating levels of melatonin in the present study. Perhaps if the dose or duration of melatonin supplementation had been increased, the results of SHBG would have been significant.

Effects on Biomarkers of Inflammation and Oxidative Stress

Our findings showed that melatonin intake after 12 weeks in women with PCOS significantly decreased serum hs-CRP and plasma MDA, and significantly increased plasma GSH and TAC levels. Moreover, melatonin consumption downregulated IL-1 and TNF- α gene expression in PBMCs of women with PCOS, but did not affect gene expression of IL-8, TGF- β , and VEGF. Previous studies have shown a promising effect of melatonin on inflammation and oxidative stress. Mistraretti et al. (45) demonstrated that 6 mg/day melatonin administration to critically ill patients significantly ameliorated TAC levels. Melatonin intake at a dosage of 5 mg/day for 8 weeks among people with metabolic syndrome reduced MDA levels and enhanced catalase activity (46). Recently, we depicted in another study that the administration of 10 mg/day melatonin for 12 weeks to diabetic people with ischemic

TABLE 4 | Metabolic profiles at baseline and after the 12-week intervention in women with polycystic ovary syndrome who received either melatonin supplements or placebo^a.


Variables	Placebo group (n = 28)		Melatonin group (n = 28)		Difference in outcome measures between melatonin and placebo groups ^a	
	Baseline	Week 12	Baseline	Week 12	β (95% CI)	p^b
Total testosterone (ng/mL)	1.1 ± 0.3	1.1 ± 0.4	1.0 ± 0.2	0.9 ± 0.2	-0.11 (-0.21, -0.02)	0.01
SHBG (nmol/L)	39.3 ± 8.4	39.2 ± 8.3	37.7 ± 7.7	38.9 ± 8.0	1.24 (-0.31, 2.80)	0.11
mFG scores	11.5 ± 2.7	11.2 ± 2.6	12.0 ± 2.4	11.2 ± 2.3	-0.47 (-0.86, -0.09)	0.01
hs-CRP (mg/L)	4.1 ± 2.1	4.3 ± 1.9	4.3 ± 1.7	3.9 ± 1.8	-0.61 (-0.95, -0.26)	0.001
NO (μ mol/L)	29.9 ± 2.9	29.5 ± 2.8	28.1 ± 2.5	28.3 ± 2.6	0.26 (-0.73, 1.26)	0.59
TAC (mmol/L)	1,011.2 ± 92.4	1,000.7 ± 100.8	1,030.0 ± 85.0	1,119.3 ± 102.7	106.07 (62.87, 149.28)	< 0.001
GSH (μ mol/L)	525.7 ± 62.6	533.4 ± 63.2	568.8 ± 133.5	655.5 ± 167.9	81.05 (36.08, 126.03)	0.001
MDA (μ mol/L)	2.6 ± 0.4	2.7 ± 0.5	2.6 ± 0.5	2.5 ± 0.5	-0.25 (-0.38, -0.11)	< 0.001

Data are means \pm SDs.

^a"Outcome measures" refers to the change in values of measures of interest between baseline and week 12. β [difference in the mean outcome measures between treatment groups (melatonin group = 1 and placebo group = 0)].

^bObtained from multiple-regression model (adjusted for baseline values of each biochemical variables, age, and baseline BMI).

GSH, total glutathione; hs-CRP, high-sensitivity C-reactive protein; mFG, modified Ferriman-Gallwey; MDA, malondialdehyde; NO, nitric oxide; SHBG, sex hormone-binding globulin; TAC, total antioxidant capacity.



heart disease had favorable impacts on inflammatory status and oxidative stress parameters (26). Pakravan et al. (47) also found that supplementation with 10 mg/day melatonin significantly decreased CRP levels in people with non-alcoholic fatty liver disease. In a recent meta-analysis conducted by Akbari et al. (48), melatonin supplementation decreased systemic inflammatory markers in patients with metabolic diseases. However, melatonin consumption (6 mg/day) in obese women for 40 days did not influence hs-CRP, TAC, and MDA levels (49). The varieties in study design, population characteristics, dosage of melatonin taken, and duration of intervention might explain the discrepancies among the results of published trials. Insulin resistance might be correlated with a specific ultrasound pattern in patients with PCOS (50). In addition, increased oxidative stress and inflammatory status are correlated with an elevated risk of CVD, diabetes mellitus, hyperandrogenism, and insulin resistance in PCOS (6, 51). Both oxidative stress and PCOS have also been involved in several aspects of female reproduction (52). Melatonin

can attenuate inflammatory status and oxidative damage *via* modulating the function of antioxidative enzymes, inhibiting nuclear factor-kappaB activation, enhancing glutathione synthesis, and directly scavenging free radicals in the inflamed tissues (53, 54).

This study had a few limitations. Due to a funding shortage, we were not able to measure serum melatonin and/or urinary 6-sulfatoxymelatonin (its main excretory metabolite) to assess precisely the compliance of study participants. This limitation has been considered in the interpretation of our findings. In addition, we were unable to determine the impacts of melatonin administration on other biomarkers of oxidative stress and inflammation. Also, we did not evaluate biomarkers of inflammation and oxidative stress in the ovary. Measurement of hirsutism as a clinical finding in the current study is undervalued; however, we did not evaluate menstrual cyclicity before and after the treatment, which is one of the most pivotal outcomes for the health and reproduction of women with PCOS.

Overall, this RCT suggests that melatonin administration for 12 weeks to women with PCOS might reduce hirsutism, total testosterone, hs-CRP, and MDA, while increasing TAC and GSH levels. In addition, melatonin administration might reduce gene expression of IL-1 and TNF- α .

ETHICS STATEMENT

This randomized double-blinded, placebo-controlled trial registered in the Iranian website for registration of clinical trials (<http://www.irct.ir: IRCT2017082733941N9>) and followed the Declaration of Helsinki and Good Clinical Practice guidelines. The study protocol was approved by the Ethics Committee of Arak University of Medical Sciences (AUMS). Written informed consent was obtained from all participants prior to the intervention.

REFERENCES

- Baptiste CG, Battista MC, Trottier A, Baillargeon JP. Insulin and hyperandrogenism in women with polycystic ovary syndrome. *J Steroid Biochem Mol Biol.* (2010) 122:42–52. doi: 10.1016/j.jsbmb.2009.12.010
- Nelson VL, Legro RS, Strauss JF, McAllister JM. Augmented androgen production is a stable steroidogenic phenotype of propagated theca cells from polycystic ovaries. *Mol Endocrinol.* (1999) 13:946–57. doi: 10.1210/mend.13.6.0311
- Gilling-Smith C, Story H, Rogers V, Franks S. Evidence for a primary abnormality of thecal cell steroidogenesis in the polycystic ovary syndrome. *Clin Endocrinol.* (1997) 47:93–9. doi: 10.1046/j.1365-2265.1997.2321049.x
- Gonzalez F, Rote NS, Minium J, Kirwan JP. Reactive oxygen species-induced oxidative stress in the development of insulin resistance and hyperandrogenism in polycystic ovary syndrome. *J Clin Endocrinol Metab.* (2006) 91:336–40. doi: 10.1210/jc.2005-1696
- Gonzalez F. Inflammation in polycystic ovary syndrome: underpinning of insulin resistance and ovarian dysfunction. *Steroids.* (2012) 77:300–5. doi: 10.1016/j.steroids.2011.12.003
- Boots CE, Jungheim ES. Inflammation and human ovarian follicular dynamics. *Semin Reprod Med.* (2015) 33:270–5. doi: 10.1055/s-0035-1554928
- van Zuuren EJ, Fedorowicz Z. Interventions for hirsutism excluding laser and therapy alone: abridged Cochrane systematic review including GRADE assessments. *Br J Dermatol.* (2016) 175:45–61. doi: 10.1111/bjd.14486
- Reiter RJ, Mayo JC, Tan DX, Sainz RM, Alatorre-Jimenez M, Qin L. Melatonin as an antioxidant: under promises but over delivers. *J Pineal Res.* (2016) 61:253–78. doi: 10.1111/jpi.12360
- Klein DC, Moore RY. Pineal N-acetyltransferase and hydroxyindole-O-methyl-transferase: control by the retinohypothalamic tract and the suprachiasmatic nucleus. *Brain Res.* (1979) 174:245–62. doi: 10.1016/0006-8993(79)90848-5
- Cajochen C, Kräuchi K, Wirz-Justice A. Role of melatonin in the regulation of human circadian rhythms and sleep. *J Neuroendocrinol.* (2003) 15:432–7. doi: 10.1046/j.1365-2826.2003.00989.x
- Maestroni GJ, Conti A, Pierpaoli W. Role of the pineal gland in immunity: circadian synthesis and release of melatonin modulates the antibody response and antagonizes the immunosuppressive effect of corticosterone. *J Neuroimmunol.* (1986) 13:19–30. doi: 10.1016/0165-5728(86)90047-0
- Tan D-X, Manchester LC, Esteban-Zubero E, Zhou Z, Reiter RJ. Melatonin as a potent and inducible endogenous antioxidant: synthesis and metabolism. *Molecules.* (2015) 20:18886–906. doi: 10.3390/molecules201018886
- Tan D, Reiter RJ, Manchester LC, Yan M, El-Sawi M, Sainz RM, et al. Chemical and physical properties and potential mechanisms: melatonin as a broad spectrum antioxidant and free radical scavenger. *Curr Top Med Chem.* (2002) 2:181–97. doi: 10.2174/1568026023394443

AUTHOR CONTRIBUTIONS

ZA contributed in conception, design, statistical analysis, and drafting of the manuscript. MJ, FF, EK, EsA, FB, VO, ElA, and MM contributed in data collection and manuscript drafting. All authors approved the final version for submission. ZA supervised the study. NM, MK, and TE have substantially contributed to the revised version, including checking all statistical analyses. All authors approved the final version for resubmission and change in authorship.

ACKNOWLEDGMENTS

The authors would like to thank the staff of Kosar Clinic (Arak, Iran) for their assistance in this project (grant no.1396.89).

- Esposito E, Cuzzocrea S. Antiinflammatory activity of melatonin in central nervous system. *Curr Neuropharmacol.* (2010) 8:228–42. doi: 10.2174/157015910792246155
- Ecklund LC, Usadi RS. Endocrine and reproductive effects of polycystic ovarian syndrome. *Obstet Gynecol Clin North Am.* (2015) 42:55–65. doi: 10.1016/j.ogc.2014.09.003
- Thessaloniki ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Consensus on infertility treatment related to polycystic ovary syndrome. *Hum Reprod.* (2008) 23:462–77. doi: 10.1093/humrep/dem426
- Cagnacci A, Volpe A. A role for melatonin in PCOS? *Fertil Steril.* (2002) 77:1089. author reply:90. doi: 10.1016/S0015-0282(02)03070-4
- Tamura H, Nakamura Y, Korkmaz A, Manchester LC, Tan DX, Sugino N, et al. Melatonin and the ovary: physiological and pathophysiological implications. *Fertil Steril.* (2009) 92:328–43. doi: 10.1016/j.fertnstert.2008.05.016
- Jain P, Jain M, Halder C, Singh TB, Jain S. Melatonin and its correlation with testosterone in polycystic ovarian syndrome. *J Hum Reprod Sci.* (2013) 6:253–8. doi: 10.4103/0974-1208.126295
- Yang HL, Zhou WJ, Gu CJ, Meng YH, Shao J, Li DJ, et al. Pleiotropic roles of melatonin in endometriosis, recurrent spontaneous abortion, and polycystic ovary syndrome. *Am J Reprod Immunol.* (2018) 80:e12839. doi: 10.1111/aji.12839
- Shreeve N, Cagampang F, Sadek K, Tolhurst M, Houldey A, Hill CM, et al. Poor sleep in PCOS; is melatonin the culprit? *Hum Reprod.* (2013) 28:1348–53. doi: 10.1093/humrep/det013
- Spinedi E, Cardinali DP. The polycystic ovary syndrome and the metabolic syndrome: a possible chronobiotic–cytoprotective adjuvant therapy. *Int J Endocrinol.* (2018) 2018:1349868. doi: 10.1155/2018/1349868
- Pacchiarotti A, Carlomagno G, Antonini G, Pacchiarotti A. Effect of myo-inositol and melatonin versus myo-inositol, in a randomized controlled trial, for improving *in vitro* fertilization of patients with polycystic ovarian syndrome. *Gynecol Endocrinol.* (2016) 32:69–73. doi: 10.3109/09513590.2015.1101444
- Unfer V, Raffone E, Rizzo P, Buffo S. Effect of a supplementation with myo-inositol plus melatonin on oocyte quality in women who failed to conceive in previous *in vitro* fertilization cycles for poor oocyte quality: a prospective, longitudinal, cohort study. *Gynecol Endocrinol.* (2011) 27:857–61. doi: 10.3109/09513590.2011.564687
- Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertil Steril.* (2004) 81:19–25. doi: 10.1016/j.fertnstert.2003.10.004
- Raygan F, Ostadmohammadi V, Bahmani F, Reiter RJ, Asemi Z. Melatonin administration lowers biomarkers of oxidative stress and cardio-metabolic risk in type 2 diabetic patients with coronary heart disease: a randomized, double-blind, placebo-controlled trial. *Clin Nutr.* (2019) 38:191–6. doi: 10.1016/j.clnu.2017.12.004

27. Jamilian M, Foroozanfard F, Bahmani F, Talaei R, Monavari M, Asemi Z. Effects of zinc supplementation on endocrine outcomes in women with polycystic ovary syndrome: a randomized, double-blind, placebo-controlled trial. *Biol Trace Elem Res.* (2016) 170:271–8. doi: 10.1007/s12011-015-0480-7
28. Hatch R, Rosenfield RL, Kim MH, Tredway D. Hirsutism: implications, etiology, and management. *Am J Obstet Gynecol.* (1981) 140:815–30. doi: 10.1016/0002-9378(81)90746-8
29. Ramezani Tehrani F, Minoos S, Azizi F. Validation of a simplified method to assess hirsutism in the Iranian population. *Eur J Obstet Gynecol Reprod Biol.* (2014) 174:91–5. doi: 10.1016/j.ejogrb.2013.12.008
30. Tatsch E, Bochi GV, Pereira Rda S, Kober H, Agertt VA, de Campos MM, et al. A simple and inexpensive automated technique for measurement of serum nitrite/nitrate. *Clin Biochem.* (2011) 44:348–50. doi: 10.1016/j.clinbiochem.2010.12.011
31. Benzie IJ, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: the FRAP assay. *Anal Biochem.* (1996) 239:70–6. doi: 10.1006/abio.1996.0292
32. Beutler E, Gelbart T. Plasma glutathione in health and in patients with malignant disease. *J Lab Clin Med.* (1985) 105:581–4.
33. Dunkley PR, Jarvie PE, Robinson PJ. A rapid Percoll gradient procedure for preparation of synaptosomes. *Nat Protoc.* (2008) 3:1718–28. doi: 10.1038/nprot.2008.171
34. Cassone VM, Natesan AK. Time and time again: the phylogeny of melatonin as a transducer of biological time. *J Biol Rhythms.* (1997) 12:489–97. doi: 10.1177/074873049701200602
35. Nakamura Y, Tamura H, Takayama H, Kato H. Increased endogenous level of melatonin in preovulatory human follicles does not directly influence progesterone production. *Fertil Steril.* (2003) 80:1012–6. doi: 10.1016/S0015-0282(03)01008-2
36. Ronnberg L, Kauppila A, Leppaluoto J, Martikainen H, Vakkuri O. Circadian and seasonal variation in human preovulatory follicular fluid melatonin concentration. *J Clin Endocrinol Metab.* (1990) 71:492–6. doi: 10.1210/jcem-71-2-493
37. Wetterberg L, Arendt J, Paunier L, Sizonenko PC, Donselaar W, Heyden T. Human serum melatonin changes during the menstrual cycle. *J Clin Endocrinol Metab.* (1976) 42:185–8. doi: 10.1210/jcem-42-1-185
38. Wickenheisser JK, Quinn PG, Nelson VL, Legro RS, Strauss JF, McAllister JM. Differential activity of the cytochrome P450 17 α -hydroxylase and steroidogenic acute regulatory protein gene promoters in normal and polycystic ovary syndrome theca cells. *J Clin Endocrinol Metab.* (2000) 85:2304–11. doi: 10.1210/jcem.85.6.6631
39. Kumar A, Woods KS, Bartolucci AA, Azziz R. Prevalence of adrenal androgen excess in patients with the polycystic ovary syndrome (PCOS). *Clin Endocrinol.* (2005) 62:644–9. doi: 10.1111/j.1365-2265.2005.02256.x
40. Dokras A, Sarver DB, Allison KC, Milman L, Kris-Etherton PM, Kunselman AR, et al. Weight loss and lowering androgens predict improvements in health-related quality of life in women with PCOS. *J Clin Endocrinol Metab.* (2016) 101:2966–74. doi: 10.1210/jc.2016-1896
41. Boztosun A, Acmaz G, Ozturk A, Muderris II. Clinical efficacy of low dose flutamide plus Diane-35 in the treatment of idiopathic hirsutism and polycystic ovary syndrome. *Ginekol Pol.* (2013) 84:258–62. doi: 10.17772/gp/1573
42. Tagliaferri V, Romualdi D, Scarinci E, Cicco S, Florio CD, Immediata V, et al. Melatonin treatment may be able to restore menstrual cyclicality in women with PCOS: a pilot study. *Reprod Sci.* (2018) 25:269–75. doi: 10.1177/1933719117711262
43. Nikmard F, Hosseini E, Bakhtiyari M, Ashrafi M, Amidi F, Aflatoonian R. Effects of melatonin on oocyte maturation in PCOS mouse model. *Anim Sci J.* (2017) 88:586–92. doi: 10.1111/asj.12675
44. Pai SA, Majumdar AS. Protective effects of melatonin against metabolic and reproductive disturbances in polycystic ovary syndrome in rats. *J Pharm Pharmacol.* (2014) 66:1710–21. doi: 10.1111/jphp.12297
45. Mistraletti G, Paroni R, Umbrello M, D’Amato L, Sabbatini G, Taverna M, et al. Melatonin pharmacological blood levels increase total antioxidant capacity in critically ill patients. *Int J Mol Sci.* (2017) 18:759. doi: 10.3390/ijms18040759
46. Kozirog M, Poliwczyk AR, Duchnowicz P, Koter-Michalak M, Sikora J, Broncel M. Melatonin treatment improves blood pressure, lipid profile, and parameters of oxidative stress in patients with metabolic syndrome. *J Pineal Res.* (2011) 50:261–6. doi: 10.1111/j.1600-079X.2010.00835.x
47. Pakravan H, Ahmadian M, Fani A, Aghaee D, Brumanad S, Pakzad B. The effects of melatonin in patients with nonalcoholic fatty liver disease: a randomized controlled trial. *Adv Biomed Res.* (2017) 6:40. doi: 10.4103/2277-9175.204593
48. Akbari M, Ostadmohammadi V, Tabrizi R, Lankarani KB, Heydari ST, Amirani E, et al. The effects of melatonin supplementation on inflammatory markers among patients with metabolic syndrome or related disorders: a systematic review and meta-analysis of randomized controlled trials. *Inflammopharmacology.* (2018) 26:899–907. doi: 10.1007/s10787-018-0508-7
49. Mesri Alamdari N, Mahdavi R, Roshanravan N, Lotfi Yaghin N, Ostadrahimi AR, Faramarzi E. A double-blind, placebo-controlled trial related to the effects of melatonin on oxidative stress and inflammatory parameters of obese women. *Horm Metab Res.* (2015) 47:504–8. doi: 10.1055/s-0034-1384587
50. Alviggi C, Conforti A, De Rosa P, Strina I, Palomba S, Vallone R, et al. The distribution of stroma and antral follicles differs between insulin-resistance and hyperandrogenism-related polycystic ovarian syndrome. *Front Endocrinol.* (2017) 8:117. doi: 10.3389/fendo.2017.00117
51. Pawelczyk M, Rosenthal J, Milla S, Liu YH, Shah B. Evaluation of the pro-inflammatory cytokine tumor necrosis factor- α in adolescents with polycystic ovary syndrome. *J Pediatr Adolesc Gynecol.* (2014) 27:356–9. doi: 10.1016/j.jpog.2014.01.104
52. Alviggi C, Cariati F, Conforti A, De Rosa P, Vallone R, Strina I, et al. The effect of FT500 Plus(R) on ovarian stimulation in PCOS women. *Reprod Toxicol.* (2016) 59:40–4. doi: 10.1016/j.reprotox.2015.10.014
53. Lin YW, Lee LM, Lee WJ, Chu CY, Tan P, Yang YC, et al. Melatonin inhibits MMP-9 transactivation and renal cell carcinoma metastasis by suppressing Akt-MAPKs pathway and NF- κ B DNA-binding activity. *J Pineal Res.* (2016) 60:277–90. doi: 10.1111/jpi.12308
54. Mohan N, Sadeghi K, Reiter RJ, Meltz ML. The neurohormone melatonin inhibits cytokine, mitogen and ionizing radiation induced NF- κ B. *Biochem Mol Biol Int.* (1995) 37:1063–70.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Jamilian, Foroozanfard, Mirhosseini, Kavossian, Aghadavod, Bahmani, Ostadmohammadi, Kia, Eftekhari, Ayati, Mahdavinia and Asemi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.