

# Versatile and Enantioselective Preparation of Planar-Chiral Metallocene-Fused 4-Dialkylaminopyridines and Their Application in Asymmetric Organocatalysis

*Kazuhiro Yoshida,<sup>\*,†</sup> Qiang Liu,<sup>‡,§</sup> Risa Yasue,<sup>†</sup> Shiro Wada,<sup>†</sup> Ryosuke Kimura,<sup>†</sup> Takuma Konishi,<sup>‡</sup> and Masamichi Ogasawara<sup>\*,‡</sup>*

<sup>†</sup> Department of Chemistry, Graduate School of Science, Chiba University, Yayoi-cho, Inage-ku, Chiba 263-8522, Japan

<sup>‡</sup> Department of Natural Science, Graduate School of Science and Technology and Research Cluster on "Innovative Chemical Sensing", Tokushima University, Tokushima 770-8506, Japan

<sup>§</sup> Graduate School of Life Science, Hokkaido University, Kita-ku, Sapporo 001-0021, Japan


**Abstract:** A series of ferrocene-fused planar-chiral *N*-tosyl-4-pyridones (*S*)-**2b-d** were prepared in enantiomerically pure forms. Starting with the chiral ferrocenyl acetals, 1-[(2*S*,4*S*)-4-methoxymethyl-1,3-dioxan-2-yl]-1',2',3',4',5'-R<sub>5</sub>-ferrocenes ((-)-**3b**, R = Me; (-)-**3c**, R = Ph; (-)-**3d**, R = Bn), *N*-tosylamino and formyl groups were introduced at the 1- and 2-positions of the ferrocene cores in (*S*)-**11b-d** with control of the planar chirality. After the reaction with

ethynylmagnesium bromide, generated propargyl alcohol derivatives (*S*)-**17** were treated with MnO<sub>2</sub> and catalytic TBAI to give the planar-chiral pyridones by the iodide-catalyzed cyclization. This method is highly practical with a shorter and higher-yield sequence without using noble metal catalysts. Planar-chiral ferrocene-pyridones (*S*)-**2b-d** were reacted with various Me<sub>3</sub>Si-NR'<sub>2</sub> to give a library of ferrocene-fused 4-dialkylaminopyridines ((*S*)-**1**, DAAPs) in high yields as single-enantiomers by the detosylative amination. The cymantrene-fused DAAPs were also prepared in the same way. The library of the chiral DAAPs were examined in the two asymmetric reactions as organocatalysts, and some newly prepared Fc-DAAPs showed better enantioselectivity than the known species.

**Keywords:** planar-chiral; ferrocene; cymantrene; enantioselective; 4-pyridone; 4-dialkylaminopyridine; organocatalyst; cyclization


## Introduction

Ferrocene is a cylindrical compound with unique three-dimensional properties.<sup>1,2</sup> Introduction of two different substituents at the 1- and 2-positions of an Fe(II)-bound η<sup>5</sup>-cyclopentadienyl ligand breaks the symmetry to induce planar chirality in the molecule (Figure 1, left).<sup>3</sup> The ferrocene-based planar chirality has been proven to be useful chiral scaffolds in asymmetric synthesis, and various such ferrocene derivatives have been utilized in a wide range of asymmetric reactions as chiral ligands<sup>4</sup> and chiral catalysts.<sup>5,6</sup>



**Figure 1.** Enantiomeric pair of 1,2-disubstituted ferrocene (left) and Fu's ferrocene-fused planar-chiral 4-dialkylaminopyridines **1** (right).

In 1990s, G. C. Fu employed this design concept for desymmetrizing flat-shaped *N*-heteroarenes, and since then various ferrocene-fused 4-dialkylaminopyridines (Fc-DAAPs; **1**; Figure 1, right) were prepared and applied in asymmetric organocatalysis. Among many chiral 4-dialkylaminopyridine derivatives reported so far,<sup>7,8</sup> Fc-DAAPs are arguably the most successful compounds.<sup>6</sup> And indeed, **1** showed excellent enantioselectivity in a variety of asymmetric transformations.<sup>9</sup> In spite of the exceptional catalytic properties of **1**, applications of these stylish molecules have been rather limited<sup>9,10</sup> probably due to their limited availability. The synthetic sequence developed by Fu is shown in Scheme 1. Two major drawbacks in this synthetic protocol are (i) necessity of the late stage chiral resolution of preformed racemic **1**, and (ii) limited diversity with respect to a substituent at the 4-position in the pyridine ring.<sup>9b,9c,11</sup> A dialkylamino group at the 4-pyridyl position, which plays important roles in controlling the catalytic activity/selectivity of chiral DAAPs,<sup>12</sup> was introduced in the middle stage of the synthetic sequences. This means that each derivative of **1** with a different 4-substituent needs to be prepared as a respective racemate through the independent reaction sequence, and, naturally, the late stage chiral resolution of each compound is essential.




**Scheme 1.** Fu's Original Synthesis of *rac*-**1** and Enantiomeric Resolution.

In 2015, we reported an enantioselective synthesis of Fc-DAAPs and related compounds, which *partly* overcame the drawbacks in the Fu's original synthesis of **1** (Scheme 2).<sup>13</sup> The three features of our synthetic strategy are: (i) introduction of proper substituents at the 1- and 2-positions of a ferrocene platform with controlling its planar chirality utilizing a chiral *ortho*-directing group (**3** to **4** in Scheme 2),<sup>14</sup> (ii) construction of the ferrocene-fused 4-pyridones by the ring-closing metathesis reaction (**8** to **2** in Scheme 2), and (iii) the late stage introduction of various substituents at the 4-position of the ferrocene-fused pyridines (**2** to **1**, **9**, and **10**). The chiral 1,3-dioxan-2-yl group in **3**, developed by Kagan and his coworkers,<sup>15-17</sup> is a powerful chiral directing group for the highly diastereoselective lithiation of **3** (>99% de). Planar-chiral 4-pyridones **2**, obtained in essentially enantiomerically pure forms, are versatile synthetic intermediates, and they could be converted to the various pyridine derivatives with retention of the planar chirality in **2**. An analogous synthetic strategy was applied to the synthesis of metallocene-fused planar-chiral phospholes recently.<sup>18</sup>

Although the synthetic method in Scheme 2 provided a library of the planar-chiral pyridine-based nucleophilic organocatalysts, including previously unknown species, without chiral resolution, it still had room for further improvement. For example, the reaction sequence in Scheme 2 is somewhat lengthy and some steps require the use of expensive noble metal catalysts, such as the Hoveyda/Grubbs-II catalyst,<sup>19</sup> with relatively high catalyst loading. Furthermore, this

"first-generation synthesis" shows severe steric hindrance. The method can be used for the preparation of the Cp- and Cp\*-derivatives ( $R' = H$  or Me), but the corresponding  $\eta^5\text{-C}_5\text{Ph}_5\text{-}$  analogues were not obtained in the same way (vide infra). It should be mentioned that the  $\text{C}_5\text{H}_5\text{-}$  derivatives of **1** are synthetically far less important due to the lower-enantioselectivity on their applications in asymmetric catalysis, and the sterically demanding  $\eta^5\text{-C}_5\text{Ph}_5\text{-}$  analogues show much better performance in many reactions.<sup>9b,d,e,m</sup>



## Scheme 2. Enantioselective Synthesis of Ferrocene-Fused 4-Pyridones **2** and Pyridine

Derivatives **1**, **9**, and **10**<sup>21</sup>: "First-Generation Synthesis" Using Ruthenium-Catalyzed Ring-Closing Metathesis for Cyclization.<sup>13</sup>

With the background mentioned above, we started the present studies with intention to overcome the limitations of the first generation synthesis of **1** (Scheme 2). The improved "second generation synthesis" developed in this study is much shorter (i. e., more effective) and much less costly without using the expensive noble metal catalysts. And more importantly, the newly

developed synthesis can provide sterically demanding analogues of Fc-DAAPs which are with an ( $\eta^5$ -C<sub>5</sub>Ph<sub>5</sub>)Fe or ( $\eta^5$ -C<sub>5</sub>Bn<sub>5</sub>)Fe substructure. These bulky Fc-DAAP derivatives are synthetically more useful showing better enantioselectivity in many cases. It should be mentioned that the ( $\eta^5$ -C<sub>5</sub>Bn<sub>5</sub>)Fe derivatives are previously unknown and prepared for the first time through this study.


In this article, we would like to report the details of the innovative second-generation enantioselective synthesis of planar-chiral Fc-DAAPs and their application in asymmetric organocatalysis.

## Results and Discussion

**Unsuccessful Attempts to Prepare  $\eta^5$ -C<sub>5</sub>Ph<sub>5</sub>-Derivatives of Fc-DAAPs by First Generation Synthesis.** As mentioned in the introduction section, our original method of enantioselective synthesis of Fc-DAAPs, shown in Scheme 2, works only for the  $\eta^5$ -C<sub>5</sub>H<sub>5</sub>- and  $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>-derivatives, and the corresponding  $\eta^5$ -C<sub>5</sub>Ph<sub>5</sub>-derivatives could not be prepared in the same way.

The results of the unsuccessful attempts at preparing the  $\eta^5$ -C<sub>5</sub>Ph<sub>5</sub>-derivatives are summarized in Scheme 3. Chiral acetal (–)-**3c** was prepared from 1-formyl-1',2',3',4',5'-pentaphenylferrocene<sup>20</sup> and (*S*)-4-methoxybutane-1,3-diol in 76% yield. Chiral (2*S*,4*S*)-4-methoxymethyl-1,3-dioxan-2-yl substituent in **3c** is an excellent chiral directing group,<sup>15,16</sup> and deprotonation of **3c** took place with high diastereoselectivity of >99.5% de. Lithiation of **3c** using <sup>t</sup>BuLi at –78 °C in THF followed by a reaction with tosyl azide gave the corresponding ferrocenyl azide, and subsequently the crude azide was reduced under a hydrogen atmosphere in


the presence of catalytic Pd/C to give ferrocenylamine (–)-**4c** in 84% yield (for two steps). The <sup>1</sup>H- and <sup>13</sup>C-NMR analyses of (–)-**4c** clarified the compound to be essentially diastereomerically pure. After *N*-tosylation of (–)-**4c** in >99%, *N*-allylation of (–)-**5c** was examined. To our disappointment, however, the *N*-allylation did not proceed under the conditions in Scheme 3 and desired **6c** could not be obtained. It was suspected that the robustness of (–)-**5c** toward the *N*-allylation might be ascribed to the steric hindrance by the bulky η<sup>5</sup>-C<sub>5</sub>Ph<sub>5</sub> ligand. To diminish the steric problems in (–)-**5c** as much as possible, the *N*-allylation was also examined after deprotection of the chiral 1,3-dioxanyl moiety, but the reaction on aldehyde (*S*)-**11c** did not afford *N*-allylated species **7c**. At this stage, we gave up to pursue the synthesis of the (η<sup>5</sup>-C<sub>5</sub>Ph<sub>5</sub>)Fe-derivatives by the method in Scheme 2.



**Scheme 3.** Unsuccessful Attempts to Prepare Pentaphenylferrocene-Fused DAAP Derivatives by First Generation Synthesis.


**Development of Transition-Metal Free Iodide-Catalyzed Cyclization Producing 4-Pyridones 2.** During our investigations on the transformation of pyridone (*S*)-**2a** to pyridine (*S*)-**1ax**,<sup>13,22</sup> we encountered the formation of unexpected "ring-opening" product (*S*)-**12** using an equimolar mixture of pyrrolidine and titanium(IV) chloride under the unoptimized conditions. The formation of (*S*)-**12** was rationalized as depicted in Scheme 4. In the presence of Lewis acidic  $\text{TiCl}_4$ , pyrrolidine attacks at the  $\beta$ -olefinic carbon in (*S*)-**2a** to give intermediate **13**, which subsequently eliminates the tosylamide anion to give the ring-opening product via a proton shift.






**Scheme 4.** Formation of Ring-Opening Product (*S*)-12 by Reaction of Pyridone (*S*)-2a with Pyrrolidine in the Presence of Titanium(IV) Chloride.

On the assumption that the reaction steps between **13** and (*S*)-12 were reversible, a transition-metal-free cyclization process producing 4-pyridones **2** was pursued (Scheme 5).<sup>23</sup> We envisioned that enone **15**, which possesses a potential leaving group "LG" on the β-sp<sup>2</sup>-carbon, might undergo a cyclization providing pyridone **2** as in Scheme 5. For the success of this pyridone formation instead of the ring-opening as in Scheme 4, the LG substituent in **16** must be a better leaving group than the tosylamide moiety. Cyclization precursor **15** could be generated *in situ* by the conjugate addition of a nucleophilic LG<sup>-</sup> to ynone **14**. Whereas LG<sup>-</sup> would be regenerated at the transformation of **16** to **2**, a substoichiometric (catalytic) amount of LG<sup>-</sup> might be sufficient.




**Scheme 5.** Strategy for Nucleophile-Catalyzed Cyclization of 1-Propynoyl-2-*N*-tosylaminoferrocene **14**.

At the outset, this idea was examined in the synthesis of pentaphenylferrocene-pyridone **2c**, which could not be prepared by our first generation synthesis as mentioned in the previous section. The results are summarized in Scheme 6. Treatment of aldehyde (*S*)-**11c**, which was prepared as in Scheme 3, with ethynylmagnesium bromide afforded propargylic alcohol (–)-**17c** in 95% yield as a diastereomeric mixture. Ynone (*S*)-**14c** could be prepared by the MnO<sub>2</sub>-oxidation of (–)-**17c** in dichloromethane, but the compound was easily polymerized into uncharacterized gummy materials and was not isolable. Due to the susceptibility of (*S*)-**14c**, a direct conversion of (–)-**17c** into (*S*)-**2c** "in one pot" was tested. That is, propargylic alcohol (–)-**17c** was reacted with manganese(IV) oxide in the presence of a catalytic nucleophile. After an extensive survey of various nucleophiles, which include pyridine, DMAP, DABCO, triethylamine, copper(I) iodide, etc., it was found that a reaction using tetrabutylammonium iodide (TBAI; 0.5 equiv. to **17c**) proceeded cleanly to produce (*S*)-**2c** in 99% yield (see the Supporting Information for details).



**Scheme 6.** Iodide-Catalyzed Synthesis of Ferrocene-Fused 4-Pyridone **2c**.


Deep red crystals of **(-)**-**2c** was grown by slow diffusion of pentane into the concentrated dichloromethane solution of **(-)**-**2c**. The single-crystal X-ray crystallography revealed that the compound was a single enantiomer. The unit cell contains two independent molecules, having slightly different conformations, and the structure of one of the two crystallographically independent molecules is shown in Figure 2.<sup>24</sup> The absolute configuration of levorotatory **2c** was determined to be **(S)**, which is consistent with the stereochemistry deduced from the diastereoselective lithiation of **(-)**-**3c**.<sup>15,16</sup> The tosyl group on N(1) takes the position opposite to the bulky  $\eta^5$ -C<sub>5</sub>Ph<sub>5</sub> ligand avoiding steric interactions with the phenyl groups. On the other hand, the space above the carbonyl moiety of the pyridone ring remains relatively open, which may be an important issue for the conversion of **2c** to dialkylaminopyridine derivatives **1** by the reactions with trimethylsilylamines (*vide infra*).



**Figure 2.** Ball-and-stick drawing of the single-crystal X-ray structure of (*S*)-(-)-**2c** with selected atom numbering.<sup>24</sup>

Scheme 7 shows a plausible reaction mechanism for the iodide-catalyzed cyclization of 1-propynoyl-2-*N*-tosylaminoferrocenes producing the ferrocene-fused *N*-tosyl-4-pyridones. The iodide anion attacks the alkynyl terminal in ynone (*S*)-**14**, which is generated *in situ* by the oxidation of propargylic alcohol (*S*)-**17** with manganese(IV) oxide, to give intermediate **A**. After the isomerization of allenol **A** into enone **B**, the intramolecular conjugate addition of the anionic tosylamide to the enone moiety affords intermediate **C**. Subsequent elimination of the iodide anion from **C** gives pyridone (*S*)-**2** to complete the catalytic cycle. For the effective synthesis of

(*S*)-**2**, the undesirable self-polymerization of (*S*)-**14** should be avoided. Ynone (*S*)-**14**, generated by the oxidation of (*S*)-**17**, needs to react with iodide anion faster than the self-polymerization. This may be a reason why relatively high catalyst-loading (i.e., 50% TBAI) is required to realize the high yields in this process. Although Scheme 7 could explain the formation of (*S*)-**2**, involvement other processes, such as a radical mechanism, could not be ruled out.




**Scheme 7.** Plausible Reaction Mechanism for Iodide-Catalyzed Cyclization Producing Ferrocene-Fused 4-Pyridones.

**Optimization of Reaction Conditions for More Efficient Synthesis of Planar-Chiral Ferrocene-Fused 4-Pyridones **2**.** With the iodide-catalyzed cyclization reaction in our hands, the enantioselective synthesis of (*S*)-**2c** was finally accomplished. The synthetic sequence

developed in this study is versatile and applicable to the synthesis of other ferrocene-fused pyridones, such as **2b** and **2d** containing the ( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)Fe and ( $\eta^5$ -C<sub>5</sub>Bn<sub>5</sub>)Fe moieties, respectively.

To enhance the practicality of this "second-generation synthesis", the reaction conditions for each step were thoroughly optimized to simplify the synthetic operations. The results are summarized in Scheme 8. The improved enantioselective synthesis of planar-chiral ferrocene-fused pyridones (*S*)-**2** starts from respective chiral acetals (–)-**3** and needs only three isolation/purification steps to (*S*)-**2**. The diastereoselective lithiation of (–)-**3** using *t*BuLi followed by a reaction with tosyl azide afforded the corresponding ferrocenyl azide, and the subsequent palladium-catalyzed reduction of the crude azide under a hydrogen atmosphere gave the ferrocenylamine, which was immediately tosylated without purification to provide *N*-tosylaminoferrocenyl acetal (–)-**5** in high yield ranging 58-84%. Acetal (–)-**5**, purified by silica gel column chromatography, was confirmed to be diastereomerically pure by the <sup>1</sup>H- and <sup>13</sup>C-NMR analyses. The acid-catalyzed hydrolysis of (–)-**5** gave the corresponding aldehyde (*S*)-**11** in >92% yield in enantiomerically pure form. The purification/isolation of (*S*)-**11** may be skipped, but we checked the enantiomeric homogeneity of (*S*)-**11** at this stage. Aldehyde (*S*)-**11** was treated with ethynylmagnesium bromide, and the crude propargylic alcohols, which were obtained by simple extraction, was applied to the iodide-catalyzed cyclization as in Schemes 6/7 to give planar-chiral 4-pyridone (*S*)-**2** in high yield ranging from 66% to 95% (Scheme 8).



**Scheme 8.** Optimized Synthetic Sequence for Enantioselective Construction of Ferrocene-Fused 4-Pyridones **2**: "Second-Generation Synthesis" Using Iodide-Catalyzed Cyclization of Propynoyl-*N*-tosylaminoferrocenes.


The second-generation synthesis of pyridones **2** developed here is much shorter with the simpler operations and higher yields compared to the first-generation synthesis. Furthermore, the second-generation synthesis can afford the sterically demanding ferrocenopyridones such as **2c** and **2d**, which are not accessible by the first-generation synthesis. While the total yield from (-)-**3b** to (*S*)-**2b** was 35% in nine-steps by the first-generation synthesis, the second-generation synthesis according to Scheme 8 provided (*S*)-**2b** in 50% total yield starting with (-)-**3b** in six-steps. The synthesis of (*S*)-**2c** was more effective in 71% total yield from (-)-**3c**. The chiral source of the chiral directing group in (-)-**3** is commercially available (*S*)-1,2,4-butanetriol, of which optical antipode, (*R*)-1,2,4-butanetriol, is also commercially available for the synthesis of (*R*)-enantiomers of planar-chiral ferrocenopyridones (*R*)-**3**.

**Preparation of Planar-Chiral Ferrocenopyridines **1** from 4-Pyridones **2**.** In the previous report, we demonstrated that planar-chiral *N*-tosyl-4-pyridones (*S*)-**2a** and (*S*)-**2b** were versatile precursors to various pyridine derivatives.<sup>13</sup> The direct conversion of (*S*)-**2b** into a series of 4-dialkylaminopyridine derivatives (*S*)-**1bw-bz** was achieved by the reaction with an appropriate *N*-trimethylsilylamine in the presence of titanium(IV) chloride in good to excellent yields (see,

Schemes 2 and 9). This detosylative amination reaction was found to be operative for the sterically demanding ferroc-pyridones such as (*S*)-**2c** and (*S*)-**2d** as well, and the results are summarized in Scheme 9. The reactions introducing dimethylamino, pyrrolidinyl, or morpholyl groups at the 4-position of the pyridine rings proceeded very cleanly and the corresponding Fc-DAAPs were obtained in excellent yields ranging 82-99%. On the other hand, the reaction with diethyl(trimethylsilyl)amine was somewhat slow and the yields of the 4-diethylaminopyridine derivatives are relatively low (47-80%). Steric influence of the  $\eta^5\text{-C}_5\text{R}'_5$  ligands on the detosylative amination reaction is minimal and three pyridones **2b-d** showed similar reactivity. The isolated yields of the  $\eta^5\text{-C}_5\text{Me}_5$  derivatives are somewhat lower, which can be ascribed to their sensitivity toward air-oxidation, since the chromatographic purification was conducted under air.


It should be emphasized that all the transformations shown in Schemes 8 and 9 are stereoretentive. Whereas (*S*)-**2b-d** obtained by our method are enantiomerically pure, planar-chiral ferroc-DAAP derivatives are also single enantiomers. Namely, we have established the divergent process preparing a library of various planar-chiral pyridine-based nucleophilic organocatalysts in enantiomerically pure forms without optical resolution. Our second-generation synthesis is fairly versatile and various substituents can be introduced both at the 4-pyridyl position as well as in the  $\eta^5\text{-C}_5\text{R}'_5$  moiety. The  $\eta^5$ -pentabenzylcyclopentadienyl derivatives (**1dw-dy**) and 4-diethylamino-/4-morpholyl-ferroc-pyridines (**1by**, **1cy**, **1dy**, **1bz**, and **1cz**) are new compounds and prepared for the first time by our synthetic method.






**Scheme 9.** Conversion of (*S*)-Ferroco-*N*-tosyl-4-pyridones **2** to Ferroco-DAAP Derivatives **1**.

Single crystals of (*S*)-(-)-**1dw** were grown from pentane/dichloromethane and isolated as deep red prisms. The X-ray crystallography revealed that the compound was single enantiomeric.<sup>24</sup> The absolute configuration of levorotatory **2c** was determined to be (*S*), which is consistent with the stereochemistry deduced from the reported diastereoselective lithiation of the closely related (-)-**3b**.<sup>15,16</sup> All five phenyl groups in the  $\eta^5\text{-C}_5\text{Bn}_5$  ligand point away from Fe(1) avoiding the steric interaction with the ( $\eta^5\text{-pyrido-Cp}$ )Fe moiety. The ferrocene core in (-)-**3b** shows essentially no distortion with the dihedral angle between the two  $\eta^5$ -ligand being 3.08°.




**Figure 3.** Ball-and-stick drawing of the single-crystal X-ray structure of (*S*)-(-)-**1dw** with selected atom numbering.<sup>24</sup>

A plausible reaction mechanism for the detosylative amination reaction is shown in Scheme 10. A reaction of a trimethylsilylamine with titanium(IV) chloride generates a titanium amide,  $\text{Cl}_3\text{Ti-NR}_2$ , that attacks the carbonyl moiety in (*S*)-**2** to form intermediate **D**. Elimination of titanium(IV) oxodichloride gives iminium species **E**, that also exists in different resonance form **F**. The subsequent reaction of *N*-tosylpyridinium intermediate **F** with  $\text{Me}_3\text{Si-NR}_2$  affords the corresponding (*S*)-**1** together with tosylamide  $\text{TsNR}_2$ , which was indeed isolated from the reaction mixture.



**Scheme 10.** Plausible Reaction Pathway of Detosylative Amination of (S)-2 Producing (S)-1.

**Enantioselective Synthesis of Planar-Chiral Cymantrene-Fused DMAP (S)-1ew and PPY (S)-1ex.** The generality of the synthetic method developed in this study was tested on preparation of DAAP derivatives fused with a metallocene other than ferrocenes. Cymantrene, ( $\eta^5$ -cyclopentadienyl)manganese(I) tricarbonyl, was chosen as a platform for this purpose, and the results are shown in Scheme 11. The synthesis began with known chiral cymantrenyl acetal (+)-**3e**,<sup>25</sup> of which sequential diastereoselective lithiation, azidation, and NaBH<sub>4</sub> reduction gave aminocymantrene (+)-**4e** in 70% yield as a single-diastereomer. After the *N*-tosylation of (+)-**4e**, the hydrolysis of the chiral acetal moiety using aqueous hydrochloric acid provided aldehyde (S)-**11e** in enantiomerically pure form in 48% yield. Treatment of (S)-**11e** in the same way as the ferrocene derivatives provided the corresponding cymantrene-fused 4-pyridone (S)-**2e** in 68% yield. Pyridone (S)-**2e** could be converted to cymantrene-fused DMAP (S)-**1ew** and PPY (S)-**1ex** in 49% and 46% yields, respectively.



**Scheme 11.** Enantioselective Synthesis of Planar-Chiral Cyantro-pyridines (*S*)-**1ew** and (*S*)-**1ex**.


Although the yields of the cyantrene derivatives were lower than those of the ferrocene derivatives, probably due to the air sensitivity of the manganese complexes, the reaction sequence in Scheme 11 demonstrated the versatility and generality of the second-generation synthesis of the planar-chiral DAAP species.

**Application of (*S*)-1 in Asymmetric Organocatalysis.** The library of the nucleophilic organocatalysts, (*S*)-**1**, obtained in this study was applied in the two prototypical asymmetric reactions and their catalytic performance was evaluated.

The first asymmetric reaction catalyzed by (*S*)-**1** is an addition reaction of 2-*t*Bu-phenol (**19**) to ethyl(*p*-tolyl)ketene (**18**), which was suggested to take place by the Brønsted acid-catalyzed

mechanism (Table 1).<sup>9i</sup> The  $\eta^5$ -C<sub>5</sub>Me<sub>5</sub> derivatives, (*S*)-**1bw-bz**, were prepared by our first-generation synthesis and their catalytic applications in this reaction were already examined in the previous report.<sup>13</sup> The reported results are included in the table for comparison (entries 1-4). Among them, ferroc-PPY (*S*)-**1bx**, which is Fu's original, showed good catalytic activity in the addition reaction to give ester **20** in 90% yield with 92% ee (entry 2). The  $\eta^5$ -C<sub>5</sub>Ph<sub>5</sub>-derivatives, (*S*)-**1cw-cz**, were poor catalysts for this transformation, and **20** was obtained in less than 8% yields with low enantioselectivities of 32% ee at best (entries 5-8). The poor performance of (*S*)-**1cw-cz**, which contain the electron-withdrawing  $\eta^5$ -C<sub>5</sub>Ph<sub>5</sub> ligand,<sup>26</sup> could be attributed to their lower basicity compared to (*S*)-**1bw-bz**. And indeed, cymantro-pyridines (*S*)-**1ew-ex**, which were also weakly basic due to the electron-withdrawing nature of the cymantrene framework, showed similar poor catalytic activities and enantioselectivities in the reaction (entries 12-13). On the other hand,  $\eta^5$ -C<sub>5</sub>Bn<sub>5</sub>-derivatives (*S*)-**1dw-dy**, which contain a bulkier and electron-donating  $\eta^5$ -pentaalkylcyclopentadienide, could be regarded as refined variants of  $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>-derivatives (*S*)-**1bw-bz** and showed excellent catalytic activities (entries 9-11). Pyridine catalysts (*S*)-**1dw-dy** provided (*R*)-**20** in excellent yields with better enantioselectivities compared to the products obtained by homologous (*S*)-**1bw-by** (entries 5-7 vs entries 9-11). Notably, newly developed (*S*)-**1dx** displayed the highest enantioselectivity of 94% ee in the present reaction and outperformed the other ferroc-/cymantro-pyridines.

**Table 1.** Enantioselective Addition of *o*-<sup>t</sup>Bu-Phenol (**19**) to Ethyl(*p*-tolyl)ketene (**18**) Catalyzed by (*S*)-**1**<sup>a</sup>




entry	( <i>S</i> )- <b>1</b>	yield (%) <sup>b</sup>	% ee of <b>20</b> <sup>c</sup>
1 <sup>d</sup>	<b>1bw</b>	92	79
2 <sup>d</sup>	<b>1bx</b>	90	92
3 <sup>d</sup>	<b>1by</b>	90	90
4 <sup>d</sup>	<b>1bz</b>	84	79
5	<b>1cw</b>	8	9
6	<b>1cx</b>	7	15
7	<b>1cy</b>	0	-
8	<b>1cz</b>	4	32
9	<b>1dw</b>	92	90
10	<b>1dx</b>	91	94
11	<b>1dy</b>	92	90
12	<b>1ew</b>	5	40
13	<b>1ex</b>	16	15

<sup>a</sup> The reaction was carried out in toluene at 23 °C in the presence of catalyst (*S*)-**1** (3 mol %). The absolute configuration of **20** was deduced by comparison with the reported results [ref. 9i]. <sup>b</sup> Isolated yield by silica gel chromatography. <sup>c</sup> Determined by HPLC analysis on a chiral stationary phase (see the Supporting Information for details). <sup>d</sup> Taken from ref. 13.

The second reaction examined was the kinetic resolution of racemic secondary alcohol *rac*-**21**. The acetylation of *rac*-**21** with acetic anhydride proceeds in an enantioselective fashion in the presence of (*S*)-**1** (4 mol %) to give ester (*S*)-**22** and recovered (*R*)-**21** (Table 2). The catalytic

properties of the  $\eta^5\text{-C}_5\text{Me}_5$  derivatives (*S*)-**1bw-bz** were already examined in the previous report<sup>13</sup> and the reported results are included in Table 2 for comparison (entries 1-4). Among the  $\eta^5\text{-C}_5\text{Me}_5$  derivatives, (*S*)-**1by** having a 4-diethylamino substituent showed the highest enantioselectivity with *s*-factor<sup>27</sup> of 6.7 (entry 3). It was reported, however, that  $\eta^5\text{-C}_5\text{Me}_5$  derivative **1bw** was not an appropriate catalyst for the analogous kinetic resolution of secondary alcohols and the corresponding  $\eta^5\text{-C}_5\text{Ph}_5$ -derivative **1cw** showed much better enantioselectivity.<sup>9b</sup> With our second-generation synthesis of the planar-chiral DAAPs, a series of  $\eta^5\text{-C}_5\text{Ph}_5$ -derivatives (*S*)-**1cw-cz** were available for this study. As we expected, indeed, (*S*)-**1cw-cz** showed excellent enantioselectivities with *s*-factors ranging 13 to 69 (entries 5-8). The best selectivity (*s* = 69) was recorded using (*S*)-**1cz**, which was prepared for the first time in this study, but its catalytic activity was decreased due to the less electron-donating ability of the 4-morpholyl substituent (entry 8). The performance of  $\eta^5\text{-C}_5\text{Bn}_5$  derivatives (*S*)-**1dw-dy** was somewhat similar to that of (*S*)-**1bw-by**, but the former showed slightly better selectivities than the latter (entries 9-11). Once again, (*S*)-**1dw-dy** can be seen as improved variants of (*S*)-**1bw-by**. Cymantrene-fused DMAP (*S*)-**1ew** and PPY (*S*)-**1ex** were also applied to the kinetic resolution of *rac*-**21**. The results were, however, unsatisfactory in terms of both catalytic activity and enantioselectivity, which can be ascribed to the less electron-donating and the sterically compact character of the cymantrene platform in (*S*)-**1ew-ex** (entries 12 and 13).

**Table 2.** Enantioselective Acetylation of *rac*-**21** Catalyzed by (*S*)-**1**<sup>a</sup>



entry	(S)-1	conv (%) <sup>b</sup>	% ee of <b>22</b> <sup>c</sup>	% ee of <b>21</b> <sup>c</sup>	<i>s</i> -factor <sup>d</sup>
1 <sup>e</sup>	<b>1bw</b>	65	41	75	5.0
2 <sup>e</sup>	<b>1bx</b>	62	43	70	5.0
3 <sup>e</sup>	<b>1by</b>	70	39	92	6.7
4 <sup>e</sup>	<b>1bz</b>	66	41	78	5.3
5	<b>1cw</b>	41	89	62	33
6	<b>1cx</b>	46	91	76	48
7	<b>1cy</b>	45	75	60	13
8	<b>1cz</b>	7	97	8	69
9	<b>1dw</b>	62	47	78	6.3
10	<b>1dx</b>	66	43	83	6.0
11	<b>1dy</b>	68	45	98	11
12	<b>1ew</b>	31	50	23	3.7
13	<b>1ex</b>	27	54	19	4.0

<sup>a</sup> The reaction was carried out in diethyl ether at 20 °C in the presence of catalyst (S)-1 (4 mol %). <sup>b</sup> Determined by <sup>1</sup>H NMR analysis. <sup>c</sup> Determined by HPLC analysis on a chiral stationary phase (see the Supporting Information for details). The absolute configurations of **22** and recovered **21** were determined by comparison with the reported results [ref. 9b]. <sup>d</sup> Calculated based on a first-order equation [ref. 27]. <sup>e</sup> Taken from ref. 13.

## Conclusions

In summary, we have developed a substantially improved method of preparing various metallocene-fused planar-chiral 4-pyridones **2**/4-dialkylaminopyridines **1** in enantiomerically



pure forms. The renewed method, "second-generation synthesis of Fc-DAAPs", is much more practical and advantageous than our previous enantioselective synthesis of **2/1**, because it is (i) with shorter and more efficient synthetic sequence, (ii) results in higher yields of **2/1**, (iii) no need for expensive noble metal catalysts, and (iv) accessible to sterically demanding Fc-DAAPs having an ( $\eta^5$ -C<sub>5</sub>Ph<sub>5</sub>)Fe or ( $\eta^5$ -C<sub>5</sub>Bn<sub>5</sub>)Fe moiety. Whereas the bulky Fc-DAAP derivatives are synthetically more useful showing better enantioselectivity in general, the last point is particularly important innovation in the present protocol. These bulky Fc-DAAPs could not be prepared by our first-generation synthesis. The  $\eta^5$ -C<sub>5</sub>Bn<sub>5</sub>-DAAP derivatives are previously unknown and prepared for the first time in the present studies. The success of the second-generation synthesis depends in large part on the discovery of the iodide-catalyzed cyclization of 1-propynoyl-2-*N*-tosylaminoferrocenes **14** giving the corresponding 4-pyridones **2** in excellent yields. One unique feature of our synthesis of Fc-DAAPs is the late stage introduction of various dialkylamino-substituents at the 4-position of Fc-DAAPs by the detosylative amination of **2**. Since the detosylative amination worked efficiently even on the bulky pyridones such as **2c** and **2d**, our method is divergent and could provide a library of planar-chiral pyridine-based nucleophilic organocatalysts. Due to the versatility of the second-generation synthesis, a new family of planar-chiral DAAP derivatives, cymantrene-fused DAAPs **1e**, were also prepared in the same way.

Finally, the library of the planar-chiral DAAPs were examined in the two prototypical asymmetric reactions as organocatalysts. Some newly prepared Fc-DAAP derivatives showed better performance than the known species in these reactions. This is a clear-cut advantage of having the library of Fc-DAAPs.

Since the development of Fc-DAAPs **1** in 1990s by Fu, these compounds have been recognized as one of the most attractive and versatile chiral nucleophilic catalysts showing applicability to a wide range of asymmetric reactions. We believe that present study should contribute further development of these unique molecules.

## **AUTHOR INFORMATION**

### **Corresponding Authors**

kyoshida@faculty.chiba-u.jp

ogasawar@tokushima-u.ac.jp

## **ASSOCIATED CONTENT**

**Supporting Information.** The Supporting Information is available free of charge on the ACS Publications website at DOI: \*\*\*\*\*.

The following files are available free of charge.

Experimental procedures and compound characterization data (PDF)

Crystallographic data for (*S*)-(-)-**2c** and (*S*)-(-)-**1dw** (CIF)

## **ACKNOWLEDGMENT**

This work was supported by JSPS KAKENHI "Precisely Designed Catalysts with Customized Scaffolding" to K.Y. (Grant #18H04236) and a Grant-in-Aid for Scientific Research (B) to M.O. (Grant #18H01979) from MEXT, Japan.

## REFERENCES

(1) (a) Kealy, T. J.; Pauson, P. L. "A New Type of Organo-Iron Compound" *Nature* **1951**, *168*, 1039-1040. (b) Wilkinson, G.; Rosenblum, M.; Whiting, M. C.; Woodward, R. B. "The Structure of Iron Bis-cyclopentadienyl" *J. Am. Chem. Soc.* **1952**, *74*, 2125-2126. (c) Fischer, E. O.; Pfab, W. "Zur Kristallstruktur der Di-cyclopentadienyl-verbindungen des Zwiwertigen Eisens, Kobalts, und Nickels" *Z. Naturforsch. B* **1952**, *7*, 377-379. (d) Dunitz, J.; Orgel, L.; Rich A. "The Crystal Structure of Ferrocene" *Acta Crystallographica* **1956**, *9*, 373-375.

(2) (a) Pauson, P. L. "Ferrocene—How It All Began" *J. Organomet. Chem.* **2001**, *637-639*, 3-6. (b) Rosenblum, M. "The Early Ferrocene Days—A Personal Recollection" *J. Organomet. Chem.* **2001**, *637-639*, 13-15. (c) Whiting, M. C. "Recollections of the Arrival of Ferrocene" *J. Organomet. Chem.* **2001**, *637-639*, 16-17. (d) Laszlo, P.; Hoffmann, R. "Ferrocene: Ironclad History or Rashomon Tale ?" *Angew. Chem. Int. Ed.* **2000**, *39*, 123-124.

(3) (a) Halterman, R. L. "Synthesis and Applications of Chiral Cyclopentadienylmetal Complexes" *Chem. Rev.* **1992**, *92*, 965-994. (b) Wagner, G.; Herrmann, R. "Chiral Ferrocene Derivatives. An Introduction" In *Ferrocenes*; Togni, A.; Hayashi, T., Eds.; VCH: Weinheim, 1995; Chap. 4, p. 173-218. (c) Togni, A. "Planar-chiral Ferrocenes: Synthetic Methods and Applications" *Angew. Chem., Int. Ed.* **1996**, *35*, 1475-1477. (d) Richards, C. J.; Locke, A. J. "Recent Advances in the Generation of Non-racemic Ferrocene Derivatives and Their Application to Asymmetric Synthesis" *Tetrahedron: Asymmetry* **1998**, *9*, 2377-2407. (e) Štěpnička, P.; Lamač, M. "Synthesis and Catalytic Use of Planar Chiral and Polydentate Ferrocene Donors" In *Ferrocenes*; Štěpnička, P., Ed.; Wiley: Chichester, 2008; Chap. 7, p. 237-277. (e) Ogasawara, M.; Watanabe, S. "Transition-metal-Catalyzed Enantioselective Synthesis of Compounds with Non-centrochirality" *Synthesis* **2009**, 1761-1785. (f) Alba, A.-N. R.; Rios, R.

"Kinetic resolution: a Powerful Tool for the Synthesis of Planar-chiral Ferrocenes" *Molecules* **2009**, *14*, 4747-4757. (g) Schaarschmidt, D.; Lang, H. "Selective Syntheses of Planar-chiral Ferrocenes" *Organometallics* **2013**, *32*, 5668-5704. (h) Arae, S.; Ogasawara, M. "Catalytic Asymmetric Synthesis of Planar-chiral Transition-metal Complexes" *Tetrahedron Lett.* **2015**, *56*, 1751-1761.

(4) (a) Hayashi, T. "Asymmetric Catalysis with Chiral Ferrocenylphosphine Ligands" In *Ferrocenes*; Togni, A.; Hayashi, T., Eds.; VCH: Weinheim, 1995; Chap. 2, p. 105-142. (b) Togni, A. "New Chiral Ferrocenyl Ligands for Asymmetric Catalysis" In *Metallocenes*; Togni, A.; Halterman, R. L., Eds.; Wiley-VCH: Weinheim, 1998; Vol. 2, Chap. 11, p. 685-721. (c) Colacot, T. J. "A Concise Update on the Applications of Chiral Ferrocenyl Phosphines in Homogeneous Catalysis Leading to Organic Synthesis" *Chem. Rev.* **2003**, *103*, 3101-3118. (d) Barbaro, P.; Bianchini, C.; Giambastiani, G.; Parisel, S. L. "Progress in Stereoselective Catalysis by Metal Complexes with Chiral Ferrocenyl Phosphines" *Coord. Chem. Rev.* **2004**, *248*, 2131-3150. (e) Arrayás, R. G.; Adrio, J.; Carretero, J. C. "Recent Applications of Chiral Ferrocene Ligands in Asymmetric Catalysis" *Angew. Chem., Int. Ed.* **2006**, *45*, 7674-7715. (f) Fu, G. C. "Applications of Planar-chiral Heterocycles as Ligands in Asymmetric Catalysis" *Acc. Chem. Res.* **2006**, *39*, 853-860. (g) Ganter, C. "Planar Chiral Phosphaferrocene-based Ligands" In *Phosphorus Ligands in Asymmetric Catalysis*; Börner, A. Ed.; Wiley-VCH: Weinheim, 2008; Chapter 4.3, p. 393-407. (h) Dai, L.-X.; Hou, X.-L., Eds. *Chiral Ferrocenes in Asymmetric Catalysis*; Wiley-VCH: Weinheim, 2010.

(5) Butsugan, Y.; Araki, S.; Watanabe, M. "Enantioselective Addition of Dialkylzinc to Aldehydes Catalyzed by Chiral Ferrocenyl Amino Alcohols" In *Ferrocenes*; Togni, A.; Hayashi, T., Eds.; VCH: Weinheim, 1995; Chap. 3, p. 143-169.

(6) (a) Fu, G. C. "Enantioselective Nucleophilic Catalysis with "Planar-chiral" Heterocycles" *Acc. Chem. Res.* **2000**, *33*, 412-420. (b) Fu, G. C. "Asymmetric Catalysis with "Planar-chiral" Derivatives of 4-(Dimethylamino)pyridine" *Acc. Chem. Res.* **2004**, *37*, 542-547. (c) Fu, G. C. "Planar-chiral Heterocycles as Enantioselective Organocatalysts" In *Asymmetric Synthesis (2nd Edition)*; Christmann, M.; Brase, S., Eds.; Wiley-VCH: Weinheim, 2008; p 195-199. (d) Marion, N.; Fu, G. C. "Applications of Aza- and Phosphoferrocenes and Related Compounds in Asymmetric Catalysis" In *Chiral Ferrocenes in Asymmetric Catalysis*; Dai, L.-X.; Hou, X.-L., Eds.; Wiley-VCH: Weinheim, 2010; p 307-335.

(7) (a) Vedejs, E.; Chen, X. "Kinetic Resolution of Secondary Alcohols. Enantioselective Acylation Mediated by a Chiral (Dimethylamino)pyridine Derivative" *J. Am. Chem. Soc.* **1996**, *118*, 1809-1810. (b) Kawabata, T.; Nagato, M.; Takasu, K.; Fujii, K. "Nonenzymic Kinetic Resolution of Racemic Alcohols through an "Induced Fit" Process" *J. Am. Chem. Soc.* **1997**, *119*, 3169-3170. (c) Spivey, A. C.; Fekner, T.; Adams, H. "Synthesis of Atropisomeric Analogs of DMAP" *Tetrahedron Lett.* **1998**, *39*, 8919-8922. (d) Kotsuki, H.; Sakai, H.; Shinohara, T. "High-pressure Organic Chemistry. Part 23. A New Facile Method for the Synthesis of 4-Dialkylaminopyridine Derivatives by High-pressure-promoted Amination of 4-Chloropyridine" *Synlett* **2000**, 116-118. (e) Naraku, G; Shimomoto, N.; Hanamoto, T.; Inanaga, J. "Synthesis of Enantiomerically Pure  $C_2$ -symmetric 4-Pyrrolidinopyridine Derivative as a Chiral Acyl Transfer Catalyst for the Kinetic Resolution of Secondary Alcohols" *Enantiomer* **2000**, *5*, 135-138. (f) Shaw, S. A.; Aleman, P.; Vedejs, E. "Development of Chiral Nucleophilic Pyridine Catalysts: Applications in Asymmetric Quaternary Carbon Synthesis" *J. Am. Chem. Soc.* **2003**, *125*, 13368-13369. (g) Yamada, S.; Misono, T.; Iwai, Y. "Kinetic Resolution of *sec*-Alcohols by a New Class of Pyridine Catalysts Having a Conformation Switch System" *Tetrahedron Lett.* **2005**, *46*,

2239-2242. (h) Dálaigh, C. Ó.; Connon, S. J. "Nonenzymatic Acylative Kinetic Resolution of Baylis-Hillman Adducts" *J. Org. Chem.* **2007**, *72*, 7066-7069. (i) Kawabata, T.; Muramatsu, W.; Nishio, T.; Shibata, T.; Schedel, H. "A Catalytic One-step Process for the Chemo- and Regioselective Acylation of Monosaccharides" *J. Am. Chem. Soc.* **2007**, *129*, 12890-12895.

(8) (a) Wurz, R. P. "Chiral Dialkylaminopyridine Catalysts in Asymmetric Synthesis" *Chem. Rev.* **2007**, *107*, 5570-5595. (b) Furuta, T.; Kawabata, T. "Chiral DMAP-type Catalysts for Acyl-transfer Reactions" In *Science of Synthesis: Asymmetric Organocatalysis I*; List, B., Ed.; Georg Thieme KG: Stuttgart, 2012; p 497-546.

(9) (a) Ruble, J. C.; Fu, G. C. "Chiral  $\pi$ -Complexes of Heterocycles with Transition Metals: A Versatile New Family of Nucleophilic Catalysts" *J. Org. Chem.* **1996**, *61*, 7230-7231. (b) Ruble, J. C.; Latham, H. A.; Fu, G. C. "Effective Kinetic Resolution of Secondary Alcohols with a Planar-Chiral Analog of 4-(Dimethylamino)pyridine. Use of the Fe(C<sub>5</sub>Ph<sub>5</sub>) Group in Asymmetric Catalysis" *J. Am. Chem. Soc.* **1997**, *119*, 1492-1493. (c) Ruble, J. C.; Fu, G. C. "Enantioselective Construction of Quaternary Stereocenters: Rearrangements of O-Acylated Azlactones Catalyzed by a Planar-chiral Derivative of 4-Pyrrolidinopyridine" *J. Am. Chem. Soc.* **1998**, *120*, 11532-11533. (d) Tao, B.; Ruble, J. C.; Hoic, D. A.; Fu, G. C. "Nonenzymatic Kinetic Resolution of Propargylic Alcohols by a Planar-chiral DMAP Derivative: Crystallographic Characterization of the Acylated Catalyst" *J. Am. Chem. Soc.* **1999**, *121*, 5091-5092. (e) Bellemin-Lapponnaz, S.; Tweddell, J.; Ruble, J. C.; Breitling, F. M.; Fu, G. C. "The Kinetic Resolution of Allylic Alcohols by a Non-enzymatic Acylation Catalyst; Application to Natural Product Synthesis" *Chem. Commun.* **2000**, 1009-1010. (f) Hodous, B. L.; Fu, G. C. "Enantioselective Staudinger Synthesis of  $\beta$ -Lactams Catalyzed by a Planar-chiral Nucleophile" *J. Am. Chem. Soc.* **2002**, *124*, 1578-1579. (g) Hodous, B. L.; Fu, G. C. "Enantioselective Addition of Amines to Ketenes

Catalyzed by a Planar-chiral Derivative of PPY: Possible Intervention of Chiral Brønsted-acid Catalysis" *J. Am. Chem. Soc.* **2002**, *124*, 10006-10007. (h) Wilson, J. E.; Fu, G. C. "Lactone Synthesis: Asymmetric Synthesis of Highly Substituted  $\beta$ -Lactones by Nucleophile-catalyzed [2+2] Cycloadditions of Disubstituted Ketenes with Aldehydes" *Angew. Chem., Int. Ed.* **2004**, *43*, 6358-6360. (i) Wiskur, S. L.; Fu, G. C. "Catalytic Asymmetric Synthesis of Esters from Ketenes" *J. Am. Chem. Soc.* **2005**, *127*, 6176-6177. (j) Schaefer, C.; Fu, G. C. "Catalytic Asymmetric Couplings of Ketenes with Aldehydes to Generate Enol Esters" *Angew. Chem., Int. Ed.* **2005**, *44*, 4606-4608. (k) Berlin, J. M.; Fu, G. C. "Enantioselective Nucleophilic Catalysis: the Synthesis of Aza- $\beta$ -lactams through [2+2] Cycloadditions of Ketenes with Azo Compounds" *Angew. Chem., Int. Ed.* **2008**, *47*, 7048-7050. (l) Dochnahl, M.; Fu, G. C. "Catalytic Asymmetric Cycloaddition of Ketenes and Nitroso Compounds: Enantioselective Synthesis of  $\alpha$ -Hydroxycarboxylic Acid Derivatives" *Angew. Chem., Int. Ed.* **2009**, *48*, 2391-2393. (m) Lee, S. Y.; Murphy, J. M.; Ukai, A.; Fu, G. C. "Nonenzymatic Dynamic Kinetic Resolution of Secondary Alcohols via Enantioselective Acylation: Synthetic and Mechanistic Studies" *J. Am. Chem. Soc.* **2012**, *134*, 15149-15153.

(10) (a) Chen, Y.-H.; McDonald, F. E. "New Chiral Synthons for Efficient Introduction of Bispropionates via Stereospecific Oxonia-Cope Rearrangements" *J. Am. Chem. Soc.* **2006**, *128*, 4568-4569. (b) Kim, J.; Ashenurst, J. A.; Movassaghi, M. "Total Synthesis of (+)-11,11'-Dideoxyverticillin A" *Science* **2009**, *324*, 238-241. (c) Francais, A.; Leyva, A.; Etxebarria-Jardi, G.; Ley, S. V. "Total Synthesis of the Anti-apoptotic Agents Iso- and Bongkrekiac Acids" *Org. Lett.* **2010**, *12*, 340-343. (d) DeLorbe, J. E.; Jabri, S. Y.; Mennen, S. M.; Overman, L. E.; Zhang, F.-L. "Enantioselective Total Synthesis of (+)-Gliocladiine C: Convergent Construction of Cyclotryptamine-Fused Polyoxopiperazines and a General Approach for Preparing

Epidithiodioxopiperazines from Trioxopiperazine Precursors" *J. Am. Chem. Soc.* **2011**, *133*, 6549-6552.

(11) Wurz, R. P.; Lee, E. C.; Ruble, J. C.; Fu, G. C. "Synthesis and Resolution of Planar-chiral Derivatives of 4-(Dimethylamino)pyridine" *Adv. Synth. Catal.* **2007**, *349*, 2345-2352.

(12) (a) Spivey, A. C.; Leese, D. P.; Zhu, F.; Davey, S. G.; Jarvest, R. L. "New Atropisomeric Biaryl Derivatives of 4-Aminopyridine - Identification of an Improved Nucleophilic Catalyst for Asymmetric Acylation of *sec*-Alcohols" *Tetrahedron* **2004**, *60*, 4513-4525. (b) Mittal, N.; Sun, D. X.; Seidel, D. "Kinetic Resolution of Amines via Dual Catalysis: Remarkable Dependence of Selectivity on the Achiral Cocatalyst" *Org. Lett.* **2012**, *14*, 3084-3087. (c) Tandon, R.; Nigst, T. A.; Zipse, H. "Inductive Effects through Alkyl Groups - How Long is Long Enough?" *Eur. J. Org. Chem.* **2013**, 5423-5430.

(13) Ogasawara, M.; Wada, S.; Isshiki, E.; Kamimura, T.; Yanagisawa, A.; Takahashi, T.; Yoshida, K. "Enantioselective Synthesis of Planar-chiral Ferrocene-fused 4-Pyridones and Their Application in Construction of Pyridine-based Organocatalyst Library" *Org. Lett.* **2015**, *17*, 2286-2289.

(14) Snieckus, V. "Directed Ortho Metalation. Tertiary Amide and O-Carbamate Directors in Synthetic Strategies for Polysubstituted Aromatics" *Chem. Rev.* **1990**, *90*, 879-933.

(15) (a) Riant, O.; Samuel, O.; Kagan, H. B. "A General Asymmetric Synthesis of Ferrocenes with Planar Chirality" *J. Am. Chem. Soc.* **1993**, *115*, 5835-5836. (b) Riant, O.; Samuel, O.; Flessner, T.; Taudien, S.; Kagan, H. B. "An Efficient Asymmetric Synthesis of 2-Substituted Ferrocenecarboxaldehydes" *J. Org. Chem.* **1997**, *62*, 6733-6745. (c) Mamane, V. "The



Diastereoselective Ortho-lithiation of Kagan's Ferrocenyl Acetal. Generation and Reactivity of Chiral 2-Substituted Ferrocenecarboxaldehydes" *Tetrahedron: Asymmetry* **2010**, *21*, 1019-1029.

(16) (a) Geisler, F. M.; Helmchen, G. "A Straightforward Synthesis of (3*S*)-4-Methoxybutane-1,3-diol and Its Use as Chiral Auxiliary for the Preparation of (p*S*)-1-(Diphenylphosphino)-2-formyl-1',2',3',4',5'-pentamethylferrocene" *Synthesis* **2006**, 2201-2205. (b) Wölfle, H.; Kopacka, H.; Wurst, K.; Ongania, K.-H.; Görtz, H.-H.; Preishuber-Pflügl, P.; Bildstein, B. "Planar Chiral Ferrocene Salen-type Ligands Featuring Additional Central and Axial Chirality" *J. Organomet. Chem.* **2006**, *691*, 1197-1215.

(17) Mroczek, A.; Erre, G.; Taras, R.; Gladiali, S. "Chiral Ferrocenyl Ligands with Bidentate Pyridine Donors. Synthesis and Application in Pd-catalyzed Asymmetric Allylic Alkylation of 1,3-Diphenylpropenyl-1-esters" *Tetrahedron: Asymmetry* **2010**, *21*, 1921-1927.

(18) Hu, H.; Wu, W.-Y.; Takahashi, T.; Yoshida, K.; Ogasawara, M. "Enantioselective Synthesis of Ferrocene- or Cymantrene-Fused Planar-chiral Phospholes" *Eur. J. Inorg. Chem.* **2017**, 325-329.

(19) (a) Garber, S. B.; Kingsbury, J. S.; Gray, B. L.; Hoveyda, A. H. "Efficient and Recyclable Monomeric and Dendritic Ru-based Metathesis Catalysts" *J. Am. Chem. Soc.* **2000**, *122*, 8168-8179. (b) Hoveyda, A. H.; Gillingham, D. G.; Van Veldhuizen, J. J.; Kataoka, O.; Garber, S. B.; Kingsbury, J. S.; Harrity, J. P. A. "Ru Complexes Bearing Bidentate Carbenes: from Innocent Curiosity to Uniquely Effective Catalysts for Olefin Metathesis" *Org. Biomol. Chem.* **2004**, *2*, 8-23.

(20) Butler, D. C. D.; Richards, C. J. "Synthesis of 1'-Substituted Derivatives of 1,2,3,4,5-Pentaphenylferrocene" *Organometallics* **2002**, *21*, 5433-5436.

(21) Dai, X.; Nakai, T.; Romero, J. A. C.; Fu, G. C. "Enantioselective Synthesis of Protected Amines by the Catalytic Asymmetric Addition of Hydrazoic Acid to Ketenes" *Angew. Chem., Int. Ed.* **2007**, *46*, 4367-4369.

(22) (a) Tamano, M.; Nagai, Y.; Koketsu, J. "Titanium Tetrachloride-mediated Reactions of Aromatic Ketones with Amines" *Nippon Kagaku Kaishi* **1988**, 1977-1983. (b) Kořínek, M.; Rybáčková, M.; Böhm, S. "The Use of *N,N*-dialkyltrimethylsilylamines in the Synthesis of Pentamethinium Salts" *Synthesis* **2009**, 1291-1296.

(23) A literature search revealed that similar synthetic methods for 4-quinolones have been reported. (a) Torii, S.; Okumoto, H.; Xu, L. H. "Palladium-catalyzed Carbonylation to form 2-Substituted 1,4-Dihydro-4-oxoquinoline" *Tetrahedron Lett.* **1991**, *32*, 237-240. (b) Torii, S.; Okumoto, H.; Xu, L. H.; Sadakane, M.; Shostakovsky, M. V.; Ponomaryov, A. B.; Kalinin, V. N. "Syntheses of Chromones and Quinolones via Palladium-catalyzed Carbonylation of *o*-Iodophenols and Anilines in the Presence of Acetylenes" *Tetrahedron* **1993**, *49*, 6773-6784. (c) Ward, T. R.; Turunen, B. J.; Haack, T.; Neuenswander, B.; Shadrack, W.; Georg, G. I. "Synthesis of a Quinolone Library from Ynones" *Tetrahedron Lett.* **2009**, *50*, 6494-6497. (d) Seppänen, O.; Muuronen, M.; Helaja, J. "Gold-Catalyzed Conversion of Aryl- and Alkyl-Substituted 1-(*o*-Aminophenyl)-2-propyn-1-ones to the Corresponding 2-Substituted 4-Quinolones" *Eur. J. Org. Chem.* **2014**, 4044-4052. (e) Seki, H.; Georg, G. I. "2,3-Dihydropyridin-4(1H)-ones and 3-Aminocyclohex-2-enones: Synthesis, Functionalization, and Applications to Alkaloid Synthesis" *Synlett* **2014**, *25*, 2536-2557. (f) Ai, W.; Wu, Y.; Tang, H.; Yang, X.; Yang, Y.; Li, Y.; Zhou, B. "Rh(III)- or Ir(III)-catalyzed Ynone Synthesis from Aldehydes via Chelation-assisted C-H Bond Activation" *Chem. Commun.* **2015**, *51*, 7871-7874.

(24) CCDC 1860884 [(*S*)-(-)-**2c**] and 1860885 [(*S*)-(-)-**1dw**] contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; +44 1223 336033.

(25) Ferber, B.; Top, S.; Jaouen, G. "Introduction of a Planar Chirality onto Steroid Substrates: Synthesis of (*S*) and (*R*)-2'-Formylcymantrenyl-17 $\alpha$ -ethynylestradiols Using (*S*)- and (*R*)-1-Formyl-2-iodo-cymantrenes" *J. Organomet. Chem.* **2004**, *689*, 4872-4876.

(26) Broadley, K.; Lane, G. A.; Connelly, N. G.; Geiger, W. E. "Electrochemical Routes to Paramagnetic Dinuclear and Mononuclear Palladium  $\pi$  Complexes Stabilized by the Pentaphenylcyclopentadienyl Ligand" *J. Am. Chem. Soc.* **1983**, *105*, 2486-2487; b) Koelle, U.; Khouzami, F. "Permethylated Electron-Excess Metallocenes" *Angew. Chem. Int. Ed. Engl.* **1980**, *19*, 640-641.

(27) (a) Kagan, H. B.; Fiaud, J. C. "Kinetic Resolution" *Top. Stereochem.* **1988**, *18*, 249-330.  
(b) Vedejs, E.; Jure, M. "Efficiency in Nonenzymatic Kinetic Resolution" *Angew. Chem., Int. Ed.* **2005**, *44*, 3974-4001.

## Table of Contents/Abstract Graphics

