Computer Charting: An Evaluation of a Respiratory Care Computer System

Robert D Andrews MS M T (ASCP), Reed M Gardner PhD, Sandy M Metcalf RRT, and Deon Simmons RRT

A respiratory care computer-charting system was developed and implemented as an addition to our hospital's computerized information system. Medical personnel charted and reviewed respiratory care procedures at nursing station computer terminals instead of using the patient's traditional paper chart. The computer automatically performed billing and provided management as well as clinical information. In an attempt to isolate specific benefits or shortcomings, we evaluated charting systems both before and after computer implementation. Four assessments were made: (1) a survey of therapists' attitudes, (2) an observation of work patterns, (3) an audit of the content of charting, and (4) an analysis of productivity statistics. Computer charting was well accepted by therapists. Charge capture was reduced from a four-step manual process to a single-step computer documentation of the procedure. Computer charting was more complete and informative. Productivity increased 18% although it remains unclear to what degree the computer was responsible. Computer charting streamlined the process of documentation and allowed more beneficial use of clinical information. (Respir Care 1985;30:695-707.)

Introduction

In efforts to increase the efficiency of medical care delivery, institutions are turning to computers as useful tools for processing and storing medical, financial, and administrative information. It has been reported that 25 to 35% of a health professional's time is spent doing paperwork,¹⁻³ and although many hospital departments have computerized information systems, the clinical information in the patient's chart remains essentially unchanged.⁴⁻⁶ This clinical information includes patient history, observations, medications, and progress notes used in diagnosis and treatment. The documentation of most procedures in respiratory care (RC) is similar in content. We report the usefulness of a computer-charting system in documenting and processing clinical information.

An Optimal System

The efficiency of any system is measured by the 'useful' work completed compared to the energy required. The most efficient RC computer system would have the following characteristics:

- No repetition of work or reporting
- Easy access for entry and review
- Accurate and descriptive documentation
- Automatic performance of many functions from a single input (i.e., billing, reporting, checking for errors, alerting, and gathering of management statistics)

Mr Andrews was a graduate student, Department of Medical Biophysics and Computing, University of Utah, Salt Lake City, when this study was performed. Dr Gardner is Professor of Medical Biophysics and Computing, LDS Hospital/University of Utah, Salt Lake City. Ms Metcalf is Associate Director and Ms Simmons is Assistant Director, Department of Respiratory Care, LDS Hospital. Work done at LDS Hospital.

Reprints: Reed M Gardner PhD, Biophysics, LDS Hospital, 325 8th Ave, Salt Lake City UT 84143.

- Exact correlation between charting and billing
- Integration of RC information with that of other hospital departments
- Availability of information for diagnostic and research purposes
- Easy implementation
- Reliability (no down time)
- Inexpensive equipment that pays for itself

Perhaps the best proof of a computer's usefulness is the degree to which people want to use it because it helps them do their jobs, not simply because its use is mandatory.

Institutional Background

LDS Hospital

LDS Hospital, a major referral center with 520 beds and 5 (4 adult, 1 newborn) intensive care units (ICUs), has been a leader in the development of computer applications in medicine. A highly developed Hospital Information System (HIS), known as HELP, integrates all patient information.^{7,8} A Tandem "non-stop" computer system (Tandem, Cupertino CA) is connected to more than 300 terminals and 95 printers. It is highly reliable and has little downtime (0.2%)⁹ because of its redundant processing and storage of data.¹⁰ The computer has an integrated central billing system. The functions of order entry, reporting, data entry, and alerting are well developed for most departments. At least four terminals are available on every nursing division (each of which handles 48 patients), as is a printer. The ICUs have a terminal at each bedside.

Respiratory Care Department

Respiratory care presented several unique problems for computer implementation. By 1982 only about a dozen RC departments in the country had reached a level of substantial computerization; an equal number of departments bad tried, but failed.¹¹ At LDS Hospital we introduced computer charting as an improvement on the written patient chart and to meet the clinical, financial, and management needs of RC.

The RC service is highly mobile. Therapists do not have a permanent workstation, as work is performed at the bedside and throughout the hospital. Therefore, entering computer information required having access to terminals En many locations or recording information on paper for later computer entry in the RC department. Thus, the logistics problems of where the data could be reviewed and how it could be entered in the patient's chart had to be solved.

Patient records vary in quality and detail because from one third to one half of them are in narrative which makes information difficult to collect and process.^{5, 12-14} Unlike computerized systems in clinical laboratories that process large amounts of numeric data, computerized RC information systems require a reporting 'vocabulary' with a wide range of descriptions. To be automated, patient records had to be converted from a narrative format to the computer's predefined vocabulary.⁶


The RC computer system was developed from a very simple concept: "Chart accurately and let the computer do the rest of the paperwork." The system was designed to maximize the efficiency of documenting procedures and thereby improve the evaluation of medical care. In addition, documentation was required for hospital accreditation¹⁵ and for verification that a procedure had been performed. The charting of clinical procedures was also used in non-medical

functions, such as management statistics and billing. Because the functions were integrated into the HIS, they became by products of the documentation process.¹⁶ As paperwork was reduced, a higher percentage of the therapist's time could be spent doing the most useful work, patient care.

Respiratory care documentation has traditionally been written into the patient's chart using specific forms-those for notes, assessments, and ventilator monitoring—with each section organized chronologically. Documentation has allowed later review so that patient care can be assessed and changed if necessary. These processes of data entry, organization, storage, and review are very similar to the operation of a computer. To permit the computer to be used for patient charting, three programming functions of the HIS were instrumental: (1) One program allowed creation of questionnaires, to be used for data entry. This program also permitted the capture of billing information. (2) Another program allowed the creation of vocabulary used in charting by assigning the medical terminology to codes that were more easily stored in the computer's files. (3) A general reporting language was used to program the reports and statistics.

Description of the RC System

The RC computer system is a subsystem of the HIS; it depends on the central computer and uses nursing division terminals for data entry and review. It avoids duplication by using existing hardware and by using information from other hospital departments, such as admission, discharge, and transfer (ADT) information. The HIS controls and processes the flow of all patient information (Fig. 1). RC charting is entered at the nursing divisions, is stored in patient



data files and can be reviewed at any nursing division terminal. A 24-hour management report provides individual and departmental productivity records, and an alert report is used for both management and patient care monitoring. Permanent copies of all RC charting are automatically

processed for delivery to Medical Records after a patient has been discharged. The HIS is integrated with a billing computer system that processes financial transactions and provides the hospital with productivity reports. Thus, all reporting and billing are extracted directly from the computerized clinical charting.

Currently the RC department is not fully computerized—order entry, workload allocation, and newborn nursery charting are still done manually. The charting of ventilator data was recently implemented, because bedside terminals axe now available in the ICUs where ventilators are used. Approximately 90% of RC charting and charge capture is now computerized.

Charting

The charting process is initiated by selecting the "Respiratory Therapy Charting" option on the computer terminal at the nursing station. Entries are made by selecting multiple-choice items from the menu, by number entry, or by typing in free text (Fig. 2). The questionnaire-entry format follows a logical sequence that corresponds with the department's charting requirements. Entries can include the charting of more than one procedure at a time, which allows procedures that are frequently done together to be charted without redundant questions and multiple data entries. To speed the process, only questions pertinent to the specific procedure are asked.




Fig. 2. Computer charting uses menu selection for entering results of RC procedures.

Follow-up questions are also specific to certain entries; this results in a highly variable pathway that allows flexibility yet decreases the time required for data entry. The only questions to which answers are mandatory are those pertaining to medical-legal or billing issues; most questions can be left unanswered, allowing the therapist to chart only that which is necessary. The therapist is responsible for complete and accurate charting. A procedure attempted but not completed is also

documented in order to verify that an attempt was made and to explain why it was not done. All entries require an employee identification number, which serves as an electronic 'signature.'

Review of Charting

The review of charting is available by using a review option on any hospital terminal. This option is on the same menu for review of laboratory, blood gas, and radiology results. Because results can be reviewed from any terminal, it is not necessary to be on a particular ward to obtain a patient's chart. The report is a text report (Fig. 3) that resembles written entries (Fig. 4).

08/18/84.17:00 -02- 02 INTERFACE: CANNULA; FLOW RATE: 5.0 L/MIN; 8 HRS ON 02;
THPST#/DUR/ENTRY: 32000/ 10/19:16
08/13/84.15:45 -MEDICATION MEBULIZERCPT- PATIENT INTERFACE: MOUTHPIECE: POSITION: DANGLE: MEDICATION: ALUPENT, 10 MG. DILUENT: NORMAL SALINE: PRE BREATH SOUNDS: RHONCHI, THROUGHOUT INSPIRATION: MECHANICAL PERCUSSION: FOR 20 MIN, LEFT/FLAT, RIGHT/FLAT, BOTH LUNDS; COUGH: WHEN ENCOURAGED, MODERATE, PRODUCTIVE: PATIENT CONDITION: ALERT, COOPERATIVE: POST BREATH SOUNDS: IMPROVED; HEART RATE: 76/ / 80 RESP RATE: 8/ /10. THPSTM/DUR/ENTRY: 46547/ 30/16:44
08/18/84.11:40 -PULMONARY EXERCISEMEDICATION NEBULIZERCPT- PATIENT INTERFACE: MOUTHFIECE: POSITION: SITTING: MEDICATION: ALUPENT, 15 MG, DILUENT: NORMAL SALINE: PRE BREATH SOUNDS: UMEEZING, THROUGHOUT INSPIRATION, BOTH LUNGS; IC: 1000 CC; 10 BREATHS DURING PLUMONARY EXERCISE: MECHANICAL PERCUSSION: FOR 20 MIN, TREMDELENBURG/KIGHT, TRENCELENBURG/LEFT, BLL: CUOGH: FREQUENT, STRONG, PRODUCTIVE; SPUTUM: SPONTAMEOUS, QUANTITY: 3 CC, YELLOW, THICY: PATIENT CONDITION: ALERT, COOPERATIVE; POST BREATH SOUNDS: IMPROVED; COMMENT: RX TOL WELL; HEART RATE: 847 847 84 RESP RATE: 247 /24 THPSTM/DUR/ENTRY: 465377 60/13:17
08/18/84.11:30 -MEDICATION NEBULIZER- PATIENT INTERFACE: MOUTHPIECE; POSITION: DANGLE; MEDICATION: ALUPENT, 15 MG, DILUENT; NORMAL SALINE; PRE BREATH SOUNDS: WHEEZING, THROUGHOUT EXPIRATION, BOTH LUNGS; COUGH: SPONTAMEOUS, STRONG, NON-PRODUCTIVE; PATIENT CONDITION: ALERT, COOPERATIVE; POST BREATH SOUNDS: UNCLANNEDE: COMMENT: RX TOL WELL; HEART RATE: 78/ / 80 THPST0/DUR/ENTRY: 46537/ 20/11:34
Fig. 3. An example of computer charting.
ng. o. An example of computer charting.
1/2/2/000 At nasally intubated on Bear-1
Vent OPAP 10 cmt, O BS Ad in haves on inspiration
Kill 3ce NS installed followed by leve yellow
patum sy.
-1-29-E4- 7:30 MEds / NE - BRAN VENT - 15 Mg Alup/NS -
1+R-96-92-94- Interved by PERCEPT TO BLL'S = DT
Side/Tarad - So See This yellow sacrations.
H-29.84 1532 mide you live to rent 15 mg Mus MM -
Ha-98 91-92: Infloored by Constant to at a
A Side Prone Mr. 8" then allow
secretions
4129/04 1945 MASS IN LINE E ISWA AL DENT MS VIA B- WENT.
14 140/136/140 74/201712 BS-1 proce & Roughing OT MAP
Bill's StT x 15 mond. 5xn = 1 Pic thick vellar sputime

Fig. 4. An example of manual charting.

Automatic Routine Reporting

Every morning at 03:00 a program automatically generates three routine reports for the RC department: (1) a complete printout of RC charting on patients discharged the previous day, (2) a 24-hour management report, and (3) an alert report. These three reports are the only hard-copy printouts that are automatically generated routinely by the RC system. This early morning

use of the computer is efficient and provides information that can be assessed by supervisors at the beginning of the day.

The 24-hour management report lists the work that has been charted for that period by each therapist (Fig. 5). The report identifies the patient, work units, and duration of each procedure. It is a record of each therapist's productivity. Supervisors review the report to confirm that assigned procedures were completed, so that missed procedures or missed charting can be identified and corrected. The management report also provides a department summary, listing a breakdown of total procedures performed and the reasons when treatments were not completed (Fig. 6). The 24-hour report provides management data extracted directly from patient charting and forms the basis for long-term individual and departmental reports.

COMPUTER CHARTING EVALUATION

RESPIRATORY CARE THERAPIST REPORT 17 AUG 1984 24 HOUR MANAGEMENT REPORT

THERAPIST #		46482	LINDA STANCHFIELD	AUG 17, 1984 00:00 - AUG	17, 1984	23:59	i			
DATE	TIME	ROOM	PATIENT	TREATMENTS	RVU'S	UNITS	DUR	PC TA	CT	ENTRY
08-17 08-17	12:00 12:00	E509 E509	PATIENT, A PATIENT, A	MEDICATION NEBULIZER NEBULIZER MONITORING HPN EQUIPMENT SET UP ON 02	23.24 2.39 20.51 .67	1 1	30 10	154 128	4.5 1.5	15:00 14:17
08-17 08-17 08-17 08-17	08:00 07:30 07:30 07:00	E509 E509	PATIENT, A PATIENT, A PATIENT, A PATIENT, B	NEDICATION NEBULIZER NEBULIZER MONITORING 02. NEBULIZER MONITORING	23.24 16.73 4.69 19.12	1 7 7 8	20 10 10 10	209 402 396 419	3.5 1.5 1.5 1.5	11:46 14.21 14:15 14:08
08-17	13:45	W623	PATIENT, C	ON 02 PULMONARY EXERCISE MEDICATION NEBULIZER	5.36 15.04 23.24	1	20	33	3.5	14:35
08-17	09:30	₩623	PATIENT, C	PULMONARY EXERCISE MEDICATION NEBULIZER	15.04 23.24	1	20	284	2.5	14:32
08-17	13:30		PATIENT, D	MEDICATION NEBULIZER	: 0	0	5 5	83 191	1.5 1.5	14:57 14:55
08-17			PATIENT, D PATIENT, D	MEDICATION NEBULIZER ## NOT DONE ## NOT ON UNIT MEDICATION NEBULIZER	23.24	1	20	225	1.5	11:44
08-17 08-17	07:00 12:30	E514	PATIENT, D PATIENT, E PATIENT, F	02 IPPB	5.36 27.34	8 1	10 35	423 108	1.5 4.5	14:12 14:49
08-17	10:30	E512	PATIENT, F	CPT IPPB CPT	24.61 27.34 24.61 27.34	1	35	224	4.5	14:45
08-17 08-17	08:30 07:30) E512) W629	PATIENT, F PATIENT, G	IPPB NEBULIZER MONITORING ON 02	27.34 19.12 5.36	1 8	20 10	355 405	4.5	14:41 14:23
08-17 08-17			PATIENT, H PATIENT, I	NEBULIZER MONITORING MEDICATION NEBULIZER	19.12 23.24		10 20	408 416	1.5 2.5	14:27 14:29
				POINTS: 14.0 TOTALS: AMERAGES:	419.19		300	0 270.2	46.0 2.6	

Fig. 5. The 24-hour management report provides a record of all procedures documented by each therapist.

The alert report (Fig. 7) is used to monitor for both management and medical errors. The listing for Patient B is an example of a management alert to an overcharge resulting from double charting. If hourly therapy, such as oxygen, is documented for more than 24 hours in a single day, an alert is printed so that the charting and billing can be corrected. A medical alert might indicate a need far closer patient assessment. If a patient is on continuous oxygen therapy for a prolonged period of time and has never had a blood gas test, an alert is printed. Alert capability will be expanded to include the monitoring of medical necessity protocols.^{17, 18}

Billing

Billing is an automatic by-product of the computer charting of a completed procedure. An example of a therapist's chart is shown in Figure 3. This documentation of oxygen therapy results in a bill for 8 hours of oxygen. The next treatment shows medications-nebulizer therapy and chest physical therapy (CPT), which are billed. Everything is charted for clinical reasons, and the program automatically bills when appropriate. Treatments ordered but not done are reported in the chart but are not billed. Thus, billing accuracy depends on the therapist's charting accuracy. Mistakes can still occur, such as charting the wrong patient or charting the same procedure twice.




Fig. 6. The 24-hour management report also provides a departmental summary of procedures performed and the reasons when procedures were not completed.

These errors can be found easily by therapists as they review the charting or by supervisors as they review the 24-hour management report, and the errors can be easily corrected by supervisory personnel. Billing accuracy is not merely of concern to the hospital and patient, but also determines RC productivity, which is used to justify the staffing requirements of the RC department. The 24-hour management report determines the individual therapist's productivity as well as that of the RC department as a whole.

```
4NB4 PATIENT, A
*** NO BLOOD GAS IN LAST 4 DAYS ***
4NB9 PATIENT, B
$$$ 2 DAYS AGO > 24 HRS 02 CHARGES / DAY $$$
*** CONTINUOUS 02 DISCONTINUED OR INTERRUPTED YESTERDAY ***
4NB7 PATIENT, C
*** NO BLOOD GASES ***
```

Fig. 7. Alert report identifies possible errors and oversights in computer charting and patient care.

Evaluation Methods

The RC computer system was evaluated in four ways: (1) therapists' appraisal, (2) observation of work patterns, (3) audit of the quality and content of charting, and (4) productivity analysis. E x h evaluation was made before computer charting (PRE) and after computer charting (POST).

Therapists' Appraisal

Questionnaires were distributed to the therapists (63 PRE and 55 POST) to be filled out anonymously 2 months before and 2 months after the establishment of computer charting (March 1984). The questionnaires were used to determine therapists' expectations, problems, suggestions, and preferences.

Work Patterns

PRE and POST individual work patterns were compared. After 2 months of computer charting, an inquiry of head nurses and ward clerks was made to obtain feedback on possible interference or congestion at nursing station terminals. The department. Managers of both Billing and Medical Records were also interviewed.

Quality and Content of Charting

We compared the quality and content of computer charting against manual documentation by auditing medications-nebulizer therapy, one of the most common RC procedures. Guided by departmental standards for this treatment, we checked documentation for inclusion of (1) therapist signature, (2) medications delivered, (3) comments (patient's condition, effects of therapy, and adverse reactions), (4) changes in breath sounds, (5) heart rate before and after treatment, (4) sputum production, (7) cough effort, and (8) patient position. Chart legibility was also evaluated. For this study, patients' charts were selected at random before and after implementation of computer charting. Five hundred manually charted procedures (performed on 22 patients by 49 therapists) were evaluated for content and quality and compared to 500 computer-charted procedures (performed on 29 patients by 51 patients). The only item that was a mandatory entry on the computer was 'therapist signature.'

Productivity

PRE and POST statistics of work volume and productivity were compared for all procedures preformed by the RC department during a 6-month period (February through July 1984). Four PRE pay periods (the 8 weeks preceding computer implementation) were compared to the first 8 pay periods (167 weeks) of POST data. Hospital data on productivity and work volume were generated from procedures billed; RC department data were generated from the supervisors' accounts of completed work assignments. These two sources were evaluated with regard to changes in productivity and work volume. An unpaired t test was used fur comparison of PRE and POST data.

Results of Evaluation

Therapists' Appraisal

Questionnaires returned by the therapists (49 PRE and 50 POST) indicated job position, location, and shift worked. Virtually all therapists were familiar with the use of computer terminals for reviewing information (96% had used a hospital. terminal before), and it took only

about 3 days for most of them to feel comfortable doing computer charting. Results of the questionnaires are presented in Table 1. Of the 50 therapists who returned the POST survey, 32 (64%) favored computer charting, compared to 10 (20%) who preferred manual charting.

Work Patterns

Computer charting reduced a four-step process- charting the procedure, filling out a charge slip, processing the charge slip and transferring it to billing, and posting the charges into the computer-to only one step--computer charting the procedure. The secretary's job was changed from that of processing charges to auditing billing mistakes and making sure that all printouts of discharged patients were delivered to Medical. Records. Shift supervisors generally had about 30 minutes added to their workload as a result of reviewing the 24-hour management report. Entering billing charges in the Kardex system was eliminated, which, according to estimates from the Industrial Engineering Department, saved each therapist 10 minutes a day. Many therapists felt that charting was faster using the computer.

Other departments affected by the computer were Nursing, Billing, and Medical Records. Access to nursing station terminals was not found to be a major problem. Occasionally problems resulted if a therapist entered several procedures at once and deprived others of access to the terminal. Because computer charting completely bypassed the Billing Department, posting RC charges was eliminated; this saved the Billing Department about 30 minutes per day. The Medical Records Department agreed to put the patient reports onto the patient's chart; this added about 30 minutes of work per day in this department. The net result of RC computer charting on other departments was one of redistribution of effort, with no major overall change.

Quality and Content of Charting

Computer charting was found to be more complete than manual charting in every case except the documentation of medication, which remained the same (Fig. 8). Both the manual and computer charts had four instances (0.8%) in which the medication was not specified. Legibility and signature were both 100% on the computer. Figures 3 and 4 illustrate the difference in legibility between computer and manual charting. It was noted that not only was there an improvement in meeting the department's requirements for charting, but often the requirements were exceeded. Computer charting was found to be more informative, concise, and compact.

Productivity

Productivity data are presented in Table 2. Significant ($P \le 0.03$) increases after computer charting was instituted are shown for both productivity and work volume. Hospital data calculated from billed procedures showed that productivity increased 18.2%; RC records showed that productivity (average workload completed per therapist) increased an average of 13.7%. Hospital data showed that work volume increased 20.9%, while RC department records showed that it increased 16.4%. The number of employees who worked during both periods was not significantly different (51.23 PRE vs. 52.40 POST).

١.	Approximat	ely how many i	minutes do you spe	end in chart	ing a treatm	ent?					
	PRE	<2 min 2 (4%)	2-5 min 27 (55%)	6- 19	10 min (39%)	n > 10 min		No Response			
	POST	11 (22%)	30 (60%)	7	(14%)	1	(2%)	1 (2%)			
	(POST) How does computer charting time compare to charting manually?										
	Faster		the same	Longer			No response				
	18 (36%) 21 (42%)			9 (18%)		2 (4%)					
	(POST) To do the same amount of work for your job, how much time has the computer saved or added?										
				# Response		Total Mir					
	Min/shift th	ne computer ha	s saved	16		414					
	Min/shift th	e computer has	s added	9		345					
•	How many	times during a sl	hift do you use a h	ospital term	inal?						
		$<\!2$ min	2-5 min	6-10 min		> 10 min		No response			
	PRE	3 (6%)	10 (20%)	19 ((39%)	14 ((29%)	3 (6%)			
	POST	2 (4%)	6 (12%)	13 ((26%)	28 ((56%)	1 (2%)			
	(POST) For	r the following a	ispects, how do yo	u feel comp	uter chartin	g compare	s to manual ch	arting?			
			Much			About		Much	No		
			Better	Better	1	The Same	Worse	e Worse	Respons		
	Quality of ti		7 (14%)	21 (42		16 (32%)	2 (4	%) 2 (4%)	2 (4%)		
	Ease of ente		12 (24%)	19 (38)		12 (24%)	5 (10	%)	2 (4%)		
	Ease of revi	ew:	8 (16%)	17 (34	121610	8 (16%)	14 (28		2 (4%)		
	Accuracy: Productivity	2	12 (24%)	14 (28)		13 (26%)	8 (16				
	Productivity		13 (26%)	24 (48)	200	9 (18%)	2 (4	%)	2 (4%)		
	How often do you have trouble getting access to a terminal on the ward?										
	Very Rarely		Occasion					s No R	esponse		
	PRE POST	17 (35%) 23 (46%)	27 (55	200	2 (4%)				(6%)		
		8 8	22 (44		2 (4%)		1 (2%)	2	(4%)		
			ave trouble getting	access to a	patient's ch	art?					
	Very Rarely	1	asionally	Often							
	9 (19%)	31	(63%) 7 (14%)			2 (4%)					
	Do you feel	computer charti	ng will make your	job any eas	sier?						
		Yes	No	N	o Response						
	PRE	17 (35%)	20 (41%)		12 (24%)						
	POST	29 (58%)	9 (18%)		12 (24%)						
	(POST) Which do you prefer, computer charting or manual charting?										
Computer Charting Manual Charting No Response											
	32 (64		10 (20%)	' 5	8 (16%						
	(POST) W	uat differance if	any do you fact a	omnutas at	uting has	ada (4					
	(POST) What difference, if any, do you feel computer charting has made in the quality of patient care? Better No Change Worse No Response										
	Dattar	NT	**************************************	T	3.7	00010200000000000000000000000000000000					

Discussion

Implementation of the computer-charting system was trouble free, and therapists learned the system quickly. Therapists' response was very positive. The preference for using the system was not only very high. but higher than anticipated. Whereas only 35% (17/49) of those who returned the PRE questionnaire felt computer charting would make their job easier, 64% (32/50) of those who returned the POST questionnaire expressed a preference for computer charting. About one third of the responding therapists reported that computer charting was faster (Table I, items 2 and 3); however, 56% of therapists returning the POST survey felt that their charting time was better spent and 74% felt productivity was better (Table 1, item 5), indicating that the computer may have been helpful in ways other than speed of charting.

The computerized clinical records were more descriptive, legible, and complete than were the manual reports (Fig. 8). Overall, computer charting was found to be 12.4% more complete than manual charting. The only item in the study that did not show a significant improvement was medication documentation, which has now been made a mandatory entry on the computer. This will ensure 100% compliance and is justified because the delivery of medication is the primary objective of medications-nebulizer therapy.

Because computer charting can be programmed so that a therapist must reply to a question in order to proceed through the entry process, an argument can be made that the answering of all questions should be mandatory, assuring 100% compliance. Although mandatory entry seems to be the ideal solution, it has the disadvantage of not allowing the therapist to exercise discretion over what is charted. Mandatory entry may force the reporting of irrelevant or incorrect information. Certainly information is better left unreported than reported incorrectly. The ultimate responsibility for complete charting is the therapist's. Computer documentation significantly improved charting without forcing the outcome.




Fig. 8. Percentages of acceptable charting from an audit of 500 procedures of manual charting and 500 procedures of computer charting. Numbers atop bars are percentages of acceptable computer charts; numbers in bars are percentages of acceptable manual charts.

Every procedure allows the entry of comments in a free-text format; therefore, a procedure can be documented entirely on comments and still be complete. However, free-text entries are not so useful as structured data (selections that are stored in the computer in coded format). As an example, if patients receiving a certain bronchodilator were to be monitored for changes in breath sounds, the computer could be programmed easily to find the data if the information was structured. If the information was free text, accurate retrieval and monitoring would not be possible. Currently, structured data accounts for more than 95% of RC charting.

An argument can be made that too much information is charted, resulting in 'information overload,' whereby irrelevant information reduces the impact of relevant information on decision making.^{8, 19-22} Just

Pay Per (2 Wee			al Data res Billed)	RC Department Data (Supervisors' Accounts)			
PRE	FTE	Productivity	Work Units	Productivity	Work Units		
1	49.96	93	221,869	87	206,834		
2	50.40	95	229,005	88	211,680		
3	49.88	91	217,955	91	215,482		
2 3 4	54.68	86	226,560	98	255,902		
Average	51.23	91.3	223,847	91.0	222,475		
POST							
5	48.82	118	275,346	92	213,832		
6	51.05	109	267,750	98	237,383		
6 7 8 9	54.44	103	268,834	103	267,845		
8	50.62	107	260,645	107	258,162		
9	50.55	112	271,524	112	271,524		
10	50.91	103	250,040	106	258,114		
11	54.88	105	277,183	104	273,302		
12	57.92	106	294,009	106	291,916		
Average	52.40	107.9	270,666	103.5	259,010		
Р	NS	0.0002	0.00004	0.0054	0.0300		
6 Increase	2.3	18.2	20.9	13.7	16.4		

Table 2. Productivity and Work-Unit Data for the 8 Weeks Preceding and the 16 Weeks Following the Implementation of Computer Charting

FTE = full-time equivalent therapists paid during pay period.

Productivity = the % of work completed compared to the amount of work expected to be completed for the number of FTEs.

Work Units = the number of minutes spent doing productive work (determined by hospital Industrial Engineering). One work unit = one productive minute of work.

P = P value from unpaired *t* test of PRE and POST results.

NS = not significant.

what information is the most useful is a question that will require further study.

Evaluation of productivity was hampered by the fact that all accounting methods and charges had been changed 8 weeks prior to computer implementation. Unfortunately, this limited the amount of useful PRE data to only four pay periods.

Because the RC department maintains a nearly constant work force, fluctuations in work volume affect the productivity of the department. The results in Table 2 show that there were increases in productivity, according to both hospital and departmental calculation, after computer implementation (18.2% and 13.7%, respectively). Work volume also increased (20.9% and 16.4%, respectively), while the number of therapists did not increase significantly.

There were three possible reasons for the apparent improvement in productivity: (1) The work volume increased, requiring the therapists to work more efficiently. (2) The computer assured that work charted was charged for, and this accuracy increased the work volume. This explanation assumes that in the PRE period, some work was done but not accounted for. We were unable from the data available to make a quantitative assessment of this factor. Nevertheless, the computer assures concordance of clinical and financial record keeping and minimizes lost charges. (3) Computer charting helped the therapists do their job more efficiently and thus allowed them to handle heavier workloads. The manual Kardex system was replaced, saving 10 minutes per therapist per shift. The therapist survey showed that 74% of the therapists thought computer charting allowed them to be more productive, but they also indicated that the timesavings was not very substantial.

We conclude that all three factors mentioned above, or a combination of them, could have been responsible for the increase in productivity, although it remains unclear to what extent each factor may have been responsible. One fact was clear: During the period when therapists were busier and 18% more productive, they were using the computer. Computer charting did not decrease productivity.

Figure 9 shows that after Pay Period 6, about 6 weeks after implementation of computer charting, procedures billed and procedures assigned became highly correlated (r = 0.96 for Pay Periods 7 though 12). These results confirmed that computer charting provided a high degree of confidence that every item billed was documented as being performed. The poor correlation for Pay Periods 5 and 6 can be partially explained. Computer charting processes billing information immediately, whereas manual charting processes billing at least a day later. Pay Period 5, the first after conversion to computer charting, reflected the billing of all procedures during that pay period, plus the carryover billing of some procedures completed in the previous pay period hence, hospital billing records and RC records differed in the work volume reported. Also, the 24-hour management report was not implemented until Pay Period 6, SO errors may have gone unnoticed before that date.




Fig. 9. Comparison of work volume from hospital records and work volume from departmental records. Computer charting started at point marked by dotted line (March 1984).

Information that is stored in the computer is used in ways that are impractical with manual methods. The alert program provides automatic quality assurance by routinely searching all current patients' records for possible needs for corrective The facet of computerized charting with the greatest potential for development is in the expansion of the automatic monitoring of patient care. Information could be incorporated into assessment protocols that automatically monitor the efficacy of treatments. Patients' assessments could be reviewed so that care could be optimized. The medical staff could be provided computer-generated reminders

for use in treatment assessment,²² The information charted could also be useful for other departments. For example, a program monitoring infectious disease could take into consideration a change in breath sounds in a patient suspected of having a pulmonary infection. RC charting is now incorporated into computerized ICU- rounds reports and patient-summary reports. These reports extract the most recent and useful data and display them in a concise format for optimal use.^{6,9,21}

The RC computer system is efficient because it has streamlined the process of documentation while extracting the most 'useful' information. Without having to provide costly cumulative paper reports, the RC system provides better access for entry and review. Overall, computer charting is preferred by therapists over manual charting, making their job easier while improving the quality of information charted. Computer charting has added a high degree of confidence that there will be good correlation of clinical, administrative, and financial records. The computerization of charting RC procedures demonstrates the advantages of using clinical information for the benefit of the therapist, the department, the hospital, and the patient.

REFERENCES

- 1. Jydstrup RA, Gross MJ. Cost of information handling in hospitals. Health Serv Res 1966;1:253-271.
- Jacobwitzk K, Strodtman L, Lomas T, Turax T. Micro- computer based data management system far nursing assessment of the diabetic patient. Proceedings of the Fifth Annual Symposium on Computers in Medical Care 1981;755-759.
- 3. Collen MF, Davis LS, Van Brunt EE, Terdiman JF. Functional goals and problems in large-scale patient record management and automated screening. Fed Proc 1974;33:2376-2379.
- 4. Bronzino JD. Computer applications for patient care, Menlo Park CA: Addison-Wesley, 1983.
- 5. Mertz SL, Ash SR, Farrell J. The CNS in the ICU: A bedside notation system for nurses. Proceedings of the Sixth Annual Symposium on Computers in Medical Care 1982;577-581.
- Barnett GO. The application of computer-based medical- record systems in ambulatory practice. N Engl J Med 1984;310:1643-1650.
- 7. Pryor TA, Gardner RM, Clayton PD, Warner HR. The HELP system. J Med Syst 1983;7:87-102.
- 8. Warner BR. Computer assisted medical decision making. NY: Academic Press, 1979.
- Gardner RM, West BJ, Pryor TA. et al. Computer based ICU data acquisition as an aid to clinical decision making. Crit Care Med 1982;10:823-830.
- 10. SerIin). Fault tolerant systems in commercial applications. Computer 1984; 17: 19-30.
- 11. Jeromin G. Computerization-Are we ready? Respir Care 1982;27:797-798.
- Garrett LE, Stead WW, Hammond WE, McDonald CJ, Buecher M. A method of handling subjective and physical data; experience with two systems. Proceedings of the Sixth Annual Symposium on Computers in Medical Care 1982;232-235.
- 13. Zielstorff RD, Roglieri JL, Marble K, et al. Experience with a computer-based medical record for nurse practitioners in ambulatory care. Comput Biomed Res 1977; 10:6 1-73.
- 14. Collen MF. Patient data acquisition. Med Instrum 1978;12-222-225.
- 15. Joint Commission on Accreditation of Hospitals. Accreditation manual for Hospitals. 1982 edition. Chicago: American Hospital Association.
- Mishelevich DJ, Robinson AL, Rogers C, DuPriest ME, Mize EI. Respiratory therapy as a component of an integrated hospital information system: The Parkland On-tine Information System (POIS). Respir Care 1982;27:846-854.
- 17. Blue Cross and Blue Shield Association. Medical necessity guidelines: Respiratory care {inpatient). Chicago: September 1982.
- 18. ACCP-NHLBI. National conference on oxygen therapy. Chest 1984;86:234-247.
- 19. Grier MR. The need for data in making nursing decisions. In: Werley HH, Grier MR, eds. Nursing information systems. New York: Springer Publishing, 198 1:15-31.
- 20. Ayers WR. The potentials and limitations-of computers in medical practice. Update: Computers in Medicine 1983;1:35-39.
- 21. Bradshaw KE, Gardner RM, Clemmer TP, Orme JF, Thomas F, West BJ. Physician decision-making--Evaluation of data used in a computerized ICU. Int J Clin Monitor & Cornput 1984;1:81-41.
- 22. McDonald CJ. Protocol-based computer reminders, the quality of care and the non-perfectibility of man. N Engl J Med 1976;295:1351-1355.
- 23. Barnett GO, Winickoff R, Dorsey JL, Morgan MM, Lurie RS. Quality assurance through automated monitoring and concurrent feedback using a computer-based medical information system. Med Care 1978;16:962-970.