

# PHYSICAL REVIEW LETTERS

VOLUME 65

10 DECEMBER 1990

NUMBER 24

## Convexity and Exponent Inequalities for Conduction near Percolation

K. Golden

*Department of Mathematics, Princeton University, Princeton, New Jersey 08544*

(Received 31 July 1989; revised manuscript received 18 June 1990)

The bulk conductivity  $\sigma^*(p)$  of the bond lattice in  $\mathbb{Z}^d$  with a fraction  $p$  of conducting bonds is analyzed. Assuming a hierarchical node-link-blob (NLB) model of the conducting backbone, it is shown that  $\sigma^*(p)$  (for this model) is convex in  $p$  near the percolation threshold  $p_c$ , and that its critical exponent  $t$  obeys the inequalities  $1 \leq t \leq 2$  for  $d=2,3$  while  $2 \leq t \leq 3$  for  $d \geq 4$ . The upper bound  $t=2$  in  $d=3$ , which is realizable in the NLB class, virtually coincides with two very recent numerical estimates obtained from simulation and series expansion.

PACS numbers: 05.60.+w, 64.60.Cn

The random-resistor network<sup>1-5</sup> is the simplest model of a disordered conductor which exhibits complex macroscopic behavior in the form of a conducting phase transition. In particular, consider the bulk conductivity  $\sigma^*(p)$  of the bond lattice in  $\mathbb{Z}^d$ , where the conductivity of the bonds is either 1 with probability  $p$ , or  $\epsilon \geq 0$  with probability  $1-p$ . When  $\epsilon=0$ ,  $\sigma^*(p)=0$  for  $p \leq p_c$ , the percolation threshold, and it is believed<sup>6</sup> that  $\sigma^*(p) \sim (p-p_c)^t$  as  $p \rightarrow p_c^+$ . In this Letter we introduce a new approach to studying  $\sigma^*(p)$  when  $\epsilon=0$ , motivated by the simple observation that in numerical simulations<sup>6-10</sup> the graph of  $\sigma^*(p)$  for bond or site models in  $d \geq 2$  is always convex near  $p_c$ . Our approach is to analyze  $d^2\sigma^*/dp^2$  and investigate the consequences for the critical exponent  $t$ , assuming a self-similar, hierarchical structure for the conducting backbone near  $p_c$ , and certain technical conditions.

The principal results of our investigation and the assumptions under which they are obtained are as follows. First, the most serious assumption is that the conducting backbone near  $p_c$  has a hierarchical node-link-blob (NLB) structure.<sup>11,12</sup> This model contains both singly and multiply connected bonds, has "loops" on arbitrarily many length scales in a self-similar fashion, and incorporates the few rigorously known features<sup>5,12</sup> about the backbone on a macroscopic scale. We further make some technical assumptions about  $\sigma^*(p)$ : It obeys the above scaling law near  $p_c$ , has at least three derivatives for all  $p > p_c$ , and obeys  $d^2\sigma^*/dp^2 + d\sigma^*/dp > 0$  at

$p=1$ , which we have verified numerically. Under these assumptions, we prove exact asymptotics for  $d^2\sigma^*/dp^2$  as  $p \rightarrow p_c^+$ . The proof employs a novel technique whereby  $d^2\sigma^*/dp^2$  for the NLB model with  $\epsilon=0$  and  $p$  near  $p_c$  is computed using perturbation theory for  $\sigma^*(p)$  (for two- and three-component resistor lattices) around  $p=1$ , with a sequence of  $\epsilon$ 's converging to 1 as one goes deeper in the hierarchy. Our asymptotics yield not only convexity near  $p_c$ , which implies  $t \geq 1$ , but delineate in which dimensions  $d^2\sigma^*/dp^2 \rightarrow 0, +\infty$ , or a positive constant as  $p \rightarrow p_c^+$ . Combining this information with the scaling law  $d^2\sigma^*/dp^2 \sim (p-p_c)^{t-2}$  yields the inequalities  $1 \leq t \leq 2$  for  $d=2,3$  and  $2 \leq t \leq 3$  for  $d \geq 4$ . The inequality  $t \leq 3$  for  $d \geq 4$  is obtained by applying a similar analysis to  $d^3\sigma^*/dp^3$  for the simpler node-link model, and can be viewed as a mean-field bound, since it is believed that  $t=3$  for  $d \geq 6$ . We stress that the convexity and inequalities are not rigorous for the actual backbone near  $p_c$  for the original lattice, but *are* rigorous for the NLB model of the backbone, under the above technical assumptions.

Our results for  $d=3$  are particularly intriguing. First, the inequality  $t \leq 2$  excludes roughly one-third of published numerical estimates of  $t$  in  $d=3$ , which have ranged from 1.5 to 2.36. Furthermore, this inequality is based on an exact calculation of  $t=2$  for one particular NLB model which provides an upper bound on  $t$  for the full class. In view of this result, it is quite striking that very recently Gingold and Lobb<sup>13</sup> have obtained for

$d=3$  the estimate  $t=2.003 \pm 0.047$  from simulation on lattices up to  $80^3$ , and Adler *et al.*<sup>14</sup> have obtained  $t=2.02 \pm 0.05$  from a thirteenth-order series expansion. In addition, our inequality is compatible with the results of an  $\epsilon=6-d$  expansion,<sup>15</sup> and the general view that “roughly  $t=2$ ”.<sup>2</sup> (We should also mention the recent work of Roman<sup>16</sup> on the ant-in-the-labyrinth problem, who indirectly obtains a value of  $t \approx 2.16$ . However, he acknowledges the inconsistency with other results, which is discussed further in Ref. 13.) The recent numerical results in Refs. 13 and 14, in conjunction with our work on the NLB model, suggest the possibility that  $t=2$  is an *exact* result for  $d=3$ . To our knowledge, the present work is the first to relate  $t$  in a direct and natural way to the number 2, rather than to other (unknown) critical exponents of percolation theory.

Before we begin, we refer the reader to Ref. 17. In addition to containing the mathematical details of the results discussed here, we obtain there numerical and rigorous results concerning the regimes in  $\epsilon$  and  $p$  of convexity of  $\sigma^*(p)$  for bond and site models, the principal rigorous results being that for the  $d=2$  bond problem, while  $\sigma^*(p)$  cannot be convex for all  $p$  when  $\epsilon=0$ , it is convex for every  $\epsilon > 0$  near  $p_c = \frac{1}{2}$ .

We now formulate the bond conductivity problem for  $\mathbb{Z}^d$ , where, for simplicity, we begin with  $d=2$ . Take an  $L \times L$  sample  $G_L$  of the bond lattice with  $M$  ( $\sim dL^d$ ) bonds. Assigned to  $G_L$  are  $M$  independent random variables  $c_i$ ,  $1 \leq i \leq M$ , the bond conductivities, which take the values 1 with probability  $p$  and  $\epsilon \geq 0$  with probability  $1-p$ . We attach perfectly conducting bus bars to two opposite edges, and let  $\sigma_L(p)$  be the effective conductivity of this network, averaged over realizations of the bond conductivities. For  $d \geq 1$ , the bulk conductivity of the lattice is defined as

$$\sigma^*(p) = \lim_{L \rightarrow \infty} L^{2-d} \sigma_L(p). \tag{1}$$

For  $\epsilon > 0$ , the infinite-volume limit in (1) has been shown to exist,<sup>18-21</sup> and for  $\epsilon=0$  the existence of  $\sigma^*$  has recently been proven in the continuum.<sup>22</sup>

The calculation of  $d^2\sigma^*/dp^2$  will require the following definition. For any graph  $B$  with bonds  $b_i$  of unit conductivity, define

$$\delta^2\sigma(B) = \sum_{\substack{b_i, b_j \in B \\ b_i \neq b_j}} [\sigma_{ij}(1,1) + \sigma_{ij}(0,0) - \sigma_{ij}(1,0) - \sigma_{ij}(0,1)], \tag{2}$$

where in (2)  $\sigma_{ij}(1,1) = \sigma(B)$  the conductivity of  $B$  measured between two vertices,  $\sigma_{ij}(0,0)$  is the conductivity of  $B$  with  $b_i$  and  $b_j$  removed, and so on. The expression in (2) represents the discrete second derivative of  $\sigma$  with respect to  $p$ , as follows. Let  $G$  be the lattice in  $d \geq 2$  with bond conductivities 1 and 0 and bulk conductivity function  $\sigma^*(p)$ . If  $B=B(p)$  is a realization of occupied

bonds of  $G$  at probability  $p$ , then<sup>17</sup>

$$p^2 \frac{d^2\sigma^*}{dp^2} = \delta^2\sigma^*(B(p)), \tag{3}$$

where  $\delta^2\sigma^*$  is the scaled infinite-volume limit of (2) and the right-hand side in (3) is appropriately averaged. [We are assuming in (3) that  $\sigma^*(p)$  is twice differentiable for  $p > p_c$  when  $\epsilon=0$ .] In (2) note that dangling bonds do not contribute, so that one may think of  $B(p)$  as a realization of the backbone at bond fraction  $p$ . For clarity, note that at  $p=1$ ,  $B(p)=G$ . We remark that analysis of simple graphs shows that, typically, positive contributions to (2) arise from series pairs, while negative contributions arise from pairs in parallel.

The idea now is to replace an actual backbone graph  $B(p)$  for  $p$  near  $p_c$  by a node-link-blob graph  $A$ , which is based on the work of Stanley<sup>11</sup> and Coniglio.<sup>12</sup> This graph is a “superlattice” constructed by replacing the bonds of the hypercubic lattice  $G$  in  $d \geq 2$  by first-order necklaces composed of strings (links) and first-order beads (blobs), and separating the nodes of  $G$  by a correlation length  $\xi$ , as in Fig. 1(a). The beads themselves have a hierarchical structure, as shown in Fig. 1(b), consisting of two second-order necklaces in parallel, and so on, in a self-similar fashion to order  $N$  for an arbitrary large integer  $N$ . We assume that any  $k$ th-order necklace has  $\beta-1$  beads on it for an arbitrary large integer  $\beta$ , and that each pair of beads is joined by a string of  $n_k$  bonds, so that there are a total of  $\beta n_k$  string bonds on each necklace. The  $\beta n_1$  string bonds on any first-order necklace are called singly connected because removal of one of them breaks the connection between nodes separated by  $\xi$ . All the rest of the bonds in the NLB graph are multiply connected, and among these it is useful to iden-




FIG. 1. Node-link-blob model of the conducting backbone near  $p_c$ . In (a), the nodes are a correlation length  $\xi$  apart, and are connected by necklaces of beads (blobs) and strings (links) with  $n_1$  bonds connecting two beads. The beads have a self-similar structure, as shown in (b), with  $n_2$  bonds connecting two beads.

tify the  $\beta n_2$  string bonds on a second-order necklace as doubly connected, since it is possible to remove two of them (in parallel) and break a connection between nodes. Based on a result of Coniglio's,<sup>12</sup> implying in our context that the number of singly and doubly connected bonds between the nodes both diverge with exponent 1 as  $p \rightarrow p_c^+$ , we assume that  $n_1 = 2\beta n_2$ . Because of self-similarity, we assume that

$$n_{j-1} = 2\beta n_j, \quad j = 2, \dots, N. \quad (4)$$

Relation (4) can be used to solve for the  $n_j$ ,  $j > 1$  in terms of  $n_1$  with  $n_2 = n_1/2\beta$ ,  $n_3 = n_1/4\beta^2$ , and so on, and we refer to the NLB graph as  $A(n_1)$ . In this model the percolation limit  $p \rightarrow p_c^+$  is characterized by the limits  $n_1, \beta, N, \xi \rightarrow \infty$ , so that the lengths of all orders of necklaces, and the numbers and sizes of all orders of blobs, diverge as  $p \rightarrow p_c^+$ .

Before we give the asymptotics of  $\delta^2 \sigma^*(A(n_1))$ , we must discuss the conditions under which they are proven. Consider  $\sigma^*(q_1, q_2)$  for the bond lattice in  $\mathbb{Z}^d$  with three conductivities 1,  $\epsilon_1$ , and  $\epsilon_2$  in proportions  $p$ ,  $q_1$ , and  $q_2$ , in addition to our standard two-component conductivity  $\sigma^*(p)$ . We require that  $\sigma^*(q_1, q_2)$  has second-order partials at  $q_1 = q_2 = 1$  for all  $\epsilon_1, \epsilon_2 \geq 0$ , and that  $\sigma^*(p)$  has two derivatives at  $p = 1$  for all  $\epsilon \geq 0$ . For  $\epsilon, \epsilon_1$ , and  $\epsilon_2 > 0$ , these conditions are satisfied by our general results<sup>17</sup> that  $\sigma^*(p)$  is analytic for all  $p \in [0, 1]$  and  $\sigma^*(q_1, q_2)$  is analytic for all  $(q_1, q_2) \in [0, 1] \times [0, 1]$ . The  $\epsilon = \epsilon_1 = 0$  requirements will be assumed, although Kozlov<sup>23</sup> has proven the existence of  $d\sigma^*/dp|_{p=1}$  for a class of continuum analogs.

The second main condition is that given the hypercubic base lattice  $G$  for  $A(n_1)$ ,

$$\kappa(G) = \left. \frac{d\sigma^*}{dp} \right|_{p=1} + \left. \frac{d^2\sigma^*}{dp^2} \right|_{p=1} > 0. \quad (5)$$

In any  $d \geq 2$ ,  $d\sigma^*/dp|_{p=1} = d/(d-1)$ ,<sup>1</sup> while  $d^2\sigma^*/dp^2|_{p=1}$ , if negative, is quite small, e.g.,  $\approx -0.21$  in

$$\delta_{11} = z_0 \left[ \beta n_1 (\beta n_1 - 1) \frac{d\sigma^*}{dp}(p=1, h_1) + (\beta n_1)^2 \frac{d^2\sigma^*}{dp^2}(p=1, h_1) \right], \quad (10)$$

$$\delta_{12} = z_0 \left[ (\beta n_1)^2 \frac{d\sigma^*}{dp}(p=1, h_1) + (\beta n_1)^2 \frac{\partial^2 \sigma^*}{\partial q_1 \partial q_2}(p=1, h_1, h_2) \right], \quad (11)$$

where  $(d\sigma^*/dp)(p=1, h_1)$ , e.g., is for  $G$  with bond conductivities 1 and  $h_1 = 0$ , with  $h_k = z_k/z_0$ . As  $k \rightarrow \infty$ ,  $h_k \rightarrow 1$ , and as  $\beta \rightarrow \infty$ ,  $h_k \rightarrow 1$  for all  $k \geq 2$ , and similarly for  $h_{jk} = z_{jk}/z_0$ . The necessary control of the  $\delta_{jk}$  is then obtained either from (5), or from perturbation theory around a homogeneous medium ( $\epsilon = 1$  or  $\epsilon_1 = \epsilon_2 = 1$ ), which establishes (6). All the details appear in Ref. 17.

We wish to make the following remarks concerning the above result. First, a result similar to (7) holds if we

$d = 2$ ,<sup>9,17</sup> indicating that  $\sigma^*(p)$  is quite straight near  $p = 1$ , so that (5) is satisfied. Condition (5) amounts to a consequence of the long-held view that effective-medium theory (giving a straight-line solution) provides an accurate description of  $\sigma^*(p)$  near  $p = 1$ , which also holds for general lattices. In fact, the asymptotics below can be proven for a variety of periodic base lattices  $G$  which satisfy (5), and presumably hold even for random lattices.

We may now state our principal result.

Under the above assumptions, for fixed, large  $n_1$ ,  $\beta$ , and  $N$ ,

$$\delta^2 \sigma(A(n_1)) = a_N \kappa(G) \beta n_1 + \sum_{i=0}^{\infty} \frac{a_i n_1 + b_i}{\beta^i}, \quad (6)$$

where  $(a_N)^{-1} = \sum_{i=0}^N (\frac{1}{4})^i$  and the series in (6) converges, so that

$$\delta^2 \sigma^*(A(n_1)) \sim \frac{a_N \kappa(G) \beta n_1}{\xi^{d-2}} > 0, \quad n_1, \beta, N, \xi \rightarrow \infty. \quad (7)$$

The idea of the proof<sup>17</sup> of (6) is first to write

$$\delta^2 \sigma(A(n_1)) = \sum_{\substack{j,k=1 \\ k \geq j}} \delta_{jk}, \quad (8)$$

where  $\delta_{jk}$  is the sum of all contributions to  $\delta^2 \sigma(A(n_1))$  in (2) arising from pairs with one bond in a  $j$ th-order string and the other in a  $k$ th-order string, which is in either the same or a different first-order necklace. Now let  $z_k$  be the conductivity of a single first-order necklace with one bond removed from a  $k$ th-order string, with  $z_0 = a_N/\beta n_1$  for no bond removed,  $z_1 = 0$ , and

$$z_k = z_0 (1 + \gamma_k / \beta^{k-1})^{-1}, \quad k \geq 2, \quad (9)$$

where  $\gamma_k \rightarrow 0$  as  $k \rightarrow \infty$  geometrically fast. There are analogous formulas for the various forms of  $z_{jk}$  with two bonds removed, say, in series or in parallel. Then through representations like (2) and (3), we obtain formulas for the  $\delta_{jk}$  in terms of derivatives of  $\sigma^*(p)$  and  $\sigma^*(q_1, q_2)$  at  $p = 1$ , such as

replace (4) by  $n_{j-1} = \eta_j \beta_j n_j$ , where the blobs of order  $j-1$  are made of  $\eta_j$  necklaces in parallel, with reasonable assumptions about  $\eta_j$  and  $\beta_j$ . Even if the blobs have a more complicated superlattice structure themselves, an analog of (7) presumably holds. Also, as noted above, (7) can be proven for a variety of base lattices  $G$ . Finally, while the principal assumption of the NLB graph replacing the actual backbone is quite serious, our proof of (6) shows that the dominant contribution to (7)

comes from  $\delta_{11}$ , which comes from macroscopic contributions in the NLB graph, where the model reflects well the actual structure. A similar result will hold for any reasonable assumption about microscopic backbone structure.

We now proceed to the implications of (7). First, its positivity establishes convexity of  $\sigma^*(p)$  for the NLB model, which implies [under our assumptions, including scaling and the existence of three derivatives of  $\sigma^*(p)$  for all  $p > p_c$  when  $\epsilon=0$ ] that  $t \geq 1$ , for any  $d \geq 2$  (the inequality  $t \geq 1$  has been previously established in a different manner in Refs. 24 and 25). Now let  $\lambda(n_1)$  be the length of a first-order necklace, so that  $\lambda(n_1) \approx \beta n_1 + \beta^2 n_2 + \dots + \beta^N n_N = \theta_N \beta n_1$ ,  $\theta_N = \sum_{i=0}^N 2^{-i}$ . By (7), we then have

$$\delta^2 \sigma^*(A(n_1)) \sim \frac{\rho_N \lambda(n_1)}{\xi^{d-2}}, \quad n_1, \beta, N, \xi \rightarrow \infty, \quad (12)$$

where  $\rho_N = a_N \kappa(G) / \theta_N$ , so that  $\rho_N \approx \frac{2}{3}$  for large  $N$  in  $d=2$ . Since all the parameters  $n_1$ ,  $\beta$ ,  $N$  and  $\xi$  are diverging as  $p \rightarrow p_c^+$ , we can define a whole class of NLB models by how fast  $\lambda(n_1)$  scales to  $\infty$  relative to  $\xi$ . By the structure of the model, clearly  $\lambda(n_1) \geq \xi$ , and typically,  $\lambda/\xi \rightarrow \infty$ . Thus as a consequence of (12) we have in  $d=2$  and 3

$$\delta^2 \sigma^*(A(n_1)) \rightarrow +\infty, \quad n_1, \beta, N, \xi \rightarrow \infty, \quad (13)$$

except in  $d=3$  when  $\lambda(n_1) = C\xi$ ,  $C \geq 1$ , in which case

$$\delta^2 \sigma^*(A(n_1)) \rightarrow \rho C > 0, \quad (14)$$

where  $\rho = \lim_{N \rightarrow \infty} \rho_N$ . In  $d \geq 4$ , if  $\lambda$  and  $\xi$  are scaled so that  $\lambda(n_1)/\xi^{d-2} \rightarrow 0^+$ , then

$$\delta^2 \sigma^*(A(n_1)) \rightarrow 0^+. \quad (15)$$

Under our assumptions, in particular, that  $d^2 \sigma^* / dp^2 \sim (p - p_c)^{t-2}$ , we then have, collecting our results

$$1 \leq t \leq 2, \quad d=2,3; \quad 2 \leq t \leq 3, \quad d \geq 4. \quad (16)$$

In (16) the last inequality  $t \leq 3$  for  $d \geq 4$  is obtained by a result that  $\delta^3 \sigma^*(A'(n_1)) \sim C \lambda^2(n_1) / \xi^{d-2}$  for a simpler node-link graph  $A'(n_1)$ , which is believed to be adequate in higher dimensions.<sup>26</sup> For models in  $d=4,5$  which satisfy  $\lambda^2(n_1)/\xi^{d-2} \rightarrow \infty$ , we have  $\delta^3 \sigma^*(A(n_1)) \rightarrow \infty$ , so that  $d^3 \sigma^* / dp^3 \sim (p - p_c)^{t-3} \rightarrow \infty$ , which gives the inequality.

It is a pleasure to thank a number of people for helpful

discussions during the course of the work: D. Bergman, J. Chayes, L. Chayes, D. Fisher, H. Kesten, C. Lobb, G. Papanicolaou, and S. Varadhan. Special thanks is owed to P. Doyle for many long and useful discussions which helped clarify certain matters in this work. This work was supported in part by NSF Grant No. DMS-8801673 and AFOSR Grant No. AFOSR-90-0203.

<sup>1</sup>S. Kirkpatrick, Rev. Mod. Phys. **45**, 574 (1973).

<sup>2</sup>D. Stauffer, *Introduction to Percolation Theory* (Taylor and Francis, London, 1985).

<sup>3</sup>G. Grimmett and H. Kesten, Z. Wahr. **66**, 335 (1984).

<sup>4</sup>H. Kesten, *Percolation Theory for Mathematicians* (Birkhäuser, Boston, 1982).

<sup>5</sup>J. T. Chayes and L. Chayes, Commun. Math. Phys. **105**, 133 (1986).

<sup>6</sup>S. Kirkpatrick, Phys. Rev. Lett. **27**, 1722 (1971).

<sup>7</sup>B. P. Watson and P. L. Leath, Phys. Rev. **B 9**, 4893 (1974).

<sup>8</sup>A. B. Harris and S. Kirkpatrick, Phys. Rev. **B 16**, 542 (1977).

<sup>9</sup>M. H. Ernst, P. F. J. van Velthoven, and Th. M. Nieuwenhuizen, J. Phys. A **20**, 949 (1987).

<sup>10</sup>Th. M. Nieuwenhuizen, P. F. J. van Velthoven, and M. H. Ernst, Phys. Rev. Lett. **57**, 2477 (1986).

<sup>11</sup>H. E. Stanley, J. Phys. A **10**, L211 (1977).

<sup>12</sup>A. Coniglio, J. Phys. A **15**, 3829 (1982).

<sup>13</sup>D. B. Gingold and C. J. Lobb, Phys. Rev. **B 42**, 8220 (1990).

<sup>14</sup>J. Adler, Y. Meir, A. Aharony, A. B. Harris, and L. Klein, J. Stat. Phys. **58**, 511 (1990).

<sup>15</sup>A. B. Harris, S. Kim, and T. C. Lubensky, Phys. Rev. Lett. **53**, 743 (1984).

<sup>16</sup>H. E. Roman, J. Stat. Phys. **58**, 375 (1990).

<sup>17</sup>K. Golden, "Exponent Inequalities for the Bulk Conductivity of a Hierarchical Model" (to be published).

<sup>18</sup>S. M. Kozlov, Dokl. Akad. Nauk. SSSR **241**, 1016 (1978).

<sup>19</sup>G. Papanicolaou and S. Varadhan, in *Colloquia Mathematica Societatis János Bolyai 27, Random Fields* (North-Holland, Amsterdam, 1982).

<sup>20</sup>K. Golden and G. Papanicolaou, Commun. Math. Phys. **90**, 473 (1983).

<sup>21</sup>R. Künnemann, Ph.D. thesis, University of Heidelberg, 1983 (unpublished).

<sup>22</sup>V. Zhikov, Math. Zametki **45**, 34 (1989).

<sup>23</sup>S. M. Kozlov, Russ. Math. Surv. **44**, 91 (1989).

<sup>24</sup>J. P. Straley, Phys. Rev. **C 15**, 2333 (1982).

<sup>25</sup>J. Chayes, L. Chayes, and H. Kesten (unpublished).

<sup>26</sup>A. B. Harris, Phys. Rev. **B 28**, 2614 (1983).