

GRASAS Y ACEITES 67 (4)

October–December 2016, e169

ISSN-L: 0017-3495

doi: <http://dx.doi.org/10.3989/gya.0457161>

Markers of quality and genuineness of commercial extra virgin sacha inchi oils

N.A. Chasquibol^a, R.B. Gómez-Coca^b, J.C. Yácono^a, Á. Guinda^b, W. Moreda^b,
C. del Aguila^b and M.C. Pérez-Camino^{b,✉}

^aFaculty of Industrial Engineering, Institute of Scientific Research, (IDIC), Universidad de Lima, Avenida Javier Prado Este N° 4600, Urb. Fundo Monterrico Chico, Santiago de Surco, Lima, Peru.

^bDepartment of Characterization and Quality of Lipids, Instituto de la Grasa, CSIC, Campus Pablo de Olavide University, Bdg. 46, 41013 Sevilla, Spain.

✉Corresponding author: mcperezcamino@ig.csic.es

Submitted: 26 April 2016; Accepted: 19 September 2016

SUMMARY: This work tackles the study of the quality and authenticity of oils labeled and commercialized as extra virgin sacha inchi oil. Major and minor components as triglycerides, fatty acid methyl esters, tocopherols, sterols and hydrocarbons are determined as well as other physicochemical parameters (density, viscosity, acidity and peroxide value). The results showed that some of the commercialized oils do not fulfill the basic requirement established in the regulation such as the content of α -linolenic acid, higher than 44.7 or 55.0% in the cases of *P. volubilis* and *P. huayllabambana*, respectively. The calculated stigmasterol/campesterol ratio for genuine sacha inchi oils should be around 4, however not all commercial oils analyzed comply with this requirement. The presence of the flavons sesamin and sesamol indicates the addition of compounds from sesame oils. Finally, some of the commercial oils showed to contain *trans* fatty acids although this was not accompanied by the sterene hydrocarbon presence.

KEYWORDS: *Authenticity; Characterization; Commercial sacha inchi oils; Purity; Quality parameters*

RESUMEN: *Marcadores de la calidad y la genuinidad de aceites de sacha inchi extra virgen comerciales.* En este trabajo se aborda el estudio de la calidad y la genuinidad de los aceites etiquetados y comercializados como sacha inchi extra virgen. Se estudian los componentes mayoritarios como los triglicéridos y los ésteres metílicos de ácidos grasos, componentes menores insaponificables (tocoferoles, esteroides e hidrocarburos) así como otros parámetros fisicoquímicos (densidad, viscosidad, acidez, peróxidos y estabilidad). Los resultados mostraron que algunos de los aceites comercializados no cumplían con el requisito básico establecido en la normativa de tener un contenido en α -linolénico superior a 44,7 o 55,0% determinado para *P. volubilis* o *P. huayllabambana* respectivamente. La relación estigmasterol/campesterol medida en aceites de sacha inchi genuinos es de alrededor de 4, y no todos los aceites comerciales analizados cumplían con este requisito. La presencia de las flavonas sesamina y sesamolina indica la adición de compuestos procedentes de aceites de sésamo. Por último, algunos de los aceites comerciales estudiados, contenían ácidos grasos *trans* aunque no se detectó en ellos la presencia de hidrocarburos esteroideos.

PALABRAS CLAVE: *Aceites de sacha inchi comerciales; Autenticidad; Caracterización; Parámetros de calidad; Pureza*

Citation/Cómo citar este artículo: Chasquibol NA, Gómez-Coca RB, Yácono JC, Guinda Á, Moreda W, del Aguila C, Pérez-Camino MC. 2016. Markers of quality and genuineness of commercial extra virgin sacha inchi oils. *Grasas Aceites* 67, e169. <http://dx.doi.org/10.3989/gya.0457161>.

Copyright: © 2016 CSIC. This is an open-access article distributed under the terms of the Creative Commons Attribution (CC-by) Spain 3.0 License.

1. INTRODUCTION

Sacha inchi oil is a singular, although globally is an almost unknown, edible oil which belongs to the *Plukenetia* genus and is native to the Peruvian Amazon. One of its most relevant peculiarities is to be one of the edible oils with the highest content of ω 3 fatty acids (44.7–55.0% α -linolenic acid (Ln), depending on the ecotype) (Gutiérrez *et al.*, 2011; Maurer *et al.*, 2012; NTP, 2010; NTP 2014; Liua *et al.*, 2014), along with a particular sterol composition (stigmasterol > campesterol), and high amounts of γ - and δ -tocopherols (>1900 mg·kg⁻¹). Sacha inchi oils are consumed as virgin oils since they have a highly appreciated ‘flowered’ taste. All those particularities justify the development of research studies on their nutritional and health benefits (Muñoz Jauregui *et al.*, 2013). Besides, in 2014 the Food and Drugs Administration (FDA) declared sachá inchi oils as safe food (GRAS). Thanks to all those facts there are presently 11 companies which export sachá inchi oil from Peru to US, which suppose 9% of their exportations. The most important part of the Peruvian exportations (40%) goes to the Netherlands. Also in 2014 four Peruvian companies got the inclusion of sachá inchi oil into the European market by getting its exclusion from the Novel Food list. The category of Novel Food prevents the entry of foods which do not have a significant history of consumption in the European Union before May 1997 into the European market. After the Netherlands, Spain is the main importing country of sachá inchi oils (32%).

Although there are 16 species of the genus *Plukenetia* described (Bussmann *et al.*, 2009), six of them growing in the Peruvian Amazon (Rodríguez *et al.*, 2010), at present commercial sachá inchi oils come almost exclusively from the *P. volubilis* ecotype. The oils are obtained through artisanal production with a cold-press solvent-free system, and the use of decanters and filters in order to eliminate suspended particles and water. Before extraction, sachá inchi seeds are partially peeled and cleaned by ventilation to remove impurities, they are then milled mechanically. The oils obtained must comply with the requirements established in the regulations (NTP, 2010; NTP 2014). The specifications of quality and safety are established for all oils of the *Plukenetia* genus, although some parameters are specific for *P. volubilis* and *P. huayllabambana* oils. Due to their positive properties and also because of their time-consuming preparation, sachá inchi oils are very expensive. Hence and, taking advantage of the laxity of its regulation, fraudulent practices can be easily undertaken.

After a complete characterization of two of the most popular varieties of sachá inchi oils, *P. volubilis* and *P. huayllabambana* (Chasquibol *et al.*, 2012, Chasquibol *et al.*, 2014) we are in position to study the quality and genuineness of the commercialized sachá inchi oils.

2. MATERIALS AND METHODS

2.1. Oil samples

Fourteen (C-1 to C-14) sachá inchi oil samples, labeled as extra virgin from different commercial branches, were purchased in specialized shops of Lima (Perú) during 2012–2015. Samples, showed an expiration date 2014–2016 and were stored in freezer until analysis. They were bottled in dark-green 250 mL glass bottles, except in the case of C-8 sample that came in a brown 125 mL glass container. In all cases, it was specified that the oils contained 48% ω 3-acid, and in most of the samples also a 36% linoleic and 9% oleic acid contents were pointed out. An acidity value below 1% was specified in C-2 and C-8 samples, and the fact of belonging to the *volubilis* ecotype was indicated in the labels of C-8, C-9 and C-12 bottles. Special reference to the absence of *trans* fatty acids was made for C-6 and C-7 samples.

2.2. Reagents and solutions

Acetone, diethyl ether, and hexane were supplied by VWR International (West Chester, PA, USA). HPLC grade propionitrile was supplied by Panreac. Si-SPE cartridges were from Varian (EA Middelburg, The Netherlands). Standards of 5- α -cholestan-3 β -ol and trilinolenin (LnLnLn) were from Sigma-Adrich Co. (St. Louis, MO, USA). Hexamethyldisilazane, pyridine, trimethylchloroxilane, and standards of eicosane and tocopherols were from Merck-España (Merck Group, Darmstadt, Germany).

2.3. Physical-chemical parameters

Physical indexes (iodine value, refractive index, density, and viscosity), physicochemical parameters (triacylglycerol, fatty acid methyl ester, tocopherol, sterol, and hydrocarbon contents), quality indexes (percentage of acidity and peroxide value), and stability parameters (Rancimat test) were determined in order to evaluate both, oil quality and authenticity.

Free fatty acid (FA) content and peroxide value were determined according to the AOCS Official Methods (AOCS, 2012, 2013).

The oxidative stability was determined following the standard Official Method Cd 12-b-92,18,19 using a Rancimat equipment (743 Rancimat Metrohm Co., Basel, Switzerland), with an air flow of 20 L·h⁻¹ at 100±1 °C (AOCS, 1997).

Viscosity and density were measured using a Stabinger SVM 3000 viscometer (Anton Paar GmbH Graz, Austria), and the refractive index was determined with a temperature-controlled Abbe refractometer (Hilger & Watts Ltd., London, U.K.) (Ourrach *et al.*, 2012).

The iodine value was determined according to UNE-EN 14111, 2003 where the unsaturated FA

composition is taken into account according to the following formula:

$$0.95 \times \%C_{16:1} + 0.8986 \times \%C_{18:1} + 1.81 \times \%C_{18:2} \\ + 2.735 \times \%C_{18:3} + 0.8175 \times \%C_{20:1} + 0.7497 \times \\ \%C_{22:1}$$

2.4. Major glyceridic components

We will include fatty acid methyl esters and triglyceride composition in this Section.

2.4.1. Fatty acid composition

The fatty acid composition was determined by gas chromatography (GC) as fatty acid methyl esters (FAME) according to the IUPAC Standard Method (IUPAC, 1987a). Oil samples (50 mg) were transesterified using 0.5 mL 2 N methanolic solution of potassium hydroxide. GC analysis was carried out using an HP 7890B gas chromatograph (Hewlett-Packard, Palo Alto, CA) equipped with a capillary column (poly (90% biscyanopropyl–10% cyanopropylphenyl) siloxane, 60 m Å~ 0.25 mm i.d.; 0.20 µm film thickness) and a automatic split injector and a flame ionization detector (FID). The carrier gas was hydrogen at a flow rate of 1 mL min⁻¹. The temperatures of the injector and detector were 225 and 250 °C, respectively. The oven was programmed at a temperature of 165 °C (10 min), which was then increased by 1.5 °C min⁻¹ up to 200 °C (10 min). The injection volume was 1 µL. The fatty acid composition was expressed as relative percentage of each fatty acid on total fatty acids.

2.4.2. HPLC-RI Triglyceride analysis

Triacylglycerols (TG) were separated by HPLC on a Lichrosphere 100 RP-18 (4µ) column (Moreda *et al.*, 2003). Previously, the oil samples were purified by silica-SPE. A solution of the oil (0.10 g in 0.5 mL of hexane) was charged into the column and pulled with 10 mL of the hexane-diethyl ether (87:13 v/v) admixture. The collected fraction was evaporated to dryness, and the residue dissolved in 2 mL acetone and injected (10 µL) onto the HPLC system using a Beckman Gold 508 auto-sampler (Beckman-Coulter, Fullerton, CA, USA). The analyses were done using a Beckman Gold 126 pumping unit (Beckman-Coulter, Fullerton, CA, USA), a refractive index detector (RI) Perkin Elmer 200 (Perkin Elmer, Norwalk, CT, USA) and a Beckman Mistral peltier column thermostat unit fixed at 20 °C (Beckman-Coulter, Fullerton, CA, USA). The elution was done with propionitrile at a flow rate of 0.6 mL·min⁻¹. The TG composition was expressed as relative percentage of each group of TG with the same Equivalent Carbon Number (ECN) on total TG.

2.5. Minor unsaponifiable compounds

As minor unsaponifiable compounds we considered sterols, hydrocarbons and tocopherols (Gómez-Coca *et al.*, 2015). Just for sterol determination it is necessary to get the unsaponifiable matter in order to isolate and determine them quantitatively.

2.5.1. Sterol composition

The sterol fraction was isolated from the unsaponifiable matter (IUPAC, 1987c) by preparative HPLC fitted with a Si-column (250 mm Å~ 4 mm i.d.; 4 µm particle size) (Cert *et al.*, 1997). An hexane:diethyl ether (1:1, v/v) solution at a flow rate of 0.8 mL·min⁻¹ was used as eluent, together with UV detection at 210 nm. The fraction of interest eluted at Rt=10–25 min and it contained both the sterols and the triterpenic dialcohols. The collected portion was evaporated until dryness, derivatized with 500 µL of the 1:3:9 (v/v/v) trimethylchloroxilane:hexamethyldisilazane: pyridine admixture and analyzed by GC. The gas chromatograph (Agilent 6890N) was equipped with a fused silica column (poly(5% diphenyl– 95% dimethyl) siloxane, 30 m, Å~ 0.25 mm i.d., Å~ 0.25 µm film thickness), and FID. The oven was programmed isothermally at 260 °C, and the split ratio was 1:50. Hydrogen was used as carrier gas at a flow rate of 1 mL·min⁻¹. The injector and detector temperature was 300° C. The quantitative determination was done starting from 5 g of the oil sample and using α-cholestanol at a concentration of 1mg·mL⁻¹ as internal standard.

2.5.2. Tocopherols

Tocopherols were determined according to the IUPAC Standard Method 2432 (IUPAC 1987b). The oil solution in hexane (10 mg·mL⁻¹) was analyzed by HPLC fitted with a silica-column (250 mm Å~ 4 mm i.d.; 4 µm particle size). A solution of hexane:2-propanol (99:1, v/v) at a flow rate of 1 mL·min⁻¹ was used as a eluent and the detection was done by fluorescence (RF-10AXL Shimadzu fluorescence) at λ=290 and 330 nm, (excitation and emission, respectively). Standards of tocopherols in hexane at concentrations of 4–6 µg·mL⁻¹ were used for quantitative determinations.

2.5.3. Hydrocarbon fractions

2.5.3.1. Aliphatic saturated hydrocarbon quantification. Aliphatic saturated hydrocarbons (HC) were determined following a procedure previously described with minor modifications (Moreda *et al.*, 2001; Gómez-Coca *et al.*, 2016). The fraction containing the HC was isolated by low-pressure column chromatography filled with 15 g of silica gel (Si-60) impregnated with AgNO₃. A sample of

0.5 g oil introduced into the column was eluted with 80 mL petroleum ether, evaporated until dryness, re-dissolved in 0.5 mL hexane, and analyzed by on-column GC. The gas chromatogram was equipped with a poly-5% diphenyl-95% dimethylsiloxane, capillary column of 12 m, 0.32 mm Å i.d., 0.1 µm film thickness. The initial oven temperature was set at 80 °C for 2 min, and the rate was established at 12 °C·min⁻¹ up to 280 °C, then at 7 °C·min⁻¹, and then up to 340 °C. The injector temperature was set at 80 °C and the FID temperature at 350 °C. Eicosane (C20:0) at a concentration of 0.05 mg·mL⁻¹ was used as internal standard.

2.5.3.2. Steroideal hydrocarbons. Sterene determination was carried out according to the method developed by Cert *et al.*, 1994, which was standardized by the IUPAC (Dobarganes *et al.*, 1999) and included in International Regulations (ISO, 1999). The unsaponifiable matter was obtained from 20 g of each oil sample (IUPAC, 1987) and isolated by means of low-pressure column chromatography using light petroleum as eluent. After discarding a first fraction, a second one is collected and analyzed by GC on a fused-silica capillary column (30 m x 0.25 mm i.d., 0.1 µm film thickness) coated with 5% diphenyl 95%-methylpolysiloxane. The oven temperature program was: 235 °C for 6 min, and then to rise to 285 °C at 2 °C·min⁻¹. Hydrogen was used as carrier gas at a flow rate of 1 mL·min⁻¹

Statistical analysis

Each analytical determination was done in triplicate and data were presented as their average and standard deviation.

3. RESULTS AND DISCUSSION

Sacha inchi from the *P. volubilis* and *P. huayllabambana* ecotypes are the most extended cultivars among their gender, although the first one is the most common for oil production and commercialization. The oils from all the *plukenetia* gender have to fulfill with the quality parameters specified in the Peruvian regulation (NTP, 2010), although specific parameters are established for the *P. volubilis* and *P. huayllabambana* ecotypes as far as the characterization parameters are concern, since it is known that both of them differ slightly in their composition.

3.1. Analytical indexes and oxidative stability

The majority of the values for the indexes commonly used to evaluate the initial quality of sachá inchi commercial oils, i.e., FA, and PV, were found within the expected ranges for oils of good quality (Table 1). All of the studied samples could be classified as extra virgin according to the acidity values included in the regulation with the exception of C-2, C-11, C-12 and C-13 samples. According to this quality parameter the mentioned samples could be

TABLE 1. Physicochemical parameters of the sachá inchi commercial oils under study.

	Acidity (%)	Peroxide Value (meqO ₂ ·kg ⁻¹ oil)	Iodine Index ^a (gI ₂ ·100g ⁻¹)	Refractive Index	Density (g·cm ⁻³)	Viscosity (mPa·s)	Rancimat (h)	Unsaponifiable Matter (%)
C-1	0.7±0.00	4.84±0.06	191.9±0.1	1.4780	0.9265±0.0000	52.3±0.26	6.6±0.1	1.03±0.12
C-2	2.0±0.05	8.38±0.08	196.4±0.2	1.4790	0.9263±0.0000	45.1±0.25	3.4±0.1	2.15±0.25
C-3	0.6±0.00	15.37±0.06	116.1±0.1	1.4710	0.9156±0.0000	65.3±0.48	3.4±0.1	1.52±0.02
C-4	0.2±0.01	6.37±0.04	203.6±0.1	1.4800	0.9264±0.0000	47.2±0.80	3.9±0.1	2.58±0.30
C-5	0.5±0.00	8.87±0.12	153.3±0.1	1.4750	0.9265±0.0000	55.9±0.41	5.4±0.1	1.62±0.09
C-6	0.9±0.01	3.86±0.09	135.6±0.1	1.4730	0.9270±0.0000	55.7±0.68	8.3±0.1	2.33±0.76
C-7	0.4±0.01	3.85±0.10	147.4±0.7	1.4750	0.9384±0.0000	53.7±0.32	10.3±0.1	1.69±0.18
C-8	0.3±0.01	7.26±0.02	198.8±0.1	1.4801	0.9293±0.0000	50.7±0.45	4.8±0.1	1.44±0.13
C-9	0.2±0.01	6.39±0.01	161.9±0.1	1.4762	0.9199±0.0000	46.8±0.25	9.2±0.1	1.48±0.05
C-10	1.6±0.02	7.2±0.01	203.4±0.1	1.4805	0.9291±0.0000	49.6±0.34	3.9±0.1	1.05±0.20
C-11	0.5±0.01	7.6±0.01	164.8±0.1	1.4751	0.9241±0.0000	58.5±0.38	7.6±0.1	1.85±0.15
C-12	1.8±0.10	4.00±0.01	202.1±0.1	1.4805	0.9288±0.0000	50.1±0.45	3.9±0.1	1.05±0.10
C-13	2.0±0.01	7.15±0.01	198.4±0.1	1.4800	0.9289±0.0000	48.8±0.35	3.8±0.1	0.82±0.06
C-14	0.7±0.02	9.70±0.01	220.8±0.1	1.4800	0.9289±0.0000	47.8±0.27	3.2±0.1	1.00±0.19
NTP ^c	<1% <2%	<10	183–199 ^b	1.478–1.481	0.926–0.931			<0.36

^aUNE-EN 14111:2003.

^bHANUS.

^cPeruvian regulation.

classified as virgin but not as extra virgin because sachu inchi regulation establishes maximum values of 1.0 and 2.0 (expressed as a percent of oleic acid) for extra and for virgin oils, respectively. One of the analyzed samples (C-3) exceed the maximum PV of $10 \text{ meq O}_2 \cdot \text{kg}^{-1}$, what meant that this sample presented an oxidation level higher than that expected for good oils.

The iodine index is a determination directly related to the presence of double bonds in the FA of the oil. We determined it taking into account a response factor for each unsaturated FA according to the normative UNE.

The refractive index, density and viscosity, at $20 \text{ }^\circ\text{C}$ were in the ranges: 1.4701–1.4805, $0.9156\text{--}0.9384 \text{ g}\cdot\text{cm}^{-3}$, and $45.1\text{--}65.3 \text{ mPa}\cdot\text{s}$, respectively. The C-3, C-7 and C-9 samples were out of the ranges established in the regulation for both, refractive index and density. The C-5, C-6, and C-11 samples complied with the density range but not with the refractive index.

In relation to the unsaponifiable matter, we believe that there must be an error since the current regulations restrict this parameter to 0.36% and important compounds such as sterols and tocopherols are included in this fraction and it has no sense to establish a limit. In fact, all the studied commercial oil samples have percentages of the unsaponifiable matter in the 1.00–2.69% range, higher than the current limit.

The oil stability measured as Rancimat hours at $100 \text{ }^\circ\text{C}$ were in the 3.2–10.3 range. The values were closely related to the unsaturation of the oil sample and thus, the higher the iodine value the lower the Rancimat hours. The degree of oxidation also influences the lost of hours of stability, for that reason C-3 sample, with the lowest iodine value, would be expected to have the highest stability although it was not the case.

3.2. Major components

The oils from the *P. volubilis* ecotype must have at least 44.7% α -linolenic acid whereas oils from the *P. huayllabambana* should contain 55.5%, according to the regulation. Our previous results, with authentic sachu inchi oils extracted by us through a mechanical process, utilizing seeds from both varieties complied with the regulation (Chasquibol *et al.*, 2012; Chasquibol *et al.*, 2014). Therefore, we are in position of expecting at least 44.7% linolenic, 32.1–24.9% linoleic, and 8.9–7.9% oleic acid contents in the authentic sachu inchi oils, assuming that the ecotypes are actually *P. volubilis* or *P. huayllabambana*. Nevertheless, as we can see in Table 2, C-3, C-5, C-6, C-7, C-9, and C-11 samples do not satisfy this requirement. Besides, four of the above mentioned samples (C-3, C-5, C-6 and C-11) contain *trans*-FA that, although is a parameter not included

in the current regulation, is determined at the same time than the *cis*-FA. Actually, one of the commercial samples specified on its label the absence of *trans*-fat since these are indicative of a heated or a bleaching process, both normal steps of the refining procedure.

In the oil extraction process from seeds, a previous step consisting of subjecting the seed to roasting by applying high temperatures ($<180 \text{ }^\circ\text{C}$) is usual. The purpose of this step is to eliminate humidity, and to facilitate the extraction by eliminating the astringent off-flavors and anti-nutritional compounds, besides increasing the oil stability (Cisneros *et al.*, 2014). We have not found documentation about the formation of *trans*-isomers in roasted seeds, but the presence of these compounds in the analyzed commercial oils could be indicative of a process where temperature is involved. In any case, the presence of steroidal hydrocarbons would confirm unmistakably presence of refined oil.

Finally, according to its FA composition (59.37% Ln, 7.99% oleic, and 26.65% linoleic acids) sample number 14 could clearly belong to the *P. huayllabambana* ecotype.

As far as TG are concern, those with ECN 38 (LnLnL), 40, and 42 (formed by combinations of Ln, L, O and P) are the main ones described for genuine sachu inchi oils in the limited publications that we have found on this subject (Fanali *et al.*, 2014; Chasquibol *et al.*, 2014). Also for genuine sachu inchi oils the TG with ECN 46 and 48 are present in percentages lower than 10%, and the TG with ECN 50 (SSO) are present in percentages below 1%. The most remarkable results are the TG with ECN 46 in the range 15.6–27.3 (Table 3) found in a few commercial oil samples (3, 5, 6, 7, 9 and 11). Also percentages of the TG with ECN 50 (3.0–5.1) higher than the 1% limit for authentic sachu inchi oils were found. The abnormal results belong to the same oils whose FA which do not match with a typical sachu inchi profile.

3.3. Minor unsaponifiable compounds

Sterols and tocopherols are the main components of the unsaponifiable matter present in the majority of the edible vegetable oils and they are known to display important health benefits. Their presence in sachu inchi oils is of great importance, qualitative and quantitatively speaking. Together with high amount of total sterols, sachu inchi oils have a particular sterol profile because, in contrast to other vegetable oils, the main sterol after β -sitosterol is stigmasterol instead of campesterol (Chasquibol *et al.*, 2014). The calculation of the stigmasterol/campesterol ratio from data available in the literature regarding authentic sachu inchi oils, drove us to a ratio equal to 4 (2.7–7.0) (Bondioli *et al.*, 2006; Castaño *et al.*, 201; Chirimos *et al.*,

TABLE 2. Fatty acid methyl esters (wt% of the total) of the sachá inchi commercial oils under study determined by GC.

Commercial sample oils	Fatty acid methyl esters (wt% of the total)														<i>trans</i>
	C 16:0	C 16:1	C 17:0	C 18:0	C 18:1	C 18:2	C 20:0	C 18:3	C 20:1	C 22:0	C 24:0				
C-1	8.83±0.23	0.70±0.04	0.08±0.02	2.06±0.03	9.91±0.40	33.17±0.08	0.52±0.05	44.73±0.07	< 0.01	nd	nd	nd	nd	nd	
C-2	6.47±0.00	0.34±0.00	0.08±0.00	1.78±0.00	12.30±0.01	33.05±0.03	0.21±0.00	45.77±0.02	< 0.01	nd	nd	nd	nd	nd	
C-3	14.06±0.11	0.99±0.01	0.08±0.01	3.73±0.03	36.64±0.02	36.80±0.02	0.34±0.00	5.71±0.03	< 0.01	0.20±0.01	0.10±0.01	1.35±0.01			
C-4	3.80±0.02	0.03±0.04	0.11±0.00	2.95±0.02	9.58±0.09	35.46±0.08	0.24±0.02	47.83±0.01	< 0.01	nd	nd	nd	nd	nd	
C-5	9.37±0.04	0.55±0.01	0.11±0.02	3.42±0.00	26.21±0.01	36.09±0.03	0.35±0.01	23.37±0.01	< 0.01	< 0.01	< 0.01	0.63±0.05			
C-6	10.81±0.03	0.09±0.00	0.12±0.00	4.64±0.03	21.39±0.10	50.99±0.02	0.78±0.01	8.79±0.01	< 0.01	0.40±0.01	0.20±0.01	1.79±0.05			
C-7	9.37±0.01	0.09±0.02	0.09±0.00	4.27±0.03	22.21±0.26	50.56±0.47	0.31±0.00	13.11±0.14	< 0.01	nd	nd	nd	nd	nd	
C-8	4.35±0.01	0.05±0.00	0.09±0.01	3.19±0.01	10.50±0.01	36.92±0.08	0.12±0.01	44.78±0.03	< 0.01	nd	nd	nd	nd	nd	
C-9	9.11±0.04	0.07±0.00	0.09±0.00	2.87±2.18	20.98±0.10	50.29±0.26	0.37±0.00	15.88±0.15	< 0.01	nd	nd	nd	nd	nd	
C-10	4.15±0.01	nd	0.05±0.00	3.15±0.03	10.05±0.09	37.05±0.19	0.20±0.01	45.05±0.05	0.30±0.01	nd	nd	nd	nd	nd	
C-11	8.65±0.05	0.04±0.01	0.04±0.00	4.05±0.04	18.05±0.07	46.05±0.16	0.65±0.02	20.71±0.03	0.15±0.01	0.35±0.01	0.15±0.01	1.11±0.05			
C-12	4.15±0.02	0.45±0.03	0.15±0.01	3.15±0.02	10.25±0.05	36.45±0.12	0.25±0.01	44.91±0.04	0.25±0.01	nd	nd	nd	nd	nd	
C-13	4.34±0.02	0.16±0.02	0.06±0.01	3.06±0.03	10.87±0.05	34.43±0.12	0.19±0.01	46.61±0.06	0.30±0.01	nd	nd	nd	nd	nd	
C-14	3.01±0.02	nd	0.16±0.06	2.31±0.02	7.99±0.03	26.65±0.13	0.25±0.02	59.37±0.46	0.26±0.03	nd	nd	nd	nd	nd	
NTP ^a	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Volubilis</i>	-	-	-	-	8.9	32.1	-	44.7	-	-	-	-	-	-	
<i>Huaylla bambana</i>	-	-	-	-	7.9	24.0	-	55.0	-	-	-	-	-	-	

^aPeruvian regulation.

TABLE 3. Main tryglicerides grouped by their ECN determined by RP-HPLC-RI

Commercial sample oils	Equivalent Carbon Number							
	ECN36	ECN38	ECN40	ECN42	ECN44	ECN46	ECN48	ECN50
C-1	12.4±0.80	22.2±1.14	23.0±0.48	22.0±0.98	9.8±0.47	4.9±0.21	5.5±2.79	0.7 ±0.29
C-2	17.7±0.40	23.5±0.45	18.7±0.41	18.7±0.37	8.6±0.14	6.3±0.58	5.9±0.62	0.6 ±0.08
C-3	1.4±0.77	2.8±0.20	5.5±0.71	13.5±0.52	21.0±0.66	27.3±0.49	25.4±2.31	3.1 ±0.16
C-4	12.3±0.11	23.2±0.03	25.1±0.07	23.7±0.32	11.0±0.16	3.6±0.18	0.8±0.02	0.3 ±0.02
C-5	6.9±0.32	13.3±0.58	16.3±0.47	19.5±0.54	19.6±0.13	15.8±0.62	5.6±0.55	3.0 ±0.98
C-6	1.4±0.01	2.8±0.08	8.0±0.21	23.7±0.15	27.1±0.53	18.4±0.10	13.5±1.33	5.1±0.08
C-7	1.9±0.12	4.8±0.23	10.2±0.38	23.6±0.51	26.4±0.07	17.3±0.57	12.7±0.45	3.1 ±0.21
C-8	10.4±0.13	19.8±1.34	23.1±1.24	23.1±0.36	13.3±0.33	6.5±0.88	3.1±1.19	0.7 ±0.43
C-9	2.6±0.05	6.5±0.14	12.2±0.02	24.9±0.30	26.5±0.23	16.3±0.19	8.9±0.03	2.1 ±0.18
C-10	11.1±0.05	22.7±0.15	24.4±0.16	23.7±0.13	11.9±0.10	4.7±0.05	1.0±0.01	0.5±
C-11	4.9±0.07	9.6±0.08	16.2±0.10	27.1±0.15	25.0±0.16	15.6±0.10	1.0±0.01	0.6±0.05
C-12	10.2±0.06	20.9±0.14	25.6±0.15	24.3±0.14	12.3±0.12	4.6±0.06	1.3±0.01	0.8±0.06
C-13	11.9±0.10	21.9±0.16	23.8±0.12	23.9±0.14	11.8±0.10	4.6±0.06	1.4±0.02	0.7±0.08
C-14	10.9±0.11	25.9±0.15	23.7±0.14	22.6±0.12	11.4±0.11	3.8±0.05	1.5±0.02	0.2±0.01

ECN (Equivalent carbon number).

2013; Chirimos *et al.*, 2015). Studying the data on Table 4, we observe that C-3, C-5, C-6, C-7, C-9, and C-11 samples are clearly out of this peculiarity. Besides, some samples of this group (C-5, C-6, C-7, C-9) show small percentages (0.2–0.4%) of Δ -5-stigmastadienol, an sterol which appears when the oil is subjected to refining.

Data presented in Table 5 show the high quantities of total tocopherols present in the studied samples. This parameter is included in the regulation where it is specified that sacha inchi oils must have at least 1900 mg·kg⁻¹ of total tocopherols. As one can observe, in spite of the high quantities, not all of the samples fulfill this requirement thus, C-3, C-5, C-6, C-7, C-9 and C-11 samples have lower values than required. Besides, usually sacha inchi oils have negligible quantities of α -tocopherol and here the commercial oils (C-3, C-6, C-7 and C-9) show quantities higher above 50 mg·kg⁻¹.

The C-8 sample fulfill the requirements: its total tocopherol concentration is 2013.6 mg·kg⁻¹, however two unknown peaks, not present in a normal tocopherol chromatogram, appear between the retention times corresponding to α - and β -tocopherols. We have approached the study of their structure by GC-MS using a mass spectrometer Polaris Q (Thermo Finnigan, Manchester, UK) coupled to a gas chromatograph TRACE fitted with a DB-e MS.

For one of the peaks, the molecular ion (M⁺) was detected at m/z 354 and the base peak at m/z 149. For this compound the most important mass ion fragments are: 204, 203, 161, 150, 135 and 121. The second peak has an M⁺ to m/z 370 and a base peak 135 and its most important mass ion fragments are: 249,

233, 203, 150, 149 and 135. To confirm the structures those compounds were analyzed by HPLC-MS-TOF, using a MICROTOF (Bruker Daltonic), allowing detect the exactly mass of both compounds being 377.0960 (M+Na)⁺ and 393.0908 (M+Na)⁺. All the above results are consistent with the spectra described for sesamin and sesaminol. These two compounds are flavons present in sesame oils. Studying all the results corresponding to the C-8 sample we can observe that there is nothing that indicates that the oil has been diluted with sesame oil or with another kind of oil or fat because all its analytical parameters are within the regulation. Therefore, we believe that this sample has been contaminated with other kind of product in the same company or the oil was intended to be improved through an additive.

Hydrocarbons are the third group of compounds belonging to the most non-polar fraction of the unsaponifiable matter studied here. Lineal saturated hydrocarbons from 8 to 35 carbon atoms are compounds with demonstrated utility in oil characterization (Moreda *et al.*, 2001). The H27:0, H29:0 and H31:0 odd elements are the most abundant in sunflower, safflower and sesame oils, whereas H23:0, H25:0, H27:0 and H29:0 are detected in virgin olive oil. Our previous studies with genuine sacha inchi oils gave us a HC profile where the highest concentrations were obtained for H23:0 and H25:0, and those with even carbon-atom, mainly H24:0, were also present.

The results presented in Table 6 show H23:0 as major HC (22–40%), with two clear exceptions: C-3 and C-11 samples, where H27:0 is the major aliphatic saturated hydrocarbon, which indicates the presence of another seed oil.

TABLE 4. Sterol composition (wt % on total) and total quantities of the Sacha inchi commercial oils determined by GC.

Samples	Cholesterol	24-Methyl-cholesterol	Campesterol	Campestanol	Stigmasterol	$\Delta 7$ -Campesterol	$\Delta 5,23$ -Stigmasteradienol	Clero-sterol	β -Sitosterol	Sitostanol	$\Delta 5$ -Avena-sterol	$\Delta 5,24$ -Stigmasteradienol	$\Delta 7$ -Stigma-sterol	$\Delta 7$ -Avena-sterol	TOTAL STEROLS (mg·kg ⁻¹)
C-1	0.3±0.0	nd	5.5±0.7	nd	23.5±1.7	0.1±0.0	0.2±0.0	0.7±0.1	63.6±2.3	1.9±0.1	2.4±0.1	0.8±0.45	0.6±0.1	0.4±0.1	2060.4±96.7
C-2	0.3±0.1	0.7±0.2	5.1±0.2	0.1±0.1	25.3±0.4	nd	nd	0.9±0.2	63.2±1.2	1.0±0.1	2.2±0.4	0.4±0.13	0.5±0.1	0.3±0.1	2001.4±10.4
C-3	0.3±0.0	0.3±0.0	12.3±0.1	0.6±0.1	10.4±0.1	nd	nd	0.9±0.1	66.2±3.3	1.4±0.1	4.3±0.6	1.0±0.22	1.1±0.2	1.2±0.2	2462.5±97.8
C-4	0.3±0.0	0.3±0.1	7.4±0.2	0.3±0.1	26.5±1.1	nd	nd	0.8±0.1	56.1±3.0	2.1±0.1	5.2±0.1	0.5±0.17	0.2±0.0	0.4±0.1	2204.5±95.3
C-5	0.3±0.0	1.5±0.1	9.6±0.2	1.0±0.1	15.8±1.0	nd	0.1±0.0	0.6±0.1	62.3±2.1	1.6±0.1	5.6±0.1	0.8±0.1	0.8±0.1	0.4±0.1	2502.0±10.8
C-6	0.4±0.1	0.6±0.1	18.9±1.3	0.8±0.1	19.3±1.0	0.3±0.1	0.3±0.0	0.8±0.1	50.8±3.1	1.6±0.1	1.4±0.1	1.5±0.1	2.3±0.1	0.9±0.1	2952.0±80.6
C-7	0.4±0.0	0.9±0.1	18.3±1.4	1.2±0.1	18.0±1.0	nd	0.4±0.0	0.9±0.1	51.4±3.0	1.7±0.1	2.8±0.1	1.2±0.1	2.1±0.1	0.8±0.1	2950.0±56.8
C-8	0.2±0.0	0.5±0.1	10.2±1.0	0.5±0.1	23.2±1.0	0.3±0.1	nd	0.9±0.1	54.7±2.1	2.6±0.1	5.6±0.1	0.7±0.1	0.3±0.0	0.3±0.0	2475.4±98.0
C-9	0.4±0.0	0.7±0.1	18.3±2.0	0.8±0.1	19.4±1.0	0.6±0.1	0.2±0.0	0.8±0.1	50.6±3.5	1.8±0.2	1.9±0.1	1.6±0.2	2.2±0.1	0.9±0.1	2469.2±79.9
C-10	0.1±0.0	nd	7.3±0.9	0.1±0.1	25.8±1.0	nd	nd	1.0±0.01	56.2±2.6	0.9±0.1	7.9±0.1	0.2±0.1	0.1±0.0	0.3±0.0	2327.8±70.5
C-11	0.3±0.0	nd	16.5±0.1	nd	19.8±1.0	nd	nd	1.8±0.1	52.6±2.4	1.2±0.2	4.2±0.1	0.3±0.0	2.0±0.1	1.2±0.1	2792.0±65.6
C-12	0.1±0.0	nd	7.6±0.1	nd	27.4±1.3	nd	nd	1.1±0.1	55.1±2.2	0.9±0.1	7.2±0.5	0.2±0.0	0.1±0.0	0.3±0.0	2323.6±68.9
C-13	0.1±0.0	nd	7.1±0.1	nd	24.7±1.1	nd	nd	1.1±0.1	56.3±2.1	1.0±0.2	6.8±0.4	0.2±0.0	0.2±0.0	0.3±0.0	2386.9±70.1
C-14	0.1±0.0	nd	7.5±0.1	nd	26.4±1.0	1.0±0.1	nd	1.1±0.1	55.5±2.4	0.6±0.2	7.1±0.5	0.3±0.0	0.1±0.0	0.3±0.1	2393.0±75.8

TABLE 5. Tocopherol composition of the sachu inchi commercial oils under study determined by normal Si-HPLC-FL.

Commercial sample oils	Tocopherol (mg·kg ⁻¹)			TOTAL (mg·kg ⁻¹)
	α -Tocopherol	β -Tocopherol	γ -Tocopherol	
C-1	48.0±1.6	nd	1565.2±30.1	2400.4±15.91
C-2	20.1±0.0	nd	1736.4±12.4	2498.2±19.40
C-3	126.5±0.30	10.5±2.1	463.8±1.0	798.4±9.30
C-4	4.2±1.1	nd	1577.5±7.5	2815.9±19.92
C-5	43.0±0.6	3.8±0.3	808.4±12.9	1438.9±4.63
C-6	91.2±2.6	11.0±1.1	846.7±5.7	1295.8±7.27
C-7	55.8±15.8	nd	723.0±43.4	1167.3±111.9
C-8	nd	nd	1170.7±19.9	2013.6±38.22
C-9	89.5±8.98	nd	1002.3±46.7	1484.1±12.66
C-10	nd	nd	1781.9±10.5	2604.4±14.5
C-11	nd	nd	618.7±5.7	968.2±10.5
C-12	nd	nd	1729.8±11.4	2565.3±15.6
C-13	nd	nd	1798.1±10.3	2671.9±10.5
C-14	nd	nd	1907.2±11.7	2614.2±10.1
NTP ^a				>1900

^aPeruvian regulation.

TABLE 6. Main aliphatic hydrocarbons of the sacha inchi commercial oils under study determined by on-column GC.

Commercial oil samples	ALIPHATIC HYDROCARBONS (wt% on total)									TOTAL (mg·kg ⁻¹)
	H21:0	H22:0	H23:0	H24:0	H25:0	H26:0	H27:0	H28:0	H29:0	
C-1	11.0±0.0	10.9±0.9	38.2±1.7	18.2±0.9	14.7±0.5	2.0±0.1	2.0±0.1	1.5±0.1	1.5±0.1	114.6±1.7
C-2	11.1±0.0	11.0±0.00	39.1±1.6	18.7±0.9	14.9±0.8	1.9±0.1	1.4±0.2	0.5±0.1	1.4±0.1	131.3±5.4
C-3	7.1±0.1	6.9±0.7	18.6±1.5	10.5±0.8	18.9±0.9	10.5±0.1	20.2±1.2	6.2±0.1	1.1±0.1	21.7±0.3
C-4	10.5±0.1	13.8±0.9	38.4±1.7	22.4±0.9	11.2±0.9	0.4±0.1	0.9±0.1	1.3±0.1	1.3±0.1	92.7±7.1
C-5	10.4±0.1	13.3±0.9	33.2±1.4	19.5±0.9	10.7±0.9	2.1±0.1	4.4±0.5	4.7±0.1	1.7±0.1	34.8±1.1
C-6	8.3±0.0	9.8±0.9	22.6±1.7	18.4±0.7	13.5±0.8	11.5±0.6	7.5±0.5	7.3±0.1	1.1±0.1	12.3±0.7
C-7	10.6±0.0	13.9±0.8	29.8±1.5	19.2±0.9	9.7±0.9	3.7±0.1	5.9±0.5	5.6±0.1	1.5±0.1	14.6±0.1
C-8	11.6±0.1	14.0±0.9	35.3±1.8	20.1±1.0	9.9±0.9	2.5±0.1	2.5±0.5	3.0±0.2	1.1±0.1	60.6±6.7
C-9	8.0±0.7	12.4±0.9	26.8±1.2	17.2±0.9	9.0±0.9	5.8±0.4	5.1±0.5	14.6±0.6	0.9±0.1	22.3±1.3
C-10	10.9±0.8	12.8±0.9	39.3±1.9	21.3±0.9	10.0±0.9	0.8±0.1	1.9±0.1	1.7±0.1	1.3±0.1	100.5±1.5
C-11	8.1±0.7	7.2±0.8	18.0±1.7	9.2±0.6	19.1±0.9	9.2±0.5	21.0±1.1	7.2±0.1	1.0±0.1	30.6±1.3
C-12	10.3±0.9	13.0±0.9	38.8±1.7	21.8±0.9	12.2±0.8	0.4±0.1	0.9±0.1	1.1±0.1	1.5±0.1	90.1±1.9
C-13	11.0±0.8	13.4±0.9	39.5±1.8	22.5±0.7	10.3±0.1	0.3±0.1	0.8±0.0	1.0±0.1	1.2±0.1	85.7±1.5
C-14	11.6±0.9	12.9±0.9	40.5±1.5	19.9±0.9	10.6±0.1	0.7±0.1	1.1±0.1	1.3±0.1	1.4±0.1	99.0±1.3

In relation to the steroidal hydrocarbons, they come from dehydration of the sterols during the different steps of the refining process, (Lanzón *et al.*, 1994). Their formation during the heating process which take place when the seeds are roasted is also documented (Crew *et al.*, 2006). In the present study no sterenes or steroidal hydrocarbons have been found. This fact guarantees the absence of refined oil in the commercialized samples under study.

4. CONCLUSIONS

The sacha inchi oils analyzed here include *P. volubilis* as the main ecotype commercialized, although the analytical parameters of one of the studied samples are clearly consistent with the analytical parameters required for *P. huayllabambana*. Probably as a consequence of their high unsaturation level, the oils are very susceptible to be degraded by hydrolysis and by oxidation thus, quality parameters such as the acidity value and the PV do not correspond to extra virgin oils as is specified in all the labels. All of the oils contain percentages of unsaponifiable matter much higher than the limit established in the regulation which, on the other hand, does not seem to be very consistent with having high content of valuable compounds as are sterols or tocopherols. The fatty acid composition alert us about the existence of six samples which clearly do not comply with the percentages required for ω -3, ω -6 and ω -9 fatty acids or with the characteristic of having greater content of stigmaterol than campesterol presented in genuine oils. Although there is no presence of steroidal hydrocarbons, some of

the samples, or more probably their seeds, may have been subjected to high temperature because *trans* isomers of fatty acids are present in quantities much higher than those expected for crude or virgin oils.

ACKNOWLEDGEMENTS

The financial support of Agencia Española de Cooperación Internacional para el Desarrollo (AECID) is gratefully acknowledge (Project AP/036672/11) and the Instituto de Investigación Científica (IDIC) de la Universidad de Lima-Perú (PI.56.006.2015). The authors would like to thank Mr. JL Rios for his assistance in the MS analysis.

REFERENCES

- AOCS. Official methods of analysis. 2012. Method Ca 5a-40. Determination of free fatty acids. Official methods and recommended practices of the AOCS; American Oil Chemists' Society: Champaign, IL, USA.
- AOCS. Official methods of analysis. 2013. Method Ja 8-87. Determination of peroxide value. Official methods and recommended practices of the AOCS; American Oil Chemists' Society: Champaign, IL, USA.
- AOCS. Official methods of analysis. 1997. Method Cd 12b-92. Sampling and analysis of commercial fats and oils: Oil stability index. Official methods and recommended practices of the AOCS; American Oil Chemists' Society: Champaign, IL, USA.
- Bondioli P, Della Bella L, Rettke P. Alpha linolenic acid rich oils. 2006. Composition of *Plukenetia volubilis* (Sacha Inchi) oil from Peru. *Riv. Ital. Sostanze Grasse* **83**, 120–123.
- Bussmann R, Téllez C, Glenn A. 2009. *Plukenetia huayllabambana* (Euphorbiaceae) from upper Amazon of Perú. *Nord. J. Bot.* **27**, 313–315. <http://dx.doi.org/10.1111/j.1756-1051.2009.00460.x>
- Castañó DL, Valencia MP. 2012. Fatty acid composition of Inca peanut (*Plukenetia volubilis* L.) *Rev. Chil. Nutr.* **39**, 45–52.

- Cert A, Moreda W, García-Moreno J. 1997. Determinación de esteroides y dialcoholes triterpénicos en aceite de oliva mediante separación de la fracción por cromatografía líquida de alta eficacia y análisis por cromatografía de gases. Estandarización del método analítico. *Grasas Aceites* **48**, 207–218. <http://dx.doi.org/10.3989/gya.1997.v48.i4.791>
- Cert A, Lanzón A, Carelli A, Albi T. 1994. Formation of stigmasta-3,5-diene in vegetable oils. *Food Chem.* **49**, 287–293. [http://dx.doi.org/10.1016/0308-8146\(94\)90173-2](http://dx.doi.org/10.1016/0308-8146(94)90173-2)
- Chasquibol N, Moreda W, Yácono JC, Pez-Camino MC. 2012. Preliminary characterization studies on sacha inchi (*Plukenetia volubilis* L.) seeds and oils grown in San Martín, Perú. 20th International symposium on plant lipids (ISPL), Sevilla, Spain. p 167.
- Chasquibol NA, del Aguila C, Yácono JC, Guinda Á, Moreda W, Gómez-Coca RB, Pez-Camino MC. 2014. Characterization of Glyceridic and Unsaponifiable Compounds of Sacha Inchi (*Plukenetia huayllabambana* L.) Oils. *J. Agric. Food Chem.* **62**, 10162–10169. <http://dx.doi.org/10.1021/jf5028697>
- Chirinos R, Zuloeta G, Pedreschi R, Mignolet E, Larondelle Y, Campos D. 2013. Sacha inchi (*Plukenetia volubilis*): A seed source of polyunsaturated fatty acid, tocopherols, phyosterols, phenolic compounds and antioxidant capacity. *Food Chem.* **171**, 1732–1739. <http://dx.doi.org/10.1016/j.foodchem.2013.04.078>
- Chirinos R, Pedreschi R, Domínguez G, Campos D. 2015. Comparison of the physic-chemical and phytochemical characteristics of the oil of two *Plukenetia* species. *Food Chem.* **173**, 1203–1206. <http://dx.doi.org/10.1016/j.foodchem.2014.10.120>
- Cisneros FH, Paredes D, Arana A, Cisneros-Zevallos L. 2014. Chemical composition, oxidative stability and antioxidant capacity of oil extracted from roasted seeds of sacha inchi (*Plukenetia volubilis* L.). *J. Agric. Food Chem.* **62**, 5191–5197. <http://dx.doi.org/10.1021/jf500936j>
- Crews C, Hough P, Brereton P, Godward J, Lees M, Guiet S, Winkelmann W. 2006. Quantitation of the Main Constituents of Some Authentic Sesame Seed Oils of Different Origin. *J. Agric. Food Chem.* **54**, 6266–6270. <http://dx.doi.org/10.1021/jf0603578>
- Dobarganes MC, Cert A, Diffenbacher A. 1999. The determination of stigmastadienes in vegetable oils. Results of collaborative studies and the standardized method. *Pure Appl. Chem.* **71**, 349–359. <http://dx.doi.org/10.1351/pac199971020349>
- Gómez-Coca RB, Pérez-Camino MC, Moreda W. 2015. Neutral lipids: Unsaponifiable matter. In *Handbook of Food Analysis*, 3rd ed., CRC Press. Taylor & Francis Group: US, pp. 459–489.
- Gómez-Coca RB, Cert R, Pérez-Camino MC, Moreda W. 2016. Determination of saturated aliphatic hydrocarbons in vegetable oils. *Grasas Aceites* **67**, e127. <http://dx.doi.org/10.3989/gya.0627152>
- Gutiérrez LF, Rosada LM, Jiménez A. 2011. Chemical composition of sacha inchi (*Plukenetia volubilis* L.) seeds and characteristics of their lipid fraction. *Grasas Aceites* **62**, 76–83. <http://dx.doi.org/10.3989/gya044510>
- International Standard Office, ISO 15788-2 (2003). Animal and vegetable fats and oils. Determination of stigmastadienes in vegetable oils.
- IUPAC Standard Method 2.302. 1987a. Standard methods for the analysis of oils, fats and derivatives. Determination of FAMES by capillary GC. Blackwell Scientific: Oxford, Great Britain.
- IUPAC Standard Method 2.432. 1987b. Standard methods for the analysis of oils, fats and derivatives. Determination of tocopherol and tocotrienols in vegetable oils and fats by HPLC. Blackwell Scientific: Oxford, Great Britain.
- IUPAC Standard Method 2.401. 1987. Standard methods for the analysis of oils, fats and derivatives. Determination of the unsaponifiable matter. Blackwell Scientific: Oxford, Great Britain.
- Lanzón A, Albi T, Cert A, Gracián J. 1994. The hydrocarbon fraction of virgin olive oil and changes resulting from refining. *J. Am. Oil Chem. Soc.* **71**, 285–291. <http://dx.doi.org/10.1007/BF02638054>
- Liu Q, Xua Y, Zhanga P, Naa Z, Tanga T, Shia Y. 2014. Chemical composition and oxidative evolution of Sacha Inchi (*Pluketia volubilis* L.) oil from Xishuangbanna (China). *Grasas Aceites* **65**, e012. <http://dx.doi.org/10.3989/gya.075713>
- Maurer NE, Hatta-Sakoda B, Pascual-Chagman G, Rodriguez-Saona LE. 2012. Characterization and authentication of a novel vegetable source of omega-3 fatty acids, sacha inchi (*Plukenetia volubilis* L.) oil. *Food Chem.* **134**, 1173–1180. <http://dx.doi.org/10.1016/j.foodchem.2012.02.143>
- Moreda W, Pérez-Camino MC, Cert A. 2001. Gas and liquid chromatography of hydrocarbons in edible vegetable oils. *J. Chromatogr. A.* **936**, 159–171. [http://dx.doi.org/10.1016/S0021-9673\(01\)01222-5](http://dx.doi.org/10.1016/S0021-9673(01)01222-5)
- Moreda W, Pérez-Camino MC, Cert A. 2003. Improved method for the determination of triacylglycerols in olive oils by high performance liquid chromatography. *Grasas Aceites* **54**, 175–179. <http://dx.doi.org/10.3989/gya.2003.v54.i2.262>
- Muñoz Jauregui A, Alvarado-Ortiz C, Castañeda B, Lizaraso F, Barnett E, Cárdenas L, Manco E. 2013. Estudio nutricional de *Plukentia huayllabambana*. *Rev. Soc. Quím. Perú* **79**, 47–56.
- NTP. 2009. Norma Técnica Peruana 151.400. Requisitos aceite sacha inchi, INDECOPI: Lima, Peru, 2010.
- NTP. 2014. Norma Técnica Peruana 151.400, amendment to NTP 151.400, 2009. Requisitos Aceite Sacha Inchi, INDECOPI: Lima, Perú, 2014.
- Ourrach I, Rada M, Pérez-Camino MC, Benaissa M, Guinda A. 2012. Detection of argan oil adulterated with vegetable oils: new markers. *Grasas Aceites* **63**, 355–364. <http://dx.doi.org/10.3989/gya.047212>
- Rodríguez A, Corazon-Guivin M, Cachique D, Mejia K, Del Castillo D, Renno J, García Dávila D. 2010. Diferenciación morfológica y por ISSR (inter simple sequencer repeats) de especies del género plukentia (Euphorbiaceae) de la amazonia peruana: Propuesta de una nueva especie. *Rev. Peru. Biol.* **17**, 325–330.
- UNE-EN 14111:2003. Fat and oil derivatives. Fatty Acid Methyl Esters (FAME). Determination of iodine value.