

OXIDATIVE STRESS IN HUMAN THYROID GLAND UNDER IODINE DEFICIENCY NODULAR GOITER: FROM HARMLESSNESS TO HAZARD DEPENDING ON COPPER AND IODINE SUBCELLULAR DISTRIBUTION

H. Falfushynska^{1,2}, L. Gnatyshyna^{1,2}, A. Shulgai¹, V. Shidlovski¹, O. Stoliar²

¹ I. YA. HORBACHEVSKY TERNOPIL STATE MEDICAL UNIVERSITY, TERNOPIL, UKRAINE

² V. HNATIUK TERNOPIL NATIONAL PEDAGOGICAL UNIVERSITY, TERNOPIL, UKRAINE

Background. *Thyroid disorders are the second most common endocrinopathies found in humans and animals. Determination of their key molecular markers presents a special interest.*

Objective. *We studied iodine and copper accumulation in nodular, paranodular and contralateral (not affected tissue by node) tissues of human thyroid gland in relation to the level of metal-binding proteins, potential antioxidants, and oxidative changes in tissue for this goal. Lower level of organificated iodine and higher level and mass fraction of inorganic iodine and copper in the nodular and paranodular tissue versus contralateral part of thyroid gland was established.*

Results. *The level of both metal-binding and apo-form of metallothioneins was higher. Content of reduced glutathione was lower in node-affected tissue compared to the contralateral part. Signs of oxidative stress (higher activity of superoxide dismutase, catalase, glutathione-transferase and level of oxyradicals) and cytotoxicity (higher cathepsin D activity, higher level of DNA strand breaks and glycolysis activation) in affected tissue were observed. The range of indice variability in paranodular tissue was smaller than in nodule compared to the parenchyma of contralateral part.*

Conclusions. *Excess of copper unbound to metallothionein in goitrous-changed tissue and high level of inorganic iodine could be the reason for elevated DNA fragmentation and increased lysosomal membrane permeability and activation of antioxidant defense. The main criterions of goiter formation were represented by low level of organificated iodine and high level of DNA damage in thyroid gland.*

KEY WORDS: iodine deficiency nodular colloidal goiter, iodine, copper, metallothioneins, oxidative stress, cytotoxicity

Introduction

Thyroid disorders are the second most common endocrinopathies found in humans and animals [1]. Determination of the key molecular markers is of considerable interest as they can be used to predict such pathologies. Iodine deficiency in thyroid pathology occurrence rate is rapidly increasing [2]. It has miscellaneous origins as a result of complex interaction of endogenous and exogenous factors and arises in the setting of high level of "nonspecific" goitrogens, among them copper, in the environment. Copper is an essential element for humans and animals, especially for antioxidant defense and the metabolism of the amino acid tyrosine, which is needed for the production of thyroid hormones [3]. Besides these ones, excessive amounts of copper in the body can pose a risk. The mechanisms underlying the acute toxicity effects of copper in humans are not well understood. It is reasonable to speculate that they probably represent a combination of significant oxidative stress at different body areas together with marked perturbations in several components of the endocrine system. It was shown

*Address for correspondence: H. Falfushynska, General Chemistry Department, I. Ya. Horbachevsky Ternopil State Medical University, m. Voli, 1, Ternopil, 46001, Ukraine
 Tel.: +380506782021;
 E-mail: halynka.f@gmail.com*

that the increase of copper level in the thyroid gland of patients with colloidal goiter was combined with the prooxidant changes in tissue [4]. It should be noted that Ternopil region with combination of moderate iodine deficiency and high scale of copper water pollution [5] presents an interest to study the relationship between copper accumulation and progress of thyroid pathology.

Free radical-mediated oxidative damage has been implicated in pathogenesis of many diseases. However, the reasons of this phenomenon are disputable under endemic nodular thyroid goiter. It is believed that the prooxidant effect of copper depends on the specific accumulation by metal binding protein, such as metallothioneins [6]. Metallothioneins (MTs) are low-molecular weight proteins of 6–7 kDa, with high content of cysteine (30 %) and complete absence of aromatic amino acids and histidine, capable of binding transition metals with high affinity. MTs synthesis can be induced by a variety of metals, cytotoxic agents, stress-producing conditions, cytokines, and glucocorticoid hormones [7]. Additionally, MTs can function as efficient scavengers of reactive oxygen species to preserve homeostasis of cells. The latest study have indicated a possible role of MTs as a tumor suppressor in papillary thyroid cancer [8] and

their part in the distribution of metals, thereby optimizing the function of thyroid gland [4].

Thus, the aim of this study was to evaluate the relation between the function of MTs and oxidative stress in node, paranodular and non-affected by node contralateral part tissue of human thyroid gland under iodine deficiency endemic nodular thyroid goiter. Molecular markers of cytotoxicity were used to assess the severity of the pathological process.

Methods

The target population in this study were the people with unilateral euthyroid iodine deficiency nodular thyroid goiter, who had lived in Ternopil Region for at least 22 years. Small samples of nodule, paranodular and contralateral (not affected by node tissue) tissues from thyroid gland were dissected in 25 patients. They were operated at the General Surgery Department of the local Ternopil Emergency Hospital. All experimental studies were conducted in accordance with the approval of the First National Congress on Bioethics (Kyiv, 2000) and the approval of the Bioethics Commission of I. Ya. Horbachevsky Ternopil State Medical University.

Tissue samples were homogenized (1/10 w/v) in 0.1 M phosphate buffer, pH 7.4, containing 100 mM KCl and 1 mM EDTA, as well as 0.1 mM phenylmethylsulfonyl fluoride (PMSF) for proteolysis inhibition for enzymatic measurements. Homogenates were centrifuged for 10 min at 6,000 *g*. Protein concentration in the supernatant (soluble protein) was measured by the method of Lowry et al. (1951), using bovine serum albumin as a protein standard. The absorbance values were measured on an UV/Vis spectrophotometer "Lomo-56" (Russia), and extinction/emission values were measured on the *f*-max fluorescence microplate reader [Molecular Device (USA)].

Reduced glutathione (GSH), 5,5'-dithio-bis(2-nitrobenzoic acid) (DTNB), glutathione reductase from baker's yeast (*S. cerevisiae*), 2-vinylpyridine, dihydrorhodamine, salmon sperm DNA, Hoescht 33342, serum albumin, phenazinemethosulfate, phenylmethylsulfonyl fluoride (PMSF), β -mercaptoethanol, NADH, β -NADPH, and EDTA were purchased from Sigma-Aldrich. All chemicals were of the analytical grade or better.

MTs were determined in parts of thyroid gland tissue after ethanol/chloroform extraction by thiol measurement with DTNB according to the method of Viarengo et al. (1997) [9]. The level of MT-related thiols (MT-SH) was calculated by using following relationship: 1 mol MT-SH = 20 mol GSH. The level of MTs was defined as μ g of MTs per gram of fresh weight (FW) tissues. To assess metal concentration in the MTs (MT-Me), they were isolated as

thermostable proteins by size-exclusion chromatography on Sephadex G-50 with necessary adjustments needed to avoid their oxidation [10], as described previously [11]. A 5 % homogenate (w/v) was prepared in ice-cold 10 mM Tris-HCl buffer, pH 8.0, containing 10 mM 2-mercaptoethanol for maintaining of the reduced conditions and 0.1 mM PMSF for the inhibition of proteolysis. Fractions of the chromatographic peak with high absorbance at 254 nm and comparative high density ratio D_{254}/D_{280} , identified as MT-containing peak [12], were pooled (total 10 mL) and applied to metal determination.

To determine copper and zinc concentration, fresh tissues (500 mg) and pooled eluate of MTs fraction after the size-exclusion chromatography (10 mL) were digested in 5 mL HNO₃ (Merck) for 3 h at 105 °C for metal analysis using hermetic acid-cleaned Teflon bomb. Concentration of metals was analyzed by the atomic absorption spectrometer with flame detector (C-115, "LOMO", Russian Federation). The metal detection limits were 0.1 μ g·g⁻¹ FW. The analytical methods were validated by external intercalibrations. Quality control was assessed by the Quality Control Sample of trace metal and method of Standard Addition (www.dentalmercury.com/245_1.pdf). Metal concentration in the tissues and MTs was presented as μ g·g⁻¹ FW and nmol·g⁻¹ FW. Subcellular distribution of iodine in the thyroid gland was determined by previously developed method [13].

Superoxide dismutase (SOD, EC 1.15.1.1) activity was measured by the method of Beauchamp and Fridovich (1971) [14] based on the aerobic reduction of NBT at 535 nm by superoxide radicals and expressed as units·mg⁻¹ soluble protein; 1 unit of SOD activity is defined as the amount of protein causing 50 % inhibition of the rate of NBT reduction.

Catalase (EC 1.11.1.6) activity was measured by monitoring decomposition of 10 mM H₂O₂ according to Aebi (1974) [15] at 240 nm ($\epsilon=40$ M⁻¹·cm⁻¹) in buffer containing 50 mM KH₂PO₄ (pH 7.0) and approximately 150 μ g of proteins. The results were related to soluble protein.

Total glutathione (GSht) concentration was quantified by the glutathione reductase recycling assay [16]. To estimate the oxidized glutathione (GSSG) level, the protein free sample was treated with 2-vinylpyridine for 60 min prior to assay at 2 % final concentration [17]. The rate of 5-thionitrobenzoic acid formation was monitored spectrophotometrically at 412 nm. Standards were prepared from GSH, and concentrations were defined as nmol per g wet weight. The redox-index of glutathione (RI GSH) was calculated as the ratio of concentrations [GSHr]/[GSH].

Evaluation of oxyradical formation in thyroid gland tissue (1/10 w/v) homogenates was determined using the non-fluorescent derivative, dihydror-

hodamine, which is converted to the fluorescent dye, rhodamine-123, while reacting with reactive oxygen species [18]. The fluorescence signal was detected by using af-max fluorescence plate-reader [excitation = 485 nm, emission = 538 nm] immediately, and in 20 min.

Glutathione-S-transferase (GST, EC 2.5.1.18) activity was measured by CDNB as the substrate [19]. Enzymatic activity was determined at 25 °C by monitoring changes in absorbance at 340 nm for 2 min. The GST activity was defined as nmol min⁻¹·mg⁻¹ soluble protein.

The activity of lactate dehydrogenase (LDH, EC 1.1.1.27) was determined using the UV assay with pyruvate and NADH [20] by determining of the amount of NADH oxidation at 340 nm. Phosphate/pyruvate solution (3 mL) (50 mM phosphate, pH 7.5, 0.63 mM pyruvate) was pipetted into cuvettes and 50 mL NADH solution (11.3 mMβ-NADH) was added. Following this, 100 mL of sample was spiked and mixed. The extinction was checked after very minute interval for a period of 4 min. A molar extinction coefficient of 6.22·10⁶ M⁻¹·cm⁻¹ was used.

DNA fragmentation was evaluated by the determination of the levels of protein-free DNA strand breaks in the digestive gland by the alkaline DNA precipitation assay [21] using Hoescht 33342. The reduction of possible interference of traces of SDS in the supernatant was measured in the presence of 0.4 M NaCl, 4 mM sodium cholate, and 0.1 M Tris (pH 9). Probe fluorescence signal was detected by using af-max fluorescence plate-reader [excitation = 360 nm, emission = 450 nm]. Quantitation was done using known amounts of salmon sperm DNA in the same mixture without addition of sample and determined as ng of protein-free DNA per mg tissue soluble proteins.

Cathepsin D (EC 3.4.23.5) activity was assayed spectrometrically as described previously [22] in incubation mixtures containing buffered haemoglobin solution [4% (w/v) in 0.25M-sodium acetate buffer, pH5.0].

All measurements were carried out using samples from 25 patients. The results were defined as means ± standard deviation (SD). Data were tested for normality and homogeneity of variance using Kolmogorov-Smirnoff and Levene's tests. Since data were not normally distributed (Lilliefors' test), non-parametric tests (Kruskall–WallisANOVA and Mann–Whitney U-test) were performed (significant at *P*<0.05). For detection of correlation, the Pearson's correlation test was also performed at a 0.05 level of significance.

Data were subjected to Multiple regression analysis and Principal component analysis (PCA) with NIPALS algorithm to differentiate the group by the set of their indices and select distinguished criterions. Classification trees were built using Classification and Regression Tree (CART) software on the basis of all determined biological characteristics. All statistical calculations were performed by means of Statistica v 8.0 and Excel for Windows 2000.

Results

The results showed the coherent activation of superoxide dismutase (by 81 %), catalase (up to two times) and glutathione-transferase (by 212 %), decrease of GSH level (by 33 %) and the increase of metallothioneins level (both MT-SH and MT-Me) in affected part of thyroid gland (Fig. 1, 2). Higher level of oxyradicals (by 21 %) and GSSG (up to 41%) has been also detected in these samples. A relation between MT-SH and oxyradical level in thyroid gland was proved. Signs of cytotoxicity, higher free cathepsin D activity (up to 84.6 % and 134.4 % in paranodular tissue and node respectively), and higher level of DNA strand breaks in node (up to 22.6 %), were observed (Fig. 3). Also, activation of glycolysis in the affected part of thyroid gland was observed. The accumulation of reactive oxygen species (*r* = 0.72, *p* < 0.01) and initiation of oxidative stress in the cell could be the reason for shifting of




Fig. 1. Subcellular distribution of iodine (A) and copper (B) in the human thyroid gland under iodine deficiency euthyroid nodular goiter, µg·g⁻¹ FW, mean ± SD (N=25). In Fig. 1 – 4, the significant differences were regarded as (*p* < 0.05); ^a, differences compare to contralateral part; ^b, differences between paranodular tissue and nodule.




Fig. 2. Oxidative stress parameters in the human thyroid gland under iodine deficiency euthyroid nodular goiter. Data for A: superoxide dismutase, B: catalase, C: glutathione transferase, D: oxyradical level are present as means \pm SD (N=25).

energy balance to anaerobiosis [23]. Range of indices variability in paranodular tissue compared to parenchyma of contralateral part was lesser than in the nodular tissue, but had generally the same trend.

The most important finding was lower level of organificated iodine (by 23 and 15 %, respectively) and higher level (by 46 and 32 %, respectively) and mass fraction (by 42% and 82% respectively) of inorganic iodine in the nodular and paranodular tissue versus contralateral part of thyroid gland (Fig. 4). The disruption of iodine organification in hyperplastic thyrocytes could be caused by elevated copper level (more than twice) in thyroid gland. Copper excess in the affected tissue partly accumulated in MTs. The binding ability of MTs with copper in the nodule was lesser than in the paranodular tissue.

According to PCA test with NIPALS algorithm (Fig. 5, A), two sets were the most distinguished: nodule was characterized by oxidative stress indices, MTs characteristics, and copper distribution, whereas non-affected contralateral part was located jointly with organificated iodine and GSH level. There was no clear characteristic related to the paranodular tissue. Furthermore, it was important to select the main distinguished criterion for pathological process progress at the nodule formation. We used CART analysis (Fig. 5, B) to achieve this. Followed by analysis of all biochemical parameters of every group, the contralateral part of thyroid gland was singled out by its ability to iodine organification. Paranodular tissue and nodule in affected part of

thyroid gland were distinguished by level of DNA fragmentation.

Discussion

It is generally known that nodules develop in pathologically affected thyroid tissue [2]. We established that hyperplastic thyroid epithelium under iodine deficiency nodular colloidal goiter sacrificed its ability for iodine organification and therefore increased of level of inorganic iodine. This pattern designated with higher copper level ($r = 0.69$, $p < 0.01$), manifestation of oxidative damage (increased of SOD, catalase and GST activity ($r = 0.73$, $r = 0.59$ and $r = 0.64$ correspondingly, $p < 0.01$ and MT-SH level ($r = 0.97$, $p < 0.001$)) and cytotoxicity (increased DNA fragmentation, $r = 0.51$, $p < 0.01$) in affected by nodule part of thyroid gland. Therefore, it can be assumed that under deprivation of iodine organification surplus of it has been created in human thyroid gland and determined further formation of iodine toxic intermediates in hyperplastic thyrocytes after its oxidation by thyroperoxidase [24]. This scenario of a stimulating effect on the antioxidant defense system was proved. Similar results, such as increased lipofuscin level, lipid peroxidation, necrosis of epithelial cells, and destabilization of mitochondrial membranes and development of autoimmune processes in the tissue of the thyroid gland were obtained after iodine application in micromolar range into human body [25].

It was proved that apoptosis was among the major determinants of pathological conditions progression [24]. Cathepsin D is a lysosomal




Fig. 3. Characteristics of metallothioneins (A-C) and glutathione (D-F) in the human thyroid gland under iodine deficiency euthyroid nodular goiter. Concentration of MT-SH (A); concentration of MT-Me (B); MT-SH/MT-Me concentration ratio (C); Levels of reduced glutathione (D), oxidized glutathione (E) and redox index of glutathione (RI GSH) (F). The values are expressed as the mean \pm SD (N=25).

endopeptidase. It involves in the processing of thyroglobulin, and belongs to mediators of IFN- γ and TNF- α -induced lysosomal programmed cell death pathway [26]. We were observing its increased level in patients with endemic nodular goiter. It was related to the increase of copper level in the thyroid tissue ($r=0.63$, $p<0.01$), and particularly, in potentially toxic, unbound with MTs form ($r=0.67$, $p<0.01$). We can speculate that copper accumulated in lysosomes and consistently caused their swelling, activation of calcium-dependent phospholipase A2 [27], and determined destabilization of lysosomal membranes. The *in vivo* release of the enzymes from the organelles, which happened consequently, has been suggested to play a fundamental role in mediating caspase activation, DNA fragmentation and apoptosis [28].

Recently, it has been shown *in vitro* that MTs and GSH form a part of the pool of cellular thiols and their function in the cell is cohered [29]. We can conclude the same regulation in human thyroid tissue ($r=-0.79$, $p<0.001$) based on our observations. MTs

might act as an effective scavenger against reactive oxygen species due to high level of SH-group under weakness of glutathione functional ability, which could indicate a compensatory mechanism. Despite the fact that the MTs level in the cell is lower than GSH, they have exhibited 50x higher antioxidant activity by the molar ratio of thiols[30]. We detected a reverse relation for the MT-SH level and oxyradicals level, which indicated a leading role of MTs in detoxification of oxyradicals: $\text{oxyradicals} = -268.4 + 88.2 \times \text{GSSG} - 4.7 \times \text{GSH} + 405.5 \times \text{RI GSH} + 6.4 \times \text{GST} - 2.1 \times \text{MT-SH}^*$; $R^2=0.45$, $F(5,12)=4.7$, $p<0.03$ (* – indicated significant contribution into the mathematical model). Similar results were obtained by comparing of goitrous-changed and intact human thyroid gland tissue [4].

Conclusions

To summarize, the combination of endemic iodine deficiency with a high environmental copper level increases the risk of node formation and progress of pathological changes. At low level of




Fig. 4. Markers of cytotoxicity in the human thyroid gland under iodine deficiency euthyroid nodular goiter. Data for A: DNA strand breaks, B: cathepsin D; C: lactate dehydrogenase are present as means ± SD (N=25).




Fig. 5. Principal component analysis with NIPALS algorithm (A) and classification tree models using CART algorithm (B) of total data from human thyroid gland under iodine deficiency euthyroid nodular goiter depending on deepness of pathology processes. Data for B showing node types, split variables, and associated split values of all attributes of tissues.

iodine organification and high copper level in goitrous tissue of thyroid gland, metallothioneins may provide a partial compensatory effect on prooxidative processes. The effects on the secretion of the cathepsin D can depend on the accumulation of copper inside thyrocyte lysosomes. Nodule formation in human thyroid gland has been followed by disruption of iodine organification by thyrocytes

(general characteristic of affected by nodule tissue) and increased level of DNA fragmentation (specific character of the nodule).

References

1. Giray B, Arnaud J, Sayek I, Favier A, Hincal F. Trace elements status in multinodular goiter. *J Trace Elem Med Biol* 2010; 24: 106–110.
2. Paschke R. Molecular pathogenesis of nodular goiter. *Langenbecks Arch Surg* 2011; 396: 1127–1136.
3. Uriu-Adams JY, Keen CL. Copper, oxidative stress, and human health. *Mol Aspects Med* 2005; 26: 268–298.
4. Фальфушинська ГІ, Гнатишина ЛЛ, Осадчук ДВ, Шідловський ВО, Столяр ОБ. Металодепонуюча функція та антиоксидантні властивості цитоподібної залози людей, хворих на йододефіцитний вузловий колоїдний зоб. *Укр біохім журн* 2011; 83: 92–97.
5. Stolyar OB, Loumbourdis NS, Falfushynska HI, Romanchuk LD. Comparison of metal bioavailability in frogs from urban and rural sites of Western Ukraine. *Arch Environ Contam Toxicol* 2008; 54: 107–113.
6. Jomova K, Valko M. Advances in metal-induced oxidative stress and human disease. *Toxicology* 2011; 283: 65–87.
7. Maret W. Redox biochemistry of mammalian metallothionein. *J Biol Inorg Chem* 2011; 16: 1079–1086.
8. Ferrario C, Lavagni P, Gariboldi M, Miranda C, Losa M, Cleris L. et al. Metallothionein 1G acts as an oncosuppressor in papillary thyroid carcinoma. *Lab Invest* 2008; 88: 474–481.
9. Viarengo A, Ponzano E, Dondero F, Fabbri R. A simple spectrophotometric method for metallothionein evaluation in marine organisms: an application to Mediterranean and Antarctic molluscs. *Mar Environ Res* 1997; 44: 69–84.
10. Suzuki KT. Purification of vertebrate metallothioneins. *Method Enzymol* 1991; 205: 252–263.
11. Falfushynska HI, Gnatyshyna LL, Stolyar OB. Population-related molecular responses on the effect of pesticides in *Carassius auratus gibelio*. *Comp Biochem Physiol* 2012; 155 C: 396–406.
12. Kagi JHR, Schaffer A. Biochemistry of metallothionein. *Biochemistry* 1988; 27: 8509–8515.
13. Шідловський ВО, Столяр ОБ, Осадчук ДВ, Шідловський ОБ, Фальфушинська ГІ. Деклараційний патент на корисну модель № 45332 (UA), МПК G09B 23/28 (2009.01). Спосіб визначення концентрації йоду в біосубстраті / (Україна). Заявл. 24.04.09; опубл. 10.11.09, Бюл. № 21.
14. Beauchamp C, Fridovich I. Superoxide dismutase: improved assay and an assay applicable to acrylamide gels. *Anal Biochem* 1971; 44: 276–287.
15. Aebi H. Catalase (in) Bergmeyer HU. (ed.) *Methods of Enzymatic Analysis*. Academic Press, London; 1974: 671–684.
16. Anderson ME. Determination of glutathione and glutathione disulfide in biological samples. *Meth Enzymol* 1985; 113: 548–555.
17. Griffith OW. Determination of glutathione and glutathione disulfide using glutathione reductase and 2-vinylpyridine. *Anal Biochem* 1980; 106: 207–212.
18. Viarengo A, Burlando B, Cavaletto M, Marchi B, Ponzano E, Blasco J. Role of metallothionein against oxidative stress in the mussel *Mytilus galloprovincialis*. *Am J Physiol* 1999; 277: 1612–1619.
19. Habig WH, Pabst MJ, Jakoby WB. Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *J Biol Chem* 1974; 249: 7130–7139.
20. Bergmeyer HU, Bernt E. Lactate-dehydrogenase, UV-assay with pyruvate and NADH(in) Bergmeyer HU. (ed.) *Methods of enzymatic analysis*. Vol 2. Academic Press, New York; 1974: 579.
21. Olive PL. DNA precipitation assay: a rapid and simple method for detecting DNA damage in mammalian cells. *Environ Mol Mutagen* 1988; 11: 487–495.
22. Dingle JT, Barrett AJ, Weston PD. Cathepsin D. Characteristics of immunoinhibition and the confirmation of a role in cartilage break down. *Biochem J* 1971; 123: 1–13.
23. Morita M, Noguchi S, Kawamoto H, Tajiri J, Tamaru M, Murakami N. Thyroglobulin and lactic dehydrogenase isozymes in cystic fluid of thyroid nodules. *Endocr J* 1994; 41: 227–233.
24. Vitale M, Di Matola T, D'Ascoli F, Salzano S, Bogazzi F, Fenzi G. et al. Iodide excess induces apoptosis in thyroid cells through a p53-independent mechanism involving oxidative stress. *Endocrinology* 2000; 141: 598–605.
25. Foley TP Jr. The relationship between autoimmune thyroid disease and iodine intake: a review. *Endocrinol Pol* 1992; 43, Suppl 1: 53–69.
26. Tsukuba T, Okamoto K, Yasuda Y, Morikawa W, Nakanishi H, Yamamoto K. New functional aspects of cathepsin D and cathepsin E. *Mol Cells* 2000; 10: 601–611.
27. Marchi B, Burlando B, Moore MN, Viarengo A. Mercury- and copper-induced lysosomal membrane destabilisation depends on $[Ca^{2+}]_i$ dependent phospholipase A2 activation. *Aquat Toxicol* 2004; 66: 197–204.
28. Persson HL. Iron-dependent lysosomal destabilization initiates silica-induced apoptosis in murine macrophages. *Toxicol Lett* 2005; 159: 124–133.
29. Hidalgo J, Garvey JS, Armario AJ. On the metallothionein, glutathione and cysteine relationship in rat liver. *Pharmacol Exp Ther* 1990; 255: 554–564.
30. Thornalley PJ, Vasak M. Possible role for metallothionein in protection against radiation-induced oxidative stress. Kinetics and mechanism of its reaction with superoxide and hydroxyl radicals. *Biochim Biophys Acta* 1985; 827: 36–44.

Received: 2014.05.11