

Available online on 25.08.2019 at <http://jddtonline.info>

# Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Research Article

## Ambient Air Quality of Nashik City 2017 (Maharashtra, India)

D.S. Khandbahahale\*, V. M. Motghare, S. C. Kollur, S. Save, V. S. Badgujar

P.G. Department of Botany, K. T. H. M. College, Shivaji Nagar, Gangapur Road, Nashik, India

### ABSTRACT

Ambient air quality of Nashik city was monitored during the year 2017. Criteria pollutants selected for the monitoring was, sulphur dioxide (SO<sub>2</sub>) & nitrogen dioxide (NO<sub>2</sub>) and PM<sub>10</sub> (Particulate Matter having aerodynamic diameter less than or equal to 10 μm) for the period of January 2017 to December 2017. Sampling was done for successive periods of about 4 hours for sulphur dioxides (SO<sub>2</sub>), nitrogen dioxide (NO<sub>2</sub>) and 8 hours for Respirable suspended particulate matter (PM<sub>10</sub>) for 24 hours. For Air Quality Monitoring, four representative sites were selected i.e. Industrial-S1-I-VIP, Commercial- S2-R-RTO and Residential-S3-C-NMC and Industrial-S4-I-UB. High volume air sampler were used to measure the concentration of nitrogen dioxide (NO<sub>2</sub>), sulphur dioxides (SO<sub>2</sub>) and Respirable suspended particulate matter (PM<sub>10</sub>). The results reported pertain to an eight hour successive continuous air sampling exercise carried out at each of the four selected locations in Nashik city. The value of PM<sub>10</sub> (Particles ≥ 10μ, PM 10μg/m<sup>3</sup>) was noted to be crossing the permissible limit and exceeded the National Ambient Air Quality Standard (NAAQS) at all locations. The concentration of SO<sub>2</sub> and NO<sub>2</sub> was within the National Ambient Air Quality Standard (NAAQS, National Ambient Air Quality Standards, November 18, 2009) at all the locations.

**Keywords:** PM<sub>10</sub>, Ambient Air Quality (AQI), National Ambient Air Quality Standard (NAAQS), Pollutant, Urban air sampling, Transport emissions.

**Article Info:** Received 21 June 2019; Review Completed 13 Aug 2019; Accepted 17 Aug 2019; Available online 25 August 2019



### Cite this article as:

Khandbahahale DS, Motghare VM, Kollur SC, Save S, Badgujar VS, Ambient Air Quality of Nashik City 2017 (Maharashtra, India), Journal of Drug Delivery and Therapeutics. 2019; 9(4-s):1111-1117 <http://dx.doi.org/10.22270/jddt.v9i4-s.3805>

### \*Address for Correspondence:

D.S. Khandbahahale, P.G. Department of Botany, K. T. H. M. College, Shivaji Nagar, Gangapur Road, Nashik, India

### INTRODUCTION:

“Clean air is considered to be a basic requirement of human health and well-being. However, air pollution continues to pose a significant threat to health worldwide” (WHO, 2005). We could go days without food and hours without water, but we would last only a few minutes without air. Ambient air is the outdoor air in which mankind and other organisms survive and respire (Daly, 2007). Ambient air quality refers to the quality of outdoor air in our surrounding environment. The term “air quality” means the state of the air round us. Good air quality refers to clean, clear, unpolluted air. Air pollution occurs when the air contains substances in quantities that could harm the comfort or health of humans and animals, or could damage plants and materials. These substances are called air pollutants and can be either particles, liquids or gases in nature (Alias, *et al.*, 2007). The atmosphere, which makes the largest fraction of the biosphere, is a dynamic system that continually absorbs a wide range of solids, liquids and gases from both natural and man-made (Anthropological) activities (Mahindra S.P and Krishnamurthy D.,2005). Various types and amounts of material enters, interact

within and with atmosphere. Polluted air, polluted space, polluted land and polluted water are the resulting undesired by-products. Awareness of air contamination and measures to monitor and control air quality are inadequate considering the rapidity of the increase in pollution levels. Many epidemic effects of air pollutants are prevalent (Brunekreef, Dockery, and Krzyzan. *et al.*, 1995, CPCB, 2012). The assessment of the Air Quality is presently linked to the Air pollution levels and to the size of populations. To protect the health, the concentrations of selected harmful air pollutants should be limited and related to given ambient air quality standards (Ahmed Haytham, 1999).

The major objectives of monitoring air pollutant levels is to provide an early warning system for pollutant levels, which may have the potential for endangering public health; to assess air quality in light of established public health and welfare standards; to track air pollution trends and changes in ambient air quality due to changes in the amount of pollutants emitted; to produce information for city planning, e.g. Location of new industries and housing and to access the environmental impact of industry and other activities.

## MATERIALS AND METHODS:

Ambient air quality was monitored during the year 2017 (January 2017 to December 2017), concentrations of sulphur dioxides (SO<sub>2</sub>), nitrogen dioxide (NO<sub>2</sub>) and PM<sub>10</sub> (Particulate Matter having aerodynamic diameter less than or equal to 10 µm) were collected over successive periods of about 4 hours and 8 hours in a day respectively. Four representative sites were selected for the study, Industrial-S1-I-VIP, Commercial-S2-R-RTO and Residential-S3-C-NMC and Industrial-S4-I-UB. High volume air sampler was used to measure the concentration of nitrogen dioxide (NO<sub>2</sub>), sulphur dioxides (SO<sub>2</sub>) and Respirable suspended particulate matter (PM<sub>10</sub>).

Respirable Suspended Particulate Matter (PM<sub>10</sub>) was measured by Cyclonic Flow Technique. Nitrogen oxides

were measured by Modified Jacobs and Hochheiser Method (Sodium Arsenite method). Sulphur oxides were measured by Modified West and Gaeke Method

About the study area: Nasik, a thickly populated poorly ventilated air basin being polluted by the ever-expanding transport, construction, and commercial activities. Nasik is amongst the fastest growing cities, it rates fourth in India and sixteenth in the world, and hence, it is necessary to forecast the impact of pollutants from the various sources located in the city for the purpose of arriving at a comprehensive air pollution scenario. Nasik is one of the four places where Kumbha Mela is held once every 12 years. For the Hindus, Nasik is one of the holiest cities. Nasik is located in the "Western Ghats" on the western edge of the "Deccan" on the banks of river "Godavari".




Figure: Map of Nasik City with Ambient Air Quality Monitoring Locations

## RESULTS AND DISCUSSION:


Concentrations of selected criteria pollutants (i.e. PM<sub>10</sub>, SO<sub>2</sub> and NO<sub>2</sub>) on selected air quality monitoring sites (i.e. S1-I-VIP, S2-R-RTO, S3-C-NMC & S4-I-U.B.) were measured and calculated for respective period.

Concentration of those pollutants was compared with National Ambient Air Quality Standards (NAAQS) for the period of January 2017 to December 2017.

Table1: Concentration of Criteria pollutants i.e. SO<sub>2</sub>, NO<sub>x</sub> & PM<sub>10</sub> on S1-I-VIP site for the period of January 2017 to December 2017 (Concentration in µg/m<sup>3</sup>)

Month	S1-I-VIP*		NO <sub>2</sub> µg/m <sup>3</sup>		PM <sub>10</sub> µg/m <sup>3</sup>	
	Average	Maximum	Average	Maximum	Average	Maximum
January	15	42	25	37	107	200
February	17	43	26	38	94	160
March	18	35	26	40	116	196
April	19	42	26	38	126	227
May	13	40	24	38	74	146
June	10	23	19	34	59	112
July	8	14	13	34	44	106
August	8	16	13	18	40	67
September	9	14	15	21	35	95
October	8	15	19	53	53	208
November	7	19	19	32	67	125
December	4	20	17	32	76	138
Annual Average	11	42	20	40	74	227

\* S1- Site 1, I-Industrial, VIP- Name of Site location i.e. VIP company, MIDC Area, Satpur Nasik



Graph 1: Concentration of Criteria pollutants i.e. SO<sub>2</sub>, NO<sub>2</sub> & PM<sub>10</sub> on S1-I-VIP site for the period of January 2017 to December 2017(Concentration in µg/m<sup>3</sup>)


Graph 1 displays the monthly concentration of criteria pollutant PM<sub>10</sub> on S1-I-VIP air quality monitoring site against NAAQS in the year 2017. PM<sub>10</sub> concentration remains more than the average NAAQS from January 2017 to May 2017, November and December 2017. In April 2017

PM<sub>10</sub> concentration was 126 µg/m<sup>3</sup>, this is 66µg/m<sup>3</sup> more than the annual NAAQS average. PM<sub>10</sub> Concentration remains lower than the NAAQS from June to October 2017. was observed from November onwards.

Table2: Concentration of Criteria pollutants i.e. SO<sub>2</sub>, NO<sub>2</sub> & PM<sub>10</sub> on S2-R-RTO site for the period of January 2017 to December 2017(Concentration in µg/m<sup>3</sup>)

Month	S2-R-RTO*					
	SO <sub>2</sub>		NO <sub>2</sub>		PM <sub>10</sub>	
	Average	Max.	Average	Max.	Average	Max.
January	17	44	28	42	102	189
February	19	48	29	43	95	132
March	21	42	30	46	112	185
April	22	46	28	42	94	178
May	13	38	26	38	80	147
June	11	22	22	36	65	143
July	9	15	15	21	59	131
August	9	16	15	27	42	60
September	9	15	16	22	44	87
October	10	19	20	59	72	205
November	8	20	22	35	95	169
December	4	14	19	46	93	202
Annual Average	13	48	22	59	79	205

\* S2- Site 2, R-Residential, RTO- Name of Site location i.e. NMC Water Tank, Near Golf Club, Nasik



Graph 2: Concentration of Criteria pollutants i.e. SO<sub>2</sub>, NO<sub>2</sub> & PM<sub>10</sub> on S2-R-RTO site for the period of January 2017 to December 2017(Concentration in µg/m<sup>3</sup>)

Table 2 shows the concentrations of criteria pollutants during the monitoring period on S2- R-RTO site for the period of January 2017 to December 2017. Annual average concentration measured during the sampling period was,  $13\mu\text{g}/\text{m}^3$  and  $22\mu\text{g}/\text{m}^3$  for  $\text{SO}_2$  and  $\text{NO}_2$  respectively, which were within the permissible limit of NAAQS. Concentration of  $\text{PM}_{10}$  was  $79\mu\text{g}/\text{m}^3$ , which is more than National Ambient Air Quality Standards. Maximum concentration for the 8 hours sampling was  $205\mu\text{g}/\text{m}^3$  in the month of April


2017.

Graph 2 displays the monthly concentration of criteria pollutant  $\text{PM}_{10}$  on S2-R-RTO air quality monitoring site against NAAQS in the year 2017.  $\text{PM}_{10}$  concentration remains more than the average NAAQS from January 2017 to June 2017, November and December 2017. In March 2017  $\text{PM}_{10}$  concentration was  $112\mu\text{g}/\text{m}^3$ , this is  $52\mu\text{g}/\text{m}^3$  more than the annual NAAQS average.  $\text{PM}_{10}$  Concentration remains lower than the NAAQS from June to October 2017.

Table 3: Concentration of Criteria pollutants i.e.  $\text{SO}_2$ ,  $\text{NO}_x$  &  $\text{PM}_{10}$  on S3-C-NMC site for the period of January 2017 to December 2017(Concentration in  $\mu\text{g}/\text{m}^3$ )

S3-C-NMC*						
Month	$\text{SO}_2$		$\text{NO}_2$		$\text{PM}_{10}$	
	Average	Max.	Average	Max.	Average	Max.
January	17	49	28	41	135	270
February	20	50	27	40	113	204
March	19	44	29	41	120	273
April	21	43	28	41	99	183
May	14	44	25	41	77	204
June	10	21	22	36	68	133
July	9	15	14	23	57	116
August	9	17	14	21	43	64
September	10	14	16	21	49	110
October	9	22	19	47	70	187
November	6	19	21	32	112	161
December	4	11	20	29	111	192
Annual Average	12	50	22	47	83	273

\* S3- Site 2, C-Commercial, NMC- Name of Site location i.e. Nashik Municipal Corporation Build. Nasik



Graph 3: Concentration of Criteria pollutants i.e.  $\text{SO}_2$ ,  $\text{NO}_2$  &  $\text{PM}_{10}$  on S3-C-NMC site for the period of January 2017 to December 2017(Concentration in  $\mu\text{g}/\text{m}^3$ )

Table 3 shows the concentrations of criteria pollutants during the monitoring period on S3- C-NMC site for the period of January 2017 to December 2017. Annual average concentration measured during the sampling period was,  $12\mu\text{g}/\text{m}^3$  and  $22\mu\text{g}/\text{m}^3$  for  $\text{SO}_2$  and  $\text{NO}_x$  respectively, which were within the permissible limit of NAAQS. Concentration of  $\text{PM}_{10}$  was  $83\mu\text{g}/\text{m}^3$ , which is more than National Ambient Air Quality Standards. Maximum concentration for the 8 hours sampling was  $273\mu\text{g}/\text{m}^3$  in the month of April

2017.

Graph 3 displays the monthly concentration of criteria pollutant  $\text{PM}_{10}$  on S3-C-NMC air quality monitoring site against NAAQS in the year 2017.  $\text{PM}_{10}$  concentration remains more than the average NAAQS from January 2017 to June 2017, November and December 2017. In January 2017  $\text{PM}_{10}$  concentration was  $135\mu\text{g}/\text{m}^3$ , this is  $75\mu\text{g}/\text{m}^3$  more than the annual NAAQS average.  $\text{PM}_{10}$  Concentration remains lower than the NAAQS from July to October 2017.

Table 4: Concentration of Criteria pollutants i.e. SO<sub>2</sub>, NO<sub>2</sub> & PM<sub>10</sub> on S4-I-UB site for the period of January 2017 to December 2017(Concentration in µg/m<sup>3</sup>)


S4-I-UB*						
Month	SO <sub>2</sub>		NO <sub>x</sub>		PM <sub>10</sub>	
	Average	Max.	Average	Max.	Average	Max.
January	16	47	28	28	106	327
February	19	49	28	45	79	229
March	20	42	28	44	96	247
April	22	47	31	42	96	210
May	15	44	26	41	74	208
June	11	24	23	37	55	171
July	10	22	15	44	42	120
August	9	18	14	21	38	98
September	9	18	16	22	43	105
October	10	28	22	82	72	229
November	7	19	23	39	91	301
December	5	20	21	102	73	202
Annual Average	13	31	23	46	72	204

\* S4- Site 4, I-Industrial, UB- Name of Site location, i.e. Udyog Bhavan, Trimbak Road, Satpur, Nasik

Table 4 shows the concentrations of criteria pollutants during the monitoring period on S4- I-UB site for the period of January 2017 to December 2017. Annual average concentration measured during the sampling period was, 13µg/m<sup>3</sup> and 23µg/m<sup>3</sup> for SO<sub>2</sub> and NO<sub>x</sub> respectively, which were within the permissible limit of NAAQS. Concentration

of PM PM<sub>10</sub> was 72µg/m<sup>3</sup>, which is more than National Ambient Air Quality Standards.

Maximum concentration for the 8 hours sampling was 204µg/m<sup>3</sup> in the month of April 2017.



Graph 4: Concentration of Criteria pollutants i.e. SO<sub>2</sub>, NO<sub>2</sub> & PM<sub>10</sub> on S3-C-NMC site for the period of January 2017 to December 2017(Concentration in µg/m<sup>3</sup>)

Graph 4 displays the monthly concentration of criteria pollutant PM<sub>10</sub> on S4-I-UB air quality monitoring site against NAAQS in the year 2017. PM<sub>10</sub> concentration remains more than the average NAAQS from January 2017 to

May 2017, November and December 2017. In January 2017 PM<sub>10</sub> concentration was 106µg/m<sup>3</sup>, this is 46µg/m<sup>3</sup> more than the annual NAAQS average. PM<sub>10</sub> Concentration remains lower than the NAAQS from June to October 2017.




Table 5: Concentration of Criteria pollutants i.e. SO<sub>2</sub>, NO<sub>2</sub> & PM<sub>10</sub> on all sites for the period of January 2017 to December 2017(Concentration in µg/m<sup>3</sup>)

City	Station Name	Class	SO <sub>2</sub> (µg/m <sup>3</sup> )		NO <sub>2</sub> (µg/m <sup>3</sup> )		RSPM (µg/m <sup>3</sup> )	
			Max	Avg.	Max	Avg.	Max	Avg.
Nashik	S1-I-VIP	I	27	11	34	20	148	74
	S2-R-RTO	R	28	13	38	23	152	79
	S9-C-NMC	C	29	12	34	20	175	87
	S4-I-U.B.	I	32	13	46	23	204	72
Annual Average			32	12.25	46	21.5	204	78

Table 5 shows the concentrations of criteria pollutants during the monitoring period on all selected site for the period of January 2017 to December 2017. Annual average concentration measured during the sampling period was, 12.25µg/m<sup>3</sup> and 21.5µg/m<sup>3</sup> for SO<sub>2</sub> and NO<sub>x</sub> respectively,

which were within the permissible limit of NAAQS. Concentration of PM<sub>10</sub> was 78µg/m<sup>3</sup>, which is more than National Ambient Air Quality Standards. Maximum concentration for the 8 hours sampling was 204µg/m<sup>3</sup> in the month of April 2017.



Graph 5: Concentration of Criteria pollutants i.e. SO<sub>2</sub>, NO<sub>2</sub> & PM<sub>10</sub> on all sites for the period of January 2017 to December 2017(Concentration in µg/m<sup>3</sup>)

Graph 5 displays the monthly concentration of criteria pollutant PM<sub>10</sub> on All air quality monitoring site against NAAQS in the year 2017. PM<sub>10</sub> concentration remains more than the average NAAQS. Maximum concentration noted at S3-C-NMC which is representing the commercial area.

**CONCLUSION**

This work represents an in-depth investigation of the status and condition of ambient air quality of Nashik city during the period of January 2017 to December 2017. The concentration of SO<sub>2</sub> and NO<sub>2</sub> observed to be within the limits of National Ambient Air Quality Standards (NAAQS) on all selected air quality monitoring sites. But, the concentration of PM<sub>10</sub> observed to be much higher than National Ambient Air Quality Standards (NAAQS) on all selected air quality monitoring sites. It is observed that PM<sub>10</sub>

concentration crosses the predicible limits during summer and winter seasons, while in rainy season, dust steeled down on the ground which lower the PM<sub>10</sub> concentration. Being Nasik is one of the fastest growing cities in Maharashtra (India), Transport related emissions are the major sources of air contamination, increasing civil construction activities also contribute to particulates. The exponential rise in the number and volume of vehicles, disadvantageous traffic flow pattern, differing driving cycle pattern, old and outdated transport vehicles and human interceptions deserve due attention.

**ACKNOWLEDGEMENT**

We are thankful to Maharashtra Pollution Control Board, Mumbai and sub regional office Nashik and Savitribai Phule Pune University, formerly University of Pune.

## REFERENCES

1. Alexopolulos, Assimacopoulos, D. and Mistsoulis, E. (1993), Model for Traffic Emissions estimation, Journal of Atmospheric Environment, 27B (4):435-446.
2. Arun Srivastava, Jain V.K., (2005), A study to characterize the suspended particulate matter in an indoor environment in Delhi, India Building and Environment, pp2046- 2052.
3. Athanasios Matzoros, (1990), Results from a model of Road traffic air pollution, featuring junction effects and vehicle operating models, Traffic Engineering and control 24:37.
4. Brunekreef, B., Dockery, D.W.and Krzyzan.M.et al. (1995), Epidemiologic studies on short-term effect of low levels of major ambient air pollution components. Environment Health Perspective.103, suppl: 2:3-13.
5. CPCB (2012), Epidemiological study on effect of air pollution on human health (Adults) in Delhi. Environmental health management series (EHMS/01/2012)
6. H.K. Bandhu, Sanjiv Puri', J.S. Shahi, D. Mehta, M.L. Garg, Nirmal Singh,P.C. Mangal, C.R. Suri, E. Swietlicki, P.N. Trehan (1996), An evaluation of the sources of air pollution in the city of Chandigarh, India: A study using EDXRF technique.Nuclear Instruments and Methods in Physics Research B 114 (1996) 341-344.
7. Health and safety authority (2005),Carbon monoxide February 2008
8. Hickman, A.J. (1982), The estimation of air pollution concentrations from road traffic, TRRL Report 1052
9. Hiromasa Ureda and Gregiry R Caramichael 1995, Formation of secondary pollutants during long range transport and its contribution to air quality in east Asia,TAO, Vol.6, No.3, 487-500 Sept 1995
10. Mahindra S.P and Krishnamurthy D., (2005), Impact of introduction of one way traffic system on ambient air quality. Pollution Research 24 (1): 213-216.
11. Michal B, Sarah H, Tracy K, Kit S L, Kira R, Kay Teschke (2002), Review of the risks association with Nitrogen dioxide and sulphur dioxide in indoor air, report of Health Canada december1 2002.
12. National Summery Report, CPCB,(2010), Air quality, emission inventory and source appointment study for Indian cites National summery report, Dec 2010:197-198.
13. Ntao S.A, (2004), Study of Ambient air quality of margao town in south Goa, Pollution research 23(4):741-744(2004).
14. Pankajam K, Harikumaran Nair R, kesavachandran .C, Reethamma.K.V and shashidhar. (2005), Effects of automobile exhaust pollution on the pulmonary functions in shop keepers. Pollution Research 24 (1): 49-54.
15. Simkhada k, Murthy K, and Khanal S.N., (2005), Assessment of ambient air quality in Bishnumati corridor, Kathmandu metropolis International Journal of Environmental Science and technology, vol.2 pp217-222.

