

***In vitro* and *in vivo* studies of the thiol:disulphide
oxidoreductase ResA from *Bacillus subtilis* and
*Streptomyces coelicolor***

Christopher T C Hodson

A thesis submitted in part fulfilment of the degree of Doctor of
Philosophy at the University of East Anglia 2010

© This copy of the thesis has been supplied on condition that anyone who
consults it is understood to recognise that its copyright rests with its author and
no quotation from the thesis, nor information derived there from, may be
published without the author's prior consent.

DECLARATION

I declare that the work contained in this thesis, submitted by me for the degree of Ph.D., is my own work, except where due reference is made to other authors, and has not been previously submitted by me for a degree at any other university.

Christopher T. C. Hodson

April 2010

PUBLICATIONS ARISING FROM WORK IN THIS THESIS

Hodson, C. T. C., Lewin, A., Hederstedt, L., and Le Brun, N. E. (2008) “The active-site cysteinyls and hydrophobic cavity residues of ResA are important for cytochrome *c* maturation in *Bacillus subtilis*” *Journal of Bacteriology* **190** 4697-4705

Lewin, A., Crow, A., Hodson, C. T. C., Hederstedt, L., and Le Brun, N. E. (2008) “Effects of substitutions in the CXXC active-site motif of the extracytoplasmic thioredoxin ResA” *Biochemical Journal* **414** 81-91

Many of the ideas discussed in this thesis are also discussed in these papers. Reference to these papers in the main text has been deliberately restricted to a minimum, not out of any wish to diminish the importance of these works, but rather, to prevent the need to cite these works recurrently.

ACKNOWLEDGEMENTS

Throughout the course of my research I have received help, advice and support, both professionally and more informally, from many people, all of whom I wish to sincerely thank here.

In particular I would like to thank my supervisor, Nick Le Brun who has somehow managed to put up with me for the last four years and always found the time to support, advise and tutor; Allison Lewin who has offered advise and tuition long after she was contractually obliged to; Lars Hederstedt (and his laboratory) for much needed help with BdbD; Matt Hutchings and Dave Widdick for guiding me through *Streptomyces* genetics; Nick Cull for technical support; all my friends and colleagues in both CAP and BIO that I've not already mentioned (Jason Crack, John Holmes, Allister Crow, Alisa Gaskil, Liang Zhou, Tamara Lawson, Chloe Singleton, Oliver Hecht, Angelo Figueiredo, Rose Marie Doyle, Kate Haynes, Gaye White, Matt Bawn, Myles Cheesman, Julea Butt, Ben Thompson and sorry to anyone I have forgotten); My girlfriend Nicola; Dad, Mum, Jim, Tim and Granny, I can not thank you enough for the love and support you have all offered me over the years to get me this far; finally I would like to thank my late Grandad, Gerald Lucian Baldit, and dedicate this thesis to his memory.

This Ph.D. thesis was funded by the Engineering and Physical Sciences Research Council (EPSRC).

ABSTRACT

Thiol:disulphide oxidoreductases (TDORs) are essential in many organisms for the correct insertion and/or removal of disulphide bonds into and from peptides and proteins. One process for which TDORs have been shown to be integral is cytochrome *c* maturation (CCM). In the Gram positive soil bacterium *Bacillus subtilis* the membrane bound TDOR ResA is involved in the removal of a disulphide bond from the CXXCH haem binding motif of apo-cytochromes *c* in order to allow correct haem insertion by ResBC. The majority of TDORs contain a CXXC active site in which the sulphur residues of the cysteine side chain shuffle between the oxidised (disulphide) and reduced (thiol) forms. It is demonstrated here that both cysteines of the ResA CXXC active site are essential for protein function and that other residues, Pro141, Glu80 and Glu75, are important for stability, recognition and maintaining the reducing power of the active site, respectively. Studies of the membrane anchor domain of ResA reveal that it is important but not essential for CCM. Further to this, a homologue of *B. subtilis* ResA found in *Streptomyces coelicolor* was shown to play a similar role *in vivo* with regard to CCM; and *in vitro* studies of a purified soluble form of the protein revealed that although it has a similar low reduction potential to *B. subtilis* ResA it also has some interesting differences. Finally, *in vivo* studies of an oxidising TDOR, BdbD, from *B. subtilis* have provided some insight to the delicate balance of the redox state of proteins on the outside of the cytoplasmic membrane as well as future perspective on how to study this protein *in vivo*.

TABLE OF CONTENTS

Section	Title	Page
	TITLE PAGE	i
	DECLARATION	ii
	PUBLICATIONS ARISING FROM WORK IN THIS THESIS	iii
	ACKNOWLEDGEMENTS	iv
	ABSTRACT	v
	TABLE OF CONTENTS	vi
	LIST OF FIGURES	xii
	LIST OF TABLES	xvii
	ABBREVIATIONS	xix

Chapter 1: Introduction

1.1	Summary	1
1.2	Thiol:disulfide oxidoreductases (TDORs)	1
1.2.1	Thioredoxin	1
1.2.2	Structure of Thioredoxin	2
1.2.3	Thioredoxin in <i>B. subtilis</i>	3
1.3	TDORs of <i>B.subtilis</i>	4
1.3.1	ResA	5
1.3.1.1	Function of ResA	5
1.3.1.2	Structure of ResA	6
1.3.1.3	Biophysical properties of ResA	10
1.3.2	BdbD/BdbC	11
1.3.2.1	Function of BdbD	12
1.3.2.2	Structure of BdbD	13
1.3.2.3	Biophysical properties of BdbD	14
1.3.3	BdbA and BdbB	15
1.3.4	StoA	18
1.3.5	CcdA	20
1.4	Cytochromes <i>c</i>	22

1.4.1	Haem C	23
1.4.2	Cytochromes <i>c</i> .	24
1.4.3	Cytochromes <i>c</i> of <i>B. subtilis</i>	25
1.4.3.1	QcrC	26
1.4.3.2	CccA	27
1.4.3.3	CccB	29
1.4.3.4	CtaC	29
1.4.4	Cytochrome <i>c</i> maturation	31
1.4.4.1	CCM System I	32
1.4.4.2	CCM System II	38
1.4.4.3	CCM System III	42
1.4.4.4	Systems IV and V	45
1.4.4.5	CCM in <i>S. coelicolor</i>	47
1.5	Project Aims	52

Chapter 2: Materials and methods

2.1	Introduction	53
2.2	Bacterial Strains	53
2.3	Growth Media.	53
2.4	Molecular biology techniques	55
2.4.1	Polymerase Chain Reaction (PCR)	55
2.4.2	Site directed PCR mutagenesis	57
2.4.3	Ligations	59
2.4.4	Preparation of chemically competent <i>Escherichia coli</i> cells	61
2.4.5	Transformation of chemically competent <i>E. coli</i> cells	61
2.4.6	Preparation and natural transformation of competent <i>B. subtilis</i> cells	62
2.4.7	Transformation of <i>E. coli</i> cells by electroporation	62
2.4.8	Blue/White screening	63
2.4.9	Isolation of DNA from cell cultures	64
2.4.10	DNA quantification	65
2.4.11	Agarose gel electrophoresis	65
2.4.12	DNA sequencing	66

2.4.13	LUGOL staining for amylase activity	66
2.5	Sample preparation	69
2.5.1	<i>B. subtilis</i> membrane preparations	69
2.5.2	<i>S. coelicolor</i> membrane preparations	70
2.5.3	Determination of protein concentrations	70
2.5.4	Gel filtration chromatography	71
2.5.5	Anion exchange chromatography	72
2.6	Analytical techniques	72
2.6.1	SDS PAGE	72
2.6.1.1	SDS PAGE – Laemmli	73
2.6.1.2	SDS PAGE - Shäggar and von Jargow (SVJ)	73
2.6.2	Haem staining	74
2.6.3	Western Blots	74
2.6.4	Fluorescence spectroscopy	75
2.6.5	Cytochrome <i>c</i> oxidase assay	78
2.6.6	TMPD staining for cytochrome <i>c</i> oxidase activity	79

Chapter 3: *In vivo* studies of the active site residues of *B. subtilis*

ResA

3.1	Introduction	81
3.2	Materials and Methods	82
3.3	Results	83
3.3.1	Construction of vectors and strains	83
3.3.2	Analysis of protein production by Western blotting	85
3.3.3	Investigation of ResA activities by TMPD staining	86
3.3.4	Cytochrome <i>c</i> oxidase assays	87
3.3.5	Haem staining of the active site variants	90
3.3.6	The wild type CCM phenotype is restored with addition of dithiothreitol (DTT) to the growth media	91
3.4	Discussion.	93

Chapter 4: *In vivo* studies of the functional importance of the hydrophobic cavity and the membrane anchor of *B. subtilis* ResA

4.1	Introduction	95
4.2	Materials and Methods	97
4.3	Results	97
4.3.1	<i>In vivo</i> studies of the key residues found in the hydrophobic cavity	97
4.3.2	Confirmation of expression of variant ResA proteins by Western Blotting	98
4.3.3	TMPD staining of ResA variant strains P141S, P141T and E80Q	100
4.3.4	Cytochrome <i>c</i> oxidase assays of strains producing ResA hydrophobic cavity variants	101
4.3.5	Haem staining of strains producing hydrophobic cavity variants	103
4.3.6	Generation, expression and production of CccA:ResA fusion protein in LUL9	104
4.3.7	Western Blot of a <i>B. subtilis</i> strain producing the CccA:ResA fusion	105
4.3.8	Investigation of CccA:ResA fusion activity by haem staining	106
4.3.9	Cytochrome <i>c</i> oxidase assays of membranes from a strain producing the CccA:ResA fusion protein	107
4.4	Discussion	109

Chapter 5: *In vivo* Studies of the TDOR BdbD from *B. subtilis*.

5.1	Introduction	114
5.2	Experimental Procedures	117
5.2.1	PCR amplification of <i>bdbD</i> gene and upstream ribosome binding site	117
5.2.2	Cloning <i>bdbD</i> into pUC18	118
5.2.3	Cloning <i>bdbD</i> into pDG148	120
5.2.4	Targeted PCR mutagenesis of <i>bdbD</i>	121

5.2.5	Transformation of <i>B. subtilis</i> LUL7 protoplasts with variant <i>bdbD</i> genes	121
5.3	Results	122
5.3.1	TMPD staining of variant BdbD producing LUL7 strains.	123
5.3.2	Western Blotting	124
5.3.3	Haem staining of membranes from BdbD variant strains	126
5.4	Discussion	128

Chapter 6: Studies of the Res system in *S. coelicolor*

6.1	Introduction	131
6.2	Experimental Procedures	131
6.2.1	Generation of <i>resA</i> and <i>res</i> operon mutants	131
6.2.1.1	Verifications of cosmid StD65 by restriction digest	132
6.2.1.2	PCR amplification of the <i>resA</i> and <i>res</i> operon disruption cassettes	134
6.2.1.3	Transformation of BW25113/pIJ790 with StD65	137
6.2.1.4	Disruption of the <i>resA</i> gene and the <i>res</i> operon on StD65 by recombineering	138
6.2.1.5	Conjugative transfer of StD65 Δ <i>resA</i> and StD65 Δ <i>resA-C</i> to <i>S. coelicolor</i> from <i>E. coli</i>	143
6.2.2.	Purification of a soluble form of <i>S. coelicolor</i> ResA	144
6.2.2.1	Purification of a soluble native form of ScResA	144
6.2.2.2	Expression of GST-ScResA in <i>E. coli</i> BL21 CodonPlus(DE3)-RP	151
6.2.2.3	Purification of GST:ScResA	153
6.3	Results and preliminary characterisation of the function and physiology of <i>S. coelicolor</i> ResA	156
6.3.1	The <i>S. coelicolor</i> M145 Δ <i>resA</i> and M145 Δ <i>resA-C</i> phenotypes	156
6.3.1.1	<i>S. coelicolor</i> produces one detectable membrane associated cytochrome <i>c</i>	157
6.3.1.2	The <i>resA</i> and <i>res</i> operon deletion strains are deficient in QcrC maturation	158

6.3.2	Characterisation of ScResA	159
6.3.2.1	Determination of the extinction coefficient of ScResA at 280 nm.	159
6.3.2.2	Reduced ScResA has only one reactive thiol group at neutral pH.	161
6.3.2.3	pH stability of ScResA	162
6.3.2.4	The reactive active site cysteine of ScResA has a pK_a of 6.9 (± 0.1)	165
6.3.2.5	Reduction potential determination for ScResA.	169
6.3.2.6	Stability studies of ScResA	173
6.4	Discussion	175
Chapter 7: General discussion and future perspectives		180
Chapter 8: References		188

LIST OF FIGURES

Figure	Title	Page
<i>Chapter 1:</i>		
Figure 1.1	Schematic depiction of the accepted mechanism of disulphide exchange.	2
Figure 1.2	Structure of thioredoxin from <i>E. coli</i> .	3
Figure 1.3	Ribbon structure of the oxidised form of <i>B. subtilis</i> ResA.	6
Figure 1.4	Structure of <i>B. subtilis</i> BdbD	15
Figure 1.5	The locations of thioether and disulfide bonds in sublancin.	17
Figure 1.6	Comparison of the structural difference between sStoA and sResA from <i>B. subtilis</i> .	19
Figure 1.7	Comparison of the predicted topology of <i>B. subtilis</i> CcdA and <i>E. coli</i> DsbD.	21
Figure 1.8	Haem attachment to cytochrome <i>c</i>	22
Figure 1.9	Haem B (protohaem IX) and haem C	23
Figure 1.10	Schematic topography of <i>B. subtilis</i> cytochromes <i>c</i>	25
Figure 1.11	Aerobic respiratory chains of <i>B. subtilis</i>	26
Figure 1.12	CCM system I from <i>E. coli</i> .	32
Figure 1.13	CCM system II from <i>B. subtilis</i> .	38
Figure 1.14	The <i>res</i> operon from <i>B. subtilis</i> .	39
Figure 1.15	CCM biogenesis system III from <i>Saccharomyces cerevisiae</i> .	42
Figure 1.16	Gene organisation of the predicted <i>S. coelicolor res</i> operon	48
Figure 1.17	Sequence alignment of BsResA and ScResA	49
Figure 1.18	The <i>qcr</i> and cytochrome <i>c</i> oxidase gene cluster of <i>S. coelicolor</i>	50
<i>Chapter 2:</i>		
Figure 2.1	Insertion of pALR12 into the amyE gene in the <i>B. subtilis</i> genome	67

Figure 2.2	Conformation of gene insertion at amy locus by LUGOL staining	69
Figure 2.3	Badan	77
Figure 2.4	A typical cytochrome <i>c</i> oxidation assay plot.	79

Chapter 3:

Figure 3.1	Crystal structures of wild type ResA and active site cysteine variants	79
Figure 3.2	Insertion of pALR12 in to the <i>amyE</i> gene in the <i>B. subtilis</i> genome	82
Figure 3.3	Confirmation of gene insertion at <i>amyE</i> locus by LUGOL staining.	83
Figure 3.4	PCR amplification of <i>amyE</i> region of the LUL9 chromosome.	84
Figure 3.5	Western blot analysis of ResA production in LUL9-derivative strains	85
Figure 3.6	TMPD stains of <i>B. subtilis</i> wild type, ResA deficient and ResA active site variant strains.	86
Figure 3.7	Cytochrome <i>c</i> oxidase assays.	88
Figure 3.8	Haem stain of cytochromes <i>c</i> present in the membranes of active site cysteine variant ResA strains	91
Figure 3.9	Haem stain of cytochromes <i>c</i> found in membranes of cysteine variant ResA strains and LUL9 grown in the presence of 1 mM DTT.	92

Chapter 4:

Figure 4.1.	Crystal structure of the active site and hydrophobic cavity region of wild type <i>B. subtilis</i> ResA obtained at pH 9.25.	98
Figure 4.2.	Comparison of the N-terminal amino acid sequences from <i>B. subtilis</i> ResA and <i>B. subtilis</i> CccA.	101
Figure 4.3	PCR amplification of <i>resA</i> insertion in to the genome of	

	<i>B. subtilis</i> LUL9 at the <i>amyE</i> locus.	104
Figure 4.4	Western blot of ResA E80Q variant	105
Figure 4.5	Western blot of proline 141 variants.	106
Figure 4.6	TMPD staining of the hydrophobic cavity variants.	106
Figure 4.7	Cytochrome <i>c</i> oxidase assays of ResA hydrophobic cavity variants.	108
Figure 4.8	Haem stain of cytochromes <i>c</i> present in the membranes of <i>B. subtilis</i> strains containing hydrophobic cavity variants of ResA.	110
Figure 4.9	Western blot analysis of CccA:ResA production in the ResA deficient strain, LUL9	111
Figure 4.10	Haem stain of <i>B. subtilis</i> cytochromes <i>c</i> in LUL9/CccA:ResA membranes	112
Figure 4.11	Cytochrome <i>c</i> oxidase assays of CccA:ResA fusion	113
Figure 4.12	Alignment of haem binding domain of <i>B. subtilis</i> cytochromes <i>c</i> .	116
 Chapter 5:		
Figure 5.1	Calcium binding site of BdbD.	116
Figure 5.2	PCR amplification of the <i>bdbD</i> gene from <i>B. subtilis</i> 1A1 genomic DNA.	118
Figure 5.3	Restriction digest confirmation of <i>bdbD</i> insertion into pUC18	119
Figure 5.4	Confirmation of successful ligation of <i>bdbD</i> into pDG148	120
Figure 5.5	TMPD staining of LUL7 expressing variant <i>bdbD</i> induced by different concentrations of IPTG.	124
Figure 5.6	Western blot showing wild type BdbD production from pCHN18 in <i>B. subtilis</i> LUL7.	125
Figure 5.7	Haem stain of membrane samples prepared from LUL7/pCHN18 grown in the presence of 10–0.1 μ M IPTG.	127

Chapter 6:

Figure 6.1	Verification of StD65 by restriction digest.	134
Figure 6.2	Schematic of primer design process to design primers to PCR amplify <i>apra</i> ^R +oriT cassette with <i>ScresA</i> flanking regions.	135
Figure 6.3	PCR amplification of the <i>resA</i> and <i>resA-C</i> disruption cassettes.	137
Figure 6.4	Schematic of λ -Red mediated double crossover.	138
Figure 6.5	<i>SacI</i> restriction digest of wild type StD65 and the <i>resA</i> and <i>resA-C</i> knock out (KO) mutants.	141
Figure 6.6	PCR confirmation of <i>resA</i> and <i>resA-C</i> disruption with <i>apra</i> ^R resistance cassette.	142
Figure 6.7	Test expressions of the pCHN12 plasmid in BL21 DE3.	145
Figure 6.8	Ammonium precipitation and anion exchange of ScResA.	147
Figure 6.9	Heat stability of ScResA.	148
Figure 6.10	PCR amplification of part of <i>ScresA</i> gene predicted to encode soluble domain.	150
Figure 6.11	Conformation of correct insertion of the gene encoding <i>ScresA</i> into pUC18.	150
Figure 6.12	Confirmation of correct ScResA encoding gene insertion into pGEX-4T-1.	151
Figure 6.13	Test expression/test purification of GST:ScResA from different <i>E. coli</i> strains:	152
Figure 6.14	Coomassie stained SDS PAGE showing different stages of ScResA purification.	154
Figure 6.15	Absorbance at 280 nm from elution from gel filtration column.	155
Figure 6.16	Coomassie stained SDS PAGE showing fractions eluted from S75 gel filtration column	156
Figure 6.17	Haem stain of membranes prepared from <i>S. coelicolor</i> M145 and M145 Δ <i>qcr</i> .	157

Figure 6.18	Haem stain of M145, M145 Δ <i>resA</i> and M145 Δ <i>resA-C</i> membranes	158
Figure 6.19	Absorbance spectra of unfolded (UF) and folded (F) ScResA used to calculate the extinction coefficient.	160
Figure 6.20	Fluorescence spectra of ScResA from pH 2 to pH 12.	163
Figure 6.21	Fluorescence at 349 nm plotted as function against pH.	164
Figure 6.22	pH unfolding of ScResA.	165
Figure 6.23	Example of Badan fluorescence between 400 and 600 nm of badan incubated with ScResA.	167
Figure 6.24	Badan fluorescence at 544 nm	168
Figure 6.25	The observed pseudo first order rate constant (k_o) values plotted against pH and the curve fitted using the Henderson Hasselbalch equation.	169
Figure 6.26	Fluorescence emission spectra of oxidised and reduced ScResA	170
Figure 6.27	Fluorescence spectra of ScResA between -330 mV and -210 mV.	171
Figure 6.28	Determination of the ScResA reduction potential	172
Figure 6.29	Unfolding and refolding of ScResA in guanidine hydrochloride.	174

LIST OF TABLES

Table	Title	Page
<i>Chapter 1</i>		
Table 1.1	Changes in active site cysteine pK_a and redox midpoint potential of ResA with the active site altered to that of <i>E. coli</i> DsbA	11
<i>Chapter 2:</i>		
Table 2.1	Bacterial strains used and/or generated throughout the course of this study	54
Table 2.2	Growth media used in this work	55
Table 2.3	Typical components of a PCR amplification reaction	56
Table 2.4	Typical conditions of a DNA amplification PCR reaction	56
Table 2.5	Typical reaction mixture used for a PCR mutagenesis reaction	57
Table 2.6	Typical conditions of a site directed PCR mutagenesis reaction	57
Table 2.7	Primers used in this study	58
Table 2.8	The components of a typical ligation reaction mixture	59
Table 2.9	Plasmids and cosmids generated and/or used throughout the course of these studies	60
<i>Chapter 3:</i>		
Table 3.1	Cytochrome <i>c</i> oxidation activities of ResA and ResA active site variants	89
<i>Chapter 4:</i>		
Table 4.1	Cytochrome <i>c</i> oxidase activity of hydrophobic variants	103
Table 4.2	Cytochrome <i>c</i> oxidase assays of CccA:ResA fusion protein	108

Chapter 6:

Table 6.1	Expected bands generated by the restriction digest of StD65 with <i>EcoRI</i> and <i>HindIII</i> .	133
Table 6.2	Primers used to generate <i>resA</i> and the <i>res</i> operon knock-out disruption cassettes by PCR	136
Table 6.3	PCR reaction conditions required to amplify the <i>AprR</i> cassette with the 5' and 3' flanking regions of the gene of interest	136
Table 6.4	Predicted <i>SacI</i> restriction digest pattern of wild type StD65 and the StD65 <i>resA</i> and <i>resA-C</i> knock out mutants	139

LIST OF ABBREVIATIONS

aa	Amino acid
Abs, A	Absorbance
Amp	Ampicillin
Badan	6-Bromoacetyl-2-dimethylaminonaphthalene
B&W	Bott and Wilson solution
BCA	Bicinchononic acid
bp	Base pairs
Bs	<i>Bacillus subtilis</i>
<i>caa</i> ₃	Cytochrome <i>c</i> oxidase
Da	Daltons
DTNB	5,5'-dithio-bis(2-nitrobenzoic acid)
DTT	Dithiothreitol
ϵ	Extinction coefficient ($M^{-1} \text{ cm}^{-1}$)
EDTA	Ethylene diamide tetraacetic acid
GST	Glutathione S-transferase
IPTG	Isopropyl- β -D-thiogalactopyranoside
Kan	Kanamycin
nt	Nucleotide
dNTPs	Deoxynucleotide 5'-triphosphates
LB	Luria-Bertani broth
LBA	Luria-Bertani agar
MG	Minimal glucose
MOPS	3-(N-morpholino)-propanesulfonic acid
NSMP	Nutrient sporulation medium with phosphate

OD	Optical density
PAGE	Polyacrylamide gel electrophoresis
PCR	Polymerase chain reaction
PDB	Protein data bank
PMSF	Phenylmethanesulphonylfluoride
Sa	<i>Staphylococcus aureus</i>
Sc	<i>Streptomyces coelicolor</i>
SDS	Sodium dodecyl sulphate
TBAB	Tryptose blood agar base
TCA	Trichloroacetic acid
TIM	Inner membrane transport machinery
TMPD	N,N,N',N'-tetramethyl-p-phenylenediamide
TOM	Outer membrane transport machinery
Tris	Tris(hydroxymethyl)aminomethane
Trx	Thioredoxin
UV	Ultraviolet
% v/v	ml per 100 ml
% w/v	grams per 100 ml
WT	Wild type
X-gal	5-bromo-4-chloro-3-indonyl- β -D-galactopyranoside