

Des. Codes Cryptogr. (2007) 44:143–150
DOI 10.1007/s10623-007-9078-1

On solvable minimally transitive permutation groups

Francesca Dalla Volta · Johannes Siemons

Received: 24 September 2006 / Revised: 15 April 2007 / Accepted: 18 April 2007 /
Published online: 14 June 2007
© Springer Science+Business Media, LLC 2007

Abstract We investigate properties of finite transitive permutation groups (G, Ω) in which all proper subgroups of G act intransitively on Ω . In particular, we are interested in reduction theorems for minimally transitive representations of solvable groups.

Keywords Solvable permutation groups · Minimal transitivity

AMS Classifications 20B05 · 20D10 · 20D15

1 Introduction

A finite permutation group (G, Ω) is *minimally transitive* if G is transitive on Ω while all proper subgroups of G are intransitive on Ω . Evidently, any transitive permutation group contains minimally transitive subgroups acting on the same set and so this concept occurs naturally in reduction arguments. Closely related is the notion of minimally irreducible linear groups, namely those linear groups which act irreducibly on a vector space V while all proper subgroups leave some proper subspace of V invariant.

Solvable minimally transitive groups were first considered by Suprunenko [8] and Kopylova [3] who studied the groups of degree pq with p and q primes. More recently Lucchini [4] studied minimal generating sets in minimally transitive groups, in connection with the asymptotic properties of permutation groups considered in Pyber [7]. In Ngo [6]

Work partially supported by M.I.U.R. and London Mathematical Society.

F. Dalla Volta
Dipartimento di Matematica e Applicazioni, Università Milano-Bicocca, Via R. Cozzi 53, 20125, Milano
Italy
e-mail: Francesca.DallaVolta@unimib.it

J. Siemons (✉)
School of Mathematics, University of East Anglia, Norwich, NR4 7TJ, UK
e-mail: j.siemons@uea.ac.uk

non-regular metabelian minimally transitive groups are investigated, and Miller–Praeger [5] mention such groups in the context of vertex transitive graphs which are not Cayley graphs. A list of minimally transitive groups up to degree 30 is available in Hulpke [2], see also Conway et al. [1].

In this paper we consider solvable groups. Here in particular it happens frequently that a group action is not faithful. Therefore we study more generally arbitrary *minimally transitive representations* which may or may not be faithful. This language requires technical detail which could detract from the main matter; wherever possible we therefore try to stay close to the language of permutation groups which may appear more natural.

Any transitive permutation group contains minimally transitive subgroups and therefore it would be unreasonable to expect full classifications in general. However, under suitable restrictions some general results can be expected. For instance, for nilpotent groups there is a simple description of all their minimally transitive representations.

For a faithful action the primes dividing the order of a solvable group must divide the degree, see Theorem 3.1. In Sects. 2 and 3 we prove some reduction theorems for subgroups and factor groups. In particular, a construction is given to reduce a general minimally transitive action to the case where the degree contains only two primes. A good result is also available for actions of square-free degree, extending the work of Suprunenko and Kopylova.

2 Minimally transitive groups

Let $G \subseteq \text{Sym } \Omega$ be a transitive permutation group on a finite set Ω . Then G is *minimally transitive* on Ω if every proper subgroup of G is intransitive on Ω . In the following we consider more generally an abstract finite group G together with all its transitive actions, faithful or not. Thus if $A \subset G$ is a subgroup of G let G act on the cosets $G:A$ of A in G . The kernel of this action is the core $K_{G:A} := \bigcap_{g \in G} A^g$ of A in G .

Thus G acts *minimally transitively* on $G:A$ if and only if every subgroup H with $K_{G:A} \subseteq H \subset G$ acts intransitively on $G:A$. It will be convenient to call such a subgroup A an *mt-stabilizer* in G ; we denote this as $A \subset_m G$. Therefore $A \subset_m G$ if and only if the following holds: Whenever $H \subseteq G$ is transitive on $G:A$ then $HK_{G:A} = G$. Evidently, if A is an arbitrary subgroup of G then $G/K_{G:A}$ always acts faithfully on $G:A$ and hence is a permutation group on $G:A$. This permutation group then is minimally transitive if and only if A is an mt-stabilizer. For instance, if $A = 1$ then G is regular on $G:A$ and so $1 \subset_m G$. For another example suppose that $|G| = pq$ with distinct primes and Sylow subgroups $A \trianglelefteq G$ and $B \not\trianglelefteq G$. Then $K_{G:A} = A$ and $A \subset_m G$ while $K_{G:B} = 1$ and $B \not\subset_m G$.

2.1 Preliminaries

We begin by listing general properties of groups with minimally transitive action. For the remainder let G be a finite group and let A be a subgroup of G . The property of being an mt-stabilizer in G is quite special as it relates to the subgroup as well as its embedding in G . Let $\mathcal{L}(G)$ denote the lattice of all subgroups of G . We will begin by describing some general properties of groups in $\mathcal{L}(G)$ which are mt-stabilizers in G . The next lemma is technical but essential; the first part we use later on without further mention.

Lemma 2.1 (i) *Let $A \subseteq G$. Then $A \subset_m G$ if and only if $AH = G$ for a subgroup $H \subseteq G$ implies that $HK_{G:A} = G$.*

(ii) Let $A \subsetneq G$ and let $B \subseteq A$. Then (a): $B \subsetneq G$ or (b): $K_{G:B} \neq K_{G:A}$, $BK_{G:A} \subsetneq G$ and there exists a subgroup $H \subseteq G$ with $HK_{G:B} \neq HK_{G:A} = G$. In particular, if $A \subsetneq G$ and $K_{G:A} \subseteq B \subseteq A$ then $B \subsetneq G$.

Proof

- (i) Suppose that $A \subseteq G$ and that also $H \subseteq G$. Then H is transitive on $G : A$ if and only if $AH = G$. Therefore by definition, if $A \subsetneq G$ and if $AH = G$ then $HK_{G:A} = G$. Conversely, if $AH = G$ implies that $HK_{G:A} = G$ then H being transitive on $G : A$ means that $AH = G$ and so $HK_{G:A} = G$. Hence $A \subsetneq G$.
- (ii) Assume that $A \subsetneq G$ and $B \subseteq A$, and suppose that $B \not\subsetneq G$. Then there exists some H such that $G = BHK_{G:B}$ but $G \neq HK_{G:B}$. As $K_{G:B} \trianglelefteq K_{G:A}$ we have $G = AHK_{G:A}$. If $HK_{G:A} \neq G$ then $A \not\subsetneq G$. Therefore $HK_{G:A} = G$ and so $K_{G:A} \neq K_{G:B}$. Next we compute the core \bar{K} of $BK_{G:A}$ in G . Evidently, $K_{G:A} \subseteq \bar{K} \subseteq BK_{G:A} \subseteq A$ so that $\bar{K} = K_{G:A}$. If $BK_{G:A} \not\subsetneq G$ then there exists a subgroup \bar{H} such that $(BK_{G:A})\bar{H} = G$ but $\bar{H}\bar{K} \neq G$. But then $A(K_{G:A}\bar{H}) = G$ and $\bar{H}K_{G:A} \neq G$, a contradiction. If $K_{G:A} \subseteq B \subseteq A$ then $K_{G:B} = K_{G:A}$ and hence the second alternative can not happen. □

When dealing with the set of all m t-stabilizers in the subgroup lattice of G the following is a useful notion: If (\mathcal{L}, \leq) is any partially ordered set then a subset \mathcal{M} of \mathcal{L} is an *order ideal* in \mathcal{L} if $X \in \mathcal{M}$ and $Y \leq X$ with $Y \in \mathcal{L}$ implies that $Y \in \mathcal{M}$.

Remark 2.2 *From the last part of the lemma we deduce that the core-free m t-stabilizers in G form an order ideal in the subgroup lattice $\mathcal{L}(G)$.*

It is therefore often sufficient to know the ‘top’ m t-stabilizers, that is those which are maximal subject to being an m t-stabilizer. For instance, if G is simple then the m t-stabilizers form an order ideal and this is described completely by its top elements. We may also ask when such top elements are maximal subgroups of G . Evidently, A is maximal in G precisely when G acts primitively on $G : A$. More generally, G acts *quasi-primitively* on $G : A$ if and only if any subgroup $N \neq K_{G:A}$ with $K_{G:A} \subseteq N \trianglelefteq G$ acts transitively on $G : A$. In particular, a transitive permutation group is quasi-primitive if all its normal subgroups $\neq 1$ are transitive.

Proposition 2.3 *Let G be quasi-primitive on $G : A$. If $A \subsetneq G$ then $G/K_{G:A}$ is simple. Equivalently, if (G, Ω) is a quasi-primitive minimally transitive permutation group then G is simple.*

Proof Suppose that $A \subsetneq G$. If $G \triangleright N \supset K_{G:A}$ then N is transitive on $G : A$ as G is quasi-primitive on $G : A$. Hence $N = G$ as $A \subsetneq G$. □

2.2 A reduction theorem

When studying minimal transitivity it is obviously useful to reduce a minimally transitive action $A \subsetneq G$ to one of a smaller group or to one of smaller degree. Minimal transitivity lends itself to good reduction arguments of this kind for normal subgroups. For this let G be an arbitrary finite group with an m t-stabilizer $A \subsetneq G$ of index n in G . Let H be a normal subgroup of G with $K_{G:A} \subset H$ and $K_{G:A} \neq H \neq G$. Then H is not transitive on $\Omega := G : A$ and the orbits of H on Ω are a system of imprimitivity for G . So these are of the shape $\Omega_1, \dots, \Omega_{n^*}$ with $|\Omega_i| = s$ and $n^* := \frac{n}{s}$. Let therefore $\Omega^* := \{ \Omega_i \mid i = 1, \dots, n^* \}$.

Let Ω_1 be such that it contains the coset $1A$ and let B be the set-stabilizer of Ω_1 . In other words, $B = AH$ and in particular $H \subseteq K_{G:B}$. Now note that $B \subsetneq G$. For if $M \subseteq G$ satisfies

$BM = G$ then $AHM = G$. As $A \subset_m G$ we have $G = MHK_{G:A}$. But by choice, $K_{G:A} \subseteq H$ so that $G = MH$. As $H \subseteq K_{G:B}$ we get $G = MK_{G:B}$ and so $B \subset_m G$. Equivalently, G acts minimally transitively on Ω^* . Therefore we have the following:

Theorem 2.4 *Let $A \subset_m G$ and suppose $H \neq G$ is normal in G with $K_{G:A} \subset H \neq K_{G:A}$. Then $A \neq AH \subset_m G$.*

It is worth to formulate this statement in terms of permutation groups. In conjunction with Proposition 2.3 we have:

Theorem 2.5 *Let G be a minimally transitive permutation group on Ω . If G is quasi-primitive on Ω then G is simple. Otherwise, if H is a proper normal subgroup of G then G acts minimally transitively on the set of H -orbits.*

In other words, a minimally transitive permutation group is either simple or otherwise induces a minimally transitive action on the orbits of any normal subgroup. Another kind of reduction occurs for the action of quotient groups, and this will be used later.

Lemma 2.6 *Let N be a normal subgroup of G and let $N \subseteq A \subseteq G$. Then $A/N \subset_m G/N$ if and only if $A \subset_m G$.*

Proof If $N \subseteq A \subseteq G$ then $K_{G/N:A/N} = K_{G:A}/N$. Suppose that $A/N \subset_m G/N$ but $A \not\subset_m G$. So there exist $H \subseteq G$ with $G = AH$ and $G \neq HK_{G:A}$. Consider $G/N = A/N \cdot HK_{G:A}/N$ and evaluate $HK_{G:A}/N \cdot K_{G/N:A/N} = HK_{G:A}/N \cdot K_{G:A}/N = HK_{G:A}/N \neq G/N$, a contradiction. Conversely, suppose that $A \subset_m G$ but that $A/N \not\subset_m G/N$. So there is a subgroup $N \subseteq H \subseteq G$ with $G/N = A/N \cdot H/N$ and $H/N \cdot K_{G:A}/N = HK_{G:A}/N \neq G/N$. So $G = AH$ with $HK_{G:A} \neq G$, a contradiction. □

3 Solvable groups

For the remainder of the paper we shall restrict ourselves to minimally transitive representations of solvable groups. If n is an integer let $\pi(n)$ be the set of primes dividing n . Similarly, $\pi(G)$ and $\pi(G:H)$ are the prime divisors in $|G|$ and $|G:H|$, respectively. Also, $|n|_p$ is the highest p -power dividing n .

The following theorem states the basic relation between $\pi(G)$ and the degree of any faithful minimally transitive action when G is solvable. For nilpotent groups it completely characterizes all minimally transitive actions.

Theorem 3.1 (i) *Let $A \subset_m G$ such that $G/K_{G:A}$ is solvable. Then $\pi(G:A) = \pi(G/K_{G:A})$. In particular, for a solvable minimally transitive permutation group G of degree n we have $\pi(G) = \pi(n)$.*

(ii) *Let $A \subset G$. If $A/K_{G:A}$ is contained in the Frattini subgroup of $G/K_{G:A}$ then $A \subset_m G$. Conversely, if $G/K_{G:A}$ is nilpotent and $A \subset_m G$ then $A/K_{G:A}$ is contained in the Frattini subgroup of $G/K_{G:A}$.*

(iii) *If $A \subset_m G$ and $|G:A| = p^i$ for some prime p then $G/K_{G:A}$ is a p -group and $A/K_{G:A}$ is contained in the Frattini subgroup of $G/K_{G:A}$.*

Proof

(i) Let H be a Hall $\pi(G:A)$ -subgroup of G . Then $AH = G$ as the left hand side has order $|G|$. As $A \subset_m G$ therefore $HK_{G:A} = G$. As $K_{G:A} \subseteq A$ therefore $\pi(G:A) = \pi(G:K_{G:A})$.

- (ii) Suppose that $A \not\subseteq_m G$. Then there exists some maximal subgroup $H' \supseteq K_{G:A}$ with $G = AH'$ and $H' \neq G$. If in addition $A/K_{G:A}$ is contained in the Frattini subgroup of $G/K_{G:A}$ we have $A \subseteq H'$, a contradiction. Conversely, if $G/K_{G:A}$ is nilpotent and if $H \supseteq K_{G:A}$ is a maximal subgroup of G then H is normal in G . Therefore $AH \supseteq K_{G:A}$ is a group and if $A \subseteq_m G$ then $AH \neq G$, and hence $A \subseteq H$. (iii) This follows from (i) and (ii). □

The next result is a general splitting principle reducing representations of non-nilpotent groups to representations of subgroups, generally involving fewer primes. We denote the Fitting subgroup of X by $F(X)$.

Theorem 3.2 *Let G be a solvable group, suppose that $A \subseteq_m G$ is core-free and that A is contained in $F = F(G)$. Let $\pi^* := \pi(G : F)$ and let Q be a Hall π^* -subgroup of G . Suppose that P is a normal Sylow p -subgroup of G , let $A_P := A \cap P$ and $A_Q := A \cap Q$. Then p does not belong to π^* . Furthermore, $A_Q \times A_P$ is core-free in Q^*P and $A_Q \times A_P \subseteq_m Q^*P$ for any conjugate Q^* of Q .*

*Conversely, let P_1, P_2, \dots, P_t be the normal Sylow p_i -subgroups of G . Suppose there exist a subgroup A_Q of $F \cap Q$ and subgroups $A_{P_i} \subseteq P_i$ such that $A_Q \times A_{P_i} \subseteq_m Q^*P_i$ is core-free in Q^*P_i , for all conjugates Q^* of Q and all $i = 1, \dots, t$. Then $A_Q \times A_{P_1} \times \dots \times A_{P_t} \subseteq_m G$ is core-free in G .*

For instance, in the simplest case when $\pi^* = \{q\}$ we may take p to be any prime in $\pi(n) \setminus \{q\}$ where $n = |G : A|$. Then $Q \cap F$ is the Sylow q -subgroup of F so that A_Q is the Sylow q -subgroup of A . Similarly, A_P is the Sylow p -subgroup of A and hence $A_Q \times A_P \subseteq_m QP$ is minimally transitive of degree $|n|_q |n|_p$. Note, for at least one choice of p the group QP is not nilpotent, and evidently, groups of this type are at the basis of any induction in this case.

Proof Evidently, as P is a normal Sylow subgroup of G we have $P \subseteq F$ and so p does not belong to π^* . Put $K := K_{QP:(A_Q \times A_P)}$. Then K is centralized by every Sylow r -subgroup of G for $r \neq p$ not dividing the order of Q . Further, it is normalized by QP and hence $K \subseteq A$ is a normal subgroup of G . As A is core-free, K is trivial. Now, suppose $A_Q \times A_P \not\subseteq_m QP$. Then there exists a subgroup $Y \subseteq QP$ such that $(A_Q \times A_P)Y = QP$ but $Y \neq QP$. Let S be the direct product of all Sylow r -subgroups of F for $r \neq p$ not dividing the order of Q . This group is characteristic in F and so normal in G . Therefore, YS is a group and $A(YS) = (AY)S \supseteq (QP)S = QF = G$. However, $YS \neq G$. This is a contradiction, since $A \subseteq_m G$. Finally note that $Q \cap F$ is normal in F . Thus if Q is replaced by Q^f then $A_{Q^f} \times A_P \subseteq_m Q^fP$. But as $A \subseteq F$ we have $A \cap Q = A \cap Q^f$.

Conversely, let $A = A_Q \times A_{P_1} \dots \times A_{P_t}$. Since $A_Q \times A_{P_i}$ are core-free in QP_i also A is core-free in G . To show that $A \subseteq_m G$ suppose that this is not the case. Let therefore Y be a subgroup such that $G = AY$ but $Y \neq G$. Thus $Y = Y^*(Y_1 \times \dots \times Y_t)$ for $Y_i := Y \cap P_i$ and Y^* a Hall π^* -subgroup of Y . Therefore $Y^* \subseteq Q^f$ for some $f \in F$.

Further, $G = (P_1 \times \dots \times P_t)Q^f$ and since $Y = (Y_1 \times \dots \times Y_t)Y^*$ we have $G = AY = (A_Q \times A_{P_1} \dots \times A_{P_t})(Y_1 \times \dots \times Y_t)Y^*$. As A_Q is a π^* -subgroup of F it centralizes all terms other than Y^* . Similarly, Y_i centralizes all terms A_{P_j} with $i \neq j$. Therefore we can rewrite this as $(P_1 \times \dots \times P_t)Q^f = (A_{P_1}Y_1) \times \dots \times (A_{P_t}Y_t)(A_QY^*)$. For order reasons we have $A_QY^* = Q^f$ and $A_{P_i}Y_i = P_i$ for $i = 1, \dots, t$. As $Y \neq G$ we have $Y \cap Q^f P_r = Y^*Y_r \neq Q^f P_r$ for at least one r , say $r = 1$. Now consider $(A_{P_1} \times A_Q)(Y^*Y_1) = (A_{P_1}Y_1)(A_QY^*) = P_1Q^f$. This is a contradiction, since $A_{P_1} \times A_Q \subseteq_m P_1Q^f$. □

3.1 Representations of square-free degree:

Next we turn to representations of square-free degree. Here we get precise information on the Fitting subgroup.

Theorem 3.3 *For a solvable group G suppose that $A \leq_m G$ is core-free and has square-free index n in G . Let F be the Fitting group of G . Then $|F|$ is coprime to $|G : F|$ and all Sylow subgroups of F are elementary abelian. In particular, G is nilpotent if and only if G is cyclic of order n , with $A = 1$.*

Let $\pi^ = \pi(n) \setminus \pi(F)$ and let n^* be the product of the primes in π^* . If C is a Hall π^* -subgroup of A and if Q is a Hall π^* -subgroup of G containing C then $|Q : C| = n^*$ and the action of Q on $Q : C$ is permutationally equivalent to the action of G on $G : AF$.*

Proof If $n = p_1 p_2 \dots p_t$ with pairwise distinct primes p_i then $\pi(G) = \{p_1, p_2, \dots, p_t\}$ by Theorem 3.1. Let $N \neq 1$ be a p -subgroup of G , say $p = p_1$, which is normal in G . We claim that N is a Sylow subgroup of G . To prove this note that N has $m := \frac{n}{p}$ orbits $\Omega_1, \dots, \Omega_s, \dots, \Omega_m$ on $\Omega := G : A$, all of length $|\Omega_s| = |N : N \cap A| = p$.

Let S be a Sylow p -subgroup of AN . As AN is the setwise stabilizer of the orbit Ω_s that contains $1A$ we have that p does not divide $|G : AN|$. Hence S is a Sylow p -subgroup of G . If Q is a Hall p' -subgroup of G then $SQ = G$ so that in particular $G = (AN)Q = A(NQ)$ for order reasons. As $A \leq_m G$ we have $(NQ)K_{G:A} = G$ but as A is core-free we have $NQ = G$. Therefore $N = S$ is a Sylow subgroup of G . For any $p \in \pi(F)$ let now N be the unique Sylow p -subgroup of F . Thus N is normal in G and hence is a Sylow p -subgroup of G . It follows that $|F|$ is co-prime to $|G : F|$. By the same argument N is characteristically simple and hence elementary abelian. Evidently, if $G = F$ is nilpotent then G is abelian, hence regular on Ω and so cyclic of order $n = |\Omega|$.

As $|F|$ is co-prime to $|G : F|$ we may assume for the remainder that Q is a π^* -subgroup of G complementing F , with the further property that $Q \cap A = C$ is a π^* -subgroup of A . Then $G = QF$ with $Q \cap F = 1$ and $AF = CF$ with $C \cap F = 1$ implies that the action of G on the n^* cosets of AF in G is permutationally equivalent to the action of Q on the cosets of C . Hence $C \leq_m Q$ by Theorem 2.4. □

Some comments are in order: (1) While the theorem could be formulated for permutation groups the resulting action of G on the cosets of AN is not faithful, and the same may be true for the action of Q on the cosets of C .

(2) As G is solvable there is at least one normal p -subgroup N , as in the proof, and for this p it is the unique normal p -subgroup. This subgroup is elementary abelian, and G acts irreducibly on it.

(3) The basis of induction for square free degrees occurs when n is the product of two distinct primes. A complete analysis of the possibilities for G can be found in Suprunenko [8] and Kopylova [3]. For the reader's benefit we collect their results here.

Theorem 3.4 (Suprunenko [8]) *The permutation group G is minimally transitive of degree pq , where $q < p$ are prime numbers with q not dividing $(p - 1)$, if and only if G is isomorphic to one of the following*

- (i) *the cyclic group of order pq ,*
- (ii) *a minimal non-abelian group $G = PQ$, where $|Q| = q$ and P is normal in G , with $|P| = p^m$ where m is the exponent of $p \pmod q$, or*
- (iii) *a minimal non-abelian group $G = PQ$ with Q is normal, $|P| = p$ and $|Q| = q^r$ where r is the exponent of $q \pmod p$.*

The remaining case where q^r with $r > 0$ is the highest power of q dividing $(p - 1)$ is analyzed in Kopylova [3]. Here a similar description is obtained and it is shown that G is (i) a group of order $q^t p$ with $0 < t \leq r$; (ii) a group of order $q^{r+1} p^q$ or (iii) a group of order pq^l where l is the exponent of $q \bmod p$.

3.2 Representations of degree involving two primes:

From the discussion so far it is clear that $\{p, q\}$ -groups and their minimally transitive representations play a special role. So let $A \subsetneq G$ be core-free with $\pi(G) = \{p, q\}$. From Theorem 2.4 it is clear that any normal subgroup N in G gives rise to a minimally transitive representation $AN/N \subsetneq G/N$ of degree $\leq |G : A|$. Our first observation is the following

Lemma 3.5 *Let G be solvable and let $A \subsetneq G$ be core-free in G . Suppose that the prime q divides $|G : A|$ to the first power only. Let N be a q -group which is normal in G . Then N is elementary abelian and is a Sylow subgroup, with G acting irreducibly on N .*

Proof Let Q be a Sylow q -subgroup containing N and let P be a q' -complement in $G = PQ$. Note that the N -orbits on $G : A$ are blocks of imprimitivity, all of equal size q and AN is the stabilizer of the orbit containing $1A$. Therefore q and $|G : AN|$ are co-prime so that AN contains some Sylow q -subgroups of G . From this we have $(AN)P = G$, for order reasons. Since $A \subsetneq G$ and $A(NP) = G$ we have that $(NP)K_{G:A} = G$ but $K_{G:A} = 1$ means $NP = G$. This says that N is a Sylow q -subgroup of G and hence $N = Q$. Next replace N by a minimal normal subgroup of G . This group is elementary abelian. By the same argument N has to be Sylow q -subgroup of G . \square

The lemma suggests that the natural choice for a normal subgroup is indeed the Fitting subgroup of G . We follow through with this process when A has index $p^i q$. In this case, if either Sylow subgroup S of G is normal then $AS \subsetneq G$ gives a minimally transitive representation of a nilpotent group, and this situation is known from Theorem 3.1.

Otherwise none of the Sylow subgroups are normal and by Lemma 3.5 $F_1 = F(G)$ is a p -group. By Theorem 2.4 we have $AF_1 \subsetneq G$ and if $K \supseteq F_1$ is the core of AF_1 in G then $AF_1/K \subsetneq G/K$ is minimally transitive, faithful of degree $p^j q$ for $j < i$ and with $|G/K|_p < |G|_p$. If F_2 is the pre-image of $F(G/F)$ in G then F_2/F_1 is a q -group. Thus, if F_2 is not contained in K then Lemma 3.5 shows that F_2K/K is a normal Sylow q -subgroup of G/K . In this case we are reduced to the nilpotent case. Otherwise $K \supseteq F_2$ so that $|G/K|_q < |G|_q$. The process stops when the group becomes nilpotent or when it is of Suprunenko–Kopylova type.

References

1. Conway JH, Hulpke A, McKay J (1998) On transitive permutation groups. LMS J Comput Math 1:1–8 (electronic)
2. Hulpke A The minimally transitive groups of degree up to 30, www.math.colostate.edu/hulpke/paper/ctglist.pdf
3. Kopylova TI (1985) Solvable minimal transitive groups of permutations of degree pq . (Russian) VestsiAkad Navuk BSSR Ser Fiz-Mat Navuk 6:54–60, 126
4. Lucchini A (1998) Generating minimally transitive groups, in *Groups and Geometries* (Siena, 1996), 149–153, Trends Math, Birkhäuser, Basel, 1998
5. Miller AA, Praeger CE (1994) Non-cayley vertex-transitive graphs of order twice the product of two odd primes. J Algeb Combin 3: 77–111
6. Ngo DT (1976) Minimal transitive permutation groups. (Russian) VestsiAkad Navuk BSSR Ser Fiz-Mat Navuk 6:5–14, 140

7. Pyber L Asymptotic results for permutation groups. *Groups and computation* (New Brunswick, NJ, 1991), 197–219, DIMACS Ser Discrete Math Theoret Comput Sci 11, Amer Math Soc, Providence, RI, 1993
8. Suprunenko DA Solvable minimal transitive permutation groups of degree pq , *Soviet Math Dokl* (1986) 27:337–340; (Russian) *Dokl Akad Nauk SSSR* 269 (1983), 2:295–298 Suprunenko DA Solvable minimal transitive permutation groups of degree pq , *Soviet Math Dokl* (1986) 27:337–340; (Russian) *Dokl Akad Nauk SSSR* 269 (1983), 2:295–298