

Quantum-Classical Correspondence in Nonrelativistic Electrodynamics

B. Ritchie and C.A. Weatherford

October 14, 1999

U.S. Department of Energy

Lawrence
Livermore
National
Laboratory

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

Work performed under the auspices of the U. S. Department of Energy by the University of California Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

This report has been reproduced
directly from the best available copy.

Available to DOE and DOE contractors from the
Office of Scientific and Technical Information
P.O. Box 62, Oak Ridge, TN 37831
Prices available from (423) 576-8401
<http://apollo.osti.gov/bridge/>

Available to the public from the
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Rd.,
Springfield, VA 22161
<http://www.ntis.gov/>

OR

Lawrence Livermore National Laboratory
Technical Information Department's Digital Library
<http://www.llnl.gov/tid/Library.html>

Quantum-classical correspondence in nonrelativistic
electrodynamics

Burke Ritchie* and Charles A. Weatherford**

*University of California, Lawrence Livermore National
Laboratory, Livermore CA 94550

**Physics Department, Florida A&M University,
Tallahassee, FL 32307

A form of classical electrodynamic field exists which gives exact agreement with the operator field of quantum electrodynamics (QED) for the Lamb shift of a harmonically bound point electron. Here it is pointed out that this form of classical theory with its physically acceptable interpretation is the result of an unconventional resolution of a mathematically ambiguous term in classical field theory. Finally a quantum-classical correspondence principle is shown to exist in the sense that the classical field and expectation value of the QED operator field are identical if retardation is neglected in the latter.

There is confusion in the literature regarding the limitations of classical electrodynamics to describe the Lamb shift of a bound electron. Usually it is assumed that the Abraham-Lorentz model [1], although physically flawed, is the natural outcome of a consistent application of classical electrodynamics to the fundamental problem of the self interaction of an electron with its own radiation field. On the other hand at least two other applications exist in the literature [2-3] in which exact agreement is obtained with quantum electrodynamics (QED) for this quantity.

In this paper it is pointed out that the confusion arises from writing down the classical field in two different mathematical forms, one of which,

$$\vec{A}(\vec{r},t) = \frac{e}{c} \int_0^\infty dt' \theta(t-t') [G^{(+)} - G^{(-)}] \vec{v}'(t') \quad (1)$$

where $\theta(t-t')$ is the theta function, $G^{(+)}$ ($G^{(-)}$) are the retarded (advanced) Green's functions,

$$G^{(\pm)} = \frac{\delta(t' \pm R/c - t)}{R} \quad (2)$$

$\vec{v}'(t') = \frac{d\vec{r}'(t')}{dt'}$ is the velocity for an electron whose trajectory is $\vec{r}'(t')$, and $R = |\vec{r} - \vec{r}'(t')|$, contains mathematical ambiguities at $R = 0$ in the second term which are resolved in favor of the vanishing of the term. The second form [2-3], however, arises from writing the field in terms of inverse Fourier integrals over the radial variable in \vec{k} space and evaluating the field at $R = 0$ *before* performing the integration over $k = \omega/c$. In other words if the k integration is performed first, then Eq. (1) is the result; however if the k integration is performed after finding the field at $R = 0$, then the result is a field whose application to the Lamb shift gives exact agreement with the QED result, where the k integral is recognizable from QED as the unbounded frequency integral which diverges linearly as ω . The familiar high-frequency divergence simply reflects the capacity of a point charged particle to radiate at infinitely high frequencies either quantum mechanically or classically. The latter problem, which appears to arise from an incompleteness of the physical model, is dealt with of course using

physical arguments to renormalize the result such that the infinite contributions are cancelled.

While this would seem to be an unsatisfactory situation from a mathematical point of view, yet mathematical physics abounds with examples in which ambiguities are resolved in favor of obtaining a result with a satisfactory physical interpretation. Here we point out that the usual choice of dropping the second term of Eq. (1), in the sense that it seems inevitably to lead to physically unsuccessful concepts such as the finite extent and charge distribution of the electron, is not as compelling physically as the prescription just outlined [2-3].

For completeness we outline below the derivation of the two forms of the classical field. The Maxwell equation for the vector field arising from the motion of a point electron is,

$$\left(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2}\right) \vec{A}(\vec{r}, t) = -\frac{4\pi}{c} e\vec{v}'(t) \delta[\vec{r} - \vec{r}'(t)] \quad (3)$$

Eq. (3) can be expeditiously solved by first Fourier transforming in space and then solving the resulting second-order differential equation in the time, namely,

$$\left(\frac{\partial^2}{\partial t^2} + \omega^2\right) \vec{a}(\vec{k}, t) = 4\pi e c v'(\vec{t}) e^{-i\vec{k} \cdot \vec{r}'(\vec{t})} = \vec{\rho}(\vec{k}, t) \quad (4a)$$

$$\omega = kc, \quad (4b)$$

whose solution is

$$\vec{a}(\vec{k}, t) = \frac{i}{2\omega} \int_0^\infty dt' e^{-i\omega(t_>-t')} \vec{\rho}(\vec{k}, t') \quad (5)$$

where $t_>$ ($t_<$) is the greater (lesser) of t and t' . Eq. (5) can be written out explicitly,

$$\vec{a}(\vec{k}, t) = \frac{i}{2\omega} \left(\int_0^t dt' e^{-i\omega(t-t')} + \int_t^\infty dt' e^{i\omega(t-t')} \right) \vec{\rho}(\vec{k}, t') \quad (6)$$

In fact the integral from t to infinity can be recast as the integral from zero to infinity minus the integral from zero to t . Then on dropping the term containing the constant integral, the result is still a solution to Eq. (4a), namely,

$$\vec{a}_1(\vec{k}, t) = \frac{i}{2\omega} \left(\int_0^t dt' e^{-i\omega(t-t')} - \int_0^t dt' e^{i\omega(t-t')} \right) \vec{\rho}(\vec{k}, t') \quad (7)$$

because the term we dropped is a solution of the homogeneous equation. One may easily verify that Eq. (7) is indeed a particular

solution of Eq. (4a) by direct substitution.

Boundary conditions have been used corresponding to the propagation of positive-frequency solutions forward in time. In the temporal inverse Fourier transformation satisfying this condition, the contour is closed in the lower half-plane with detour including (excluding) the positive (negative) real axis. These boundary conditions are discussed in detail for the solution of the Dirac equation in Bjorken and Drell [4].

The final expression for the field follows on taking the spatial inverse Fourier transformation of Eq. (7),

$$\begin{aligned}
 \vec{A}(\vec{r}, t) &= \frac{1}{(2\pi)^3} \int d\vec{k} e^{i\vec{k}\vec{r}} \vec{a}_1(\vec{k}, t) \\
 &= \frac{ie}{\pi c} \int_0^t dt' \int_0^\infty d\omega \frac{[e^{-i\omega(t-t')} - e^{i\omega(t-t')}] \sin(\frac{\omega R}{c})}{R} v'(t) \\
 &= \frac{e}{c} \int_0^t dt' \frac{[\delta(t-t'-R/c) - \delta(t-t'+R/c)]}{R} v'(t)
 \end{aligned} \tag{8}$$

where the identity,

$$\frac{1}{\pi} \int_0^\infty d\omega \cos\omega(t - t' \pm R/c) = \delta(t - t' \pm R/c) \tag{9}$$

has been used. On recognizing that the integral from zero to t can be replaced by the integral from zero to infinity if the term in brackets on the right side of Eq. (8) is weighted by the theta function $\theta(t-t')$, we can immediately write Eq. (8) in the compact form given by Eq. (1). On the other hand the other prescription for writing the field [2-3] is simply to defer the integration over ω until the field, as given by the second line on the right side of Eq. (8), is evaluated at $R = 0$ and used in the Lamb shift application, whereupon the ω integral is the familiar divergent integral over emitted frequencies.

In summary the result depends critically on the order in which the limits are taken: $\omega \rightarrow \infty$ followed by $R \rightarrow 0$ or $R \rightarrow 0$ followed by $\omega \rightarrow \infty$. The second choice leads to agreement with QED, while the first choice leads to the well known models of radiation reaction which are plagued by unphysical behavior (run-away and acausality).

Finally we demonstrate the classical-QED correspondence, which was first stated in [3]. First however we require the projection of the classical field (as given by the second line of Eq. (8) evaluated at $R = 0$) onto the unit vector $\hat{\epsilon}$ in the direction of polarization of the

emitted radiation, namely,

$$\langle \vec{A} \rangle = \frac{1}{4\pi} \int d\Omega 2\hat{\epsilon}(\hat{\epsilon} \cdot \vec{A}) = -\frac{ie}{4\pi c^2} \int d\Omega \int_{-\infty}^{\infty} d\omega \int_0^t dt' (e^{i\omega(t-t')} - e^{i\omega(t'-t)}) \hat{\epsilon}[\hat{\epsilon} \cdot \vec{v}'(t')] \quad , (10)$$

where the factor of 2 accounts for the two possible polarization states. On the other hand the QED operator field is written for a two-level model of an atom [5],

$$\vec{A}_Q(0,t) = -\frac{ie}{4\pi c^2} \int d\Omega \int_{-\infty}^{\infty} d\omega \int_0^t dt' (e^{i\omega(t-t')} - e^{i\omega(t'-t)}) \hat{\epsilon}[\hat{\epsilon} \cdot \vec{p}_H/m] \quad (11)$$

where Eq. (11) is evaluated at the center of the atom such that retardation is neglected and where \vec{p}_H is the Heisenberg momentum operator, where, in order to compare with the form given in [5], we have used the operator identity,

$$i \frac{e\vec{p}_H}{m} = [R_+ - R_-] \omega_0 \vec{d} \quad , \quad (12)$$

where ω_0 and d are the transition frequency and dipole moment of the two-level atom, respectively, and R_+ , R_- are the raising and lowering operators. Note that Eq. (10) is indeed the expectation

value of Eq. (11); thus within the principal approximation of Eq. (11), neglect of retardation, the correspondence principle is established. A result identical to Eq. (11) has also been presented [3] when the model is that for the linear oscillator rather than a two-level atom.

It is not surprising that the correspondence exists only when retardation is neglected in the QED field because here the distinction between a classical point particle and a quantum point particle described by a spatially distributed wave function is made. What may be surprising however to a community which believes that quantum field theory is necessary to describe effects such as the Lamb shift is the mathematical similarity of the classical electrodynamic and QED descriptions once the decision is made in the former to represent the field as a mode sum over continuously distributed frequencies of the emitted radiation. This decision effectively lifts the mathematical ambiguity at $R = 0$ of Eq. (1) in favor of an unbounded frequency model, in agreement with QED but in disagreement with the model which diverges Coulombically as R^{-1} as R goes to zero.

Acknowledgements. This work was performed under the auspices of the U. S. Department of Energy by Lawrence Livermore National Laboratory under Contract No. W-7405-ENG-48. The author is grateful to Michael D. Feit for encouragement and helpful discussion throughout the course of this work and for confirming his derivation of the classical field. The author is also grateful to John Garrison for valuable suggestions on the presentation of the work.

References

1. E. J. Moniz and D. H. Sharp, Phys. Rev. D **10**, 1133 (1974).
2. J. M. Wylie and J. E. Sipe, Phys. Rev. A **32**, 2030 (1985). See their Eqs. (3.21)-(3.30).
3. B. Ritchie and C. M. Bowden, J. Mod. Optics **41**, 1739 (1994).
4. J. D. Bjorken and S. D. Drell, **Relativistic Quantum Mechanics** (McGraw Hill, New York, 1964), p. 94.
5. Jay R. Ackerhalt, P. L. Knight, and J. H. Eberly, Phys. Rev. Lett. **30**, 456 (1973).