

GAMMA RAY BURSTS AND RADIO LOUD ACTIVE GALACTIC NUCLEI

Gabriele Ghisellini

Osservatorio Astronomico di Brera, via Bianchi 46 I-23807 Merate Italy

Abstract. We believe that the radiation we receive from Gamma-Ray Bursts (GRBs) and radio loud Active Galactic Nuclei (AGNs) originates from the transformation of bulk relativistic motion into random energy. Mechanisms to produce, collimate and accelerates the jets in these sources are uncertain, and it may be fruitful to compare the characteristics of both class of sources in search of enlightening similarities. I will present some general characteristics of radio loud AGNs and GRBs such as their bulk Lorentz factors and the power of their jets. I will also discuss the way in which the energy in bulk relativistic motion can be transformed into beamed radiation, and consider the possibility that both classes of sources can work in the same way, namely by an intermittent release of relativistic plasma at the base of the jet: shells ejected with slightly different velocities collide at some distance from the central engine, dissipating part of their kinetic energy, and keeping the rest to power the extended radio lobes (in AGNs) or to produce the afterglow (in GRBs).

INTRODUCTION

Radio loud Active Galactic Nuclei (AGNs) and Gamma Ray Bursts (GRBs) have very little in common, at first sight. GRB are flashes of γ -ray radiation, likely flagging the birth of a stellar size black hole, while radio-loud AGNs, even if remarkable for their rapid variability, live for hundreds millions years, producing spectacular and Mpc-size jets and radio lobes, and are powered by supermassive black holes. On the other hand, in both classes the emitting plasma is moving at relativistic bulk speeds, and the radiation we see is likely the result of the transformation of part of this well ordered kinetic energy into random energy and then into radiation. Furthermore, there are strong evidences that also GBRs have collimated jets. And finally, consider that the dynamical timescale for a GRB should be of the order the light travel time to cross the gravitational radius, i.e. $R_g/c \sim 10^{-4} M_1$ seconds, where M_1 is the mass of the black hole in units of tens of solar masses. A burst with a duration of 10 seconds therefore lasts for 10^5 dynamical times: it can be a quasi steady process (for a 10^9 solar mass black hole, this time is equivalent to 30 years). What we naively consider an "explosion" is instead a long event.

In both classes of sources we have non-thermal particles and magnetic field, suggesting that non-thermal radiation processes are the main contributors to the radiation we see. This radiation, being produced by plasma in relativistic motion, is strongly beamed in the velocity direction, and we have evidences that also in GRBs the emitting fireball is collimated in a cone, i.e. a "jet". For these reasons it is instructive to compare them looking for similarities and differences, to see if their physics is similar. In the following I will briefly discuss some of the basic facts of blazars and GRBs, and discuss the possibility that, at the origin of their phenomenology, there is a common engine.




FIGURE 1. The blazar sequence: note the shift of the peak energies as the total power changes. From Fossati et al. (1998) and Donato et al. (2002).

BLAZARS

Bulk Lorentz factor — The best evidences for bulk relativistic motion in blazars come from VLBI observations of knots of radio emission moving superluminally. Apparent speeds up to $\beta_{\text{app}} = 30h^{-1}$ are measured (see e.g. Jorstad et al. 2001), at least in those blazars that are powerful γ -ray emitters, indicating even larger Lorentz factors Γ (note that Γ must always be larger than β_{app}). Indications of a large degree of beaming come also from spectral fitting, especially of low powerful, TeV emitting, blazars where beaming factors $\delta > 20$ are derived (Tavecchio et al. 1998). There are also indications that the jet could be structured, with a fast “spine” surrounded by a slower “layer” (Laing 1993; Chiaberge et al. 2001; Giroletti et al. 2003), explaining for example why the non-thermal radiation of the core of radio-galaxies is not as faint as predicted if the plasma is moving at the same large speed and if they are observed at large angles with respect to the jet axis.

The blazar sequence — Fossati et al. (1998), collecting data from three complete sample of blazars, demonstrated that the SED is controlled by the bolometric observed luminosity, with both peaks shifting at smaller frequencies when increasing the luminosity (see Fig. 1). Furthermore, the dominance of the high energy peak increases when increasing the bolometric luminosity (but this latter inference was based on the few low

power BL Lacs detected by EGRET). This “blazar sequence” can be explained by a different degree of radiative cooling: in powerful blazars electrons cool faster, producing a break in the electron distribution functions at smaller and smaller energies when increasing the total (radiation plus magnetic) energy density in the comoving frame (Ghisellini et al. 1998).

Jet power — The radio lobes of radio–galaxies and blazars are a sort of calorimeter: the power required to feed them can be calculated dividing the total energy of a radio lobe by its lifetime (estimated from spectral aging or from advance motion). This estimate has been done, among others, by Rawling & Saunders (1991): they find an average power ranging from 10^{43} – 10^{44} erg s⁻¹ for FR I radiogalaxies to 10^{46} – 10^{47} erg s⁻¹ for FR II radiogalaxies and radio–loud quasars.

One can also calculate the power carried by the jet by inferring its density through modeling the observed SED and requiring that the jet carries at least the particles and the magnetic field necessary to make the radiation we see. This has been done on the pc scale by Celotti & Fabian (1993), on sub–pc scale (the γ –ray emitting zone) by Celotti & Ghisellini (2003, see also Ghisellini 2003), and on the hundreds of kpc scale (the X–ray jets seen by Chandra) by Celotti, Ghisellini & Chiaberge (2001) and Tavecchio et al. (2000). These studies suggest large values of the power transported by the jet and require the presence of a dynamically dominating proton component (see also arguments by Sikora & Madejski 2000).

GAMMA RAY BURSTS

The main breakthrough in GRB science was the precise localization of some of them made possible by the coded mask of the wide field camera of *BeppoSAX*, which in turn made possible the prompt follow up in X–rays, optical and radio. Then it was possible to measure the redshift and end a decade long discussion about the galactic or cosmological origin of GRBs. Up to now, about 30 redshifts of GRBs have been measured. Apart from the controversial case of GRB 980425, possibly associated with the nearby SN 1998bw (at $z \sim 0.008$), all other redshifts are within the 0.17–4.5 range. A particularly useful updated link with all the relevant information about bursts with good localization is maintained by Jochen Greiner at: www.aip.de/~jcj/grbgen.html .

Duration — The majority of GRBs lasts for more than 2 seconds, while about one third is shorter. All information derived from the precise localization of GRBs refer to *long* bursts. The bimodality of the distribution of their duration is confirmed by the associated spectral shape, since short bursts, on average, appear harder than long GRBs. The light curve of GRBs is erratic and sometimes highly variable: spikes as short as a fraction of a millisecond have been detected (see Shaefer & Walker 1999). The extremely short timescales we observe demand large Lorentz factors, and the fact that the spikes at early and late times of the prompt emission have similar timescales (i.e. their duration does not increase) are major proofs against external shocks (see below) causing the prompt emission of GRBs (Fenimore, Ramirez–Ruiz & Wu 1999).

Spectra of the prompt emission — The spectra of GRBs are very hard, with a peak

(in a E - EF_E plot) at an energy E_{peak} of a few hundreds keV. Some bursts have been detected at very large γ -rays energies (> 100 MeV) by the EGRET instrument (see the review by Fishman & Meegan 1995, and references therein).

The GRB–Supernova connection — GRB 030329 is certainly associated with the supernova 2003dh (see e.g. Stanek et al. 2003). This burst, exceptionally bright because close by ($z \sim 0.17$), will probably be a Rosetta stone for GRB science. We now have quite a secure confidence that the progenitors of GRBs are massive stars, most likely exploding as SN Ic. What remains to do is to find if there is a lag between the SN and the GRB explosion in some bursts, as envisaged by the SupraNova scenario (Vietri & Stella 1998).

Iron lines — For a few bursts, there are evidence for large amounts of X-ray line emitting material around the site of the explosion. The detection of emission features (albeit with relatively low significance) in the afterglow spectra of GRBs some hours after the GRB event poses strong constraints on the properties of the line-emitting material (see Lazzati 2003 for a recent review).

Jet breaks — Assume that the burst is collimated within a cone of semiaperture θ . Assume also that, initially, the bulk Lorentz factor of the fireball is such that $1/\Gamma < \theta$. Because of relativistic aberration, the observer (which is within the cone defined by θ) will receive light only from a cone of aperture angle $1/\Gamma$. This leads to the estimate of how the received flux varies in time. If the fireball is spherical, this will continue as long as the motion is relativistic. But if the fireball is collimated, there is a time when $1/\Gamma$ becomes comparable to θ . After this time the observed solid angle will remain constant, and then there will be a change in the slope of the light curve. An *achromatic* break is predicted (“jet break”), This break allows to estimate θ and then to correct the isotropic values of the energetics of GRBs. Frail et al. (2001) in this way obtain the remarkable result that despite the “isotropic” energy values differ by some orders of magnitude, the corrected values are all very similar and cluster around a value of a few times 10^{51} erg.

Polarization — GRB 021206 was serendipitously observed by RHESSI (a satellite for solar studies), showing a prompt emission (in the hard X-rays) linearly polarized at the extraordinary level of $(80 \pm 20)\%$ (Coburn & Boggs 2003). Polarimetric observations were performed for several afterglows in the optical, finding always a moderate (but non-zero) linear polarization at the 1–3% level (see Covino et al. 2003).

Bulk Lorentz factors — If the source is moving relativistically, then the observed photon energies are blueshifted, and the typical angles (as observed in the lab frame) between photons are smaller, decreasing the probability for them to interact. Bulk Lorentz factors $\Gamma > 100$ are required to avoid strong suppression of high energy γ -rays due to photon–photon collisions.

There is a second argument demanding for strong relativistic motion, concerning the very fast observed variability. In fact the size associated with one millisecond is $R \sim 3 \times 10^7$ cm, which is much too small to be optically thin. To match the observed timescales with the size at which the fireball becomes transparent ($R_t \sim 10^{13}$ cm) we need a Doppler contraction of time given approximately by $ct_{var} \sim R_t(1 - \beta)$, yielding $\Gamma \sim 400$.




FIGURE 2. The “standard” fireball model for GRBs.

The fireball

If there is a huge release of energy in a small volume, no matter in which form the energy is initially injected, a quasi-thermal equilibrium (at relativistic temperatures) between matter and radiation is reached, with the formation of electron-positron pairs accelerated to bulk relativistic speeds by the high internal pressure. This is a *fireball* (Cavallo & Rees 1978). See Fig. 2. When the temperature of the radiation (as measured in the comoving frame) drops below ~ 50 keV the pairs annihilate faster than the rate at which they are produced. But the presence of even a small amount of barions, corresponding to only $\sim 10^{-6} M_{\odot}$, makes the fireball opaque to Thomson scattering: the internal radiation thus continues to accelerate the fireball until most of its initial energy has been converted into bulk motion. After this phase the fireball expands at a constant speed and at some point becomes transparent.

If the central engine does not produce a single pulse, but works intermittently, it can produce many shells (i.e. many fireballs) with slightly different Lorentz factors. Late but faster shells can catch up early slower ones, producing shocks which give rise to the observed burst emission. In the meantime, all shells interact with the interstellar medium, and at some point the amount of swept up matter is large enough to decelerate the fireball and produce other radiation which can be identified with the afterglow emission observed at all frequencies.

This is currently the most accepted picture for the burst and afterglow emission, and it is called the internal/external shock scenario (Rees & Mészáros 1992; Rees & Mészáros 1994; Sari & Piran 1997). According to this scenario, the burst emission is due to collisions of pairs of relativistic shells (internal shocks), while the afterglow is generated by the collisionless shocks produced by shells interacting with the interstellar medium (external shocks).

INTERNAL SHOCKS FOR BLAZARS

Rees (1978) was the first to propose the internal shocks idea to transport energy from the nucleus to the outer jet of M87 (a famous AGN). As mentioned above, this idea then became “standard” for GRBs. But it can work equally well, if not better, for blazars. For powerful blazars, in fact, we require that most of the power is not dissipated and transformed into radiation, but kept to feed the huge radio lobes. Internal shocks are not very efficient in transforming bulk into random energies (the shells are both relativistic: it is only the *relative* kinetic energy that can be dissipated). This is a problem for GRBs, due to the theoretical desire to limit the total energetic amount of GRBs, but it is welcome in the blazar field. Therefore Ghisellini (1999) and Spada et al. (2001) applied these ideas to blazars, finding that the SED and the general behavior of blazars could be well reproduced by this scenario. In addition, one can find a simple explanation of why the jet is dissipationless in its first ~ 100 Schwarzschild radii: this is the minimum distance required by a shell to catch up the previous one.

With respect to GRBs, there is an important difference: in the case of blazars the shell cannot be accelerated, initially, by its internal radiation pressure: this would cause some visible effects on their SED, which are not observed: the shells must be initially cold.

MATTER OR MAGNETICALLY DOMINATED JETS?

The internal shock scenario requires the jets to be matter dominated. The role of the Poynting flux, at the scales of the dissipation regions, should be minor. On the other hand Blandford (2003) and Lyutikov & Blandford (2003) propose an alternative scenario both for GRBs and radio loud AGNs, in which the jet is magnetically dominated at all scales. The radiation could be produced, in this case, by reconnection. Since the magnetic field is the dominant jet energy carrier, in this scenario the matter is less important, and the jet could even be made by electron–positron pairs required to produce the radiation we see, with a negligible proton component.

We have seen above that the bolometric luminosity of **blazars** is often dominated by the high energy peak: if this is due to the inverse Compton process, then the relative importance of the synchrotron component must be minor. Therefore this “Compton dominance” implies a modest role of the magnetic field in the emitting region, to limit the synchrotron emitted power. This conclusion relies on the assumption that the high energy peak of blazars is due to inverse Compton. It could be, instead, again due to the synchrotron process, by ultrarelativistic electrons and positrons, resulting from electromagnetic cascades involving relativistic protons. But in this case the cascading process would result also in an overproduction of X-rays, at the “valley” between the two peaks (see Ghisellini 2003).

For **GRBs** the issue is much more controversial (and interesting) because of two recent observational results: 1) in some burst the spectrum of the prompt emission can be fitted by a blackbody, at least for the first seconds (Ghirlanda et al. 2003); 2) the prompt emission of GRB 021206 was strongly linearly polarized (Coburn & Boggs 2003). Some blackbody radiation is expected in the standard, hot fireball, scenario: the

radiation responsible for the acceleration of the fireball can escape when the fireball becomes transparent, and it can well be blackbody in shape, in close analogy with the fossil cosmic background radiation. But the very large polarization, instead, would point towards a very important, dominant, and very well ordered magnetic field.

THE SAME ENGINE?

The primary energy source of GRBs and blazars could well be the same. The main store of energy is the spin of the black hole: for a maximally rotating Kerr hole, one can extract the 29% of its total mass–energy, amounting to $\sim 5 \times 10^{53} M/M_{\odot}$ erg. The problem is how to extract this energy sufficiently fast (especially in the case of GRBs). One promising way to extract this energy is the Blandford & Znajek (1977) process, in which the rotational energy of a Kerr black hole can be extracted by a magnetic field surrounding the hole providing a source of power:

$$L_{\text{BZ}} \sim 10^{51} \left(\frac{a}{m}\right)^2 \left(\frac{M_{\text{BH}}}{10M_{\odot}}\right)^2 \left(\frac{B}{10^{14}\text{G}}\right)^2 \text{ erg s}^{-1} \sim \left(\frac{a}{m}\right)^2 (3R_s)^2 U_{\text{BC}} \quad (1)$$

where (a/m) is the specific black hole angular momentum (~ 1 for maximally rotating black holes), R_s is the Schwarzschild radius and $U_B = B^2/(8\pi)$. The duration of a typical GRB could be associated to the duration of the accretion process (a few seconds for long bursts). To sustain such large magnetic fields, the torus surrounding them should be very dense, but this is only natural, given the fact that it is the core of the progenitor star not yet collapsed into the hole.

Even if this mechanism is purely magnetic, suggesting that the shells are born “cold” also in GRBs, it is not clear if the shell can remain cold: the shell could be initially purely magnetic and cold, but soon a fraction of the transported energy could be converted into hot pairs and trapped radiation, forming a classical hot fireball. On the other hand, for blazars, we do have observational constraints suggesting that this does not occur, but we also have other convincing evidence that the shells in blazars have magnetic fields below equipartition. The situation is rather puzzling, and this is an open issue.

One should also consider that short GRBs could be powered by the same basic mechanism even if their progenitors are two merging neutron stars (with the possible help of $\sim 10^{51}$ ergs in neutrinos).

CONCLUSIONS

GRBs last for tens of thousands of dynamical times, and are not single explosions, as supernovae, even if the association between GRBs and supernovae is now certain. It is very likely that they are collimated, and their radiation is certainly beamed. Their power can exceed $10^{50} \text{ erg s}^{-1}$ in γ -rays even accounting for collimation, and the total emitted energy is of the order of 10^{52} ergs. Being so luminous, albeit for a short time, they are the best torchlights we have to illuminate the far universe. Since they are associated with massive stars, there is the hope to study, through them, Pop III stars and the re-ionization

phases of the universe, at redshifts as large as 15–20. The basic physics of GRBs may be similar to the physics of relativistic jets in general, and therefore share many aspects with blazars, even if they are obviously more extreme. For both classes of sources we may see, at action, the more efficient engine that nature invented to produce radiation, more efficient than accretion. That this is the case is already clear considering those blazars that although having powerful jets, do not show any sign of thermal emission coming from accretion, such as lineless BL Lacs. But that jets can be orders of magnitude more powerful than accretion becomes dramatically evident with GRBs.

REFERENCES

1. Blandford R.D., 2003, in *The Physics of Relativistic Jets in the CHANDRA and XMM Era*, Bologna, in press.
2. Cavallo G. & Rees M.J., 1978, MNRAS, 183, 359
3. Celotti, A. & Fabian, A.C. 1993, MNRAS, 264, 228
5. Celotti A., Ghisellini G. & Chiaberge M., 2001, MNRAS, 321, L1
5. Celotti, A. & Ghisellini, 2003, in prep.
6. Chiaberge M., Celotti A., Capetti A. & Ghisellini G., 2000, A&A, 358, 104
7. Coburn W. & Boggs S., 2003, Nature, 423, 415
8. Covino S. et al., 2003, in *Gamma Ray Bursts in the afterglow era– 3rd Workshop*, ASP Conf. series, in press
9. Donato D. et al. 2001, A&A, 375, 739
10. Fenimore E.E., Ramirez–Ruiz E. & Wu B., 1999, ApJ, 518, L73
11. Fishman G.J. & Meegan C.A., 1995, ARA&A, 33, 415
12. Fossati G. et al. 1998, MNRAS, 299, 433
13. Frail D.A. et al., 2001, ApJ, 562, L55
14. Ghirlanda G., Celotti A & Ghisellini G., 2003, A&A, 406, 879
15. Ghisellini G. et al., 1998, MNRAS, 301, 451
16. Ghisellini G., 1999, *Astronomische Nachrichten*, 320, p. 232
17. Ghisellini G., 2003, in *The Restless High-Energy Universe* Nucl. Physics B. Suppl. Ser., in press (astro-ph/0308526)
18. Giroletti M. et al., 2003, ApJ, in press (astro-ph/0309285)
19. Jorstad S.G., Marscher A.P., Mattox J.R., Wherle A.E., Bloom S.D. & Yurchenko A.V., 2001, ApJS, 134, 181
20. Laing, R.A. 1993, in *Astrophysical Jets*, Burgarella D., Livio M., O’Dea C., eds., Cambridge University Press (Cambridge), 95
21. Lazzati D., 2003, in *Beaming and Jets in Gamma Ray Bursts*, Copenhagen (astro-ph/0211174)
22. Lyutikov M. & Blandford R.D., 2003, in *Beaming and Jets in Gamma Ray Bursts*, in press (astro-ph/0210671)
23. Rawlings, S.G. & Saunders, R.D.E., 1991, Nature, 349, 138
24. Rees M.J., 1978, MNRAS, 184, 61
25. Rees M.J. & Mészáros P., 1992, MNRAS, 258, P41
26. Rees M.J. & Mészáros P., 1994, ApJ, 430, L93
27. Sari R. & Piran, T., 1997, MNRAS, 287, 110
28. Schaefer B.E. & Walker K.C., 1999, ApJ, 511, L89
29. Sikora M. & Madejski G., 2000, ApJ, 534, 109
30. Spada M. et al., 2001, MNRAS, 325, 1559
31. Stanek K.Z. et al., 2003, ApJ, 591, L17
32. Tavecchio F., Maraschi L. & Ghisellini G., 1998. ApJ, 509, 608
33. Tavecchio F. et al., 2000, ApJ, 544, L23
34. Vietri M. & Stella L., 1998, ApJ, 507, L45