



Dietary pomegranate by-product improves oxidative stability of lamb meat

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Natalello, A., Priolo, A., Valenti, B., Codini, M., Mattioli, S., Pauselli, M., Puccio, M., Lanza, M., Stergiadis, S. and Luciano, G. (2020) Dietary pomegranate by-product improves oxidative stability of lamb meat. *Meat Science*, 162. 108037. ISSN 0309-1740 doi:
<https://doi.org/10.1016/j.meatsci.2019.108037> Available at
<http://centaur.reading.ac.uk/88016/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.meatsci.2019.108037>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1
2
3
4 **1 Dietary pomegranate by-product improves oxidative stability of lamb meat**

5
6
7 2

8 3 Antonio Natalello^{a,*}, Alessandro Priolo^a, Bernardo Valenti^b, Michela Codini^c, Simona
9
10 4 Mattioli^b, Mariano Pauselli^b, Mario Puccio^a, Massimiliano Lanza^a, Sokratis Stergiadis^d,
11
12 5 Giuseppe Luciano^a
13
14 6

15
16 7 ^a *University of Catania, Dipartimento di Agricoltura, Alimentazione e Ambiente (Di3A), Via*
17
18 8 *Valdisavoia 5, 95123, Catania, Italy*

19
20 9 ^b *University of Perugia, Dipartimento di Scienze Agrarie, Alimentari e Ambientali (DSA3), Borgo XX*
21
22 10 *Giugno 74, 06123, Perugia, Italy*

23
24 11 ^c *University of Perugia, Dipartimento di Scienze Farmaceutiche, Via Ariodante Fabretti 48, 06123,*
25
26 12 *Perugia, Italy.*

27
28 13 ^d *University of Reading, Department of Animal Sciences, School of Agriculture, Policy and*
29
30 14 *Development, PO Box 237, Earley Gate, Reading RG6 6AR, United Kingdom*

31
32 15

33
34 16 *Corresponding author. *E-mail address:* antonio.natalello@unict.it
35
36 17

Highlights

- The inclusion of 20% whole pomegranate by-product (WPB) in lamb diet was evaluated
- WPB is rich in bioactive compounds (e.g. conjugated FA, vitamin E and phenols)
- WPB-lamb meat contained higher concentration of vitamin E and PUFA
- Lipid oxidation and metmyoglobin formation of meat were reduced by WPB treatment
- Dietary WPB increased the meat antioxidant capacity in the lipophilic fraction

60
61
62 **18 ABSTRACT:**
63

64
65 19 This study investigated the effect of including whole pomegranate by-product in lamb diet on
66
67 20 meat oxidative stability. Seventeen lambs were assigned to two experimental treatments and
68
69 21 fed a cereal-based concentrate (CON) or the same concentrate where 200 g/kg DM of cereals
70
71 22 were replaced by whole pomegranate by-product (WPB). Meat from WPB-fed lambs had a
72
73 23 greater concentration of vitamin E (α - and γ -tocopherols), polyunsaturated fatty acids (PUFA),
74
75 24 highly peroxidizable PUFA and a higher peroxidability index ($P < 0.05$). Feeding WPB limited
76
77 25 the formation of metmyoglobin ($P = 0.05$) and reduced lipid oxidation (TBARS values) after
78
79 26 7 days of storage for raw meat ($P = 0.024$) or 4 days for cooked meat ($P = 0.006$). Feeding
80
81 27 WPB increased meat antioxidant capacity (ORAC assay) in the lipophilic fraction ($P = 0.017$),
82
83 28 but not in the hydrophilic. These results suggest that vitamin E in the pomegranate by-product
84
85 29 contributed to the higher antioxidant capacity of meat from the WPB-fed lambs.
86
87

88 30
89
90 31 **KEYWORDS:** pomegranate by-product; lipid oxidation; meat quality; vitamin E;
91
92 32 antioxidants; phenolic compounds.
93
94 33
95
96 34
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118

119
120
121 **35 1. Introduction**
122

123 36 The agro-industrial by-products have long been used in ruminant feeding as an effective
124 37 strategy to reduce the cost of the diet. Furthermore, the use of alternative feeds that do not
125 38 compete with human foods is currently one of the primary objectives for the scientific
126 39 community and the re-use of agro-industrial wastes is pivotal to mitigate their impact on the
130 40 environment (Salami et al., 2019). Among the many agro-industrial wastes available,
131 41 pomegranate (*Punica granatum* L.) by-products are gaining enormous interest due to the global
132 42 increase in consumption of pomegranate juice or ready-to-eat arils, also linked to the
133 43 recognition of its potential health properties. Pomegranate fruit contains numerous bioactive
134 44 compounds, such as peculiar conjugated fatty acids, phenolic compounds and vitamins, which
135 45 possess antioxidant, antimicrobial, anti-inflammatory, antitumoral and immunomodulatory
136 46 properties (Viuda-Martos, Fernández-López, & Pérez-Álvarez, 2010; Johanningsmeier &
137 47 Harris, 2011).

148 48 The pomegranate by-products have a high nutritional value as ruminant feeds, and can be
149 49 effectively used in ruminant diets to replace cereals. In a previous study, dietary administration
150 50 of fresh pomegranate peels to beef calves increased feed intake and the concentration of α -
151 51 tocopherol in plasma (Shabtay et al., 2008). Subsequent studies reported evidences of the
152 52 beneficial effects of dietary pomegranate seed pulp on goat kids' antioxidant status (Emami,
153 53 Ganjkhanlou, Fathi Nasri, Zali, & Rashidi, 2015) and meat fatty acid composition (Emami,
154 54 Fathi Nasri, Ganjkhanlou, Rashidi, & Zali, 2015). Recently, we also observed a desirable
155 55 increment of the polyunsaturated fatty acids (PUFA) in meat (Natalello et al., 2019) and milk
156 56 (Valenti, Luciano, et al., 2019) when sheep diets were supplemented with a dried whole
157 57 pomegranate by-product (WPB), which contained both seeds and peels. Nevertheless, although
158 58 a high concentration of PUFA in meat is considered desirable from a human health perspective,
159 59 the higher susceptibility of PUFA to peroxidation may drastically reduce meat shelf-life
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177

178
179
180 60 (Bekhit, Hopkins, Fahri, & Ponnampalam, 2013). Few studies have investigated the shelf-life
181
182 61 of meat from ruminants fed with pomegranate by-products. Emami, Fathi Nasri, Ganjkhanlou,
183
184 62 Zali, & Rashidi (2015) observed a greater lipid and colour stability of meat when kids were fed
185
186 63 with 150 g/ kg (dry matter basis) of pomegranate seed pulp. These authors mainly attributed
187
188 64 the reduction in lipid oxidation and colour discoloration to the phenolic compounds present in
189
190 65 the by-product. Nevertheless, the concentration of total phenols in pomegranate seed is not
191
192 66 high, especially if compared to other portions of the fruit, such as peels (Pande & Akoh, 2009;
193
194 67 Orak, Yagar,& Isbilir, 2012; Natalello et al., 2020). Moreover, as stated above, pomegranate
195
196 68 fruits and the by-products residual after juice extraction contain other bioactive compounds,
197
198 69 such as vitamin E, which could play a major role in determining the stability of meat to
199
200 70 oxidative deterioration (Bellés, Campo, Roncalés, & Beltrán, 2019).

201
202 71 To our knowledge, no other studies have tested the dietary inclusion of whole pomegranate by-
203
204 72 product on meat oxidative stability. Therefore, the aim of the present study was to investigate
205
206 73 the effect of feeding lambs with WPB on the resistance of meat to oxidation. We hypothesized
207
208 74 that the diverse bioactive molecules present in the WPB could delay the oxidative deterioration
209
210 75 even in meat with a high PUFA content. To test this hypothesis, we used the same animals
211
212 76 from the experiment by Natalello et al. (2019) and we evaluated the vitamin E and the
213
214 77 antioxidant capacity in lipophilic and hydrophilic fraction in the muscle, as well as the colour
215
216 78 and lipid stability in meat preserved in common retail conditions.
217
218
219
220

221 79

222 80 **2. Materials and methods**

223 81 *2.1. Whole pomegranate by-product*

224
225 82 The experimental feeding trial is described in detail by Natalello et al. (2019). Briefly, fresh
226
227 83 pomegranate fruits, from Wonderful variety, were processed in a local juice manufacturing
228
229 84 company (Catania, Sicily, Italy) by mechanically halving and squeezing the fruit. After
230
231
232
233
234
235
236

237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295

85 processing, the residual part containing peels, seeds, membranes and portion of arils was
86 collected and dried in a ventilated oven set at 40 °C for approximately 36 hours until constant
87 weight. Chemical composition and antioxidant capacity of the dried residual part are presented
88 in Table 1.

90 *2.2. Animals and experimental treatments*

91 The experimental procedures were approved by the University of Catania (approval:
92 015CT325). The animals were raised at the university’s experimental farm (Catania, Italy;
93 37°24’35.3’’N 15°03’34.9’’E) and handled by specialized personnel following the European
94 Union Guidelines (2010/63/ EU Directive). As described by Natalello et al. (2019), the trial
95 involved seventeen Comisana male lambs, born within an interval of 10 days in the same
96 commercial farm. At 60 days of age, animals were transported to the university facilities,
97 weighed (average body weight 14.82 kg ± 2 kg) and allocated indoors in individual pens (1.5
98 m² each). Lambs were randomly assigned to two dietary treatments, balanced for bodyweight,
99 and adapted to the experimental diet over 8 days, during which the pre-experimental
100 concentrate was gradually replaced with the experimental diets. After this adaptation period,
101 lambs were fed *ad libitum* for 36 days with a barley-corn based concentrate diet (CON, 8 lambs)
102 or a concentrate diet containing 200 g/kg dry matter (DM) of whole pomegranate by-product
103 to partially replace barley and corn (WPB, 9 lambs). Ingredients and chemical composition of
104 the experimental diets are reported in Table 1. All the ingredients were ground (5-mm screen),
105 mixed and pelleted (at 40 °C) using a pelleting machine (CMS-IEM - Colognola ai Colli,
106 Verona, Italy) to avoid selection. Lambs had free access to fresh water throughout the
107 experiment. Every day, the amount of offered and refused diet was recorded in order to
108 calculate the dry matter intake (DMI). Lambs were weighed every week from the beginning to
109 the end of the trial to calculate average daily gain (ADG).

296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354

110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134

2.3. Slaughter procedure and samplings

At the end of the trial, all animals were slaughtered on the same day at a commercial abattoir according to the European Union welfare guidelines (Council Regulation no. 1099/2009). Lambs were firstly stunned by a captive bolt and exsanguinated. Each carcass was immediately weighted and stored at 4° C for 24 h. Then, carcasses were halved and the entire *longissimus thoracis et lumborum* muscle (LTL) was excised from both sides. The right LTL was immediately vacuum-packed and stored at -80 °C until analysis of intramuscular fatty acid composition, antioxidant vitamins and antioxidant capacity. The left LTL was firstly used to measure the muscle pH by a pH-meter (HI-110; Hanna Instruments, Padova, Italy), then was aged vacuum-packaged for 3 days at 4 °C, after which it was used for oxidative stability measurements.

2.4. Feedstuffs analyses

Samples of the experimental diets were collected at the beginning, middle, and end of the trial, vacuum-packed and stored at -30 °C. Feed sample for analysis was obtained by mixing equal amounts the above subsamples collected during the trial. Neutral detergent fibre (NDF), acid detergent fibre (ADF) and acid detergent lignin (ADL) were determined according to Van Soest, Robertson, & Lewis (1991). Furthermore, crude protein, crude fat (ether extract) and ash were analysed according to the AOAC methods (1995).

Total phenolic compounds and total tannins in the feeds were extracted and determined as described by Natalello et al. (2019). Briefly, finely ground feeds (200 mg) were extracted sequentially in a sonicating water-bath with acetone 70% (v/v) followed by methanol 80% (v/v). The combined supernatants were evaporated and the residue was dissolved in methanol 70% (v/v). Total phenolic compounds were quantified by reaction of the extract with the Folin-

355
356
357 135 Ciocalteu reagent (1N) and sodium carbonate 20% (w/v), after which the absorbance at 725
358
359 136 nm was measured using a double-beam spectrophotometer (model UV-1601; Shimadzu
360
361 137 Corporation, Milan, Italy). Non-tannin phenolics were determined with the same procedure,
362
363 138 after removal of the tannins from the extract with insoluble polyvinylpyrrolidone (PVPP). The
364
365 139 concentration of total tannins was calculated as difference between total phenols and total non-
366
367 140 tannin phenols. Standard solutions of tannic acid (TA) were used to prepare an external
368
369 141 calibration curve in order to quantify phenolic compounds, which were expressed as g TA
370
371 142 equivalents/100 g dry matter. Additionally, total condensed tannins were determined by the *in*
372
373 143 *situ* thiolysis assay according to the method described by Gea, Stringano, Brown, and Mueller-
374
375 144 Harvey (2011) with slight modifications. In short, 200 mg of ground feedstuffs were weighed
376
377 145 into a screw-top glass tube, and a reagent containing 2 mL of MeOH, 1 mL of 3.3% HCl in
378
379 146 MeOH, and 100 μ L of benzyl mercaptan (BM) was added. The tubes were heated at 40 °C for
380
381 147 2 h under vigorous stirring. Then, 9 mL of 1% formic acid in water was added, and the tubes
382
383 148 were subsequently vortex mixed and centrifuged for 5 min. The supernatant was transfer to 2-
384
385 149 mL vials and analysed within 48 h by liquid chromatography-mass spectrometry (LC-MS;
386
387 150 Agilent 1100 series, Agilent Technologies, Waldbronn, Germany) with taxifolin as an external
388
389 151 standard. The concentration of condensed tannins was expressed as g/100 g dry matter.
390
391
392 152 Tocopherols from feedstuffs were extracted and analysed as described by Valenti et al. (2018).
393
394 153 Briefly, feed samples were homogenized with an ethanolic butylated hydroxytoluene (BHT)
395
396 154 solution (0.06%, w/v) and saponified with KOH (60%, w/v) at 70 °C for 30 min. Tocopherols
397
398 155 were extracted three times using hexane/ethyl acetate (9/1, v/v), dried under N₂ and dissolved
399
400 156 with acetonitrile. A 50 μ L volume was injected in a HPLC system (pump model Perkin Elmer
401
402 157 series 200), equipped with an autosampler (model AS 950-10, Tokyo, Japan) and a Synergy
403
404 158 Hydro-RP column (4 μ m, 4.6 \times 100 mm; Phenomenex, Bologna, Italy). A mobile phase
405
406 159 consisting of acetonitrile/methanol/tetrahydrofuran/1% ammonium acetate (68/22/7/3, v/v/v/v)
407
408
409
410
411
412
413

414
415
416 160 was used and the flow rate was set at 2 mL/min. The tocopherols were identified using a
417
418 161 fluorescence detector (model Jasco, FP-1525) set at excitation and emission wavelengths of
419
420 162 295 nm and 328 nm, respectively and were quantified by using external calibration curves of
421
422
423 163 commercial standard compounds (Sigma Aldrich, Steinheim, Germany).

424
425 164 The feed antioxidant capacity was determined using the oxygen radical absorbance capacity
426
427 165 (ORAC). The hydrophilic and lipophilic fractions were extracted from 1 g finely powdered
428
429 166 feed using 10 ml of either hexane or phosphate buffer (pH 7.2), for the lipophilic and
430
431 167 hydrophilic fractions, respectively. Both fractions were extracted by vortex-mixing the samples
432
433 168 for 1 min, followed by centrifugation at 4000 x g for 30 min at 25 °C. The supernatant (2 ml)
434
435 169 was stored at -80 °C prior to analysis. The ORAC assays were carried out on a FLUOstar
436
437
438 170 OPTIMA microplate fluorescence reader (BMG LABTECH, Offenburg, Germany) following
439
440 171 the procedure previously described by Valenti, Luciano, et al. (2019). In short, 2,20-azobis (2-
441
442 172 methylpropionamide) dihydrochloride (AAPH; Sigma-Aldrich) was used as peroxy radical
443
444 173 generator, Trolox was used as the reference antioxidant standard and fluorescein was used as a
445
446 174 fluorescent probe. A 100 µL volume of diluted sample, blank or Trolox calibration solution
447
448 175 (10–80µmol) was mixed with 1 mL of fluorescein (80 nM); then, 200 µL of each mixture was
449
450 176 placed in a well of the microplate. The microplate was placed in the reader and pre-incubated
451
452 177 for 20 min at 37 °C, after which AAPH (60 µL) was automatically added in each well to initiate
453
454 178 the reaction. The fluorescence was recorded every 1.9 min, using excitation and emission
455
456 179 wavelengths of 485 nm and 520 nm, respectively. The area under the fluorescence decay curve
457
458
459 180 was measured for each sample and compared to that obtained with the Trolox standard
460
461 181 solutions in order to express the data as µmol Trolox equivalents (TE) / g of sample. All the
462
463 182 reaction mixtures were prepared in duplicate, and at least three independent assays were
464
465 183 performed for each sample.

466
467 184
468
469
470
471
472

473
474
475 185 *2.5. Myoglobin concentration, fatty acids, antioxidant vitamins and antioxidant capacity of*
476
477 186 *meat*

479 187 As described by Krzywicki (1982), myoglobin (Mb) was extracted by homogenization of
480
481 188 muscle samples with phosphate buffer, followed by centrifugation at $6800 \times g$ at $4\text{ }^{\circ}\text{C}$ and
482
483 189 filtration through Whatman 541 paper. The filtered supernatant was scanned in a UV/VIS
484
485 190 spectrophotometer (UV-1601; Shimadzu Co., Milan, Italy) and the absorbance at 525 nm was
486
487 191 used to calculate Mb concentration, expressed as mg/g of fresh tissue.

490 192 Intramuscular fatty acid composition from the same muscles used here was determined by
491
492 193 Natalello et al. (2019). In this previous investigation, the detailed fatty acid profile was reported
493
494 194 as g/100 g of total fatty acids according to the purpose of that study. In the present study, the
495
496 195 main classes of fatty acids, such as saturated, monounsaturated and polyunsaturated fatty acids
497
498 196 (SFA, MUFA and PUFA, respectively), n-3 and n-6 PUFA are expressed as mg/g of muscle.
499
500 197 In addition, the susceptibility of fatty acids to oxidation was estimated by the amount of the
501
502 198 highly peroxidizable polyunsaturated fatty acids (HP-PUFA) with unsaturation degree ≥ 3 and
503
504 199 the peroxidability index was calculated according to Valenti, Natalello, et al. (2019).

507 200 The concentration of vitamin E in muscle (α - and γ -tocopherols) was analysed as described by
508
509 201 Luciano et al. (2017). Briefly, 2 g of sample was homogenized with aqueous BHT (0.06%),
510
511 202 saponified with ethanolic KOH (60%) at $70\text{ }^{\circ}\text{C}$ for 30 min and extracted with hexane/ethyl
512
513 203 acetate (9/1, v/v). The extracted solution was dried under nitrogen and resuspended with in
514
515 204 acetonitrile. The HPLC analysis of tocopherols was performed as described above for feeds.

517 205 The muscle antioxidant status was measured on the lipophilic and hydrophilic fraction by the
518
519 206 ORAC assay, using the conditions described above for feed samples.

522 207

524 208 *2.6. Meat oxidative stability*
525
526
527
528
529
530
531

532
533
534 209 Oxidative stability was measured in fresh and cooked meat over aerobic storage, as described
535
536 210 by Valenti, Natalello, et al. (2019). Briefly, six slices (2 cm thickness) of each left LTL muscle
537
538 211 were cut from the 9 to the 13 ribs using a knife. Three slices were packed under vacuum and
539
540 212 cooked for 30 min at 70 °C in a water bath. One of these was used immediately for
541
542 213 measurement of lipid oxidation (day 0), whereas the other two slices were placed in polystyrene
543
544 214 trays, over-wrapped with 3-layers of domestic cling film and stored in the dark at 4 °C for 2
545
546 215 and 4 days. The other three raw slices were immediately placed in polystyrene trays, covered
547
548 216 as cooked meat and stored at 4 °C in dark for 0 (after 2 hours of blooming), 4 and 7 days. At
549
550 217 the end of the respective storage time, each slice of raw meat was used for measuring colour
551
552 218 stability by a Minolta CM 2022 spectrophotometer (d/8° geometry; Minolta Co. Ltd. Osaka,
553
554 219 Japan) set to operate in the specular components excluded (SCE) mode and to measure with
555
556 220 the illuminant A and 10° standard observer. Two measurements were taken on the meat surface
557
558 221 and the mean value was calculated. The colour descriptors L* (lightness), a* (redness), b*
559
560 222 (yellowness), C (saturation) and h_{ab} (hue angle) were measured in the CIE L* a* b* colour
561
562 223 space. The reflectance spectra from 400 to 700 nm wavelength were recorded for calculation
563
564 224 of metmyoglobin percentage (MMb) formation as described by Krzywicki (1979).
565
566 225 For both raw and cooked slices, lipid oxidation was determined by measuring the 2-
567
568 226 thiobarbituric acid reactive substances (TBARS) at the end of each storage time, as described
569
570 227 by Valenti, Natalello et al. (2019). Meat samples (2.5 g) were homogenized with 12.5 mL of
571
572 228 distilled water using a Heidolph Diax 900 tissue homogenizer (Heidolph Elektro GmbH & Co.
573
574 229 KG, Kelheim, Germany) operating at 9500 rpm. During the homogenization, samples were put
575
576 230 in a water/ice bath. Subsequently, 12.5 mL of 10% (w/v) trichloroacetic acid (TCA) was added
577
578 231 to precipitate proteins, after which samples were filtered through Whatman No. 1 filter paper.
579
580 232 The clear filtrate (4 mL) was added to 1 mL of 0.06M aqueous thiobarbituric acid into pyrex-
581
582 233 glass tubes. The tubes were incubated in a water bath at 80 °C for 90 min and the absorbance
583
584
585
586
587
588
589
590

591
592
593 234 of each sample was read at 532 nm using a Shimadzu UV/vis spectrophotometer (UV-1601;
594
595 235 Shimadzu Corporation, Milan, Italy). The assay was calibrated with solutions of TEP (1,1,3,3,-
596
597 236 tetraethoxypropane) in distilled water ranging from 0 to 65 nmoles/4mL. Results were
598
599 237 expressed as mg of malonaldehyde (MDA)/kg of meat.
600

601
602 238

603 239 *2.7. Statistical Analysis*

604
605
606 240 Data on animal performances and intakes, as well as on fatty acid classes, myoglobin,
607
608 241 tocopherols and antioxidant capacity of meat were analysed using the general linear model
609
610 242 (GLM) to test the effect of the dietary treatment. Data on oxidative stability measured in raw
611
612 243 and cooked meat were analysed using a mixed model to test the effect of the dietary treatment
613
614 244 and of the time of storage, as well as of their interaction as the fixed factors, while individual
615
616 245 lamb was considered a random effect. Differences between means were assessed using the
617
618 246 Tukey's Honest Significant Difference test. Significance was declared at $P \leq 0.05$, while trends
619
620 247 toward significance were considered when $0.05 < P \leq 0.10$. Statistical analyses were performed
621
622 248 using Minitab, version 16 (Minitab Inc., State College, PA, USA).
623
624

625 249

626 627 250 **3. Results**

628 629 251 *3.1. Feed composition, animal performances and intakes*

630
631 252 As shown in Table 1, the partial replacement of barley and corn with the whole pomegranate
632
633 253 by-product produced slight variations in the nutrient composition parameters of the diet mostly
634
635 254 related to the fibre fractions, with higher NDF and ADF in the WPB diet compared to CON.
636
637 255 The whole pomegranate by-product contained almost 10% DM of total phenolic compounds,
638
639 256 mostly represented by tannins (98.21% of total phenols). Consequently, the WPB diet had a
640
641 257 greater content of total phenolic compounds and total tannins compared to the CON diet.
642
643 258 Regarding vitamin E, α -tocopherol represented the main compound when compared with γ -
644
645
646
647
648
649

650
651
652 259 tocopherol in the pomegranate by-product. Both compounds were found at a greater
653
654 260 concentration in the WPB diet when compared with CON. The hydrophilic fraction accounted
655
656 261 for most of the antioxidant capacity (ORAC) in all the experimental feeds analysed and the
657
658 262 WPB diet exhibited a greater antioxidant capacity (ORAC) of both hydrophilic and lipophilic
659
660 263 fractions compared to CON.

662
663 264 The dietary treatment did not affect ($P > 0.05$) the performance parameters of lambs, measured
664
665 265 as final bodyweight, carcass weight, average daily gain, voluntary feed intake and feed
666
667 266 conversion ratio (Table 2). The above differences in the concentration of phenolic compounds
668
669 267 between the diets, led to a greater intake of total phenolic compounds and tannins by lambs fed
670
671 268 the WPB diet ($P < 0.001$). Similarly, feeding the WPB diet increased the daily intake of α - and
672
673 269 γ -tocopherols compared to the CON treatment ($P < 0.001$).

674
675
676 270

677 271 *3.2. Myoglobin, vitamin E, fatty acids and antioxidant capacity of meat.*

678 272 As shown in Table 3, the dietary treatment did not affect the ultimate pH of meat, the
679
680 273 concentration of myoglobin and the intramuscular fat content ($P > 0.05$). Feeding the WPB
681
682 274 diet increased the concentration of vitamin E in meat (α - and γ -tocopherols; $P < 0.001$).
683
684 275 Regarding the fatty acid composition of the intramuscular fat, the dietary treatment did not
685
686 276 affect the concentration of saturated and monounsaturated fatty acids (SFA and MUFA,
687
688 277 respectively; $P > 0.05$), while a greater concentration of polyunsaturated fatty acids (PUFA)
689
690 278 was found in meat from the WPB-fed lambs ($P < 0.05$). Particularly, compared to CON, feeding
691
692 279 WPB increased the concentration of highly peroxidizable (HP) PUFA with at least three double
693
694 280 bonds and the peroxidability index of intramuscular fatty acids ($P < 0.05$). Lastly, as shown in
695
696 281 Figure 1, the WPB diet increased the antioxidant capacity (ORAC) of the lipophilic fraction of
697
698 282 meat ($P < 0.05$).

699
700
701 283

709
710
711 284 *3.3. Meat oxidative stability*
712

713 285 The effect of the dietary treatment and time of storage on the oxidative stability parameters
714
715 286 measured in raw and cooked meat are reported in Table 4. The time of storage affected some
716
717
718 287 of the colour parameters measured in raw meat, with a* values decreasing over the 7 days of
719
720 288 storage, while b* and h_{ab} values were increased ($P < 0.001$). Also, the percentages of
721
722 289 metmyoglobin (MMb%) increased over time, indicating meat browning ($P < 0.001$). The
723
724 290 dietary treatment did not affect any of the colour parameters overall measured in meat during
725
726 291 storage, except for L* values which were lower in meat from the CON group ($P < 0.01$).
727
728 292 Additionally, feeding the WPB diet reduced the average MMb% measured in meat across the
729
730 293 7-day storage period ($P = 0.05$). In both raw and cooked meat, lipid oxidation (TBARS values)
731
732 294 increased over storage duration ($P < 0.001$) and the WPB diet reduced the extent of lipid
733
734 295 oxidation overall measured in meat over time ($P < 0.05$). A significant diet \times time interaction
735
736 296 was found for TBARS values measured in raw meat. Specifically, compared to day 0, while
737
738 297 the TBARS values increased already after 4 days in meat from CON-fed animals, lipid
739
740 298 oxidation increased in WPB meat after 7 days ($P < 0.05$; Figure 2). Raw meat from lambs in
741
742 299 the WPB treatment had lower TBARS values compared to the CON group after 7 days of
743
744 300 storage ($P < 0.05$). Statistically comparable results were observed between fresh meat (day 0)
745
746 301 from control group and the WPB meat stored over 7 days ($P > 0.05$).
747
748
749
750 302

751
752 303 **4. Discussion**
753

754 304 The resistance of meat to oxidation depends upon the complex balance between pro-oxidant
755
756 305 factors, such as readily oxidizable substrates and catalysts, and various antioxidant defences,
757
758 306 including both endogenous systems and exogenous antioxidants of dietary origin (Bekhit et al.,
759
760 307 2013). Among the pro-oxidant factors, heme iron in myoglobin has been demonstrated to
761
762 308 promote the initiation of lipid oxidation (Baron & Andersen, 2002), so that a greater content of
763
764
765
766
767

768
769
770 309 myoglobin in muscle might increase its susceptibility to lipid peroxidation. Nevertheless, in
771
772 310 the present study, we did not observe differences in the concentration of myoglobin between
773
774 311 meat from CON- and WPB-fed lambs. The fatty acid composition of the intramuscular fat is
775
776 312 another factor determining meat oxidative stability. Particularly, polyunsaturated fatty acids
777
778 313 (PUFA) are the primary target for lipid oxidation (Bekhit et al., 2013) and their susceptibility
780
781 314 to oxidation increases with increasing degree of unsaturation (Johnson & Decker, 2015).
782
783 315 Therefore, feeding strategies aimed at increasing the PUFA concentration in muscle may
784
785 316 present the drawback of impairing the oxidative stability of meat if not balanced by adequate
786
787 317 antioxidant interventions (Bekhit et al., 2013). In the present study, we found that the
788
789 318 concentration of PUFA was higher (+38%) in meat from lambs fed WPB compared to the
790
791 319 control. In a previous study, Natalello et al. (2019) investigated the effect of the whole
792
793 320 pomegranate by-product on the fatty acid metabolism of lambs and reported the detailed fatty
794
795 321 acid composition of muscle, liver and ruminal digesta from the same animals used in the present
796
797 322 study. As reported in that study, the greater concentration of total PUFA observed in meat from
798
799 323 the WPB-fed animals was due to the combined effect of both PUFA and bioactive substances
800
801 324 (such as tannins, able to alter the ruminal lipid metabolism) present in the pomegranate by-
802
803 325 product. Consequently, compared to CON, meat from the WPB-fed lambs contained specific
804
805 326 PUFA derived from the whole pomegranate by-product (i.e., conjugated linolenic acid
806
807 327 isomers). Also, other PUFA, such as rumenic acid, were more abundant in meat from animals
808
809 328 in the WPB treatment compared to the CON group (Natalello et al., 2019). In the present study,
810
811 329 caused by the greater amount of highly unsaturated PUFA, we also observed a greater
812
813 330 peroxidability index in meat from the WPB-fed lambs. Therefore, a possible higher
814
815 331 susceptibility to lipid oxidation could be expected in meat from the WPB-fed animals.
816
817 332 Nevertheless, no reduction of meat shelf-life was observed; on the contrary, the formation of
818
819 333 the secondary lipid oxidation products (TBARS) was reduced in raw meat. Furthermore, meat
820
821
822
823
824
825
826

827
828
829 334 from the WPB-fed lambs displayed a greater resistance to lipid oxidation even under more
830
831 335 stressful oxidative conditions, such as cooking. These results demonstrate that feeding the
832
833 336 whole pomegranate by-product improved the antioxidant capacity of meat.
834
835
836 337 In the present study, we measured the overall antioxidant capacity of meat using the ORAC
837
838 338 (oxygen radical absorbance capacity) assay which was adopted because it first offers a high
839
840 339 sensitivity compared to other common assays (Cao & Prior, 1998). Moreover, while most of
841
842 340 the other tests only measure antioxidants in the hydrophilic fraction, the ORAC assay has been
843
844 341 successfully adopted to separately determine the antioxidant capacity of the lipophilic and
845
846 342 hydrophilic fractions of the sample (Prior et al., 2003; Huang, Ou, Hampsch-Woodill,
847
848 343 Flanagan, & Deemer, 2002). In the case of dietary phenolic compounds, this approach can be
849
850 344 useful to assess their antioxidant activities in the diet and their possible antioxidant effects in
851
852 345 animal tissues, as most of these compounds and their metabolites have hydrophilic nature.
853
854 346 Therefore, due to the greater concentration of phenolic compounds in the WPB diet compared
855
856 347 to the CON, it is not surprising that the former displayed a much higher ORAC value in the
857
858 348 hydrophilic fraction and this finding agree with previous reports on the antioxidant capacity of
859
860 349 pomegranate fruits (Elfalleh et al., 2011; Valenti, Luciano et al., 2019).
861
862 350 Nevertheless, although the animals fed WPB ingested a higher quantity of polyphenols than
863
864 351 those fed the CON diet control, the antioxidant capacity of the hydrophilic fraction of meat was
865
866 352 not affected by the dietary treatment. Even if the bioavailability of phenolic compounds was
867
868 353 not tested, this result might lead to suppose that pomegranate phenolic compounds did not
869
870 354 contribute to improving meat oxidative stability with a mechanism that involved their intestinal
871
872 355 absorption and transfer to the muscle as previously suggested. Kotsampasi et al. (2014)
873
874 356 reported that a greater content of phenolic compounds in meat from lambs fed with a silage
875
876 357 pomegranate by-product. However, the Folin-Ciocalteu method used by the authors is biased
877
878 358 by several interfering substances, some of which present in muscle, with a consequent
879
880
881
882
883
884
885

886
887
888 359 erroneous measure of the phenolic compounds (Georgé, Brat, Alter, & Amiot, 2005). Also,
889
890 360 among the heterogeneous class of phenolic compounds, tannins are considered to be poorly
891
892 361 bioavailable in animals (Vasta & Luciano, 2011; López-Andrés et al., 2013). In this context, it
893
894 362 is of note that phenolic compounds in the WPB used in the present study were almost
895
896 363 exclusively represented by tannins (approximately 98%), in agreement with previous reports
897
898 364 (Seeram, Lee, Hardy, & Heber, 2005; Mphahlele, Fawole, Mokwena, & Opara, 2016; Natalello
899
900 365 et al., 2020).

903 366 Differently from the hydrophilic fraction, our results demonstrate that feeding the whole
904
905 367 pomegranate to lambs increased the antioxidant capacity in the lipophilic fraction of muscle.
906
907 368 This result might be explained by the greater concentration of Vitamin E (α - and γ -tocopherols)
908
909 369 in muscle from lambs in the WPB treatment, as tocopherols react in the ORAC assay (Huang
910
911 370 et al., 2002). The composition of the diets can directly explain the results found on the
912
913 371 concentration of vitamin E in meat. Indeed, vitamin E is highly bioavailable as it is largely
914
915 372 hydrolysed in the intestine and then absorbed in combination with lipid micelles. For this
916
917 373 reason, the concentration of vitamin E in muscle responds to its content in the diet, as
918
919 374 extensively demonstrated in different animal species (Bellés et al., 2019; Sales & Koukolová,
920
921 375 2011). Therefore, in the present study, the greater deposition of vitamin E in meat from the
922
923 376 WPB-fed lambs could derive from their higher intake of tocopherols, especially α -tocopherol
924
925 377 which is the most bioavailable among the vitamin E isoforms (Bellés et al., 2019). Moreover,
926
927 378 compared to the CON diet, the WPB diet contained a greater amount of lipids, which can
928
929 379 further increase the intestinal absorption of vitamin E (Lodge, Hall, Jeanes, & Proteggente,
930
931 380 2004). Finally, other possible effects of feeding whole pomegranate by-product on the
932
933 381 concentration of vitamin E in meat cannot be excluded. For example, it has been suggested that
934
935 382 dietary phenolic compounds could exert indirect antioxidant effects. Among these, polyphenols
936
937 383 could protect and/or regenerate other antioxidant compounds, such as vitamin E, in the
938
939
940
941
942
943
944

945
946
947 384 gastrointestinal tract or in the animal tissues for the most bioavailable compounds (Halliwell,
948
949 385 Rafter, & Jenner, 2005; Iglesias, Pazos, Torres, & Medina, 2012). In agreement with this
950
951 386 observation, recent studies demonstrated a greater concentration of tocopherols in meat and
952
953 387 milk from sheep fed diets supplemented with phenolic compounds (Lobón, Sanz, Blanco,
954
955 388 Ripoll, & Joy, 2017; Ortuño, Serrano, & Bañón, 2015; Valenti, Natalello et al., 2019).
956
957 389 Therefore, it can be supposed that the polyphenols contained in the pomegranate by-product
958
959 390 used in the present study might have contributed to the greater deposition of vitamin E in meat
960
961 391 from the WPB-fed animals and future studies would be necessary to investigate this possible
962
963 392 effect.

964
965
966 393 Vitamin E has been extensively shown to be one of the main determinants of meat oxidative
967
968 394 stability (Bellés et al., 2019). Indeed, it has been demonstrated that pro-oxidant factors, such
969
970 395 as intramuscular polyunsaturated fatty acids and hem iron content, play a less important role
971
972 396 when feeding strategies promote an adequate deposition of vitamin E in meat (Ponnampalam,
973
974 397 Norn, Burnett, Dunshea, Jacobs, & Hopkins, 2014). Therefore, in the present study, it is
975
976 398 plausible that vitamin E exerted a considerable role in reducing the extent of lipid oxidation in
977
978 399 raw and cooked meat from lambs fed the WPB diet. This is the first study demonstrating that
979
980 400 vitamin E could contribute to the antioxidant capacity of dietary pomegranate by-products as,
981
982 401 to our knowledge, no previous studies have determined the vitamin E concentration, or the
983
984 402 lipophilic antioxidant capacity, in muscle from animals fed with pomegranate by-products.
985
986 403 Therefore, it is not possible to fully compare our results with the existing literature and further
987
988 404 studies would be necessary to confirm these results. Although comparisons between studies
989
990 405 should always be made with caution due to the different experimental conditions, it is possible
991
992 406 that vitamin E contributed to the improvement of meat oxidative stability observed also in
993
994 407 previous studies where lambs and kids were fed with pomegranate by-products. For example,
995
996 408 Emami, Fathi Nasri, Ganjkanlou, Zali, et al. (2015) used a by-product mostly composed of
997
998
999
1000
1001
1002
1003

1004
1005
1006 409 pomegranate seeds and attributed the observed antioxidant effects mainly to the phenolic
1007
1008 410 compounds. Nevertheless, it has been shown that tocopherols occur at higher concentrations in
1009
1010 411 pomegranate seeds than in other parts of the pomegranate fruit, such as peels and pulp (Pande
1011
1012 & Akoh, 2009). Furthermore, tocopherols were shown to be the main contributors to the
1013 412 antioxidant capacity of pomegranate seed oil, while phenolic compounds are associated to the
1014
1015 413 antioxidant capacity of other portions of the fruits such as peels (Elfalleh, et al. 2011). Similar
1016
1017 414 considerations may apply to the results provided by other studies in which dietary pomegranate
1018
1019 415 seed pulp or other by-products containing seeds reduced lipid oxidation and improved
1020
1021 416 antioxidant activity in meat (Emami, Fathi Nasri, Ganjkhanelou, Rashidi, et al., 2015;
1022
1023 417 Kotsampasi et al., 2014).
1024
1025
1026 418
1027
1028 419 Regarding meat colour, it is not easy to propose a plausible explanation for the higher L* values
1029
1030 420 found, in the present study, in meat from lambs given the WPB diet. Indeed, factors potentially
1031
1032 421 affecting meat lightness, such as ultimate pH, intramuscular fat and the concentration of
1033
1034 422 myoglobin in meat did not differ between treatments. It is possible to suppose an effect of
1035
1036 423 tannins from pomegranate in increasing L* values, as it was reported in studies where lambs
1037
1038 424 were fed with different tannin-containing feeds, such as acacia foliage, fresh sulla or carob pulp
1039
1040 425 (Priolo, Waghorn, Lanza, Biondi, and Pennisi, 2000; Priolo, Ben Salem, Atti, and Nefzaoui,
1041
1042 426 2002; Priolo et al., 2005). On the other colour parameters, it is known that meat browning,
1043
1044 427 caused by the redox conversion of myoglobin forms, can be evaluated by the instrumental
1045
1046 428 colour measurement, with some descriptors being particularly relevant (Mancini & Hunt,
1047
1048 429 2005). Specifically, the decrease of a* values and the increase of b* and h_{ab} values over time
1049
1050 430 of storage observed in the present study are consistent with several shelf-life studies on lamb
1051
1052 431 meat stored in comparable conditions (Aouadi et al., 2014; Valenti, Natalello, et al., 2019;
1053
1054 432 Luciano et al., 2019). Although the saturation index (C values) has often been reported to
1055
1056 433 decrease following meat browning, C values did not change over time in the present study. In
1057
1058
1059
1060
1061
1062

1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121

434 agreement with our results, Emani, Fathi Nasri, Ganjkhanlou, Zali, et al. (2015) studied the
435 effects of feeding kids with pomegranate seed pulp on meat oxidative stability and reported
436 that, despite the variation of a*, b* and h_{ab} values, the saturation index was not affected by the
437 storage time. Lastly, it has been reported that strategies to reduce the extent of lipid oxidation
438 often improve the stability of myoglobin to oxidation (Faustman, Sun, Mancini, & Suman,
439 2010). Therefore, in the present study, the ability of dietary pomegranate by-product to reduce
440 lipid oxidation might explain the reduction of metmyoglobin accumulation observed in meat
441 from WPB-fed animals compared to the CON treatment. However, it should be stressed that
442 the accumulation of metmyoglobin over time was overall numerically small, albeit significant.
443 This could partially explain the absence of differences in the colour descriptors between
444 treatments, as well as their negligible numerical variation over time.

5. Conclusions

447 The results of this study provided evidence that a high amount of conventional cereal grains in
448 the diet of lambs can be replaced by whole pomegranate by-product without negative effects
449 on animal performances. Furthermore, this dietary strategy led to a reduction of lipid oxidation
450 in fresh and cooked meat during refrigerated storage, despite the greater concentration of
451 polyunsaturated fatty acids, while the formation of metmyoglobin was also slightly reduced.
452 These results could be linked to the higher concentration of vitamin E in muscle from animals
453 fed whole pomegranate by-product, which was associated to the higher antioxidant capacity
454 measured in the lipophilic fraction of muscle. Therefore, these findings suggest for the first
455 time vitamin E from dietary pomegranate by-products as a main factor contributing to improve
456 meat oxidative stability.

Acknowledgements

1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180

459 This study was supported by the project “Ricerca di Base di Ateneo DSA3/2015”, University
460 of Perugia, Department DSA3 and was conducted under the 2-year collaborative research
461 program (2016-2018) established between the Departments Di3A (University of Catania) and
462 DSA3 (University of Perugia). Authors would also like to thank Mr. Chris Humphrey
463 (University of Reading) for his help with the profiling of condensed tannins in animal feed.

465 **References**

466 AOAC (1995). *Association of Official Analytical Chemists. Official methods of analysis* (16th
467 ed.). Washington, DC:AOAC.

468 Aouadi, D., Luciano, G., Vasta, V., Nasri, S., Brogna, D.M.R., Abidi, S., Priolo, A., & Ben
469 Salem, H. (2014). The antioxidant status and oxidative stability of muscle from lambs
470 receiving oral administration of *Artemisia herba alba* and *Rosmarinus officinalis essential*
471 *oils. Meat Science, 97, 237-243*

472 Baron, C.P., & Andersen, H.J. (2002). Myoglobin-induced lipid oxidation. A review. *Journal*
473 *of Agricultural and Food Chemistry, 50, 3887–3897*

474 Bekhit, A. E. D. A., Hopkins, D. L., Fahri, F. T., & Ponnampalam, E. N. (2013). Oxidative
475 processes in muscle systems and fresh meat: Sources, markers, and remedies.
476 *Comprehensive Reviews in Food Science and Food Safety, 12(5), 565-597.*

477 Bellés, M., del Mar Campo, M., Roncalés, P., & Beltrán, J. A. (2019). Supranutritional doses
478 of vitamin E to improve lamb meat quality. *Meat Science, 149, 14-23*

479 Cao, G., & Prior, R. L. (1998). Comparison of different analytical methods for assessing total
480 antioxidant capacity of human serum. *Clinical Chemistry, 44(6), 1309-1315.*

481 Elfalleh, W., Tlili, N., Nasri, N., Yahia, Y., Hannachi, H., Chaira, N., Ying, M., & Ferchichi,
482 A. (2011). Antioxidant capacities of phenolic compounds and tocopherols from Tunisian
483 pomegranate (*Punica granatum*) fruits. *Journal of Food Science, 76(5), C707-C713.*

1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239

484 Emami, A., Fathi Nasri, M. H., Ganjkhanlou, M., Rashidi, L., & Zali, A. (2015). Dietary
485 pomegranate seed pulp increases conjugated-linoleic and-linolenic acids in muscle and
486 adipose tissues of kid. *Animal Feed Science and Technology*, 209, 79-89.

487 Emami, A., Fathi Nasri, M. H., Ganjkhanlou, M., Zali, A., & Rashidi, L. (2015). Effects of
488 dietary pomegranate seed pulp on oxidative stability of kid meat. *Meat Science*, 104, 14-
489 19.

490 Emami, A., Ganjkhanlou, M., Fathi Nasri, M. H., Zali, A., & Rashidi, L. (2015). Pomegranate
491 seed pulp as a novel replacement of dietary cereal grains for kids. *Small Ruminant*
492 *Research*, 123(2-3), 238-245.

493 Faustman, C., Sun, Q., Mancini, R., & Suman, S. P. (2010). Myoglobin and lipid oxidation
494 interactions: Mechanistic bases and control. *Meat science*, 86(1), 86-94.

495 Gea, A., Stringano, E., Brown, R. H., & Mueller-Harvey, I. (2011). In situ analysis and
496 structural elucidation of sainfoin (*Onobrychis viciifolia*) tannins for high-throughput
497 germplasm screening. *Journal of Agricultural and Food Chemistry*, 59(2), 495–503.

498 Georgé, S., Brat, P., Alter, P., & Amiot, M. J. (2005). Rapid determination of polyphenols and
499 vitamin C in plant-derived products. *Journal of Agricultural and Food Chemistry*, 53(5),
500 1370-1373.

501 Halliwell, B., Rafter, J., & Jenner, A. (2005). Health promotion by flavonoids, tocopherols,
502 tocotrienols, and other phenols: direct or indirect effects? Antioxidant or not? *The*
503 *American journal of clinical nutrition*, 81(1), 268S-276S.

504 Huang, D., Ou, B., Hampsch-Woodill, M., Flanagan, J.A., & Deemer, E.K. (2002).
505 Development and validation of oxygen radical absorbance capacity assay for lipophilic
506 antioxidants using randomly methylated beta-cyclodextrin as the solubility enhancer.
507 *Journal of Agricultural and Food Chemistry*, 50,1815-1821

- 1240
1241
1242
1243 508 Iglesias, J., Pazos, M., Torres, J.L., & Medina, I. (2012). Antioxidant mechanism of grape
1244
1245 509 procyanidins in muscle tissues: redox interactions with endogenous ascorbic acid and α -
1246
1247 510 tocopherol. *Food Chemistry*, *134*, 1767–1774.
- 1248
1249 511 Johanningsmeier, S. D., & Harris, G. K. (2011). Pomegranate as a functional food and
1250
1251 512 nutraceutical source. *Annual review of Food Science and Technology*, *2*, 181-201.
- 1252
1253 513 Johnson, D. R., & Decker, E. A. (2015). The role of oxygen in lipid oxidation reactions: a
1254
1255 514 review. *Annual Review of Food Science and Technology*, *6*, 171-190.
- 1256
1257 515 Kotsampasi, B., Christodoulou, V., Zotos, A., Liakopoulou-Kyriakides, M., Goulas, P.,
1258
1259 516 Petrotos, K., Natas, P., & Bampidis, V.A. (2014). Effects of dietary pomegranate
1260
1261 517 byproduct silage supplementation on performance, carcass characteristics and meat quality
1262
1263 518 of growing lambs. *Animal Feed Science and Technology*, *197*, 92-102.
- 1264
1265
1266 519 Krzywicki, K. (1979). Assessment of relative content of myoglobin, oxymyoglobin and
1267
1268 520 metmyoglobin at the surface of beef. *Meat Science*, *3(1)*, 1–10.
- 1269
1270 521 Krzywicki, K. (1982). The determination of haem pigments in meat. *Meat Science*, *7(1)*, 29-
1271
1272 522 36.
- 1273
1274 523 Lobón, S., Sanz, A., Blanco, M., Ripoll, G., & Joy, M. (2017). The type of forage and
1275
1276 524 condensed tannins in dams' diet: Influence on meat shelf life of their suckling lambs. *Small*
1277
1278 525 *Ruminant Research*, *154*, 115–122
- 1279
1280
1281 526 Lodge, J. K., Hall, W. L., Jeanes, Y. M., & Proteggente, A. R. (2004). Physiological factors
1282
1283 527 influencing vitamin E biokinetics. *Annals of the New York Academy of Sciences*, *1031(1)*,
1284
1285 528 60-73.
- 1286
1287 529 López-Andrés, P., Luciano, G., Vasta, V., Gibson, T. M., Biondi, L., Priolo, A., & Mueller-
1288
1289 530 Harvey, I. (2013). Dietary quebracho tannins are not absorbed, but increase the antioxidant
1290
1291 531 capacity of liver and plasma in sheep. *British Journal of Nutrition*, *110(4)*, 632-639.
- 1292
1293
1294
1295
1296
1297
1298

- 1299
1300
1301 532 Luciano, G., Natalello, A., Mattioli, S., Pauselli, M., Sebastiani, B., Niderkorn, V., Copani, G.,
1302
1303 533 Benhissi, H., Amanpour, A., & Valenti, B. (2019). Feeding lambs with silage mixtures of
1304
1305 534 grass, sainfoin and red clover improves meat oxidative stability under high oxidative
1306
1307 535 challenge. *Meat Science*, 156, 59-67.
- 1309
1310 536 Luciano, G., Roscini, V., Mattioli, S., Ruggeri, S., Gravador, R. S., Natalello, A., Lanza, M.,
1311
1312 537 De Angelis, A., & Priolo, A. (2017). Vitamin E is the major contributor to the antioxidant
1313
1314 538 capacity in lambs fed whole dried citrus pulp. *Animal*, 11(3), 411–417.
- 1316 539 Mancini, R. A., & Hunt, M. (2005). Current research in meat color. *Meat Science*, 71(1), 100-
1317
1318 540 121.
- 1320 541 Mphahlele, R. R., Fawole, O. A., Mokwena, L. M., & Opara, U. L. (2016). Effect of extraction
1321
1322 542 method on chemical, volatile composition and antioxidant properties of pomegranate
1323
1324 543 juice. *South African Journal of Botany*, 103, 135-144.
- 1326 544 Natalello, A., Luciano, G., Morbidini, L., Valenti, B., Pauselli, M., Frutos, P., Biondi, L.,
1327
1328 545 Rufino-Moya, P. J., Lanza, M., & Priolo, A. (2019). Effect of feeding pomegranate by-
1329
1330 546 product on fatty acid composition of ruminal digesta, liver and muscle in lambs. *Journal*
1331
1332 547 *of Agricultural and Food Chemistry*, 67, 4472-4482.
- 1333 548 Natalello, A., Hervás, G., Toral, P.G., Luciano, G., Valenti, B., Mendoza, A.G., Pauselli, M.,
1334
1335 549 Priolo, A., & Frutos, P. (2020). Bioactive compounds from pomegranate by-products
1336
1337 550 increase the in vitro ruminal accumulation of potentially health promoting fatty acids.
1338
1339 551 *Animal Feed Science and Technology*, 259, 114355
- 1342 552 Orak, H. H., Yagar, H., & Isbilir, S. S. (2012). Comparison of antioxidant activities of juice,
1343
1344 553 peel, and seed of pomegranate (*Punica granatum* L.) and inter-relationships with total
1345
1346 554 phenolic, tannin, anthocyanin, and flavonoid contents. *Food Science and*
1347
1348 555 *Biotechnology*, 21(2), 373-387.
- 1350
1351
1352
1353
1354
1355
1356
1357

1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416

556 Ortuño, J., Serrano, R., & Bañón, S. (2015). Antioxidant and antimicrobial effects of dietary
557 supplementation with rosemary diterpenes (carnosic acid and carnosol) vs. vitamin E on
558 lamb meat packed under protective atmosphere. *Meat Science*, *110*, 62–69

559 Pande, G., & Akoh, C. C. (2009). Antioxidant capacity and lipid characterization of six
560 Georgia-grown pomegranate cultivars. *Journal of agricultural and food chemistry*, *57*(20),
561 9427-9436.

562 Ponnampalam, E.N., Norn, S., Burnett, V.F., Dunshea, F.R., Jacobs, J.L., & Hopkins, D.L.
563 (2014). The synergism of biochemical components controlling lipid oxidation in lamb
564 muscle. *Lipids*, *49*, 757–766

565 Priolo, A., Bella, M., Lanza, M., Galofaro, V., Biondi, L., Barbagallo, D., Ben Salem, H., &
566 Pennisi, P. (2005). Carcass and meat quality of lambs fed fresh sulla (*Hedysarum*
567 *coronarium* L.) with or without polyethylene glycol or concentrate. *Small Ruminant*
568 *Research*, *59*, 281–288

569 Priolo, A., Ben Salem, H., Atti, N., & Nefzaoui, A. (2002). Polyethylene glycol in concentrate
570 or feedblocks to deactivate condensed tannins in *Acacia cyanophylla* Lindl. Foliage 2.
571 Effects on meat quality of Barbarine lambs. *Animal Science*, *75*, 137-140.

572 Priolo, A., Waghorn, G. C., Lanza, M., Biondi, L., & Pennisi, P. (2000). Polyethylene glycol
573 as a means for reducing the impact of condensed tannins in carob pulp: Effects on lamb
574 growth performance and meat quality. *Journal of Animal Science*, *78*, 810–816.

575 Prior, R. L., Hoang, H., Gu, L., Wu, X., Bacchiocca, M., Howard, L., Hampsch-Woodill, M.,
576 Huang, D., Ou, B., & Jacob, R. (2003). Assays for hydrophilic and lipophilic antioxidant
577 capacity (oxygen radical absorbance capacity (ORAC_{FL})) of plasma and other biological
578 and food samples. *Journal of Agricultural and Food Chemistry*, *51*(11), 3273–3279.

- 1417
1418
1419 579 Salami, S. A., Luciano, G., O'Grady, M. N., Biondi, L., Newbold, C. J., Kerry, J. P., & Priolo,
1420
1421 A. (2019). Sustainability of feeding plant by-products: a review of the implications for
1422 580
1423 ruminant meat production. *Animal Feed Science and Technology*, 251, 37-55
1424 581
1425
1426 582 Sales, J., & Koukolová, V. (2011). Dietary vitamin E and lipid and color stability of beef and
1427
1428 583 pork: Modeling of relationships. *Journal of Animal Science*, 89, 2836–2848
1429
1430 584 Seeram, N., Lee, R., Hardy, M., & Heber, D. (2005). Rapid large scale purification of
1431
1432 585 ellagitannins from pomegranate husk, a by-product of the commercial juice
1433
1434 586 industry. *Separation and Purification Technology*, 41(1), 49-55.
1435
1436 587 Shabtay, A., Eitam, H., Tadmor, Y., Orlov, A., Meir, A., Weinberg, P., Weinberg Z. G., Chen,
1437
1438 588 Y., Brosh, A., Izhaki, I., & Kerem, Z. (2008). Nutritive and antioxidative potential of fresh
1439
1440 589 and stored pomegranate industrial byproduct as a novel beef cattle feed. *Journal of*
1441
1442 590 *Agricultural and Food Chemistry*, 56(21), 10063-10070.
1443
1444
1445 591 Valenti, B., Luciano, G., Morbidini, L., Rossetti, U., Codini, M., Avondo, M., Priolo, A., Bella,
1446
1447 592 M., Natalello, A., & Pauselli, M. (2019). Dietary pomegranate pulp: effect on ewe milk
1448
1449 593 quality during late lactation. *Animals*, 9(5), 283.
1450
1451 594 Valenti, B., Luciano, G., Pauselli, M., Mattioli, S., Biondi, L., Priolo, A., Natalello, A.,
1452
1453 595 Morbidini, L., & Lanza, M. (2018). Dried tomato pomace supplementation to reduce lamb
1454
1455 596 concentrate intake: Effects on growth performance and meat quality. *Meat Science*, 145,
1456
1457 597 63-70.
1458
1459
1460 598 Valenti, B., Natalello, A., Vasta, V., Campidonico, L., Roscini, V., Mattioli, S., Pauselli, M.,
1461
1462 599 Priolo, A., Lanza, M., & Luciano, G. (2019). Effect of different dietary tannin extracts on
1463
1464 600 lamb growth performances and meat oxidative stability: comparison between mimosa,
1465
1466 601 chestnut and tara. *Animal*, 13(2), 435–443.
1467
1468
1469
1470
1471
1472
1473
1474
1475

1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534

602 Van Soest, P. J., Robertson, J. B., & Lewis, B. A. (1991). Methods for dietary fiber, neutral
603 detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of*
604 *Dairy Science*, 74(10), 3583–3597.

605 Vasta, V., & Luciano, G. (2011). The effects of dietary consumption of plants secondary
606 compounds on small ruminants' products quality. *Small Ruminant Research*, 101(1-3),
607 150-159.

608 Viuda-Martos, M., Fernández-López, J., & Pérez-Álvarez, J. A. (2010). Pomegranate and its
609 many functional components as related to human health: a review. *Comprehensive*
610 *Reviews in Food Science and Food Safety*, 9(6), 635-654.

Table 1.

Ingredients, chemical composition and antioxidant capacity of the experimental diets and whole pomegranate by-product.

	Whole pomegranate by-product	Experimental diet ¹	
		CON	WPB
<i>Ingredients, g/100 g dry matter (DM)</i>			
Corn		22.6	11.6
Barley		22.6	11.6
Alfalfa hay		19.8	19.8
Wheat bran		20.0	20.0
Soybean meal		12.0	14.0
Whole pomegranate by-product		-	20.0
Molasses		0.9	0.9
Mineral premix ²		2.1	2.1
<i>Chemical composition, g/100 g DM</i>			
DM, g/100 as fed	90.0	88.7	89.2
Crude Protein	6.52	17.6	17.8
NDF ³	28.8	23.3	26.3
ADF ³	20.7	12.9	15.5
ADL ³	5.52	2.98	2.70
Ash	3.52	5.87	4.40
Crude Fat	3.99	2.11	2.51
<i>Phenolic compounds, g/100g DM</i>			
Total phenols ⁴	9.51	0.30	1.89
Total tannins ⁴	9.34	0.14	1.70
Condensed tannins	0.80	0.10	0.20
<i>Tocopherols, mg/kg DM</i>			
γ -Tocopherol	11.1	0.74	2.04
α -Tocopherol	48.3	7.82	16.8
<i>Antioxidant capacity (ORAC), μmol TE/g DM⁵</i>			
Hydrophilic fraction	684	103	342
Lipophilic fraction	27.3	21.1	31.1

¹ CON: control barley-corn based concentrate diet. WPB: diet including 20% of whole pomegranate by-product.

² Containing: 25% calcium carbonate, 25% sodium bicarbonate, 25% bicalcic phosphate and 25% sodium chloride.

³ NDF: neutral detergent fibre; ADF: acid detergent fibre; ADL: acid detergent lignin (ADL).

⁴ Expressed as tannic acid equivalents.

⁵ ORAC: oxygen radical absorbance capacity; TE: trolox equivalents.

Table 2.

Effect of the dietary treatment on lamb performances and intakes.

	Dietary treatment ¹		SEM ²	P-value
	CON	WPB		
<i>Performances</i>				
Final body weight, kg	23.6	23.1	0.524	0.637
Carcass weight, kg	11.1	10.2	0.330	0.179
DMI ³ , g/day	821	882	19.50	0.125
ADG ³ , g/day	234	235	7.200	0.921
FCR ³ , g DMI/g ADG	3.56	3.79	0.101	0.278
<i>Intakes of phenolic compounds g/day</i>				
Total phenols	2.49	16.7	1.700	<0.001
Total tannins	1.16	15.0	1.660	<0.001
Condensed tannins	0.08	0.18	0.012	<0.001
<i>Intakes of tocopherols, mg/day</i>				
γ-Tocopherol	0.61	1.80	1.444	<0.001
α-Tocopherol	6.42	14.8	1.300	<0.001

¹ CON: control barley-corn based concentrate diet. WPB: diet including 20% of whole pomegranate by-product.

² SEM: standard error of the mean.

³ DMI: dry matter intake; ADG: average daily gain; FCR: feed conversion ratio.

Table 3.

Effect of the dietary treatment on the muscle pH, myoglobin, antioxidant vitamins and oxidisable fatty acids in the intramuscular fat.

	Dietary treatment ¹		SEM ²	P-value
	CON	WPB		
pH	5.72	5.78	0.036	0.427
Myoglobin, mg/g of muscle	2.68	2.46	0.093	0.259
Intramuscular fat, g/100 g muscle	1.88	2.01	0.156	0.690
<i>Tocopherols, ng/g of muscle</i>				
γ-Tocopherol	13.0	32.0	2.610	<0.001
α-Tocopherol	162	309	22.00	<0.001
<i>Fatty acids classes and oxidizable fatty acids, mg/g of muscle</i>				
Saturated	6.39	6.27	0.637	0.928
Monounsaturated	6.62	6.15	0.566	0.702
Polyunsaturated (PUFA)	1.30	1.80	0.121	0.041
PUFA <i>n</i> -3	0.13	0.16	0.011	0.091
PUFA <i>n</i> -6	1.08	1.34	0.086	0.137
HP-PUFA ³	0.39	0.57	0.041	0.024
Peroxidability index ⁴	2.07	2.85	0.191	0.042

¹ CON: control barley-corn based concentrate diet. WPB: diet including 20% of whole pomegranate by-product.

² SEM: standard error of the mean.

³ Highly peroxidizable-PUFA, calculated as the sum of PUFA with three or more unsaturated bonds.

⁴ Calculated as: peroxidability index = (\sum dienoic \times 1) + (\sum trienoic \times 2) + (\sum tetraenoic \times 3) + (\sum pentaenoic \times 4) + (\sum hexaenoic \times 5).

Table 4.

Effect of the dietary treatment and time of storage on the oxidative stability parameters of meat.

	Dietary treatment (D) ¹		Time of storage (T) ²			SEM ³	P-values ⁴		
	CON	WPB	0	1	2		D	T	D × T
<i>Colour descriptors and metmyoglobin % of raw meat</i>									
L* (lightness)	46.99	49.02	48.57	48.65	47.48	0.296	0.005	0.068	0.533
a* (redness)	11.56	11.60	12.24 ^a	11.58 ^{ab}	10.93 ^b	0.166	0.929	<0.001	0.181
b* (yellowness)	11.21	11.39	10.15 ^b	11.87 ^a	11.93 ^a	0.194	0.663	<0.001	0.159
C (saturation)	16.14	16.29	15.91	16.59	16.19	0.214	0.789	0.474	0.168
h _{ab} (hue angle)	44.08	44.29	39.55 ^c	45.61 ^b	47.45 ^a	0.494	0.620	<0.001	0.087
MetMb, % of Mb	47.60	46.03	39.52 ^c	49.16 ^b	51.25 ^a	0.762	0.050	<0.001	0.498
<i>Lipid oxidation (TBARS values), mg/kg meat</i>									
Raw meat	0.95	0.56	0.25 ^b	0.80 ^a	1.08 ^a	0.080	0.024	<0.001	0.013
Cooked meat	3.74	3.07	1.69 ^c	3.60 ^b	4.70 ^a	0.192	0.006	<0.001	0.266

¹ CON: control barley-corn based concentrate diet. WPB: diet including 20% of whole pomegranate by-product.² Time of storage 0, 1, 2 correspond to: days 0, 4, 7 (raw meat); days 0, 2, 4 (cooked meat)³ SEM: standard error of the mean.⁴ P-values for the effects of the dietary treatment, time of storage and of the Diet × Time interaction^{a, b, c} Within row, different superscript letter indicates differences ($P < 0.05$) between times of storage tested using the Tukey's Honest Significant Difference test.

1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811

Fig 1. Effect of the dietary treatment (CON: control barley-corn based concentrate diet. WPB: diet including 20% of whole pomegranate by-product) on raw meat antioxidant capacity of (a) hydrophilic fraction and (b) lipophilic fraction. Values presented are the estimated least squares means and standard error bars.

1812
1813
1814
1815 **Fig 2.** Effect of the dietary treatment (CON: control barley-corn based concentrate diet.
1816 WPB: diet including 20% of whole pomegranate by-product) and time of storage (days 0, 4
1817 and 7) on TBARS (**thiobarbituric acid reactive substances**) values of raw meat over aerobic
1818 storage at 4 °C. Values presented are the estimated least squares means and standard error
1819 bars.

1820 **MDA: Malondialdehyde**

1821 ^{a,b,c}Indicate differences between mean values ($P < 0.05$) tested using the Tukey's Honest
1822 Significant Difference test.



