

**Manuscript version: Author's Accepted Manuscript**

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

**Persistent WRAP URL:**

<http://wrap.warwick.ac.uk/131253>

**How to cite:**

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk).

## Turnip yellows virus in winter oilseed rape

**The incidence of *Turnip yellows virus* in oilseed rape crops (*Brassica napus* L.) in three different regions of England over three consecutive growing seasons and the relationship with the abundance of flying *Myzus persicae*.**

Elvis Asare-Bediako<sup>1</sup>, Julie E. Jones<sup>1</sup>, Angela Hambidge<sup>1</sup>, Mark Stevens<sup>2</sup>, Andrew Mead<sup>3</sup>, Carol E. Jenner<sup>1</sup> and John A. Walsh\*<sup>1</sup>

<sup>1</sup> School of Life Sciences, University of Warwick, Wellesbourne, Warwick, CV35 9EF, UK

<sup>2</sup> British Beet Research Organisation, Innovation Centre, Norwich Research Park, Norwich, NR4 7GJ, UK

<sup>3</sup> Rothamsted Research, Harpenden, AL5 2JQ, UK

\* [john.walsh@warwick.ac.uk](mailto:john.walsh@warwick.ac.uk)

## Turnip yellows virus in winter oilseed rape

**Abstract**

Turnip yellows virus (TuYV) is the most important virus infecting oilseed rape in the UK. The incidence and spatial distribution of TuYV in winter oilseed rape (WOSR) crops in three regions of England was determined over three growing seasons. Leaf samples were collected from three fields in each region, in autumn (November-December) and spring (April) of the three crop seasons and tested for virus presence by enzyme-linked immunosorbent assay. Infection was detected in all fields except one. Higher TuYV incidences were recorded in the 2007-8 ( $\leq 89\%$ ) and 2009-10 ( $\leq 100\%$ ) crop seasons than in 2008-9 ( $\leq 24\%$ ). Highest incidences were recorded in Lincolnshire ( $\leq 100\%$ ), followed by Warwickshire ( $\leq 88\%$ ), with lowest incidences in Yorkshire (1-74%). There was a significant increase in incidence detected between autumn and spring sampling in eight fields, a significant decrease in one field and no significant change in 18 fields. Rothamsted Insect Survey suction trap data for the aphid *Myzus persicae* in Lincolnshire, Warwickshire and Yorkshire revealed two peaks of flight activity in most years (2007-2009). The second peak (September-November) coincided with emergence of WOSR. The highest cumulative (August-November) trap catches in the three regions during the three crop seasons occurred in Lincolnshire and the lowest in Yorkshire; catches in autumn 2009 were highest and lowest in autumn 2008. Regression analysis revealed a highly significant association between the cumulative numbers of *M. persicae* caught in the suction traps closest to the crops between August and November each year and the incidence of TuYV detected in the WOSR crops in the autumn of each year. Results are discussed in the light of factors affecting the spread of TuYV and future possibilities for control.

**Keywords:** *Brassica napus* (oilseed rape), Turnip yellows virus incidence, aphids, *Myzus persicae* abundance

## 1 INTRODUCTION

*Turnip yellows virus* (syn. *Beet western yellows virus*), a species in the *Polyomavirus* genus of the *Luteoviridae* family is the most common virus infecting oilseed rape in the UK (Hardwick et al., 1994; Stevens et al., 2008). It has a wide host range and infects more than 150 species in 23 dicotyledonous families, including economically important crops such as lettuce, spinach, field beans, radish, cabbage and oilseed rape (Graichen, 1996).

TuYV infection of oilseed rape crops in England was first reported in 1980 (Gilligan et al., 1980) but the widespread incidence of the virus in the UK was first reported by (Smith & Hinckes, 1985). Results from a UK survey in 1983, covering 80 autumn-sown oilseed rape crops from Aberdeenshire to Essex showed that 97% were extensively infected with TuYV (Smith & Hinckes, 1985). Varying incidences of TuYV infection in oilseed rape crops have since been reported in the UK, ranging from less than 10% to 85% (Hardwick et al., 1994; Hill et al., 1989; Jay et al., 1999; Walsh et al., 1989). A survey conducted in 2009 covering oilseed rape crops on 80 farms from the south coast of England to Scotland showed TuYV infection of up to 70% (Clark & Stevens, 2009). Infection of oilseed rape crops by TuYV has also been reported in other countries including Germany (Schröder, 1994), France (Kerlan, 1991), the Czech Republic (Polak & Majkova, 1992), Austria (Graichen et al., 2000), Iran (Shahraeen et al., 2003), Serbia (Jasnic & Bagi, 2007; Milošević et al. 2015) and Australia (Coutts et al., 2006). The virus has also been reported in lettuce in the UK (Walkey & Pink, 1990) and faba bean in Egypt and Morocco (Abraham et al., 2008).

## Turnip yellows virus in winter oilseed rape

1  
2  
3  
4 TuYV, like other members of the *Luteoviridae* is mostly restricted to the phloem tissue of host  
5  
6 plants and is transmitted by aphid vectors from plant to plant in a persistent (acquisition times  
7  
8 range from 30 min to hours, the vector cannot immediately transmit the virus and retains its  
9  
10 ability to transmit through the moult and often for the remainder of its life), circulative (passes  
11  
12 from the insect gut to the salivary glands via the haemolymph for transmission) and non-  
13  
14 propagative (does not replicate in vector tissues) manner. The green peach aphid, *Myzus persicae*  
15  
16 (Sulzer) is the principal vector in Europe (Schliephake et al., 2000) with a transmission efficiency  
17  
18 of about 90% (Schliephake et al., 2000). Enzyme-linked immunosorbent assay (ELISA) on  
19  
20 samples of *M. persicae* caught in yellow water traps and Rothamsted Insect Survey Suction traps  
21  
22 have shown that 2-72% of *M. persicae* were carrying TuYV (Stevens et al., 2008).  
23  
24  
25  
26  
27  
28

29 TuYV infection is thought to be one of the major reasons why oilseed rape crops do not attain  
30  
31 their potential yield in England (Stevens et al., 2008), estimated at 6.5 t/ha (Berry & Spink,  
32  
33 2006), compared to the current yield of 3.5 t/ha (Defra, 2018). TuYV infection seriously affects  
34  
35 all components of yield, including the number of pods per plant, number of seeds per pod, and  
36  
37 the oil content per seed. Plants infected with TuYV also have a reduced leaf area per plant and  
38  
39 produce fewer primary branches (Jay et al., 1999). The effect of TuYV on the yield of oilseed  
40  
41 rape depends on the incidence of virus infection and the crop variety (Walsh et al., 1989). One  
42  
43 estimate of yield losses due to TuYV infection is up to 30% (Home-Grown Cereals Authority,  
44  
45 2009). Experimental plots of oilseed rape with 100% TuYV- infection yielded approximately  
46  
47 10% less seed and 13.4% less oil than plots with 18% virus infection (Smith and Hinckes, 1985).  
48  
49 Plot experiments in Australia on spring oilseed rape, showed that a site with 96% infection  
50  
51 suffered a yield loss of up to 46% (Jones et al., 2007) and in Germany, plots of winter oilseed  
52  
53 rape with 90-100% TuYV infection yielded between 12% and 34% lower than plots that were  
54  
55  
56  
57  
58  
59  
60

## Turnip yellows virus in winter oilseed rape

1  
2  
3  
4 almost virus-free (Graichen & Schliephake, 1999). When oilseed rape plants were co-infected  
5  
6 with a mix of TuYV, cauliflower mosaic virus (CaMV) and turnip mosaic virus (TuMV), the  
7  
8 yields of plants with severe virus symptoms were reduced by an estimated 70-79% (Hardwick et  
9  
10 al., 1994).

11  
12  
13  
14  
15 Effective control of TuYV in oilseed rape is necessary to improve yields. However, for the most  
16  
17 part, farmers growing oilseed rape in the UK seem not to be aware of the presence, let alone the  
18  
19 incidence of TuYV in their crops. For this reason and as an important prerequisite for developing  
20  
21 effective approaches to controlling the disease it is important to investigate the incidence of  
22  
23 TuYV infection of oilseed rape crops at various locations, in different years and the type of  
24  
25 disease pattern and spread in the fields. The quantitative relationship between the flight activities  
26  
27 of the aphid vectors is also important for the timing and development of control strategies.  
28  
29  
30  
31

32  
33 Here the incidence and distribution of TuYV in fields of winter oilseed rape in three regions of  
34  
35 England in the autumn and spring of three consecutive crop seasons are described. It has been  
36  
37 said that the spread of brassica viruses is related to the abundance and movement of aphid vectors  
38  
39 (Walsh & Tomlinson, 1985). In Germany, high levels of TuYV infection were detected in winter  
40  
41 oilseed rape crops during 1995-96 following high levels of flight activity of aphids during the  
42  
43 autumn of 1995 (Graichen & Schliephake, 1999). We have investigated these relationships in the  
44  
45 context of *M. persicae* (said to be the most important vector of TuYV in Europe) and quantified  
46  
47 them by relating the number of flying *M. persicae* caught in the Rothamsted Insect Survey  
48  
49 suction traps located nearest to sampling sites to the incidences of TuYV at the sampling sites in  
50  
51 the three regions, over the three consecutive growing seasons.  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Turnip yellows virus in winter oilseed rape

**2 MATERIALS AND METHODS****2.1 Surveys of winter oilseed rape crops**

Using a line transect sampling method (Buckland et al., 2001), upper leaves from 100 plants were sampled from three crop fields in each of three oilseed rape-growing regions of England, Eastern (Lincolnshire), Northern (Yorkshire) and Midlands (Warwickshire) in autumn (November-December; 2-6 true leaves) and the following spring (April; starting to flower) during the 2007-8, 2008-9 and 2009-10 crop seasons (27 fields in all). Aphids were very rarely seen on the sampled plants, consequently no records were taken. Samples from each plant were placed in separate polythene bags to register sampling positions within fields. The Lincolnshire fields were near Long Sutton, the Warwickshire fields were near Gaydon, and the Yorkshire sites were near Allerton, Little Ouseburn, Aberford, Green Hammerton and Whixley. The locations of the fields were determined using a Global Positioning System (Garmin E-Trex HGPS Receiver, Garmin Corporation Olathe, KS, USA) (Supporting information, Table S1).

The sampling procedure involved estimating the length and breadth of each field in order to divide the field into ten equally-spaced transects with ten equally-spaced-samples collected per transect. Where fields were exceptionally large, only a proportion of the fields were sampled. The mean field size was 20.8 hectares (range from 4.3 to 40.0 ha).

**2.2 Sampling sites**

Farms sampled rotate oilseed rape crops every 3-6 years with crops such as wheat, potato, sugar beet, peas and may also practice land fallowing. Winter oilseed rape is normally sown in England between late August and early September, overwinters, flowers in the spring (April-May) and is harvested in July/August. Seed sowing in field three in Warwickshire during the 2008-9 crop

1  
2  
3  
4  
5  
6 season was delayed by the prevailing weather conditions until late September. Where the  
7  
8 information was available, the cultivars in the fields sampled are given in Table S1.  
9

### 10 11 12 13 **2.3 Detection of TuYV in the samples**

14  
15 The presence of TuYV was tested in all sampled plants by standard triple antibody sandwich  
16  
17 enzyme-linked immunosorbent assay (TAS-ELISA) using paired wells in microtitre plates (96-  
18  
19 well Nunc Maxisorp; Nunc, Roskilde, Denmark) as described by Hunter et al. (2002).

20  
21 Absorbance values ( $A_{405\text{nm}}$ ) were measured with a Biochrom Anthos 2010 microplate reader  
22  
23 (Biochrom Ltd., Cambridge, U.K.). Values for 10 uninfected leaf samples were also measured. A  
24  
25 sample was deemed to be positive when the absorbance was greater than the mean absorbance of  
26  
27 ten healthy samples on each ELISA plate, plus  $2.262 \times$  standard deviation of the mean of the ten  
28  
29 healthy samples (where  $2.262 =$  Inverse of Student's  $t$ -distribution at 5% probability level with 9  
30  
31 degrees of freedom).  
32  
33  
34  
35  
36  
37

### 38 **2.4 Data analysis**

39  
40 TuYV incidence data were analysed using a generalised linear model (GLM) (Nelder &  
41  
42 Wedderburn, 1972). Differences between county and crop season means and their interactions  
43  
44 were compared using the approximate least significant difference (LSD) calculated from the  
45  
46 analyses.  
47  
48  
49

50  
51 A two-sample binomial test (Armitage & Berry, 1994) was used to determine whether there was  
52  
53 a significant change in TuYV incidences in each field between autumn and spring. The overall  
54  
55 correlation between autumn and spring incidences of TuYV infections was assessed using the  
56  
57



## Turnip yellows virus in winter oilseed rape

Pearson correlation coefficient. All statistical analyses were carried out using GenStat (GenStat Release version 12.1) (Payne et al., 2009).

The cumulative numbers of *M. persicae* caught monthly in the Rothamsted Insect Survey suction traps closest to the sampling sites (Kirton in Lincolnshire [22-26 km from fields sampled], Askham Bryan in Yorkshire [13-17 km from fields sampled] and Wellesbourne in Warwickshire [10-11 km from fields sampled]) were determined from the weekly suction trap aphid catches obtained between August (crop sowing) and November for each year. A non-linear regression analysis fitting an exponential function was used to explore the relationship between mean TuYV incidences in autumn and the cumulative aphid counts.

A Black-White (BW) join-count statistic (Cliff & Ord, 1969) was calculated using GenStat to assess spatial autocorrelation within each field for each sampling occasion, comparing a null hypothesis of random distribution of infected samples with an alternative hypothesis of spatial clustering. The statistic measures the number of neighbour pairs containing both an infected and a healthy plant, where neighbours were defined here to be adjacent samples in vertical or horizontal directions (each non-edge sample has four neighbours, i.e. “rook case”). The significance of the join-count statistic is achieved by computing a standard normal deviate, called a Z-score which is given by the formula:

$$Z(BW) = \frac{\text{Observed (BW)} - \text{Expected (BW)}}{\sigma_{BW}}$$

where  $\sigma_{BW}$  is the standard deviation for BW joins. The expected number of BW neighbours is calculated based on the overall proportion of infected plants and represents the likely pattern under a random distribution. As the aim was to detect evidence of clustering, a one-sided test for

Turnip yellows virus in winter oilseed rape  
negative values of the BW join-count statistic was appropriate, negative values indicating a  
positive spatial autocorrelation between infected plants and probability levels of  $P \leq 0.05$   
indicating significant spatial autocorrelation.

### 3 RESULTS

#### 3.1 Incidence of TuYV infection in winter oilseed rape crops

The overall mean TuYV incidences recorded in the autumns of 2007 ( $36.67 \pm 4.56\%$ ) and 2009 ( $48.67 \pm 3.62\%$ ) were not significantly different from each other but were significantly higher ( $P < 0.001$ ) than recorded in the autumn of 2008 ( $6.11 \pm 2.33\%$ ) (Table 1). Highly significant differences ( $P < 0.001$ ) in the mean percentage autumn virus incidences were found between the counties (Table 1). Lincolnshire had the highest mean TuYV incidence ( $55.00 \pm 3.69\%$ ), followed by Warwickshire ( $23.78 \pm 3.94\%$ ) whilst Yorkshire had the lowest ( $12.67 \pm 3.18\%$ ) (Table 1; Fig 2b). The interaction effects between the counties and crop seasons for autumn virus incidences were significant ( $P < 0.05$ ). The highest mean incidence ( $94.00 \pm 4.08\%$ ) was recorded in Lincolnshire during the autumn of 2009, whilst the lowest ( $2.33 \pm 2.60\%$ ) was recorded in Yorkshire during the autumn of 2008.

The highest overall mean TuYV incidence in spring was recorded in the 2007-8 crop ( $55.67 \pm 6.91\%$ ), followed by the 2009-10 crop ( $53.78 \pm 5.86\%$ ); the 2008-9 crop had the lowest ( $8.67 \pm 4.11\%$ ) (Table 2). Lincolnshire had the highest overall mean spring incidence ( $56.67 \pm 5.17\%$ ), followed by Warwickshire ( $42.00 \pm 6.43\%$ ) and Yorkshire had the lowest ( $19.44 \pm 5.55\%$ ). The GLM analysis did not indicate a significant interaction between the regions and crop seasons for the spring incidences of TuYV ( $P = 0.163$ ). However, the highest mean incidence ( $94.00 \pm$

1 Turnip yellows virus in winter oilseed rape  
2 6.04%) was recorded in Lincolnshire in spring of the 2009-10 crop, whilst the lowest mean  
3  
4 incidence ( $3.67 \pm 4.79\%$ ) was recorded in Yorkshire during 2008-9 crop season (Table 2).  
5  
6  
7

8 The comparisons of autumn and spring incidences of TuYV in the individual 27 fields in the  
9  
10 three regions over the three crop seasons using the two-sample binomial test are shown in Table  
11  
12 S1. Large differences in the incidences of TuYV in the various fields, counties and crop seasons  
13  
14 surveyed, ranging from 0% (recorded in autumn and spring in Warwickshire in 2008-9) to 100%  
15  
16 (recorded in Lincolnshire in the autumn of 2009) were found. There were significant ( $P < 0.05$ )  
17  
18 changes in the proportions of plants infected with TuYV between autumn and spring in five of  
19  
20 the nine fields in 2007-8 (all 3 fields in Warwickshire and one each in Lincolnshire and  
21  
22 Yorkshire), three fields in 2008-9 (two in Lincolnshire and one in Warwickshire) and only one  
23  
24 field in 2009-10 (Yorkshire) (Table S1). With the exception of one of the fields in Lincolnshire in  
25  
26 2008-9, all significant changes involved an increase in TuYV incidence. Overall, there was a  
27  
28 significantly high correlation ( $r = 0.89$ ,  $P < 0.001$ , d.f. = 25) between autumn and spring  
29  
30 incidences of TuYV in the oilseed rape crops surveyed.  
31  
32  
33  
34  
35  
36  
37

### 38 **3.2 Within field virus distribution**

39 The results of the analysis of spatial distribution of TuYV-infected plants are given in Table S1.  
40  
41 Where possible, Black-White join-count statistics were calculated for each of the 27 fields  
42  
43 surveyed in the three-crop seasons and then tested as standard normal deviates ( $Z$ -scores). Most  
44  
45 of the fields (17 of the 25 analysed) showed positive, but non-significant spatial autocorrelation  
46  
47 (negative  $Z$ -scores,  $P > 0.05$ ) when sampled in autumn, indicating that most of the infected plants  
48  
49 showed a slightly aggregated pattern of distribution. For a one-sided test at a probability of 0.05,  
50  
51 values less than -1.645 (large negative  $Z$ -score) indicated that the number of observed join-counts  
52  
53  
54  
55  
56  
57  
58  
59  
60

Turnip yellows virus in winter oilseed rape was significantly less than expected, an indication of clustering of the infected plants. Lincolnshire field 1 was the only field showing significant positive autocorrelation ( $Z = -1.922$ ,  $P < 0.05$ ) between the infected plants in autumn (2007), indicating significant clustering (Fig. 1a). Spring sampling revealed a slightly more random pattern with half of the fields (13 of the 26 analysed) showing negative spatial autocorrelation (positive Z-scores) (e.g. Warwickshire field 2 in spring 2010,  $Z = 0.561$ ,  $P = 0.288$ , Fig. 1b). Warwickshire field 1 showed a significant positive autocorrelation ( $Z = -2.177$ ,  $P < 0.05$ ) between the infected plants in spring 2008, indicating clustering.

### 3.3 Numbers of *Myzus persicae* caught in Rothamsted Insect Survey suction traps

Rothamsted Insect Survey suction trap catches showed two peaks of flight activity of *M. persicae* in Lincolnshire, Warwickshire and Yorkshire in most years (2007-2009) (Fig. 2a). The first peak occurred between June and July and the second occurred between September and November in each year, the latter coinciding with the emergence of oilseed rape crops. Crops were sown in August or September. The highest cumulative (August to November) trap catches of *M. persicae* during the three crop seasons occurred in Lincolnshire and the lowest in Yorkshire; catches in autumn 2009 were highest and those in autumn 2008 were lowest.

### 3.4 Relationship between flying *Myzus persicae* and TuYV incidence

There was a clear relationship between TuYV incidence and aphid numbers (Fig. 2). Regression analysis revealed a highly significant association between the numbers of *M. persicae* caught in the Rothamsted Insect Survey suction traps closest to the crops between August and November each year and the incidence of TuYV in the oilseed rape crops (d.f. = 2, 8;  $F = 24.2$ ;  $P < 0.001$ ) in autumn of each year (Fig. 3).

## Turnip yellows virus in winter oilseed rape

**4 DISCUSSION**

In this study, TuYV was prevalent in winter oilseed rape fields in three regions of England over a three year period with incidences within individual fields ranging from 0-100%. The virus was detected in 26 of the 27 oilseed rape fields sampled from three regions (Lincolnshire, Warwickshire and Yorkshire). This finding corroborates the previous reports of widespread occurrence of TuYV in oilseed rape crops in the UK (Smith and Hinckes 1985; Walsh et al., 1989; Hill et al., 1989; Hardwick et al., 1994; Jay et al., 1999; Stevens et al., 2008) where incidences of 0-100% were also reported. The only field where no infection was recorded (Warwickshire field three in the 2008-9 crop season) was sown late (27<sup>th</sup> September, 2008) relative to the other two Warwickshire fields (sown 26-27<sup>th</sup> August, 2008) and the very few plants that had emerged were very small when sampled on 15<sup>th</sup> December, 2008. It is clear from Fig. 2a that aphid numbers in Warwickshire were very low by late September 2008 and this, combined with the late planting date and the sparsity of plants, is very likely to be the reason for the lack of infection. We have not done any detailed assessments of TuYV incidence in winter oilseed rape in the UK since 2010, however, the limited testing of commercial oilseed rape crops that we have done, has revealed incidences of 0-100% in the 2018-19 crop, as for the 2007-2010 study reported here. That said, many crops tested early in 2019 had incidences of 100%, reflecting the very high numbers of *M. persicae* caught in Rothamsted Insect Survey suction traps between August and November 2018.

The analysis of the autumn incidences of TuYV in the oilseed rape crops in the different regions sampled were clearly associated with the cumulative numbers of *M. persicae* caught in the suction traps in these regions between August and November. This indicates that the significantly higher incidences of TuYV recorded in the 2007-8 and 2009-10 crops compared with those in the 2008-9 crops were due to the increased flight activity of *M. persicae* between August and

Turnip yellows virus in winter oilseed rape  
November in 2007 and 2009, relative to 2008 (Fig. 2). Graichen and Schliephake (1999) also  
demonstrated that a high incidence of TuYV in winter oilseed rape appeared to be closely related  
to the flight activity of *M. persicae* vectors in Germany. Clark and Stevens (2011) indicated a  
close correlation between aphid numbers caught in the autumn and the amount of virus present in  
oilseed rape in the UK. The timing and intensity of the spring and summer *M. persicae* aphid  
flights in the Columbia basin in the USA were associated with heat unit accumulation (day  
degrees) (Thomas et al., 1997). The differences between the accumulated day degrees in the first  
part (January to May) of 2007, 2008 and 2009 for the site of the Warwickshire Rothamsted Insect  
Survey suction trap do not appear to account for the abundance of *M. persicae* caught in this  
suction trap between August and November in these years and hence the incidence of TuYV. It is  
possible that the accumulated day degrees later in these years might account for the differences,  
however other factors including rainfall, predation, parasites, pathogens, availability of suitable  
host plants etc. are also likely to have an effect. Mild autumn conditions favour the development  
of the aphid vectors and encourage TuYV spread (Stevens et al., 2008). The low numbers of  
aphids in 2008 accounting for the low incidence of TuYV, were said to be due to wet and windy  
weather and an abundance of natural enemies.

In general, the incidence of TuYV within oilseed rape crops is considered to increase from initial  
autumn infection to a maximum level in the following spring (Stevens et al., 2008). Our data  
showed significant increases in the incidence of TuYV in eight fields between autumn and  
spring, but no significant change in 18 fields and a significant decrease in only one field. Most of  
the increases in incidence between autumn and spring occurred in the 2007/8 crop. All the fields  
in Warwickshire showed a significant increase in TuYV incidence between autumn 2007 and  
spring 2008 and the cumulative day degrees at our site in Warwickshire (10-11km from the  
Warwickshire fields sampled) were greater between January and May 2007 than for the same

1 Turnip yellows virus in winter oilseed rape  
2 months in 2008 and 2009. This indicates that in warmer winters, there could be yield/financial  
3 benefits from applying control measures such as effective insecticides at some point between  
4 November and April, depending up on regulations (numbers of spray and spray intervals  
5 permitted) and costs.  
6  
7  
8  
9

10  
11  
12 The significant regional differences in TuYV incidence, where highest levels of infection were  
13 observed in Lincolnshire and lowest in Yorkshire are likely to be attributable to a number of  
14 factors. There was much increased flight activity of *M. persicae* in Lincolnshire in the autumns of  
15 2007 and 2009 relative to Yorkshire and Warwickshire. Also, the large area of vegetable  
16 brassicas grown in Lincolnshire is likely to be a reservoir of TuYV and source for aphids over the  
17 summer months, when no oilseed rape crops are present. Between 1986 and 1989, surveys in  
18 England and Wales detected lower TuYV incidences in the north and east (Hill et al. 1989). In  
19 1992 and 1993, the incidence of TuYV was higher in Wales and the midlands, western and south  
20 western regions of England than in the eastern, south eastern or northern regions of England  
21 (Hardwick et al. 1994). Our data over three years and that of others (Blake, 2009; Clark and  
22 Stevens, 2009) suggest that there has been a change in prevalence, in that the highest levels of  
23 TuYV have been in Lincolnshire, close to the Wash and on the south coast.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 The spatial autocorrelation analysis revealed that TuYV-infected oilseed rape plants showed  
44 either random or slightly aggregated pattern of distribution within individual fields, with most  
45 fields showing slightly aggregated patterns during autumn. This finding agrees with that of  
46 Bourdon (1987) who reported that crop plants infected with viruses can show random or  
47 aggregated distributions, with aggregated distributions more common in vector-borne viruses.  
48  
49 There is very little data on the spatial incidence of TuYV, however Raybould et al. (1999) also  
50 reported plants infected by TuYV and other viruses (CaMV, TuMV and turnip yellow mosaic  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 Turnip yellows virus in winter oilseed rape  
2 virus) were distributed randomly, or were very weakly aggregated within wild populations of  
3 *Brassica oleracea*. Most of the oilseed crops showed positive autocorrelation (slight aggregated  
4 pattern) during autumn but a more random pattern during the following spring. Data on a non-  
5 persistently transmitted virus showed that aggregation of infected plants appears where there is  
6 limited spread of the virus from the initial (primary) foci of infection (Eckel and Lampert, 1993).  
7 This suggests that autumn infection (tending towards aggregation) is mostly due to primary  
8 infection with some, probably limited, secondary infection. The reduction in aggregation between  
9 autumn (18 out of 25 had negative Z-scores and only seven had positive Z-scores) and the  
10 following spring (13 out of 26 had negative Z scores and 13 had positive Z-scores) indicates that  
11 the infection of plants between the autumn sampling dates and May of the following years was  
12 mostly due to secondary spread of the virus within the fields, rather than further primary infection  
13 coming from outside the fields. This is consistent with the lack of flying aphid vectors caught in  
14 suction traps during this period (Anon, 2011) and the characteristic spatial spread of persistently  
15 transmitted viruses (Thresh, 1976). All the crops we sampled were in arable crop growing  
16 vicinities and the most common crops surrounding the fields sampled were winter wheat and  
17 winter barley, neither of which are known to be important hosts for TuYV.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 Oilseed rape is not reaching its full yield potential in the UK. No doubt there are a number of  
42 reasons for this, however, there is compelling evidence for TuYV being a major contributor to the  
43 shortfalls. TuYV is probably the most widespread and common disease of oilseed rape in the UK.  
44 The high levels of infection found, in years where there have been wet and not particularly warm  
45 summers and autumns, indicates that in warmer, drier years predicted under climate change  
46 scenarios (Parry, 1990), TuYV is likely to be an even greater problem. Higher incidences of  
47 TuYV were recorded for the major OSR growing countries of Europe (Poland, Germany and  
48 France) (Newbert 2016), than those outlined here for a three year period in England. This  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1 Turnip yellows virus in winter oilseed rape  
2 indicates that losses across parts of mainland Europe must be much higher than in the UK and  
3 control measures are needed. The most effective insecticides for *M. persicae* control, the  
4 neonicotinoids, are now banned in the EU due to adverse effects on pollinating insects (Dewar,  
5 2017). Varieties of oilseed rape with partial resistance to TuYV are now available, however, all  
6 varieties possess the same resistance source. This has created high selection pressure for  
7 resistance-breaking strains of TuYV. A further alternative source of resistance has been reported  
8 recently (Hackenberg et al. 2019).  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

## 22 ACKNOWLEDGEMENTS

23  
24 The authors wish to thank Dick Evenden and Simon Carr for help in identifying fields to sample,  
25 David Brightman and Adrian Howell for allowing us to sample their crops. We also thank  
26 Philippa Hallsworth, Sue Parker, Richard Harrington, Jinhua Mu, Shujiang Zhang and Rosemary  
27 Collier for their help. This work was supported by the Government of Ghana Educational Trust  
28 Fund and Biotechnology and Biological Sciences Research Council, Crop Improvement Research  
29 Club (CIRC) Project BB/I017410/1. We are grateful to collaborators and staff who have  
30 contributed data from the Rothamsted Insect Survey, a National Capability, is funded by the  
31 Biotechnology and Biological Sciences Research Council under the Core Capability Grant  
32 BBS/E/C/000J0200.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

## 48 REFERENCES

- 49  
50  
51 Abraham A. D., Varrelmann M., Vetten H. J. (2008) Molecular evidence for the occurrence of  
52 two new luteoviruses in cool season food legumes in Northeast Africa. *African Journal of*  
53 *Biotechnology*, 7, 414-420.  
54 Anon (2011) Aphid bulletin: Bulletin Archives. *Rothamsted Insect Survey* [Online]. Available:  
55 [http://www.rothamsted.bbsrc.ac.uk/insect-](http://www.rothamsted.bbsrc.ac.uk/insect-survey/STAphidBulletinArchive.php?Year=2011)  
56 [survey/STAphidBulletinArchive.php?Year=2011](http://www.rothamsted.bbsrc.ac.uk/insect-survey/STAphidBulletinArchive.php?Year=2011) [Accessed 26 July, 2019].  
57  
58  
59  
60

## Turnip yellows virus in winter oilseed rape

- 1 Armitage P., Berry G. (1994) *Statistical methods in medical research*, Oxford: Blackwell  
2 Science.
- 3 Berry P. M., Spink J. (2006) A physiological analysis of oilseed rape yields: Past and future.  
4 *Journal of Agricultural Science*, **144**, 381-392.
- 5 Blake A. (2009) *Turnip yellows virus threatens oilseed rape* [Online]. Reed Business  
6 Information. Available: [http://www.fwi.co.uk/Articles/2009/06/04/115896/Turnip-](http://www.fwi.co.uk/Articles/2009/06/04/115896/Turnip-yellow-virus-threatens-oilseed-rape.htm)  
7 [yellow-virus-threatens-oilseed-rape.htm](http://www.fwi.co.uk/Articles/2009/06/04/115896/Turnip-yellow-virus-threatens-oilseed-rape.htm) [Accessed 26 July, 2019].
- 8 Bourdon J. J. (ed.) (1987) *Diseases and plant population biology*, Cambridge, UK: Cambridge  
9 University Press.
- 10 Buckland S. T., Anderson D. R., Burnham K. P., Laake J. L., Borchers D. L., Thomas L. (2001)  
11 *Introduction to Distance Sampling*, Oxford: Oxford University Press.
- 12 Clark B., Stevens M. (2009) Strategies to reduce the impact of *Turnip yellows virus* on oilseed  
13 rape production in the UK. In: *HGCA Research and Development, Annual Report, 2009*  
14 *results*. Stoneleigh Park, Kenilworth, UK: Home-Grown Cereal Authority. Available:  
15 <https://cereals.ahdb.org.uk/media/160420/3498-annual-project-report-2009.pdf>  
16
- 17 Clark B., Stevens M. (2011) Strategies to reduce the impact of Turnip yellows virus on oilseed  
18 rape production in the UK. In: *HGCA Research and Development, Annual Report, 2011*  
19 *results*. Stoneleigh Park, Kenilworth, UK: Home-Grown Cereal Authority. Available:  
20 <https://cereals.ahdb.org.uk/media/160414/3498-annual-project-report-2011.pdf>  
21
- 22 Cliff A. D., Ord J. K. (eds.) (1969) *The problem of spatial autocorrelation*, London: Pion.
- 23 Coutts B. A., Hawkes J. R., Jones R. A. C. (2006) Occurrence of *Beet western yellows virus* and  
24 its aphid vectors in over-summering broad-leaved weeds and volunteer crop plants in the  
25 grainbelt region of south-western Australia. *Australian Journal of Agricultural Research*,  
26 **57**, 975-982.
- 27 Department for Environment Food and Rural Affairs (2018) National Statistics. Farming  
28 Statistics, Provisional crop areas, yields and livestock populations, At June 2018 - United  
29 Kingdom. Available:  
30 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/747210/structure-jun2018prov-UK-11oct18.pdf)  
31 [data/file/747210/structure-jun2018prov-UK-11oct18.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/747210/structure-jun2018prov-UK-11oct18.pdf) (Accessed 26.7.19).
- 32 Dewar, A. M. (2017) The adverse impact of the neonicotinoid seed treatment ban on crop  
33 protection in oilseed rape in the United Kingdom. *Pest Management Science* **73**, 1305-  
34 1309.
- 35 Eckel R. V. W., Lampert E. P. (1993) Spatial and temporal analysis of tobacco etch virus  
36 distribution and its relationship to aphid (Homoptera:Aphidae) vectors in flue-cured  
37 tobacco. *Journal of Economic Entomology*, **86**, 1534-1545.
- 38 Gilligan C. A., Pechan P. M., Day R., Hill S. A. (1980) *Beet western yellows virus* on oilseed  
39 rape (*Brassica napus* L). *Plant Pathology*, **29**, 53-53.
- 40 Graichen K. (1996) Rape is not a source of infection of *Beet western yellows virus*. *L'Informatore*  
41 *Agrario*, **52**, 79-81.
- 42 Graichen K., Rabeinstein F., Kurtz E. (2000) The occurrence of *Turnip yellows virus* in winter  
43 oilseed rape in Austria. *Pflanzenschutzberichte*, **59**, 35-46.
- 44 Graichen K., Schliephake E. (1999) Infestation of winter oilseed rape by *Turnip yellows*  
45 *luteovirus* and its effect on yield in Germany. In: *10th International Rapeseed Congress*,  
46 pp. 31-136. Canberra, ACT: International Consultative Group for Rapeseed Research.
- 47 Hackenberg, D., Asare-Bediako, E., Baker, A., Walley, P., Jenner, C., Greer, S., Bramham, L.,  
48 Batley, J., Edwards, D. Delourme, R., Barker, G., Teakle, G., Walsh, J. (2019)  
49 Identification and QTL mapping of resistance to Turnip yellows virus (TuYV) in oilseed  
50 rape, *Brassica napus*. *Theoretical and Applied Genetics* [https://doi.org/10.1007/s00122-](https://doi.org/10.1007/s00122-019-03469-z)  
51 [019-03469-z](https://doi.org/10.1007/s00122-019-03469-z)

## Turnip yellows virus in winter oilseed rape

- 1 Hardwick N. V., Davies J. M., Wright D. M. (1994) The incidence of three virus diseases in  
2 England and Wales in the 1991/92 and 1992/93 seasons. *Plant Pathology*, **43**, 1045-1049.
- 3 Hill S. A., Lane A., Hardwick N. V. (1989) The incidence and importance of *Beet western*  
4 *yellows virus* in oilseed rape. *Aspects of Applied Biology*, **23**, 311-318.
- 5 Hunter P. J., Jones J. E., Walsh J. A. (2002) Involvement of *Beet western yellows virus*,  
6 *Cauliflower mosaic virus* and *Turnip mosaic virus* in internal disorders of stored white  
7 cabbage. *Phytopathology*, **92**, 816-826.
- 8 Jasnic S., Bagi F. (2007) Viral diseases of oilseed rape. *Plant Doctor*, **35**, 458-461.
- 9 Jay C. N., Rossall S., Smith H. G. (1999) Effects of *Beet western yellows virus* on growth and  
10 yield of oilseed rape (*Brassica napus*). *Journal of Agricultural Science, Cambridge*, **133**,  
11 131-139.
- 12 Jones R. A. C., Coutts B. A., Hawkes J. (2007) Yield-limiting potential of *Beet western yellows*  
13 *virus* in *Brassica napus*. *Australian Journal of Agricultural Research*, **58**, 788-801.
- 14 Kerlan C. (1991) Les viroses, une etude de longue haleine. *Le dossier, Oleoscop no.5, Septembre*,  
15 **91**, 6-7.
- 16 Milošević, D., Marjanović-Jeromela, A., Ignjatov, M., Jovičić, D., Stanković, I., Bulajić, A.,  
17 Krstić, B. (2015) First Report of Turnip yellows virus on Oilseed Rape in Serbia. *Plant*  
18 *Disease* **99**, 1869.
- 19 Nelder J. A., Wedderburn R. W. M. (1972) Generalised linear models. *Journal of Royal*  
20 *Statistical Society, Series A*, **135**, 370-384.
- 21 Newbert, M. J. (2016) The genetic diversity of Turnip yellows virus in oilseed rape (*Brassica*  
22 *napus*) in Europe, pathogenic determinants, new sources of resistance and host range.  
23 PhD thesis, University of Warwick, 241pp.
- 24 Parry M. L. (1990) *Climate change and world agriculture*, London: Earthscan Publications.
- 25 Payne R. W., Murray D. A., Harding S. A., Baird D. B., Soutar D. M. (eds.) 2009. *GenStat for*  
26 *Windows: Introduction*, Hemel Hempstead: VSN International.
- 27 Polak J., Majkova L. (1992) Winter oilseed rape as a likely source and reservoir of *Beet western*  
28 *yellows virus*. *Ochrana Rostlin-UVTIZ*, **28**, 191-196.
- 29 Raybould A. F., Maskell L. C., Edwards M. L., Cooper J. I., Gray A. J. (1999) The prevalence  
30 and spatial distribution of viruses in natural populations of *Brassica oleracea*. *New*  
31 *Phytologist*, **141**, 265-275.
- 32 Schliephake E., Graichen K., Rabenstein F. (2000) Investigations on the vector transmission of  
33 the *Beet mild yellowing virus* (BMVYV) and the *Turnip yellows virus* (TuYV). *Zeitschrift*  
34 *Fur Pflanzenkrankheiten Und Pflanzenschutz-Journal of Plant Diseases and Protection*,  
35 **107**, 81-87.
- 36 Schroder M. (1994) Investigations on the susceptibility of oilseed rape (*Brassica napus* L. ssp  
37 *napus*) to different virus diseases. *Zeitschrift Fur Pflanzenkrankheiten Und*  
38 *Pflanzenschutz-Journal of Plant Diseases and Protection*, **101**, 576-589.
- 39 Shahraeen N., Farzadfar S., Lesemann D. E. (2003) Incidence of viruses infecting winter oilseed  
40 rape (*Brassica napus* ssp. *oleifera*) in Iran. *Journal of Phytopathology*, **151**, 614-616.
- 41 Smith H. G., Hinckes J. A. (1985) Studies on *Beet western yellows virus* in oilseed rape (*Brassica*  
42 *napus* ssp *oleifera*) and sugar beet (*Beta vulgaris*). *Annals of Applied Biology*, **107**, 473-  
43 484.
- 44 Stevens M., McGrann G., Clark B. (2008) *Turnip yellows virus* (syn *Beet western yellows virus*):  
45 an emerging threat to European oilseed rape production? *HGCA Research Review*, **69**, 1-  
46 36. Retrieved from <https://cereals.ahdb.org.uk/media/269200/rr69.pdf>
- 47 Thomas P. E., Pike, K.S. and Reed, G.L. (1997) Role of Green peach aphid flights in the  
48 epidemiology of *Potato leaf roll* disease in the Columbia Basin. *Plant Disease*, **81**, 1311-  
49 1316.

## Turnip yellows virus in winter oilseed rape

1 Thresh J. M. (1976) Gradients of plant virus diseases. *Annals of Applied Biology*, **82**, 381-406.

2 Walkey D. G. A., Pink D. A. C. (1990) Studies on resistance to *Beet western yellows virus* in  
3 lettuce (*Lactuca sativa*) and the occurrence of field sources of the virus. *Plant Pathology*,  
4 **39**, 141-155.

5  
6 Walsh J. A., Perrin R. M., Miller A., Laycock D. S. (1989) Studies on beet western yellows virus  
7 in winter oilseed rape (*Brassica napus* ssp. *oleifera*) and the effect of insecticidal  
8 treatment on its spread. *Crop Protection*, **8**, 137-143.

9 Walsh J. A., Tomlinson J. A. (1985) Viruses infecting winter oilseed rape (*Brassica napus* ssp.  
10 *oleifera*). *Annals of Applied Biology*, **107**, 485-495.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## Turnip yellows virus in winter oilseed rape

**TABLE TITLES AND FIGURE LEGENDS**

**TABLE 1** Mean autumn percentage incidence of turnip yellows virus (TuYV) in winter oilseed rape crops in Lincolnshire, Warwickshire and Yorkshire in the 2007-8, 2008-9 and 2009-10 crop seasons.

**TABLE 2** Mean spring percentage incidence of turnip yellows virus (TuYV) in winter oilseed rape crops in Lincolnshire, Warwickshire and Yorkshire in the 2007-8, 2008-9 and 2009-10 crop seasons.

**TABLE S1** Incidence and spatial analysis of turnip yellows virus (TuYV) in oilseed rape crops sampled in Lincolnshire, Warwickshire and Yorkshire in the 2007-8, 2008-9 and 2009-10 crop seasons.

**FIGURE 1a** Spatial distribution of turnip yellows virus-infected plants in oilseed rape field 1 in Lincolnshire in autumn of the 2007-8 crop season showing significant clustering (Rook's case connection of Black-White join-count statistic was used to test for the spatial autocorrelation of the TuYV-infected plants. In this field  $Z = -1.922$ ,  $P < 0.05$ ).

**FIGURE 1b** Spatial distribution of turnip yellows virus-infected plants in oilseed rape field 2 in Warwickshire in spring of the 2009-10 crop season showing random distribution (Rook's case connection of Black-White join-count statistic was used to test for the spatial autocorrelation of the TuYV-infected plants. In this field  $Z = 0.561$ ,  $P = 0.288$ ).

## Turnip yellows virus in winter oilseed rape

1  
2  
3  
4 **FIGURE 2.** (a) Rothamsted Insect Survey catches of *Myzus persicae* in suction traps located in  
5 Lincolnshire, Warwickshire and Yorkshire between June 2007 and April 2010 (Source: Aphid  
6 Bulletin, Rothamsted Insect Survey, Rothamsted Research, U.K.). (b) Mean incidences of turnip  
7 yellows virus (TuYV) from three fields in each of Lincolnshire, Warwickshire and Yorkshire in  
8 the autumns of 2007, 2008 and 2009.  
9  
10  
11  
12  
13  
14  
15  
16

17 **FIGURE 3** Relationship between cumulative numbers of *Myzus persicae* caught in the  
18 Rothamsted Insect Survey suction traps located in Lincolnshire, Warwickshire and Yorkshire  
19 between August and November in 2007, 2008 and 2009 and mean percentage turnip yellows  
20 virus incidence in oilseed rape crops in the autumn of each year in the three regions (d.f. = 2, 8; F  
21 = 24.2;  $P < 0.001$ ).  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**TABLE 1** Mean autumn percentage incidence of turnip yellows virus (TuYV) in winter oilseed rape crops in Lincolnshire, Warwickshire and Yorkshire in the 2007-8, 2008-9 and 2009-10 crop seasons

Region	Mean TuYV incidence (%) in autumn of crop season			Means <sup>1</sup>
	2007-8	2008-9	2009-10	
Lincolnshire	58.00 ± 8.51 b <sup>2,3</sup>	13.00 ± 5.80 d, e	94.00 ± 4.08 a	55.00 ± 3.69 a
Warwickshire	27.33 ± 7.69 c, d	3.00 ± 2.94 e	41.00 ± 8.48 b, c	23.78 ± 3.94 b
Yorkshire	24.67 ± 7.43 c, d	2.33 ± 2.60 e	11.00 ± 5.40 d, e	12.67 ± 3.18 c
Means <sup>4</sup>	36.67 ± 4.56 a	6.11 ± 2.33 b	48.67 ± 3.62 a	

<sup>1</sup> Means in the same column followed by different letters are significantly different from each other ( $P < 0.001$ ).

<sup>2</sup> Mean and standard error (mean ± SE).

<sup>3</sup> Region-crop season incidence interaction means followed by different letters are significantly different from each other ( $P < 0.05$ ).

<sup>4</sup> Means in the same row followed by different letters are significantly different from each other ( $P < 0.001$ ).

Analysis of the TuYV incidence data was carried out using a generalised linear model (GLM).

**TABLE 2** Mean spring percentage incidence of turnip yellows virus (TuYV) in winter oilseed rape crops in Lincolnshire, Warwickshire and Yorkshire in the 2007-8, 2008-9 and 2009-10 crop seasons

Region	Mean TuYV incidence (%) in spring of crop season			Means <sup>1</sup>
	2007-8	2008-9	2009-10	
Lincolnshire	66.00 ± 12.08 <sup>2,3</sup>	10.00 ± 7.65	94.00 ± 6.04	56.67 ± 5.17 a
Warwickshire	68.33 ± 11.86	12.33±8.38	45.33±12.69	42.00 ± 6.43 a
Yorkshire	32.67 ± 11.96	3.67 ± 4.79	22.00 ± 10.56	19.44 ± 5.55 b
Means <sup>d</sup>	55.67 ± 6.91 a	8.67 ± 4.11 b	53.78 ± 5.86 a	


<sup>1</sup> Means in the same column followed by different letters are significantly different from each other ( $P = 0.002$ ).

<sup>2</sup> Mean and standard error (mean ± SE).


<sup>3</sup> The interaction effect between county and crop season is not significant ( $P = 0.163$ ).

Analysis of the TuYV incidence data was carried out using a generalised linear model (GLM).







**FIGURE 1a** Spatial distribution of turnip yellows virus-infected plants in oilseed rape field 1 in Lincolnshire in autumn of the 2007-8 crop season showing significant clustering (Rook's case connection of Black-White join-count statistic was used to test for the spatial autocorrelation of the TuYV-infected plants. In this field  $Z = -1.922$ ,  $P < 0.05$ ).



**FIGURE 1b** Spatial distribution of turnip yellows virus-infected plants in oilseed rape field 2 in Warwickshire in spring of the 2009-10 crop season showing random distribution (Rook's case connection of Black-White join-count statistic was used to test for the spatial autocorrelation of the TuYV-infected plants. In this field  $Z = 0.561$ ,  $P = 0.288$ ).



**FIGURE 2.** (a) Rothamsted Insect Survey catches of *Myzus persicae* in suction traps located in Lincolnshire, Warwickshire and Yorkshire between June 2007 and April 2010 (Source: Aphid Bulletin, Rothamsted Insect Survey, Rothamsted Research, U.K.). (b) Mean incidences of turnip yellows virus (TuYV) from three fields in each of Lincolnshire, Warwickshire and Yorkshire in the autumns of 2007, 2008 and 2009.



**FIGURE 3** Relationship between cumulative numbers of *Myzus persicae* caught in the Rothamsted Insect Survey suction traps located in Lincolnshire, Warwickshire and Yorkshire between August and November in 2007, 2008 and 2009 and mean percentage turnip yellows virus incidence in oilseed rape crops in the autumn of each year in the three regions (d.f. = 2, 8;  $F = 24.2$ ;  $P < 0.001$ ).