

Scientific note

**First record of ectoparasitic isopods on the invasive lionfish
Pterois miles (Bennett, 1828)**

(Crustacea, Cymothoidae and Teleostei, Scorpaenidae)

Charalambos Antoniou, Periklis Kleitou, Fabio Crocetta & Maurizio Lorenti

The lionfish *Pterois miles* (Bennett, 1828) is a benthic coastal ray-finned fish native to the western Indo-Pacific region, that recently invaded the Mediterranean basin (Kletou et al. 2016). We first document the occurrence of six *Nerocila* Leach, 1818 parasites collected from four adult lionfish host specimens from the southern coasts

of Cyprus (Table 1). Five isopods were collected on the skin, while one was found attached in the inner branchial cavity, in the approximate location of the vestibular labyrinth (Table 2; Fig. 1). The isopods specimens consisted of four transitional individuals, possibly young females (Table 2: specimens 1–4), a juvenile male



Fig. 1. *Nerocila bivittata* found parasitizing *Pterois miles* in Cyprus. A–B. Larvigerous female. C. “Aster” form. D. Male.

Charalambos Antoniou (corresponding author) & Periklis Kleitou, Marine and Environmental Research (MER) Lab Ltd., Limassol, Cyprus; e-mail: cantoniou@merresearch.com

Fabio Crocetta & Maurizio Lorenti, Stazione Zoologica di Napoli Anton Dohrn, Department of Integrative Marine Ecology, Center of Villa Dohrn – Benthic Ecology, Ischia (NA), Italy

Table 1. Localities where parasitized lionfishes were found, with date, depth (in meters), and environmental features.

Locality	Date	Coordinate	Depth	Substrate
Limassol major port	12/07/2017	34°38.677'N 33°01.251' E	3	Port breakwaters
Moni	18/07/2017	34°41.698'N 33°11.885' E	26.5	Hard substrate with sparse <i>Posidonia oceanica</i> meadows
Cavo Greco	03/08/2017	34°59.038'N 34°04.667' E	20	Hard substrate
Cavo Greco	03/08/2017	34°59.091'N 34°04.632' E	22	Hard substrate

Table 2. Lionfish and isopod morphometric, and location of infestation. Abbreviations used: BL, body length (in mm); BW, body width (in mm); TL, total length (in cm); W, weight (in grams).

<i>Nerocila bivittata</i> specimen	Figure	Location of infestation	<i>Pterois miles</i>		<i>Nerocila bivittata</i>	
			TL	W	BL	BW
1		Caudal peduncle	22.5	178	14.5	7.8
2	1C	Anal fin	22.5	178	10.5	4.5
3		Caudal peduncle	19.8	108	12.9	6.9
4		Abdomen	22.8	164	13.33	7.11
5	1D	Gill cavity	30.3	528	8.43	3.75
6	1A, 1B	Between the pelvic fins	30.3	528	26.22	13.82

(Table 2: specimen 5; Fig. 1D), and a large larvigerous female (Table 2: specimen 6; Fig. 1), belonging to *Nerocila bivittata* (Risso, 1816) (Crustacea, Isopoda, Cymothoidea, Cymothoidea). The brooding female and the male were found co-occurring in different positions on the same fish, namely pelvic fins and gill cavity (Table 2; Figs 1A–B,D). One individual (Table 2: specimen 2; Fig. 1C) also featured a peculiar extension of pereonites and coxae reminiscent of the “aster” form of the Pacific-Atlantic *Nerocila acuminata* (Brusca, 1981). However, among the other features, the morphology of uropods in the present specimen (endopod lacking an acute point or a distinct distomedial tooth or sinuosity) looks different from that figured for *N. acuminata* (Brusca 1981, Bruce 1987), and therefore we assigned that specimen to *N. bivittata* as well.

Our findings constitute the first records of a *Pterois miles*–*Nerocila bivittata* association. In addition, the occurrence of different stages of *N. bivittata* on lionfish shows that the isopod has the potential of completing its life cycle on this introduced host, which points to a stability of the interactions between the two species.

References

- Bruce, N. L. 1987. Australian species of *Nerocila* Leach, 1818, and *Creniola* n. gen. (Isopoda: Cymothoidea), crustacean parasites of marine fishes. Records of the Australian Museum 39(6): 355–412.
- Brusca, R. C. 1981. A monograph on the Isopoda Cymothoidea (Crustacea) of the eastern Pacific. Zoological Journal of the Linnean Society 73(2): 117–199.
- Kletou, D., Hall-Spencer, J. M. & Kleitou, P. 2016. A lionfish (*Pterois miles*) invasion has begun in the Mediterranean Sea. Marine Biodiversity Records 9(1): 46.