



THE CENTRAL AFRICAN JOURNAL OF MEDICINE

Vol. 40, No. 8

CONTENTS

August, 1994

ORIGINAL ARTICLES

- Neoplasms involving the spinal cord in Zimbabweans: an analysis of 262 cases PI Garrido, S Laher-Mooncey, NL Murphree, N Jonker, LF Levy, S Makarawo 201
- Termination of pregnancy using extra-amniotic prostaglandin F₂_α – experience in a peripheral hospital A Krolkowski, KA Janowski, JV Larsen 204
- Epidemiological study of drug use in urban and rural secondary schools in Zimbabwe SW Acuda, AH Eide 207
- A comparison of three types of "M" traps for sampling tsetse fly (Diptera: Glossinidae) populations at South Luangwa Game Management Area, Zambia MM Ziba, M Mukunyandela, S Siziya, JS Chipipa, L Mwanza 212
- Increased susceptibility of erythrocyte membrane lipids to peroxidation in sickle cell disease EU Essien 217

CASE REPORTS

- Intracranial Burkitt's lymphoma with extension into orbital spaces resulting in bilateral blindness K Ramesh, LD Gahukamble, DB Gahukamble ... 220
- Munchausen syndrome by proxy – a case report and review of literature OA Oyelami, AM Alhaj, IK Airede 222
- Hypothyroxinemia in Acquired Immune Deficiency Syndrome (AIDS) CC Ezeala, E Chukwurah 226

REVIEW ARTICLES

- Advances in electrodiagnosis in Zimbabwe. Part II: "Evoked Potentials" O Parry, LF Levy, J Mielke 229

SUPPORTED BY CIMAS MEDICAL AID SOCIETY

Increased susceptibility of erythrocyte membrane lipids to peroxidation in sickle cell disease

EU ESSIEN

SUMMARY

Erythrocytes from normal subjects and from cases of sickle cell disease were exposed to hydrogen peroxide and the extent of membrane lipid peroxidation studied. Significantly increased peroxidation was observed in intact erythrocytes, isolated lipids from intact erythrocytes or their ghosts compared with normal controls. However, the extent of lipid peroxidation was less for isolated lipids of intact erythrocytes or their ghosts. This difference may be due to loss of factors enhancing peroxidation in intact erythrocytes in the extraction procedure. It is likely that damage to erythrocyte membranes caused by increased susceptibility to lipid peroxidation may be a factor in some of the haematological features of sickle cell disease.

INTRODUCTION

Although sickle cell disease (SCD) is well understood as an abnormality of the haemoglobin molecule, the complete mechanism by which this defect is expressed at the clinical level is not fully understood^{1,2} even though secondary alterations in red cell metabolism and membrane structure and function have been described.³ A considerable body of evidence suggests that defective function of the sickle cell membrane may contribute to some of the manifestations of the disease. The membrane defect is demonstrated in the permanently deformed morphology of the irreversibly sickled cell, binding of haemoglobin to the erythrocyte

*Department of Biochemistry
College of Medical Sciences
University of Calabar
Calabar, Nigeria*

membrane, easy rupturing of the sickle cell, decreased filterability of sickle erythrocytes and increased cation leak from sickle cells when they are deoxygenated.^{2,4} Equally, it has been established that the circulation of membrane damaged, rigid and irregularly shaped erythrocytes resulting in vascular occlusion are involved in this disease.²

However, the genesis of the membrane defect remains not fully known and relatively few examinations of the factors precipitating the collapse of the membrane have been made. Increased susceptibility of the membrane to peroxidant threats may be one of those factors. It is known that the erythrocyte membrane is rich in polyunsaturated fatty acids and that these molecules are themselves highly susceptible to peroxidation.

During its life span in the circulation the erythrocyte is exposed to small amounts of hydrogen peroxide (H₂O₂) generated as a result of normal metabolic process to greater amounts resulting from interactions of certain oxidant drugs and their metabolites with oxyhaemoglobin.⁵ Intracellular H₂O₂ may cause peroxidation of membrane lipids with the formation of breakdown products, including oxygen radicals and malonyldialdehyde (MDA).⁶ Membrane lipid peroxidation has been recognized as a deleterious process leading to cell damage in various tissues.⁷ MDA itself has been found capable of reacting with aminophospholipids and proteins of membrane to produce cross linking of membrane constituents which can cause rigidity and decreased deformability of red blood cells.⁶

The present study was undertaken to examine the susceptibility of sickle cell membrane to lipid peroxidation and the possible role of lipid peroxidation in the damage to the membrane and reduced erythrocyte survival in sickle cell disease.

MATERIALS AND METHODS

Blood was drawn from healthy adult male volunteers or sickle cell patients who were in a steady state uncomplicated by infection, clinical or haematological crises, into EDTA vials and the plasma separated. Red blood cells were routinely washed in cold normal saline and suspended in phosphate-saline medium (1 vol. of potassium phosphate buffer, 0,1 M, pH 7,3 mixed with 9 vol. of 0,15 M NaCl). Saline washed, packed red cells were lysed in four times their volume of cold distilled water.

An aliquot of the lysate was used to determine the haemoglobin (Hb) content by the cynmethaemoglobin method described by Dacie and Lewis.⁸ The lipids in the remainder of the lysate were extracted according to the method of Rose and Oklander.⁹ Peroxidation of intact erythrocytes, isolated lipids of intact erythrocytes or their ghosts was carried out by exposure to hydrogen peroxide which was added directly to the medium as described by Kurien and Iyer.¹⁰

Erythrocyte ghosts were prepared according to Dodge *et al*.¹¹ The extent of peroxidation of the erythrocyte and its membrane lipids was measured by estimating the MDA formed using the thiobarbituric acid (TBA) reaction described by Stocks and Dormandy.¹² Data were analyzed by using student's t-test.

RESULTS

As seen in Table I, a significantly higher level of MDA formation was observed when intact red cells from SCD patients were exposed to H₂O₂ added directly to the red cell suspension, compared to that of erythrocytes from normal subjects, (p < 0,001). The same was true of isolated lipids from intact red cells or their ghosts, but the amount of MDA formed was less. The formation of MDA, a secondary fragmentation product of polyunsaturated fatty acid peroxides, is a specific, sensitive and extensively used measure of lipid peroxidation.¹²

Table I: Lipid peroxidation in normal and sickle erythrocytes (n moles of MDA/g Hb).¹

Lipid Source	n moles MDA (13)*	p
Intact red cell normal		
Normal	198,6 + 24	
Sickle cell	408,7 + 33	< 0,01
Isolated lipids from intact red cells		
Normal	78,5 + 11	
Sickle cell	145,8 + 12	< 0,01
Isolated lipids from red cell ghosts		
Normal	63,9 + 10	
Sickle cell	187,8 + 14	< 0,01

¹ Values are expressed as means ± SEM.

* Number of cases studied given in parenthesis.

DISCUSSION

Certain abnormalities that occur in SCD are difficult to relate directly to the presence of HbS or the sickling phenomenon.¹ It is now known that a combination of intrinsic erythrocyte abnormalities and other factors are likely to be epiphenomenologic. In this study it is shown that the susceptibility to peroxidation of intact erythrocyte and isolated lipids of intact erythrocytes or their ghosts is appreciably increased in SCD compared to normal subjects. This is suggestive that in SCD the erythrocyte membrane is more vulnerable to lipid peroxidation and that this increased vulnerability may be a feature which contributes to the easy collapse of the erythrocyte membrane in SCD.

The increased vulnerability may be due to disorganization of the Hb molecule, which as a haem-protein, is a powerful catalyst of lipid peroxidation,¹³ distortion of normal Hb-membrane and membrane-membrane relations, and plasma and erythrocyte vitamin E depletion.¹⁴ In fact, increased TBA – reactive materials have been demonstrated to precede haemolysis of sickle erythrocytes¹⁵ and abnormal cation leaks that develop in SCD damaged membranes.^{16,17}

It appears from the above consideration that in SCD H_2O_2 may be generated in increased amounts and not physiologically well managed. If its behaviour in this study is a guide, H_2O_2 is a potent oxidant initiating the production of a steady supply of free radicals which degrade membrane phospholipids, followed by the collapse of the membrane.

The disparity in the extent of peroxidation between lipids of intact erythrocytes compared to isolated lipids of intact erythrocytes or their ghosts as measured by MDA formation (Table I) may be related to loss of unsaturated lipids in the extraction procedure. This observation that the extent of peroxidation was significantly higher for intact erythrocytes than for lipids isolated from intact erythrocytes or their ghosts is consistent with the report of Stocks and Dormandy¹² that isolated lipids yield much less MDA compared to lipids of intact red cells. However, MDA formation was still higher for isolated lipids from sickle erythrocytes compared to normal erythrocytes.

From the above consideration it is plausible that in SCD, the primary event of crystallization of HbS is aided by damage to the cell membrane caused by increased peroxidation of membrane constituents among other factors. Excessive and symmetric fragmentation

and loss of cell membrane due to peroxidation may precipitate irreversible sickling of erythrocytes containing HbS. These cells will not return to a normal configuration even upon oxygenation. In fact, these cells can be lysed and the solubilised Hb removed, and the membrane will remain in the sickled configuration² indicating permanent damage to the membrane.

REFERENCES

1. Burns ER, Wilkinson WH, Nagel RL. Adherence properties of sickle erythrocytes in dynamic flow systems. *J Lab Clin Med* 1985; 105: 673–8.
2. Brewer GJ. Inherited erythrocyte metabolic and membrane disorders. *Med Clin North Am* 1980; 64:579–96.
3. Beutler E. The sickle cell disease and related disorders. In: Haematology, 3rd ed. Williams WJ, Beutler E, Erslev AJ, et al New York: McGraw-Hill Book Co, 1986;583–608.
4. Noguchi CT, Schechter AN. The intracellular polymerisation of sickle haemoglobin and its relevance to sickle cell disease. *Blood* 1981; 58:1057–68.
5. Tudhope GR, Hopkins J. Lipid peroxidation in human erythrocyte in tocopherol deficiency. *Acta haemat* 1975;53:98–104.
6. Mead JF, Alfin-Slater RB, Howton DR, Popjak G. Lipids: chemistry, biochemistry and nutrition, New York: Plenum Press, 1986.
7. Pelisseir MA, Boisset M, Albrecht R. The effects of vitamin A nutritional status on microsomal lipid peroxidation and alpha-tocopherol level in rat liver. *Experientia* 1989;45:342–3.
8. Dacie JV, Lewis SM. Practical Haematology. 6th ed. Edinburgh: Churchill Livingstone, 1984;175–90.
9. Rose HG, Oklander M. Improved procedure for the extraction of lipids from human erythrocytes. *J Lipid Res* 1965;6:528–31.
10. Kurien M, Iyer GYN. Lipid peroxidation in erythrocytes from different animal species. *Ind J Biochem Biophys* 1976;13:67–70.
11. Dodge JT, Mitchell C, Hanahan DG. The preparation and chemical characteristics of haemoglobin-free ghosts of human erythrocytes. *Archs Biochem Biophys* 1963;100:119–25.

12. Stocks J, Dormandy TL. The autoxidation of human red cell lipids induced by hydrogen peroxide. *Br J Haematol* 1971;20:95-111.
13. Devlin TM. Textbook of biochemistry with clinical correlations. 2nd ed. New York: John Wiley and Sons, 1986;355-65.
14. Sergent GR. Sickle cell disease. Oxford: Oxford University Press, 1988;51-8.
15. Jacob HS, Lux SE. Degradation of membrane phospholipids and thiols in peroxide hemolysis: studies in vitamin E deficiency. *Blood* 1968; 32:549-68.
16. Hebbel RP, Mohandas N. Reversible deformation-dependent erythrocyte cation leak. Extreme sensitivity conferred by minimal peroxidation. *Biophys J* 1991;60:712-5.
17. Van der Berg JJ, Kuypers FA, Lubin BB, Roelofsen BB, Roelofsen B, Op den Kamp JA. Direct and continuous measurement of hydroperoxide-induced oxidative stress on the membranes of intact erythrocytes. *Free Radic Biol Med* 1991;11:255-61.
18. Sagihara T, Rawicz W, Evans EA, Hebbel RP. Lipid hydro peroxides permit deformation dependent leak of monovalent cation from erythrocytes. *Blood* 1991;77:2757-63.
19. Adelekan DA, Thurnhan DI, Adekile AD. Reduced antioxidant capacity in paediatric patients with homozygous sickle cell disease. *Eur J Clin Nutr* 1989;43:609-14.
20. Jain SK, Ross JD, Levy GJ, Little RL, Duett J. The accumulation of malonyldialdehyde, an end product of membrane lipid peroxidation, can cause potassium leak in normal and sickle red blood cells. *Biochem Med* 1989;42:60-5.



This work is licensed under a
Creative Commons
Attribution – NonCommercial - NoDerivs 3.0 License.

To view a copy of the license please see:
<http://creativecommons.org/licenses/by-nc-nd/3.0/>

This is a download from the BLDS Digital Library on OpenDocs
<http://opendocs.ids.ac.uk/opendocs/>